

Counting the sum of cubes for Lucas and Gibonacci Numbers

Wamiliana^{1*}, Suharsono¹, P. E. Kristanto²

¹Dept. of Mathematics, Universitas Lampung, Jl. Soemantri Brojonegoro No. 1, Bandarlampung, Indonesia

²SMP Xaverius 1, Jl. Sultan Hasanuddin No. 20, Telukbetung, Bandarlampung, Indonesia

*Corresponding author: wamiliana.1963@fmipa.unila.ac.id

Abstract

Lucas and Gibonacci numbers are two sequences of numbers derived from a wellknown numbers, Fibonacci numbers. The difference between Lucas and Fibonacci numbers only lies on the first and second elements. The first element in Lucas numbers is 2 and the second is 1, and nth element, $n \geq 3$ determined by similar pattern as in the Fibonacci numbers, i.e : $L_n = L_{n-1} + L_{n-2}$. Gibonacci numbers $G_0, G_1, G_2, G_3, \dots$; $G_n = G_{(n-1)} + G_{(n-2)}$ are generalized of Fibonacci numbers, and those numbers are nonnegative integers. If $G_0=1$ and $G_1=1$, then the numbers are the wellknown Fibonacci numbers, and if $G_0=2$ and $G_1=1$, the numbers are Lucas numbers. Thus, the difference of those three sequences of numbers only lies on the first and second of the elements in the sequences. For Fibonacci numbers there are quite a lot identities already explored, including the sum of cubes, but there have no discussions yet about the sum of cubes for Lucas and Gibonacci numbers. In this study the sum of cubes of Lucas and Gibonacci numbers will be discussed and showed that the sum of cubes for Lucas numbers is $\sum_{i=0}^n (L_i)^3 = \frac{L_n(L_{n+1})^2 + 5(-1)^n L_{n-1} + 19}{2}$ and for Gibonacci numbers is $\sum_{i=0}^n (G_i)^3 = \frac{[G_n(G_{n+1})^2 + (G_1 - G_0)^3 - 3G_0^2 G_1 + 4G_0^3 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2) G_{n-1}]}{2}$

Keywords

Fibonacci numbers, Lucas numbers, Gibonacci numbers, identities, sum of cubes

Received: 11 Februari 2019, Accepted: 8 April 2019

<https://doi.org/10.26554/sti.2019.4.2.31-35>

1. INTRODUCTION

Fibonacci number was originally stated by Leonardo Pisano or Fibonacci in his book, Liber Abaci in 1202. In his book he stated a problem about the growth of pairs rabbits. A young pair of rabbits (male and female) is placed on an island. A pair of rabbits does not breed until they are two months old. After they are two month old, each pair of rabbits produces another pair each month. Assuming that there are no rabbits ever die then the total number of pair of rabbits form a sequence 1,1,2,3,5,8,13,...and this sequence of numbers is known as Fibonacci numbers. Lucas number was introduced by François Edouard Anatole Lucas in 1877 by setting the first number is 2 and the second is 1, and the other numbers are following the pattern as on Fibonacci numbers so that it form a sequence of numbers 2, 1, 3, 4, 7, 11, 18, 29, 47,... Gibonacci numbers $G_0, G_1, G_2, G_3, \dots; G_n = G_{n-1} + G_{n-2}$ are generalized of Fibonacci numbers, and those numbers are nonnegative integers. If $G_0 = 1$ and $G_1 = 1$, then the numbers are the wellknown Fibonacci numbers, and if $G_0 = 2$ and $G_1 = 1$, the numbers are Lucas numbers. Thus, the difference of those three sequence of numbers only lies on the first and second of the elements in the sequences.

There are many identities already developed that related among Gibonacci, Fibonacci, and Lucas numbers. Posamentier

et al. (2007) showed that Fibonacci, Lucas, and Gibonacci numbers related to Pascal triangle, art and golden ratio. There are some researchers already investigated identities for Fibonacci Lucas and Gibonacci numbers such as Carlitz et al. (1972a,b,c,d), and Benjamin et al. (1999, 2000, 2009, 2003). Krishna (1980), investigated generalized Fibonacci sequence and William (1972) discussed how to get Fibonacci numbers from Pascal's triangle. Dunlap (2003) investigated about the golden ratio and Generalized Fibonacci numbers, including the application in biology and crystallography. Benjamin et al. (2003) proved more than one hundred Fibonacci identities by combinatorial arguments and Benjamin et al. (2009) proposed the identity for finding the sum of cubes for Fibonacci numbers. Frontczak (2018) discussed about the relationship of sum of powers for Fibonacci and Lucas numbers for sum of first, second, third, and fourth powers; and Arangala et al. (2016) investigated the relationship of the sum of power for Fibonacci and Lucas numbers for the power of square, fourth, sixth, and eighth. However, until recently there have no discussion about the sum of cubes for Lucas numbers where in the formula just contains the Lucas numbers itself, or the sum of cubes for Gibonacci numbers where in the formula just contains the Gibonacci numbers itself. Therefore, in this study we will discuss about the sum of cubes for Gibonacci numbers as an

extension identity of [Koshy \(2001\)](#), and [Benjamin et al. \(1999\)](#). In Section 2 and 3 we will present the results from [Koshy \(2001\)](#), [Benjamin et al. \(1999\)](#), and [Dunlap \(2003\)](#), and in Section 4 we will present the results.

2. Some Identities for Lucas and Gibonacci Numbers

2.1 Some Identities for Lucas Number

The Identity 1, 2, and 3 are proposed by [Koshy \(2001\)](#), and the Identity 4 is proposed by [Benjamin et al. \(1999\)](#).

Identity 1

$$L_0 + L_1 + L_2 + \dots + L_n = L_{n+2} - L_1$$

Proof. Using the definition of Lucas number we get

$$L_{n-2} = L_n - L_{n-1} \text{ then,}$$

$$L_0 = L_2 - L_1$$

$$L_1 = L_3 - L_2$$

$$L_2 = L_4 - L_3$$

$$L_3 = L_5 - L_4$$

.

.

.

$$L_{n-1} = L_{n+1} - L_n$$

Take the sum on both sides simultaneously we get :

$$L_0 + L_1 + L_2 + L_3 + \dots + L_n = L_{n+2} - L_1$$

Identity 2

$$L_0 + L_1 + L_2 + L_3 + \dots + L_{2n} = L_{2n+1} - L_1$$

Proof. By definition, $L_{n-1} = L_n - L_{n-2}$ then,

$$L_0 = L_2 - L_1$$

$$L_2 = L_4 - L_1$$

$$L_4 = L_6 - L_3$$

$$L_6 = L_8 - L_5$$

.

.

.

$$L_{2n-2} = L_{2n-1} - L_{2n-3}$$

$$L_{2n} = L_{2n+1} - L_{2n-1}$$

Take the sum on both sides simultaneously we get :

$$\begin{aligned} L_0 + L_1 + L_2 + L_3 + \dots + L_{2n} &= L_{2n+1} - L_1 + L_0 \\ &= L_{2n+1} - 1 + 2 \\ &= L_{2n+1} + 1 \\ &= L_{2n+1} L_1 \end{aligned}$$

Identity 3

$$L_1 + L_3 + L_5 + \dots + L_{2n-1} = L_{2n} - L_0$$

Proof. By definition, $L_{n-1} = L_n - L_{n-2}$ then,

$$L_1 = L_2 - L_0$$

$$L_3 = L_4 - L_2$$

$$L_5 = L_6 - L_4$$

.

.

.

$$L_{2n-3} = L_{2n-2} - L_{2n-4}$$

$$L_{2n-1} = L_{2n} - L_{2n-2}$$

Take the sum on both sides simultaneously we get :

$$L_1 + L_3 + L_5 + \dots + L_{2n-1} = L_{2n} - L_0$$

□

Identity 4

$$L_{n+1} L_{n-1} = L_n^2 - (-1)^n \cdot 5$$

Proof. Using mathematical induction, and we leaven the basic induction and just do the induction step. Asumme that the statement is true for $n = k$, $L_{k+1} L_{k-1} = L_k^2 - (-1)^k \cdot 5$

For $n = k + 1$, then :

$$L_{k+1}^2 = L_{k+1} \cdot L_{k+1}$$

$$L_{k+1}^2 = (L_{k+2} - L_k) L_{k+1}$$

$$L_{k+1}^2 = L_{k+2} \cdot L_{k+1} - L_k \cdot L_{k+1}$$

$$L_{k+1}^2 = L_{k+2} - (L_k + L_{k-1}) \cdot L_k \cdot L_{k+1}$$

$$L_{k+1}^2 = L_{k+2} L_k + L_{k-1} + L_{k+2} \cdot L_{k-1} - L_k \cdot L_{k+1}$$

$$L_{k+1}^2 = L_{k+2} L_k + (L_{k-1} + L_k) L_{k-1} \cdot L_k L_{k+1}$$

$$L_{k+1}^2 = L_{k+2} L_k + L_{k+1} L_{k-1} + L_k \cdot L_{k-1} - L_k \cdot L_{k+1}$$

$$L_{k+1}^2 = L_{k+2} L_k + L_{k+1} \cdot L_{k-1} + L_k (L_{k+1} - L_{k-1})$$

$$L_{k+1}^2 = L_{k+2} L_k + L_{k+1} L_{k-1} + L_k (-L_k)$$

$$L_{k+1}^2 = L_{k+2} L_k + L_{k+1} L_{k-1} - L_k^2$$

Since, $L_{k-1} L_{k+2} = -L_k^2 - (-1)^k \cdot 5$, then

$$L_{k+1}^2 = L_{k+2} \cdot L_k - (-1)^k \cdot 5$$

$$L_{k+1}^2 = L_{k+2} \cdot L_k + (-1)^{k+1} \cdot 5$$

$$L_{k+2} \cdot L_k = L_{k+1}^2 + (-1)^{k+1} \cdot 5$$

□

2.2 Some Identities for Gibonacci Number

The Identity 5, 6, and 7 are proposed by [Dunlap \(2003\)](#), Identity 8 is proposed by [Benjamin et al. \(2009\)](#), while Identity 9 derived from Definition 8. Those Identity will be used in the process to determine the sum of cubes for Gibonacci numbers.

Identity 5

$$G_0 + G_1 + G_2 + \dots + G_n = G_{n+2} - G_1$$

Proof. Using the definition of Gibonacci numbers, we get

$$G_{n-2} = G_n - G_{n-1} \text{ then,}$$

$$G_0 = G_2 - G_1$$

$$G_1 = G_3 - G_2$$

$$G_2 = G_4 - G_3$$

$$G_3 = G_5 - G_4$$

.

.

.

$$G_{n-1} = G_{n+1} - G_n$$

$$G_n = G_{n+2} - G_{n+1}$$

Take the sum of both sides simultaneously we get:

$$G_0 + G_1 + G_2 + \dots + G_n = G_{n+2} - G_1$$

□

Identity 6

$$G_0 + G_1 + G_2 + G_3 + \dots + G_{2n} = G_{2n+1} + G_0 - G_1$$

Proof. Again, using the definition of Gibonacci numbers, we get

$$G_{n-1} = G_n - G_{n-2},$$

$$G_0 = G_0$$

$$G_2 = G_3 - G_1$$

$$G_4 = G_5 - G_3$$

$$G_6 = G_7 - G_5$$

.

.

.

$$G_{2n-2} = G_{2n-1} - G_{2n-3}$$

$$G_{2n} = G_{2n+1} - G_{2n-1}$$

Take the sum of both sides simultaneously we get :

$$G_0 + G_1 + G_2 + G_3 + \dots + G_{2n} = G_{2n+1} + G_0 - G_1$$

Identity 7

$$G_1 + G_3 + G_5 + \dots + G_{2n-1} = G_{2n} - G_0$$

Proof. $G_{n-1} = G_n - G_{n-2}$ then,

$$G_1 = G_2 - G_0$$

$$G_3 = G_4 - G_2$$

$$G_5 = G_6 - G_4$$

.

.

.

$$G_{2n-3} = G_{2n-2} - G_{2n-4}$$

$$G_{2n-1} = G_{2n} - G_{2n-2}$$

Take the sum on both sides simultaneously we get :

$$G_1 + G_3 + G_5 + \dots + G_{2n-1} = G_{2n} - G_0$$

Identity 8

$$G_{n+1}G_{n-1} = G_n^2 - (-1)^n \cdot \{G_1^2 - G_1G_0 - G_0^2\}$$

Proof. By using mathematical induction.

For $n = k + 1$, then :

$$G_2 \cdot G_0 = G_n^2 - (-1)^n \cdot \{G_1^2 - G_1G_0 - G_0^2\}$$

$$(G_1 + G_0)G_0 = G_1^2 - G_1^2 - G_1G_0 - G_0^2$$

$$G_1G_0 + G_0^2 = G_1G_0G_0^2$$

Assume that the statement is true for $n = k$, then

$$G_{k+1}^2 = G_{k+1} \cdot G_{k+1}$$

$$G_{k+1}^2 = (G_{k+2} - G_k)G_{k+1}$$

$$G_{k+1}^2 = G_{k+2} \cdot G_{k+1} - G_k \cdot G_{k+1}$$

$$G_{k+1}^2 = G_{k+2} - (G_k + G_{k-1}) \cdot G_k \cdot G_{k+1}$$

$$G_{k+1}^2 = G_{k+2}G_k + G_{k-1}L_{k+2} \cdot G_{k-1} - G_k \cdot G_{k+1}$$

$$G_{k+1}^2 = G_{k+2}G_k + (G_{k-1} + G_k)G_{k-1} \cdot G_kG_{k+1}$$

$$G_{k+1}^2 = G_{k+2}G_k + G_{k+1}G_{k-1} + G_k \cdot G_{k-1} - G_k \cdot G_{k+1}$$

$$G_{k+1}^2 = G_{k+2}G_k + G_{k+1} \cdot G_{k-1} + G_k(G_{k+1} - G_{k-1})$$

$$G_{k+1}^2 = G_{k+2}G_k + G_{k+1}G_{k-1} + G_k(-G_k)$$

$$G_{k+1}^2 = G_{k+2}G_k + G_{k+1}G_{k-1} - G_k^2$$

Because, $G_{n+1}G_{n-1} = G_n^2 - (-1)^n \cdot \{G_1^2 - G_1G_0 - G_0^2\}$ then

$$G_{k+1}^2 = G_{k+2} \cdot G_k - (-1)^k \cdot \{G_1^2 - G_1G_0 - G_0^2\}$$

$$G_{k+1}^2 = G_{k+2} \cdot G_k + (-1)^{k+1} \cdot \{G_1^2 - G_1G_0 - G_0^2\}$$

$$\text{We get } G_{k+2} \cdot G_k = G_{k+1}^2 + (-1)^{k+1} \cdot \{G_1^2 - G_1G_0 - G_0^2\}$$

The following identity was derived from Identity 8

Identity 9

$$2G_n^2 = G_{n+1}^2 - G_nG_{n-1} - (-1)^n(G_1^2 - G_1G_0 - G_0^2)$$

Proof. $G_n^2 = G_n \cdot G_n$

$$G_n^2 = G_n \cdot (G_{n+1} - G_{n-1})$$

$$G_n^2 = G_n \cdot G_{n+1} - G_n \cdot G_{n-1}$$

$$G_n^2 = (G_{n+1} - G_{n-1}) \cdot G_{n+1} - G_n \cdot G_{n-1}$$

$$G_n^2 = G_{n+1}^2 - G_{n+1} \cdot G_{n-1} - G_n \cdot G_{n-1}$$

Using Identity 4

$$G_{n+1}G_{n-1} = G_n^2 - (-1)^n \cdot \{G_1^2 - G_1G_0 - G_0^2\}$$

Then:

$$G_n^2 = G_{n+1}^2 \{G_n^2 - (-1)^n \cdot \{G_1^2 - G_1G_0 - G_0^2\}\} \cdot G_{n-1} - G_n \cdot G_{n-1}$$

$$2G_n^2 = G_{n+1}^2 - G_nG_{n-1} - (-1)^n(G_1^2 - G_1G_0 - G_0^2)$$

3. Sum of Cubic for Lucas and Gibonacci Numbers

Based on the Definition above, we can derive an Definition for finding the sum of cubes for Lucas and Gibonacci number as follow:

Identity 10

The sum of cubic of Lucas numbers is

$$\sum_{i=0}^n (L_i)^3 = \frac{L_n(L_{n+1})^2 + 5(-1)^n L_{n-1} + 19}{2}$$

Proof. Using Identity 5: $2L_n^2 = L_{n+1}^2 - L_nL_{n-1} - (-1)^n \cdot 5$, then

$$2L_0^3 = 2L_0^3$$

$$2L_1^3 = 2L_1L_1^2$$

$$2L_1^3 = L_1(L_2^2L_1L_0 + (-1)^15)$$

$$2L_1^3 = L_1L_2^2 - L_1^2 \cdot -5L_1$$

$$2L_2^3 = 2L_2L_2^2$$

$$2L_2^3 = L_2(L_3^2L_2L_1 + (-1)^15)$$

$$2L_2^3 = L_2L_1^2 - L_2^2 \cdot -5L_1$$

$$2L_3^3 = 2L_3L_3^2$$

$$2L_3^3 = L_3(L_4^2L_3L_2 + (-1)^15)$$

$$2L_3^3 = L_3L_4^2 - L_3^2 \cdot -5L_1$$

$$2L_4^3 = 2L_4L_4^2$$

$$2L_4^3 = L_4(L_5^2L_4L_3 + (-1)^15)$$

$$2L_4^3 = L_4L_3^2 - L_4^2 \cdot -5L_1$$

$$2L_5^3 = 2L_5L_5^2$$

$$2L_5^3 = L_5(L_6^2L_5L_4 + (-1)^15)$$

$$2L_5^3 = L_5L_4^2 - L_5^2 \cdot -5L_1$$

$$2L_6^3 = 2L_6L_6^2$$

$$2L_6^3 = L_6(L_7^2L_6L_5 + (-1)^15)$$

$$2L_6^3 = L_6L_5^2 - L_6^2 \cdot -5L_1$$

.

.

.

$$2L_n^3 = 2L_nL_n^2$$

$$2L_n^3 = L_n(L_{n+1}^2L_nL_{n-1} + (-1)^15)$$

$$2L_n^3 = L_nL_{n-1}^2 - L_n^2 \cdot -5L_1$$

Take the sum on both sides simultaneously we get :

$$2(L_0^3 + L_1^3 + L_2^3 + L_3^3 + \dots + L_n^3)$$

$$2(L_0^3 - L_1^2L_0 - 5(L_1 - L_2 + L_3 - L_4 + L_5 - L_6 + \dots - (-1)^n L_n) L_n L_{n+1}^2$$

$$L_n L_{n+1}^2 - 5(L_1 - L_2 + L_3 - L_4 + L_5 - L_6 + \dots - (-1)^n L_n) 2(L_0^3 - L_1^2L_0$$

Next, we need to find the sum of

$L_1 - L_2 + L_3 - L_4 + L_5 - L_6 + \dots - (-1)^n L_n$ By using Identity 2 and 3, we get:

$$-L_2 - L_4 - L_6 + \dots L_{2n} = -(L_{2n+1} + L_1 - L_0) \quad (1)$$

$$-L_1 - L_3 - L_5 + \dots L_{2n-1} = -(L_{2n} - L_0) \quad (2)$$

From equation (3), adding both sides with $L_{2n} - L_0$, equation (3) becomes:

$$-L_2 - L_4 - L_6 + \dots L_{2n} + L_{2n} - L_0 = -(L_{2n+1} + L_1 - L_0) - (L_{2n} - L_0) \quad (3)$$

By using equation (4), equation (5) becomes :

$$\begin{aligned} & -L_2 - L_4 - L_6 + \dots L_{2n} + (-L_1 - L_3 - L_5 + \dots L_{2n-1}) \\ & = -(L_{2n+1} + L_1 - L_0) - L_{2n} - L_0 \\ & L_1 - L_2 + L_3 - L_4 + L_5 - L_6 + \dots + L_{2n-1} - L_{2n} \\ & = -L_{2n+1} + L_1 - L_0 - L_{2n} - L_0 \\ & = -L_{2n+1} + L_1 - L_{2n} \\ & = -L_{2n-1} - L_{2n} \end{aligned} \quad (4)$$

Adding L_{2n+1} to both sides of equation (5), we get :

$$\begin{aligned} & L_1 - L_2 + L_3 - L_4 + L_5 - L_6 + \dots L_{2n-1} - L_{2n} + L_{2n+1} \\ & = -L_{2n-1} - L_1 + L_{2n+1} \\ & L_1 - L_2 + L_3 - L_4 + L_5 - L_6 + \dots L_{2n-1} - L_{2n} + L_{2n+1} \\ & = L_{2n+1} - L_{2n-1} - L_1 \\ & \text{Thus,} \\ & L_1 - L_2 + L_3 - L_4 + L_5 - L_6 + \dots L_{2n-1} - L_{2n} + L_{2n+1} \\ & = L_{2n} - L_1 \end{aligned} \quad (5)$$

By equation (6) and (7), we can conclude that :

$$L_1 - L_2 + L_3 - L_4 + L_5 - L_6 + \dots - (-1)^n L_n = -(-1)^n L_{n-1} - L_1 \quad (6)$$

Therefore, by using equation (8), equation (3) becomes:

$$\begin{aligned} & 2(L_0^3 + L_1^3 + L_2^3 + L_3^3 + \dots + L_n^3) \\ & = L_n L_{n+1}^2 - 5[(-1)^n L_{n-1} - L_1] 2(L_0^3 - L_1^2 L_0) \\ & = L_{n+1}^2 + 5(-1)^n L_{n-1} + 5L_1 + 2L_0^3 - L_1^2 L_0 \end{aligned} \quad (7)$$

By using definition that $L_0 = 2$ and $L_1 = 1$, then:

$$\begin{aligned} & 2(L_0^3 + L_1^3 + L_2^3 + L_3^3 + \dots + L_n^3) = \\ & L_n L_{n+1}^2 - 5[(-1)^n L_{n-1} + 5 + 16 - 2] \\ & = L_n L_{n+1}^2 - 5[(-1)^n L_{n-1} + 19] \end{aligned}$$

$$\text{Therefore : } \sum_{i=0}^n (L_i)^3 = \frac{L_n (L_{n+1})^2 + 5(-1)^n L_{n-1} + 19}{2}$$

□

Example : Given Lucas sequence :

2, 1, 3, 4, 7, 11, 18, 29, 47, 76, 123, 199, ..., then

$$\begin{aligned} & 2(L_0^3 + L_1^3 + L_2^3 + L_3^3 + \dots + L_{10}^3) = \\ & \frac{L_{10}(L_{11})^2 + 5(-1)^{10} L_9 + 19}{2} = \\ & \frac{123(199)^2 + 5(-1)^{10} 76 + 19}{2} = \\ & \frac{123(39601) + 5(-1)^{10} 76 + 19}{2} = \\ & \frac{4870923 + 380 + 19}{2} = \\ & 2435661 \end{aligned}$$

By using direct summation we get :

$$\begin{aligned} & 2(L_0^3 + L_1^3 + L_2^3 + L_3^3 + \dots + L_{10}^3) = \\ & 2^3 + 1^3 + 3^3 + 4^3 + 7^3 + 11^3 + 18^3 + 29^3 + 47^3 + 76^3 + 123^3 = \\ & 8 + 1 + 27 + 64 + 343 + 1331 + 5832 + 24389 + 103823 + \\ & 438976 + 1860867 = \\ & 2435661 \end{aligned}$$

Identity 11

$$\sum_{i=0}^n (G_i)^3 = \frac{G_n(G_{n+1})^2 + G_1 - G_0^3 - 3G_0^2 G_1 + 4G_0^3 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2) G_{n-1}}{2}$$

Proof. By using Identity 9:

$$\begin{aligned} & 2G_n^2 = G_{n+1}^2 - G_n G_{n-1} - (-1)^n (G_1^2 - G_1 G_0 - G_0^2) \\ & 2G_0^3 = 2G_0^3 \\ & 2G_1^3 = 2G_1 G_1^2 \\ & 2G_1^3 = G_1 (G_2 - G_1 G_0 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2)) \\ & 2G_1^3 = G_1 G_2^2 - G_2^2 \cdot G_0 + (G_1^2 - G_1 G_0 - G_0^2) G_1 \\ & 2G_2^3 = 2G_2 G_2^2 \\ & 2G_2^3 = G_2 (G_3 - G_2 G_1 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2)) \\ & 2G_2^3 = G_2 G_3^2 - G_3^2 \cdot G_0 + (G_1^2 - G_1 G_0 - G_0^2) G_2 \\ & 2G_3^3 = 2G_3 G_3^2 \\ & 2G_3^3 = G_3 (G_4 - G_3 G_2 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2)) \\ & 2G_3^3 = G_3 G_4^2 - G_4^2 \cdot G_0 + (G_1^2 - G_1 G_0 - G_0^2) G_3 \\ & 2G_4^3 = 2G_4 G_4^2 \\ & 2G_4^3 = G_4 (G_5 - G_4 G_3 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2)) \\ & 2G_4^3 = G_4 G_5^2 - G_5^2 \cdot G_0 + (G_1^2 - G_1 G_0 - G_0^2) G_4 \\ & 2G_5^3 = 2G_5 G_5^2 \\ & 2G_5^3 = G_5 (G_6 - G_5 G_4 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2)) \\ & 2G_5^3 = G_5 G_6^2 - G_6^2 \cdot G_0 + (G_1^2 - G_1 G_0 - G_0^2) G_5 \\ & 2G_n^3 = 2G_n G_n^2 \\ & 2G_n^3 = G_n (G_{n+1} - G_n G_{n-1} - (-1)^n (G_1^2 - G_1 G_0 - G_0^2)) \\ & 2G_n^3 = G_n G_{n+1}^2 - G_n^2 \cdot G_{n-1} - (-1)^n (G_1^2 - G_1 G_0 - G_0^2) G_n \end{aligned}$$

By taking the sum of both sides simultaneously we get:

$$2(G_0^3 + G_1^3 + G_2^3 + G_3^3 + \dots + G_n^3) = 2G_0^3 - G_1^2 \cdot G_0 + (G_1^2 - G_1 G_0 - G_0^2)(G_1 - G_2 + G_3 - G_4 + G_5 - G_6 + \dots - (-1)^n G_n) + G_n G_{n+1}^2 \quad (8)$$

Next, we need to find

$(G_1 - G_2 + G_3 - G_4 + G_5 - G_6 + \dots - (-1)^n G_n) + G_n G_{n+1}^2$ By using Identity 2, we get :

$$-G_2 - G_4 - G_6 \dots - G_{2n} = -(G_{2n+1} - G_1) \quad (9)$$

and Identity 7:

$$G_1 + G_3 + G_5 \dots + G_{2n-1} = G_{2n} - G_0 \quad (10)$$

From equation (11), add both sides of the equation with $G_{2n} - G_0$ then equation (12) becomes :

$$-G_2 - G_4 - G_6 \dots - G_{2n} + G_{2n} - G_0 = (G_{2n+1} - G_1) + G_{2n} - G_0 \quad (11)$$

By equation (12), equation(13) will be :

$$\begin{aligned} -G_2 - G_4 - G_6 \dots - G_{2n} + G_{2n} + G_1 + G_3 + G_5 \dots + G_{2n-1} \\ = (G_{2n+1} - G_1) + G_{2n} - G_0 \\ G_1 - G_2 + G_3 - G_4 + G_5 + G_6 + \dots + G_{2n-1} - G_{2n} \\ = G_{2n+1} + G_{2n} + G_1 - G_0 \quad (12) \end{aligned}$$

From equation (13), add both sides with G_{2n+1} , we get:

$$\begin{aligned} G_1 - G_2 + G_3 - G_4 + G_5 + G_6 + \dots + G_{2n-1} - G_{2n} + G_{2n+1} \\ = G_{2n+1} + G_1 - G_0 + G_{2n+1} \\ = G_{2n+1} - G_{2n-1} + G_1 - G_0 \end{aligned}$$

By using the definition we get

$G_{2n} = G_{2n+1} - G_{2n-1}$ thus,

$$\begin{aligned} G_1 - G_2 + G_3 - G_4 + G_5 + G_6 + \dots + G_{2n-1} - G_{2n} + G_{2n+1} \\ = G_{2n} + G_1 - G_0 \quad (13) \end{aligned}$$

From equations (14) and (15), we can conclude that :

$$\begin{aligned} G_1 - G_2 + G_3 - G_4 + G_5 - G_6 + \dots - (-1)^n G_n \\ = -(-1)^n G_{n-1} + G_1 - G_0 \quad (14) \end{aligned}$$

By using equation (16), equation (17) become:

$$\begin{aligned} 2G_0^3 + G_1^3 + G_2^3 + G_3^3 + \dots + G_n^3 \\ = 2G_0^3 - G_1^2 \cdot G_0 + (G_1^2 - G_1 G_0 - G_0^2) \\ [-(-1)^n G_{n-1} + G_1 - G_0] + G_n G_{n+1}^2 \\ = G_n G_{n+1}^2 + 2G_0^3 - G_1^2 G_0 - \\ (-1)^n + G_1^2 - G_1 G_0 - G_0^2 G_{n-1} + G_1 - G_0 \\ = G_n G_{n+1}^2 + 2G_0^3 - G_1^2 G_0 + (G_1^2 - G_1 G_0 - G_0^2) \\ [-(-1)^n G_{n-1} + (G_1^2 - G_1 G_0 - G_0^2)(+G_1 - G_0)] \\ = G_n G_{n+1}^2 + 2G_0^3 - G_1^2 G_0 + (G_1^2 - G_1 G_0 - G_0^2) \\ (+G_1 - G_0) - (-1)^n (G_1^2 - G_1 G_0 - G_0^2) G_{n-1} \\ = G_n G_{n+1}^2 + 3G_0^3 - G_1^3 3G_1^2 G_0 - \\ (-1)^n (G_1^2 - G_1 G_0 - G_0^2) G_{n-1} \\ = G_n G_{n+1}^2 + (G_1 - G_0)^3 + 4G_0^3 - 3G_0^2 G_1 - \\ (-1)^n (G_1^2 - G_1 G_0 - G_0^2) G_{n-1} \quad (15) \end{aligned}$$

Therefore:

$$G_0^3 + G_1^3 + G_2^3 + \dots + G_n^3 = \frac{[G_n(G_{n+1})^2 + (G_1 - G_0)^3 - 3G_0^2 G_1 + 4G_0^3 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2) G_{n-1}]}{2} \quad \square$$

Example : For $G_0 = 3$ and $G_1 = 2$, the Gibonacci sequence is 3, 2, 5, 7, 12, 19, 31, 50, 81, 131, 212, ...

$$\begin{aligned} G_0^3 + G_1^3 + G_2^3 + \dots + G_9^3 = \\ \frac{[G_9(G_{10})^2 + (G_1 - G_0)^3 - 3G_0^2 G_1 + 4G_0^3 - (-1)^9 (G_1^2 - G_1 G_0 - G_0^2) G_8]}{2} \\ = \frac{[131(212)^2 + (2-3)^3 - 3 \cdot 3^2 \cdot 2 + 4 \cdot 3^3 - (-1)^9 (2^2 - 2 \cdot 3 - 3^2) 81]}{2} \end{aligned}$$

$$G_0^3 + G_1^3 + G_2^3 + \dots + G_9^3 = \frac{[(131)(44944) - 1 - 54 - 108 + (-1)81]}{2}$$

$$G_0^3 + G_1^3 + G_2^3 + \dots + G_9^3 = \frac{[5887664 - 1\lambda 54 + 108 - 891]}{2} = 2943413$$

By using direct summation we get :

$$\begin{aligned} G_0^3 + G_1^3 + G_2^3 + \dots + G_9^3 \\ = 3^3 + 2^3 + 5^3 + 7^3 + 12^3 + 19^3 + 31^3 + 50^3 + 81^3 + 131^3 \\ = 9 + 8 + 125 + 343 + 1728 + 6859 + \\ 29791 + 125000 + 531441 + 2248091 = 2943413 \end{aligned}$$

4. CONCLUSIONS

From the discussion above we can conclude that the sum of cubic for Lucas number is $L_0^3 + L_1^3 + L_2^3 + L_3^3 + \dots + L_n^3 = \sum_{i=0}^n (L_i)^3 = \frac{L_n(L_{n+1})^2 + 5(-1)^n L_{n-1} + 19}{2}$ and for Gibonacci numbers is

$$\begin{aligned} G_0^3 + G_1^3 + G_2^3 + G_3^3 + \dots + G_n^3 = \\ \sum_{i=0}^n (G_i)^3 = \frac{[G_n(G_{n+1})^2 + (G_1 - G_0)^3 - 3G_0^2 G_1 + 4G_0^3 - (-1)^n (G_1^2 - G_1 G_0 - G_0^2) G_{n-1}]}{2} \end{aligned}$$

REFERENCES

- Arangala, C., B. Hrovat, and J. Kelner (2016). Sum of Powers of Fibonacci and Lucas Numbers. *The Minnesota Journal of Undergraduate Mathematics*, 2(1); 1-12
- Benjamin, A. T, T. A. Carnes, and B. Cloitre (2009). Recounting the Sums of Cubes of Fibonacci Numbers. *Congressus Numerantium*, Proceedings of the Eleventh International Conference

- on Fibonacci Numbers and their Applications. *William Webb, ed*, **194**; 45–51
- Benjamin, A. T, and J. J. Quinn (1999). Recounting Fibonacci and Lucas Identities. *College Mathematics Journal*, **30**(5); 359 – 366
- Benjamin, A. T, and J. J. Quinn (2003). *Proofs That Really Count : The Art of Mathematical Association of America*. Washington DC
- Benjamin, A. T, J. J. Quinn, and F. E. Su (2000). Phased Tilings and Generalized Fibonacci. *The Fibonacci Quarterly*, **38**(3); 282–288
- Carlitz, L., R. Scoville, and V. E. Hoggatt (1972a). Fibonacci Representations. *Fibonacci Quarterly*, **10**(1); 1 – 28
- Carlitz, L., R. Scoville, and V. E. Hoggatt (1972b). Fibonacci Representations. *Fibonacci Quarterly*, **10**(1); 29 – 42
- Carlitz, L., R. Scoville, and V. E. Hoggatt (1972c). Fibonacci Representations of Higher Order. *Fibonacci Quarterly*, **10**(1); 71 – 80
- Carlitz, L., R. Scoville, and V. E. Hoggatt (1972d). Fibonacci Representations of Higher Order. *Fibonacci Quarterly*, **10**(1); 43 – 70
- Dunlap, R. A. (2003). *The Golden Ratio and Fibonacci Numbers*. World Scientific Press, Singapore
- Frontczak, R. (2018). Sums of powers of Fibonacci and Lucas numbers: A new bottom-up approach. *Notes on Number Theory and Discrete Mathematics*, **24**(2); 94 –103
- Koshy, T. (2001). *Fibonacci and Lucas Numbers with Application*. John Wiley & Sons Inc., New York
- Krishna, H. V. (1980). Identities of A Generalized Fibonacci Sequence. *The Fibonacci Association*, **18**; 65 – 66
- Posamentier, A. S, and I. Lehmann (2007). *The Fabulous Fibonacci Numbers*. Prometheus Books, New York
- William, H. C. (1972). Fibonacci Numbers Obtained from Pascal's Triangle with Generalization. *Fibonacci Quarterly*, **10**(4); 405 – 412