

CONFERENCE PROCEEDING



Faculty of Agricultural Technology
in Collaboration with
Indonesian Association for Microbiology



2nd ICGAB 2018
International Conference on
Green Agro-Industry and Bioeconomy



FKPT-TPI



**Indonesian Biotechnology
Consortium**

The selected papers will be published conference proceeding indexed by Scopus
IOP Publishing **IOP Conference Series**
Earth and Environmental Science



Sweet potato greens ‘neglected vegetables rich in bioactive compounds’ (part I): radical scavenging activity, inhibitory effect on α - amylase, total phenolic and flavonoid contents of local sweet potato (*Ipomoea batatas*) leaves

S Nurdjanah¹, N Yuliana¹, O Nawansih¹, R Dewi²

¹Department of Agriculture Product Technology, Faculty of Agriculture, University of Lampung Indonesia

²Department of Agronomy Politeknik Negeri Lampung, Indonesia

E-mail: siti.nurdjanah@fp.unila.a.c.id

Abstract. Sweet potato greens or leaves are commonly consumed by rural people in some part of Indonesia, Malaysia, The Philippines, and other part of the world. In addition to their nutritive value, sweet potato leaves have been reported to content an appreciable amount of bioactive compounds such as polyphenol including flavonoids beneficial for health. Over the years, phenolic including flavonoids are intensely studied for their physiological health benefits as functional foods. However information on bioactive contents of sweet potato greens grown locally in Lampung are inexistent, therefore this paper reports some components believed to be potential for human health. The leaves of locally grown three sweet potato genotypes used in this study were harvested six weeks after planting. The results showed methanol extract of sweet potato leaves had 63-79% DPPH radical scavenging activity 55-59 % inhibitory effect on α amylase , 0.81-1.75 g GAE/100 g dry weight of total phenolic content, and 7.57-10.51mg QE/100 g dry weight of flavonoids.

1. Introduction

Sweet potatoes are cultivated widely around the world primarily for their underground tubers utilized as a raw material for starch production, staple and snack foods . They can be harvested up to three times a year. In addition to be regarded as biomass waste after harvesting the tubers, sweet potato leaves are regularly harvested and consumed in some Asian countries mainly because it has a high content of protein, mineral substances, vitamins and bioactive compounds as well as soluble and non-soluble dietary fiber. Sweet potato leaves contain up to 15-27% protein on dry matter basis depending on the varieties [1] , and are a valuable source of iron, vitamins E, beta carotene lutein and polyphenol [2]. These compounds have been reported to have antioxidant properties, inhibit α glucosidase and α amylase activities [3] which is very potential to manage blood glucose level in diabetic patients.

Bioactive components, namely polyphenol, of sweet potato leaves have been intensively studied because of their ability to protect against free radicals. The over production of free radicals and nitrogen species could result in oxidative stress that increases the onset of chronic non-communicated diseases (NCDs) such as some types of cancers, neurological problem, and cardiovascular[4,5]. The main polyphenol contents of sweet potato leaves are caffeoylquinic acid (CQA) derivatives[6]. The amount of polyphenol in leaves is higher than those of in the tuber flesh and the peel[7]. Eugenio et al.[8] reported sweet potato leaves contain high amount of phenolic compounds mainly quercetin, chlorogenic acid, and rosmarinic .

In Lampung areas, sweet potatoes are grown widely among small growers. However, despite of their valuable content of protein and bioactive compounds , sweet potato leaves have no economic value, they are either fed to cattle or utilized as compost. Therefore it is important to provide information on various health-promoting biological activities of sweet potato leaves to encourage people to utilize sweet potato leaves as part of their healthy diet. This study revealed the antioxidant property, α amylase inhibitor, flavonoid and total phenolic contents of three sweet potato genotypes namely LPG-01, LPG-06, and LPG-07 grown at Politeknik Negeri Lampung Experimental Field. These sweet potato genotypes are still on-going experiment and have not been grown commercially.

2. Material and Methods

2.1 Materials

Three sweet potato genotypes namely LPG-01, LPG-06, and LPG-07 grown at Politeknik Negeri Lampung Experimental Field were used in this study. Sweet potato leaf tips used in this study had the desirable attribute for human consumption. They had small stems and petioles they were glabrous, tender and showed some vein purple color. The 15 cm long tips were harvested 6 weeks after transplanting. For each genotype, harvested fifteen sweet potato tip stems were rinsed with tap water, then steam blanched for 3 min, and cooled at room temperature. They were then cut into small pieces, put into plastic bags, and frozen at -15°C until analysis was performed. The samples were evaluated for their DPPH radical scavenging activity , α amylase inhibition, total phenolic, and total flavonoid contents. The chemicals such as Folin-Ciocalteau reagent, DPPH, methanol were purchased from Sigma-Aldrich via local supplier.

2.2 Sweet Potato Leaf Extraction

Samples of frozen blanched sweet potato leaves (50g) were macerated in 100 ml of 60% methanol in a closed container for 24 h at 4°C in the dark room, then filtered through Whatman No. 42 filter paper. The remaining residues were washed with 100 ml of methanol, both filtrates were collected as the crude extract filtrate. The methanol in the crude extract was evaporated by the rotary evaporator (0.1 MPa, 40°C) until the volume was 50 mL, kept in a dark brown bottle and stored at -20°C until further analysis.

2.3 Scavenging Radical Activity

DPPH (2,2-diphenyl-1-picrylhydrazyl) radical-scavenging activity of leaf extract was determined using method described by Chung et al. [9] with slight modification. Two μ L of the methanol extracted leaves (as described in above procedure) were mixed with 2.0 mL of 2×10⁻⁴ M DPPH in methanol. The mixture was shaken vigorously and left in the dark at 25°C for 30 min. The absorbance of mixture was read immediately at 517 nm using a GENESYS 10S S UV-Visible spectrophotometer (A_{sample}). The mixture of 95% methanol (2 mL) and sample (2 mL) serve as blank (A_{blank}). The control solution was prepared by mixing methanol (2 mL) and DPPH radical solution (2 mL) ($A_{control}$). The ability of sweet potato leaf extract to scavenge DPPH radical or antioxidant activity was calculated by the following equation:

$$\text{Scavenging Radical Activity} = \left[100 - \frac{A_{sample} - A_{blank}}{A_{control}} \right] \times 100 \%$$

2.4 Invitro α -amylase Inhibition Study

The α -amylase inhibitory activity was determine using method described by Zengin[10] with modifications. Briefly, 250 μ l of leaf extracts with varying concentrations (125–500 μ g/ml) and 250 μ l of 0.02 M sodium phosphate buffer (pH 6.9 with 0.006 M sodium chloride) containing α -amylase (from *Aspergillus oryzae* , Merck) solution (0.5 mg/ml) were incubated for 10 min at 25°C. After preincubation, 250 μ l of 1% starch solution in 0.02 M sodium phosphate buffer (pH 6.9 with 0.006 M sodium chloride) was added to each tube at 5 s intervals. The reaction mixtures were then incubated at 25°C for 10 min. The reaction was stopped with 500 μ l dinitrosalicylic acid (Sigma, St. Louis, USA) color reagent. The tubes were then incubated in a boiling water bath for 5 min and cooled to room

temperature. The reaction mixture was then diluted by adding 5 ml of distilled water, and absorbance was read at 540 nm using a GENESYS 10S S UV-Visible spectrophotometer. the α -amylase inhibitory activity was calculated as percentage inhibition, using the formula.

$$\% \text{ Inhibition} = ((\text{Abs}_{\text{control}} - \text{Abs}_{\text{samples}}) / \text{Abs}_{\text{control}}) \times 100$$

2.5 Total Phenolic

Total phenolic content in sweet potato leaf extract was determined by colorimetry method using Folin-Ciocalteu reagent assay as modified from Singleton and Rossi [11]. SP leaf extract (1ml) was mixed with 1ml of Folin-ciocalteu's phenol reagent and allowed to react for 3 minutes. Then, 0.8ml of 7.5% (w/v) sodium carbonate was added. The mixture was agitated and allowed to stand for further 30 minutes in the dark. The absorbance of leaf extracts and a prepared blank were read at 765 nm using GENESYS UV-Visible spectrophotometer. Gallic acid with the concentration range from 10 $\mu\text{g}/\text{mL}$ to 100 $\mu\text{g}/\text{mL}$, was used for the standard curve and the total phenolic content was expressed as mg gallic acid equivalents per gram of dried sample (g GAEs/100g dry weight).

2.6 Total Flavonoids

Total flavonoid content was determined using the method described by Vuong et al. [12] with slight modification. Extract (0.5 mL) was added 2 mL of deionized (DI) water and 0.15 mL of 5% (w/v) NaNO_2 , kept at 25°C for 6 min. Then, 0.15 mL of 10% (w/v) of AlCl_3 was added and allowed to stand for 6 min, followed by the addition of 2 mL of 4% (w/v) NaOH and 0.7 mL of DI water. The mixture was mixed thoroughly and left at 25°C for 15 min. The absorbance was read at 380 nm using GENESYS UV-Visible spectrophotometer. Quercetin with the concentration range from 10 $\mu\text{g}/\text{mL}$ to 100 $\mu\text{g}/\text{mL}$ was used as a standard and the total flavonoid content was expressed as mg quercetin equivalents per gram of dried sample (mg QEs/100g dry weight).

2.7 Statistical Analysis

All the analysis were conducted in triplicates for each genotype, and the data were reported as means \pm standard deviation

3. Results and Discussion

The results of scavenging activity, enzyme inhibition, total phenolic and total flavonoid contents are summarised in Table 1.

3.1 Scavenging Radical Activity

The DPPH radical scavenging activity of the three sweet potato genotypes can be ranked as LPG-7 > LPG-6 > LPG-2. The difference in total antioxidant activity among sweet potato varieties grown under similar condition might be influenced by the plants tolerance against viruses and fungi [13]. DPPH is a free radical generating compound used to measure the scavenging ability of sweet potato leaf extract. The radical scavenger of the leaf extract will decolorize the color of DPPH in methanol solution (purple) to yellow color because of reduction of the stable DPPH radicals to diphenyl-pricrylhydrazine in the presence of H-donating antioxidant . In this study, the antioxidant property of the sweet potato leaf extract was determined using only single method (DPPH assay). The decision was based on its availability and simplicity, furthermore Steed and Truong [14] reported that the antioxidant activities assayed by DPPH and ORAC (oxygen radical absorbance capacity) showed a significant correlation.

3.2 α -Amylase Inhibition

The inhibitory effect on α -amylase of three genotypes sweet potato leaf extracts were between 55 to 60%. LPG-7 leaf extract showed the strongest inhibitory effect , followed by LPG-2 and LPG-6. The variation of inhibitory effect on α -amylase could be due to variation in the phenolic content. It was claimed that the higher levels of chlorogenic acid (one type of phenolic) may contribute to α -amylase and α -glucosidase inhibitory effect [15].

3.3 Total Phenolic

The total phenolic contents among leaf extract from three genotypes sweet potato were between 0.82 and 1.75g GAEs/100 g extract. Phenolic compounds contribute to the ability to scavenge the reactive oxygen species (ROS)^[12], and therefore are important to keep healthy body. Phenolics also has important role in modulating the carbohydrate absorption [16], which may be used to manage blood glucose in diabetic patients.

3.4 Total Flavonoids

The flavonoids contents were between 7.57 – 10.51 mg QEs/100g dry weight. The highest was found in LPG-6 leaf extract, followed by those of LPG-2 and LPG-7. Flavonoids have been intensely studied mainly because of their function as antioxidant . The mechanism of flavonoids as antioxidant or in removing free radicals can be divided in to two, which is scavenging and chelating [17].

Table 1. Antioxidant ability, enzyme inhibition, total phenolic and total flavonoid of sweet potato leaf extracts

Sampel	DPPH Scavenging radical activity (%)	α -amylase inhibition (%)	Total Phenolic (g GAEs/100g dry weight)	Total Flavonoid (mg QEs/100g dry weight)
LPG-2	79.68±0.13	57.95±0.05	1.75±0.27	8.23±0.51
LPG-6	75.72±0.24	54.92±0.03	0.81±0.09	10.51±0.72
LPG-7	63.39±0.36	59.84±0.15	1.49±0.05	7.57± 0.18

4. Conclusion

The methanol extract of sweet potato leaves from 3 genotypes have shown to contain considerable level of antioxidant properties, enzyme inhibition, total phenolic and total flavonoid. Based on these findings, sweet potato leaves are very potential to be promoted as promising natural source of antioxidant, and α -amylase inhibition that could be used to overcome hyperglycemia problem.

References

- [1] Nkongho G O, Achidi AU, Ntonifor1 N N, Numfor FA, Dingha B N, Louis E N, Jackai and Bonsi C K 2014 Sweet potatoes in Cameroon: Nutritional profile of leaves and their potential new use in local foods *Afric J of Agric. Res.* **9(18)**1371-1377
- [2] Chandrika UG, Basnayake BM, Athukorala I, Colombagama PW, Goonetilleke A 2010 Carotenoid content and in vitro bioaccessibility of lutein in some leafy vegetables popular in Sri Lanka. *J Nutr Sci Vitaminol (Tokyo)* **56** 203–207
- [3] Hue S-M, Boyce AN and Somasundram C 2012 Antioxidant activity, phenolic and flavonoid contents in the leaves of different varieties of sweet potato (*Ipomoea batatas*) *Austral. J. of Crop Sci.* **6(3)** 375-380
- [4] Valko M, Rhodes C J, Moncol J, Izakovic M, Mazur M 2006 Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-Biological Interactions* **160(1)** 1-40.
- [5] Moreira P I, Santos M S, Oliveira C R, Shenk J C, Nunomura A, Smith M A, Zhu X, Perry G 2008 Alzheimer disease and the role of free radicals in the pathogenesis of the disease. *CNS & Neurological Disorders - Drug Targets* **7.3** –10. 3
- [6] Islam M S, Yoshimoto M, Yahara S, Okuno S, Ishiguro K, Yamakawa O 2002 Identification and characterization of foliar polyphenolic composition in sweetpotato (*Ipomoea batatas* L.) genotypes *J Agric Food Chem* **50** 3718–3722

- [7] Truong V D, McFeeters R F, Thompson R T, Dean L L, Shofran B 2007 Phenolic acid content and composition in leaves and roots of common commercial sweetpotato (*Ipomea batatas* L.) cultivars in the United States *J Food Sci.* **72** C343–C349
- [8] Eugenio M H A, Pereira R G F A, César de Abreu W and Pereira M C A 2017 Phenolic compounds and antioxidant activity of tuberous root leaves *Intl. J of Food Properties* **20**(12) 2966-2973
- [9] Chung Y C, Chen S J and Hsu C K 2005 Studies on the antioxidative activity of *Graptophetalum paraguayense* E. Walther. *Food Chem.* 91419-424
- [10] Zengin G 2016 A study on in vitro enzyme inhibitory properties of Asphodeline Anatolica: New Sources of Natural Inhibitors for Public Health Problems *Ind. Crops Prod.* **83** 39–43
- [11] Singleton V L and Rossi J A Jr 1965 Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents *Am J Enol Vitic.* **16** 144-158
- [12] Vuong Q V Hirun S, Roach P D, Bowyer M C, Phillips P A, Scarlett C J 2013 Effect of extraction conditions on total phenolic compounds and antioxidant activities of *Carica papaya* leaf aqueous extracts. *J. Herb. Med.* 3: 104–111.
- [13] Osiru MO, Olanya OM, Adipala E, Kapinga R, Lemaga B. 2009. Yield stability analysis of *Ipomoea batatas* L. cultivars in diverse environments. *Aust J Crop Sci.* 3(4): 213-220
- [14] Steed, L. E., and V. D. Truong. 2008. Anthocyanin content, antioxidant activity and selected physical properties of flowable purple-fleshed sweet potato purees *J Food Sci.* **73** 215–221
- [15] Zengin G, Menghini, L., Malatesta, L., De Luca, E., Bellagamba, G., Uysal, S., Aktumsek A, Locatelli M 2016 Comparative study of biological activities and multicomponent pattern of two wild Turkish species: Asphodeline Anatolica and Potentilla Speciosa. *J Enzyme Inhib Med Chem.* 31 sup1203–208
- [16] Jeng T O, Chiang Y C, Lai C C, Liao C T, Lin Y S, Lin C T and Sung J M 2015 Sweet potato leaf extract inhibits the simulated in vitro gastrointestinal digestion on native starch *J Food and drug Anal.* **23** 399-406
- [17] Cook N C and Samman S 1996 Flavonoids- chemistry, metabolism, cardioprotective effects, and dietary sources. *Nutr Biochem.* 7 66-76