

Jurnal TPCJ, Hendri Busman agust 2018

By Hendri Busman

WORD COUNT

2159

TIME SUBMITTED

19-JUN-2019 11:28AM

PAPER ID

48001455



Antispermatic Effect of Seeds Extract of Papaya (*Carica papaya* L.) in Mice

Nuning Nurcahyani, Hendri Busman, Sutyarso, Pratami Dwi Rahmawati, Mohammad Kanedi*

6

Department of Biology, Faculty of Maths and Sciences, University of Lampung, Bandar Lampung, Indonesia

31

Abstract To enrich new evidence about the anti-fertility activity of seeds extract of papaya (*Carica papaya* L.) in test animals, especially mice. Male mice (n=20) were grouped into four. The first group received only distilled water containing 1% CMC as the control. Group 2, 3 and 4 consecutively received papaya seeds extract of 2, 4 and 8 mg/40 g body weight. All treatment administered orally using stomach sonde once daily for 35 days. Study parameters assessed were the number spermatogonia, spermatocyte and spermatid cell counts, diameter and epithelial thickness of seminiferous tubules of the testis. There is no significant different in the number of spermatogonia between treatment groups. Spermatocyte and spermatid cells significantly decreased by the extract, especially at the highest dose. Additionally, the diameter and epithelial thickness of seminiferous tubules of testis are significantly reduced by the extract at the dose of 8 mg/40g body weight. Papaya seeds extract as potential to be used as anti fertility agent.

Keywords: Carica papaya, contraceptives, papaya seeds, antispermatic, antifertility

1. Introduction

There are many medicinal plants that are reported to have potential as anti-fertility agents. However, to be acceptable the ingredient must be eligible, among others causing azoospermia (cement), easy to use, no side effects and toxic, not interfere with libido or sexual behavior and is reversible [1].

Papaya (*Carica papaya* L.) is one of tropical plants worth considering to be developed as a contraceptive. In Langur monkey seeds extract of papaya revealed to cause tubules of testis shrunken, vacuolation of Sertoli cells and abnormalities of germ cells, and inhibition of steroidogenic function of Leydig cells [2]. In Wistar rats, seeds extract of this plant at the dose of 50-200 mg/kg caused hypertrophy of pituitary gonadotrophs, gradual degeneration of germ cells, Sertoli cells and Leydig cells as well as germinal epithelium, progressive collapse dan shrank villi of seminal vesicles [3]. In mice, treatment of seed extract of papaya for 7 days caused the decrease in sperm concentration, motility and viability [4].

Although the results of research on the antifertility activity of papaya seed extract showed a positive trend, but in-depth verification of the effects of these plant preparations on test animals is still required as recommended by the World Health Organization [5]. This study is intended to enrich new evidence about the anti-fertility activity of papaya seed extracts in test animals, especially mice. In the study reported Wiryawan, giving extract only for 7 days. In this research, papaya seed extract was given for 35 days, based on the duration of one cycle of spermatogenesis.



2. Materials and Methods

Plant Materials

Seed samples of papaya (*Carica papaya*) used in the study were originated from ripen fruit collected from a local papaya farmer in suburb of Bandar Lampung, Indonesia. The sun dried seeds were milled into powder and then macerated using methanol. After vacuum evaporation, the viscous extract then suspended in 1% CMC in distilled water in accordance with the prescribed treatment doses.

20

Animals and Treatment

Twenty male Swiss albino mice, aged 4 months and weighing 25-30 g from Lampung Veterinary Center, Indonesia, were used and divided into four groups. The first group received only distilled water containing 1% CMC as the control. Group 2, 3 and 4 consecutively received papaya seeds extract of 2, 4 and 8 mg/40 g body weight. All treatment administered orally using stomach sonde once daily for 35 days. On day 36 all mice were sacrificed and the testes were taken for histological preparation using formalin as a fixative and stained using hematoxylin-eosin. Microscopic slides of the testes were then observed using light microscope at 400x magnification.

Study Parameters and Data Analyses

Study parameters assessed were spermatogonia cell counts, spermatocyte and spermatid cell counts, diameter and epithelial thickness of seminiferous tubules of the testis. To detect intergroup differences one-way ANOVA was performed, followed by Fisher's least significant difference (LSD) test to detect differences between individual groups at the $p < 0.05$.

3. Results

Spermatogenic cell count of mice after daily treatment of papaya seeds extract for 35 days are shown in Table 1. There is no significant different in the number of spermatogonia between treatment groups. However, both spermatocyte and spermatid cells significantly affected by the extract, especially at the highest dose. Table 2 depict the effect of papaya seeds extract on the diameter and epithelial thickness of seminiferous tubules of testis. The data showed that at the dose of 8 mg/40g body weight papaya seeds extract significantly decreased diameter of seminiferous tubules. Furthermore, in comparison to control, all treatment doses significantly decreased the epithelial thickness of the tubules. The histological changes in diameter and epithelial thickness of the seminiferous tubules of mice of different treatment are visualized in Fig.1.

Table 1: Spermatogenic cell counts of mice after papaya seeds extract treatment for 35 days

Papaya seeds extract	Spermatogonia	Spermatocytes	Spermatids
0 (mg/40g bw)	59.60 ± 3.78 ^a	185.40 ± 23.00 ^a	176.60 ± 28.43 ^a
2 (mg/40g bw)	53.80 ± 4.32 ^a	138.80 ± 30.06 ^b	152.40 ± 18.03 ^a
4 (mg/40g bw)	51.00 ± 13.09 ^a	108.60 ± 24.79 ^{bc}	137.20 ± 14.42 ^{ab}
8 (mg/40g bw)	49.80 ± 7.66 ^a	90.04 ± 14.63 ^b	127.20 ± 21.34 ^b

Data are presented as mean±SD. Values in the same column followed by the same superscript are not significantly different at $\alpha < 0.05$ by LSD test

Table 2: Diameter and epithelial thickness of seminiferous tubules of mice after papaya seeds extract treatment for 35 days

Papaya seeds extract	Diameter of Seminiferous Tubule (µm)	Thickness of Seminiferous epithelium (µm)
0 (mg/40g bw)	192.60±8.26 ^a	45.30±5.46 ^a
2 (mg/40g bw)	179.80±7.59 ^a	26.50± 7.00 ^b
4 (mg/40g bw)	177.80±9.8 ^{ab}	24.60± 4.70 ^b
8 (mg/40g bw)	176.20±18.68 ^b	21.10± 2.43 ^b

Data are presented as mean±SD. Values in the same column followed by the same superscript are not significantly different at $\alpha < 0.05$ by LSD test






Figure 1: Micrographs (400x) of cross section of seminiferous tubules of mice treated with papaya seeds extract: 1) Control group; 2) 2 mg/40 g; 3) 4 mg/40 g; 4) 8 mg/40 g body weight.

4. Discussion

The data in Tables 1 and 2 as well as the photographs in Fig.1 clearly imply that the papaya seeds extract has anti-spermatogenic activity. Phytochemical screening studies showed that papaya seeds are rich in alkaloids, tannins, saponins, phenols, terpenoid, flavonoid and sterol [6-8]. There are many studies that confirm the antifertility effects of chemical compounds similar to those of papaya seeds. *Citrullus colocynthis* is revealed to contain flavonoids, saponins and steroid [9]. Methanolic extract of this plant is suggested to have contraceptive properties because proven to decrease in testicular weight, sperm concentration, level of testosterone, LH and FSH, in addition to cause impairment of spermatogenesis in rats [10]. Another plant species that also indicates antifertility properties is *Chenopodium album*. In rats treated with this plant seeds extract is known to induce instantaneous immobilization of spermatozoa allegedly due to disintegration of plasma membrane and dissolution of acrosomal cap [11]. This plant, as indicated, also contained chemicals similar to papaya seeds extract mainly flavonoids [12].

How flavonoids can affect spermatogenesis? Apparently, these substances have inhibitory activity on aromatase, an enzyme that plays an important role in catalyzing the conversion of androgens into estrogens. Due to inhibition of the enzymes, testosterone levels increase and affect pituitary function via a negative feedback mechanism so that the secretion of FSH and LH is suppressed that lead to inhibition of spermatogenesis [13]. Saponin is another instance of phytochemicals that also contained in papaya seeds extract. Saponins, mainly steroids and triterpenoids, allegedly also give effects on the hypothalamus-pituitary-testis axis in laboratory mammals that potentially lead to decrease spermatocyte and spermatid counts [14].

Data in Table 2 and Fig.1 clearly showed that papaya seeds extract affect diameter and epithelial thickness of seminiferous tubules of testis. These phenomena may also associate with function of the hypothalamus-pituitary-testis axis. Decreases in GnRH and gonadotropin (FSH and LH) will inhibit the process of spermatogenesis due to degeneration of the seminiferous tubules resulting in decreased diameter and epithelial thickness of the tubules [15].



Although significantly affecting the process of spermatogenesis, papaya seed extract did not show any significant change in the number of spermatogonia. This may be related to the testicular immune properties, especially spermatogonial stem cells, which are less sensitive to injury, capable of self-renewal [16-17].

5. Conclusion

Given proven capable of reducing the number of spermatocyte and spermatid cells and reducing the diameter and epithelial thickness of seminiferous tubules of mice testes, it can be inferred that papaya seeds extract can be used as anti fertility agent.

10

References

1. Shivabasavaiah. Antifertility Effects of Madhuca Indica Leaves in Male Swiss Albino Rats. 2011; *Journal of Pharmacy Research* 2011; 4(2): 323-326.
2. Lohiya NK, Manivannan B, Mishra PK, Pathak N, Sriram S, Bhande SS, Panneerdoss S. Chloroform extract of *Carica papaya* seeds induces long-term reversible azoospermia in Langur monkey. *Asian J Androl.* 2002. 4 (1): 17-26
3. Udoh P, Essien I, and Udoh F. Effects of *Carica papaya* seed extract on the morphology of pituitary-gonad axis of male Wistar rats. *Phytoth Res.* 2005. 19(14): 65-1068.
4. Wiryawan R.A., I'tishom R. and Purwaningsih S. Papaya Seed Extract Lowers Sperm Concentrations, Fertility and Viability in Male Mice. *Folia Medica Indonesiana*, 2015; 51(4): 252-256.
5. Sharma P., Sharma A., Agarwal M. and Joshi S.C.. A Review on Antifertility Efficacy of Plants in Males. *Int J Pharm Bio Sci* 2013 Oct; 4(4): (P) 413 - 428.
6. Tariq M.H., Ghaffar B., Ahmed T., Sultan A., Irfan M. and Farrukh M.J. Phytochemical and Microbiological Evaluation of Different Chemical Extracts of Papaya Seeds on Clinical Isolates of (Fgsh Hospital) Islamabad. *International Journal of Pharmacy*, 2015; 5(1):122-126.
7. Hayatie L., Biworo A. and Suhartono E. Aqueous Extracts of Seed and Peel of *Carica Papaya* Against *Aedes Aegypti*. *Journal of Medical and Bioengineering*, 2015; 4(5): 417-421. doi: 10.12720/jomb.4.5.417-421.
8. Dada F.A., Awaji F.O., Esan A.M., Oyeleye S.I. and Adegbola V.B. Phytochemical And Antioxidant Analysis Of Aqueous Extracts Of Unripe Pawpaw (*Carica Papaya* Linn.) Fruit's Peel And Seed. *IJRRAS*, 2016; 27 (3): 68-71.
9. Nora N.B., Hamid K., Snouci M., Boumediene M. and Abdellah M. Phytochemical and antibacterial screening of *Citrullus colocynthis* of South-west Algeria. *J. Chem. Pharm. Res.*, 2015, 7(5):1344-1348.
10. Sharma A., Sharma P., Chaturvedi M. and Joshi S.C. Contraceptive Efficacy of *Citrullus colocynthis* Methanolic Extract in Male Rats. *American Journal of Phytomedicine and Clinical Therapeutics*, 2014; 2(2):229-241.
11. Kumar S, Chatterjee R, Dolai S, Adak S, Kabir SN, Banerjee S and Mondal NB. Chenopodium album seed extract-induced sperm cell death: exploration of a plausible pathway. *Contraception* 2008; 77(6): 456-462.
12. Arora S. and Itankar P. Extraction, isolation and identification of flavonoid from *Chenopodium album* seed parts. *Journal of Traditional and Complementary Medicine* (2018), <https://doi.org/10.1016/j.jtcme.2017.10.002>.
13. Hasim, B.S., Lalithamma, A., Lakshman, J., and Changamma, C. Antifertility Effect Of *Carica Papaya* Linn. Seed Extract On Hormones In Male Albino Rats. *International Journal of Biological & Pharmaceutical Research*. 2013. 4(12): 859-861.
14. Nwaehujor C.O., Ode J.O., Ekwere M.R. dan Udegbunam R.I.. Anti-fertility effect of fraction from *Carica papaya* (PawPaw) Linn. methanol rootextract in male Wistar rats. *Arabian Journal of Chemistry* (2014), <http://dx.doi.org/10.1016/j.arabjc.2014.10.018>.



15. Turek P.J.(2005). Hypothalamic-Pituitary Gonadal (HPG) Axis and Control of spermatogenesis. In: Endocrine evaluation. Male Reproductive. Laboratory Department of Urology University of California at Francisco. San Francisco, California. 2005. Page 1220.
16. Heckert L.L. and Grisworld M.D.. The Expression of the Follicle stimulating Hormone Receptor in Spermatogenesis. *Reprod Prog Horm Res*. 2002; 57:129-48.
17. Hassanin A.M. and Ayad E. The impact of chronic testicular inflammatory infiltration on spermatogenesis in azoospermic men, evidence-based pilot study. *Middle East Fertility Society Journal*, 2016; 21(1): 31-35.



25%

SIMILARITY INDEX

PRIMARY SOURCES

- 1** Peter M. Eze, Dominic O. Abonyi, Chika C. Abba, Peter Proksch, Festus B. C. Okoye, Charles O. Esimone. 36 words — 2%
"Toxic, but beneficial compounds from endophytic fungi of *Carica papaya*", *The EuroBiotech Journal*, 2019
Crossref

- 2** Peter M. Eze, Joy C. Nnanna, Ugochukwu Okezie, Happiness S. Buzugbe et al. 28 words — 1%
"Screening of metabolites from endophytic fungi of some Nigerian medicinal plants for antimicrobial activities", *The EuroBiotech Journal*, 2019
Crossref

- 3** scholar.sun.ac.za 28 words — 1%
Internet

- 4** Euis Julaeha, Yunita Permatasari, Tri Mayanti, Ajeng Diantini. 27 words — 1%
"Antifertility Compound from the Seeds of *Carica papaya*", *Procedia Chemistry*, 2015
Crossref

- 5** Tuti Nuraini, Dadang Kusmana, Efy Afifah. 27 words — 1%
"Injection of *Carica papaya* L. Seed Extract of Cibinong Variety to *Macaca fascicularis* L. and its Effect to Quality of Spermatozoa and Level of Testosterone Hormone", *Makara Journal of Health Research*, 2012
Crossref

- 6** article.sciencepublishinggroup.com 27 words — 1%
Internet

- 7** Masahiro Itoh. 26 words — 1%
"Chapter 3 Human Testicular Autoimmunity as a Result of Breakdown of Testicular Immune Privilege", Springer Nature, 2017

-
- 8 etheses.uin-malang.ac.id 25 words — 1%
Internet
-
- 9 e-journal.unair.ac.id 25 words — 1%
Internet
-
- 10 www.ijcps.com 25 words — 1%
Internet
-
- 11 Meilin Xu, Lu Ran, Ning Chen, Xiaowei Fan, Dabing Ren, Lunzhao Yi. "Polarity-dependent extraction of flavonoids from citrus peel waste using a tailor-made deep eutectic solvent", Food Chemistry, 2019 21 words — 1%
Crossref
-
- 12 theses.ulaval.ca 20 words — 1%
Internet
-
- 13 Wajid Zaman, Mushtaq Ahmad, Muhammad Zafar, Hafiza Amina et al. "The quest for some novel antifertility herbals used as male contraceptives in district Shangla, Pakistan", Acta Ecologica Sinica, 2019 20 words — 1%
Crossref
-
- 14 doaj.org 19 words — 1%
Internet
-
- 15 W S Adayani, R Subahar, Fatmawaty, Slamet. " Effect of treatment using seed extract with Ag–TiO nanocomposite on the mortality of larvae ", Journal of Physics: Conference Series, 2018 17 words — 1%
Crossref
-
- 16 www.degruyter.com 16 words — 1%
Internet
-
- 17 Saran, Parmeshwar, Ishwar Solanki, and Ravish Choudhary. "Papaya as Medicaments", Papaya, 2015. 16 words — 1%
Crossref

-
- 18 Lewu, F.B., D.S. Grierson, and A.J. Afolayan. "Extracts from Pelargonium sidoides. Inhibit the Growth of Bacteria and Fungi", *Pharmaceutical Biology*, 2006.
Crossref 14 words — 1%
-
- 19 www.ncbi.nlm.nih.gov
Internet 14 words — 1%
-
- 20 Kumpawat, K.. "Genotoxic effect of raw betel-nut extract in relation to endogenous glutathione levels and its mechanism of action in mammalian cells", *Mut.Res.- Genetic Toxicology and Environmental Mutagenesis*, 20030708
Crossref 14 words — 1%
-
- 21 S. Carter Fox, Bin Li, Daiqiang Xu, Kevin J. Edgar. "Regioselective Esterification and Etherification of Cellulose: A Review", *Biomacromolecules*, 2011
Crossref 12 words — 1%
-
- 22 ecc.isc.gov.ir
Internet 12 words — 1%
-
- 23 www.ejpmr.com
Internet 9 words — < 1%
-
- 24 link.springer.com
Internet 9 words — < 1%
-
- 25 Samar Sayed Ibrahim, Nagwa Yousef Salem. "Insecticidal efficacy of nano zeolite against *Tribolium confusum* (Col., Tenebrionidae) and *Callosobruchus maculatus* (Col., Bruchidae)", *Bulletin of the National Research Centre*, 2019
Crossref 9 words — < 1%
-
- 26 chalapathipharmacy.in
Internet 8 words — < 1%
-
- 27 "Reproductive Endocrinology", *Springer Nature*, 2009
Crossref 8 words — < 1%

28 Scott A. Walper, Guillermo Lasarte Aragonés, Kim E. Sapsford, Carl W. Brown et al. "Detecting Biothreat Agents: From Current Diagnostics to Developing Sensor Technologies", ACS Sensors, 2018 8 words — < 1%
Crossref

29 www.indexintelligence.org 8 words — < 1%
Internet

30 Ganiyu Oboh, Ayodeji A. Olabiyi, Ayodele J. Akinyemi, Adedayo O. Ademiluyi. "Inhibition of key enzymes linked to type 2 diabetes and sodium nitroprusside-induced lipid peroxidation in rat pancreas by water-extractable phytochemicals from unripe pawpaw fruit (*Carica papaya*)", Journal of Basic and Clinical Physiology and Pharmacology, 2014 7 words — < 1%
Crossref

31 Olawuyi, O.J., A.C. Odebode, I.O. Oyewole, A.O. Akanmu, and O. Afolabi. "Effect of arbuscular mycorrhizal fungi on *Pythium aphanidermatum* causing foot rot disease on pawpaw (*Carica papaya* L.) seedlings", Archives of Phytopathology and Plant Protection, 2014. 6 words — < 1%
Crossref

32 Saran, Parmeshwar, Ishwar Solanki, and Ravish Choudhary. "References", Papaya, 2015. 6 words — < 1%
Crossref

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE MATCHES OFF