#### PAPER • OPEN ACCESS

# The Potency Study of Organotin(IV) 3-Nitrobenzoate Compounds as Antimalarial Agents

To cite this article: Sutopo Hadi et al 2019 J. Phys.: Conf. Ser. 1338 012012

View the article online for updates and enhancements.



## IOP ebooks<sup>™</sup>

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

### The Potency Study of Organotin(IV) 3-Nitrobenzoate **Compounds as Antimalarial Agents**

Sutopo Hadi<sup>1,\*</sup>, Noviany<sup>1</sup>, Hardoko Insan Qudus<sup>1</sup> and Pakorn Wattana-Amorn<sup>2</sup>

<sup>1</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Lampung, Bandar Lampung, Indonesia 35145 <sup>2</sup>Department of Chemistry, Faculty of Sciences, Kasetsart University, Bangkok Thailand sutopo.hadi@fmipa.unila.ac.id

Abstract. This paper reports the potency study of organotin(IV) 3-nitrobenzoate as antimalarial agent. The organotin(IV) 3-nitrobenzoate compounds were synthesized by reacting the intermediate products dibutyltin(IV) oxide, diphenyltin(IV) dihydroxide and triphenyltin(IV) hydroxide, with 3-nitrobenzoic acid. The antimalarial activity study was performed against *Plasmodium falcifarum* and the result of  $IC_{50}$  of the three compounds tested, i.e. dibutyltin(IV) di-3-nitrobenzoate, diphenyltinIV) di-3-nitrobenzoate and triphenyltin(IV) 3-nitrobenzoate were 8.1 x  $10^{-3}$ , 5.2 x  $10^{-2}$  and 7.2 x  $10^{-3} \mu g/mL$ , respectively, while the chloroquine gave IC<sub>50</sub> 2 x  $10^{-3}$  µg/mL). The IC<sub>50</sub> of the compounds tested were little bit higher than the positive control, however these organotin(IV) compounds synthesized are not resistent to Plasmodium, thus the possible development as antimalarial agent is widely opened.

Keywords: antimalaria activity, IC<sub>50</sub>, organotin(IV) 3-nitrobenzoate

#### **1. Introduction**

Malaria is a disease that has been known since a century ago. This disease is caused due to the bite of Plasmodium. Nowadays, it continues to be a major public health problem in Indonesia and other tropical countries. Malaria has large effect to human being thus WHO pays attention to this disease by a program called Roll Back Malaria (RBM). Few points of this program were immediate diagnoses and exact treatment to eradicate malaria [1,2]. In Indonesia, the malaria cases between the periods of 1997 -2010 increased sharply. In the provinces of Java and Bali the malaria cases were increased by ten times, while outside these provinces were increased 4-5 times. These cases were also followed by resistance cases toward standard drugs used in the malaria treatments, the chloroquine and the sulfadoxine-pirimetamine. In some provinces, there were more than 25% resistance cases which cause the use of these standard drugs to be much more limited, therefore efforts to find new potent antimalarial drugs are urgently needed [3].

The organotin(IV) compounds are part of organometallic compounds which recently attract high attention from many chemists. This is because of the strong effect of organotin(IV) in many biological tests [4, 5]. The biological activities of organotin(IV) are mainly determined by the number and the type of organic groups bound to Sn center [6]. The present and the nature of the anionic groups bound to organotin(IV) compound is only as a secondary factor [6]. The various researches on organotin compounds have produced some new organotin(IV) carboxylates compound which have shown some interesting biological activities such as antimicrobial [6-8], antitumor and anticancer [7-10], antifungal activity [6,11,12], anticorrosion inhibitor [13-15], and the latest development of these compounds has led the new finding of organotin(IV) as antimalarial agent, therefore the research on organotin(IV) as possible antimalarial agent is still very challenging, thus more attentions for future application as antimalarial drug are still needed [16, 17].

Because of promising result for organotin(IV) compounds as antimalarial agent, in this paper, we reported the potency antimalarial activity study of some organotin(IV) 3-nitrobenzoates against P. falcifarum.

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

**IOP** Publishing

#### 2. Experimental

#### 2.1. Materials

All reagents used were AR grade. Dibutyltin(IV) dichlorides ([ $(n-C_4H_9)_2Cl_2$ ]), diphenyltin(IV) dichloride ([ $(C_6H_5)_2Cl_2$ ]), triphenyltin(IV) chloride ([ $(C_6H_5)_3Cl$ ]), 3-nitrobenzoic acid, RPMI were obtained from Sigma, water HPLC grade, sodium hydroxide (NaOH) and methanol (CH<sub>3</sub>OH) were JT Baker products, and were used without further purification.

#### 2.2. Characterization and instrumentations

<sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker AV 600 MHz NMR (600 MHz for <sup>1</sup>H and 150 MHz for <sup>13</sup>C). All experiments were run in DMSO-D<sub>6</sub> at 298K. The number of runs used for <sup>1</sup>H experiments were 32 with reference at DMSO signal at 2.5 ppm, while the <sup>13</sup>C were 1000-4000 scans with the reference DMSO signal at 39.5 ppm. Microelemental analyses (CHNS) were performed on Fision EA 1108 series elemental analyser. IR spectra in the range of 4000-400 cm<sup>-1</sup> were recorded on a Bruker VERTEX 70 FT-IR spectrophotometer with KBr discs. The UV spectra were recorded in the UV region and were measured using a UV- Shimadzu UV-245 Spectrophotometer. Measurements were performed in 1 mL quartz-cells. Solutions were prepared using methanol as the solvent with concentration of  $1.0 \times 10^4$  M.

#### 2.3. Preparation of organotin(IV) nitrobenzoates

The organotin(IV) 3-nitrobenzoates used in this work were prepared based on the procedure previously reported [9-12,14-16] and was adapted from the work by Szorcsik *et al.* [18]. An example procedure in the preparation of dibutyltin(IV) di-2-nitrobenzoate was as follows:

To 3.44 g (0.01 mol)  $[(C_6H_5)_2SnCl_2]$  (4) in 50 mL methanol was added 0.8 g (0.02 mol) NaOH. The reaction mixtures were stirred for about 60 minutes. Compound  $[(C_6H_5)_2Sn(OH)_2]$  (5) was precipitated out as white solid, filtered off and dried in vacuo till they are ready for analysis and further reaction. The average yield was 2.92 g (95 %).

To 0.4605 g (1.5 mmol) [  $(n-C_4H_9)_2$ SnO] (2) in 50 mL of methanol was added with 2 mole equivalents of 3-nitrobenzoic acid (0.501 g) and was refluxed for 4 hours at 60 – 62°C. After removal of the solvent by rotary evaporator, the produced compounds [( $C_6H_5$ )\_2Sn(3-OOCC\_6H\_4(NO\_2)\_2] were dried in vacuo until they are ready for analysis and further use for in vitro antimalarial activity. The yield was 1.67 g (92%). A similar procedure was also adapted in the preparation of dibutyltin(IV) and triphenyltin(IV) derivatives, [( $n-C_4H_9$ )\_2Sn(3-OOCC\_6H\_4NO\_2)] (3) and [( $C_6H_5$ )\_3Sn(3-OOCC\_6H\_4NO\_2)] (9), respectively. For triphenyltin(IV) only one mole equivalent of the 3-nitrobenzoic acid was added.

#### 2.4. In vitro antimalarial bioactivity assays

The in vitro antimalarial assays were performed in the Institute of Tropical Disease, Universitas Airlangga, Surabaya Indonesia. The malaria parasite *P. falciparum* 3D7 clone was essentially propagated according to the previously published procedure [3]. Briefly, parasite cultures were propagated in tissue culture flasks containing RPMI-1640 medium supplemented with 25  $\mu$ g/mL gentamycin, 50  $\mu$ g/mL hypoxanthine, 25 mM Hepes buffer, 25 mM sodium bicarbonate, 10% AB+ human serum, 5% haematocrit and human erythrocytes with the pH maintained at 7.4. Each compound tested was first dissolved in DMSO and diluted to different concentration by adding complete malaria medium. Chloroquine was used as a positive control. To determine the antiplasmodial activity of each isolated compound, parasites were placed in a 24-well culture plate in the presence of a wide concentration range of each compound. The parasite growth was monitored by making a blood smear that was fixed with methanol and stained with Giemsa. Total parasitaemia was calculated as the number of parasites-observed, divided by the total erythrocyte multiplied by 100%. The concentration response parasite growth data were calculated by a linear regression provided by SYSTAT Sigma Plot, using the 50% inhibitory concentration relative to untreated control.

#### **3. Results and Discussion**

The synthesis of the organotin(IV) 3-nitrobenzoates i.e. dibutyltin(IV) di-3-nitrobenzoate,  $[(n-C_4H_9)_2Sn(3-OOCC_6H_4NO_2)_2]$  (3), diphenyltin(IV) di-3-nitrobenzoate  $[(C_6H_5)_2Sn(3-OOCC_6H_4NO_2)_2]$  (6) and triphenyltin(IV) 3-nitrobenzoate,  $[(C_6H_5)_3Sn(3-OOCC_6H_4NO_2)]$  (9), were successfully prepared from their chlorides  $[(n-C_4H_9)_2SnCl_2]$  (1),  $[(C_6H_5)_2SnCl_2]$  (4) and  $[(C_6H_5)_3SnCl]$  (7), respectively. All of these reactions were conducted via  $[(n-C_4H_9)_2SnO]$  (2),  $[(C_6H_5)_2Sn(OH)_2]$  (5) and  $[(C_6H_5)_3SnOH]$  (8) respectively similar to the procedure previously reported [9-12,14-16]. Figure 1 shows an example of the reaction occurred in the synthesis of dibutyltin(IV) di-3-nitrobenzoate.

$$[(C_4H_9)_2SnCl_2] \xrightarrow[Stirred for 60 mins]{(C_4H_9)_2Sn(OH)_2]} \xrightarrow[Stirred in MeOH]{+ HOOCC_6H_4NO_2} [(C_4H_9)_2Sn(OOCC_6H_4(NO_2)_2)]$$

$$1 \qquad 2 \qquad 3$$

Fig. 1. The scheme of preparative route of the organotin(IV) di-3-nitrobenzoate

The data of microanalysis for all compounds synthesized are tabulated in Table 1, in general, all values obtained are very good and are close to the calculated values.

Table 1. The microanalytical data of the organotin(IV) compounds synthesized

| Compound                                    | Elemental analysis found (calculated) |            |            |  |
|---|---------------------------------------|------------|------------|--|
| Compound                                    | С                                     | Н          | Ν          |  |
| $[(n-C_4H_9)_2SnCl_2](1)$                   | 31.4 (31.6)                           | 6.0 (5.9)  |            |  |
| $[(n-C_4H_9)_2SnO](2)$                      | 38.6 (38.6)                           | 7.1 (7.3)  |            |  |
| $[(n-C_4H_9)_2Sn(3-OOCC_6H_4(NO_2))_2]$ (3) | 46.8 (46.7)                           | 4.7 (4.6)  | 4.8(4.96)  |  |
| $[(C_6H_5)_2SnCl_2](4)$                     | 41.7 (41.9)                           | 2.8 (2.9)  |            |  |
| $[(C_6H_5)_2Sn(OH)_2](5)$                   | 46.5 (46.9)                           | 3.8 (3.9)  |            |  |
| $[(C_6H_5)_2Sn(3-OOCC_6H_4(NO_2)_2](6)$     | 51.3 (51.6)                           | 3.1 (2.98) | 4.5 (4.63) |  |
| $[(C_6H_5)_3SnCl]$ (7)                      | 55.8 (56.1)                           | 4.0 (3.9)  |            |  |
| $[(C_6H_5)_3Sn(OH)](8)$                     | 58.4 (58.9)                           | 4.3 (4.4)  |            |  |
| $[(C_6H_5)_3Sn(3-OOCC_6H_4(NO_2))]$ (9)     | 57.2 (58.1)                           | 3.8 (3.68) | 2.6 (2.71) |  |

Table 2. The characteristic and important IR bands of the organotin(IV) compounds (cm<sup>-1</sup>) synthesized

| Compound             | 3      | 6                | 9                | References |
|----------------------|--------|------------------|------------------|------------|
| Sn-O                 | 434.4  | 594.0            | 735.42           | 800-400    |
| Sn-O-C               | 1029.1 | 1238.2           | 1243.4           | 1050-900   |
| Sn-Bu                | 674.4  | -                | -                | 740-660    |
| CO <sub>2</sub> asym | 1419.1 | 1531.7           | 1557.5           | 1600-1400  |
| CO <sub>2</sub> sym  | 1558.1 | 1659.3           | 1630.4           | 1700-1550  |
| C-H                  | 2954 – |                  |                  | 2960 -     |
| aliphatic            | 2860   | -                | -                | 2850       |
| Phenyl               | -      | 1467.0;<br>750.7 | 1428.4;<br>729.1 | 1450, 730  |

Table 2 lists some important FT-IR data of the compound synthesized. The starting materials (1, 4, 7) gave characteristic strong stretching band of Sn-Cl bond at  $390 - 310 \text{ cm}^{-1}$ . For example, in 1 the Sn-Cl bond appeared at  $334.2 \text{ cm}^{-1}$ . The other characteristic bands of 1 is the present of stretching band from butyl ligands at 1069 cm<sup>-1</sup>. The bending vibration of C-H aliphatic stretch of the butyl appeared at 2956 – 2865 cm<sup>-1</sup>.Once compound 1 is reacted with NaOH to produce compound 2, the

band of Sn–Cl disappeared and a new strong band of Sn–O bond at of 417.4 cm in compound **2** appeared in the spectrum. The other stretching band due to the presence of butyls and their bending vibrations as expected although the frequencies have little bit shifted are still present. The formation of dibutyltin (IV) di-3-nitrobenzoate compounds,  $[(n-C_4H_9)_2Sn(3-OOCC_6H_4(NO_2))_2]$ , (**3**) is confirmed by the strong asymmetric stretching bands of the carboxylate groups which occurred at ca. 1400 cm<sup>-1</sup> and the symmetric stretch at ca. 1600 cm<sup>-1</sup> as well as the present of Sn-O stretching of the acid at 435 cm<sup>-1</sup>, and the appearance of these bands is confirming the success of the substitution reaction [9-12,14-16].

The UV-Vis spectroscopy analyses of all the compounds have also been utilized to obtain  $\lambda_{max}$  andtThe data obtain presented in Table 3. The  $\lambda_{max}$  for each compound in any steps of the reaction have been observed to have shifted. The compound **1** has  $\lambda_{max}$  of 210.7 nm, while compound **2** has  $\lambda_{max}$  of 202.9 nm. This information gave an indication that there was a shift to a shorter  $\lambda_{max}$  value when the conversion of compound **1** to **2** took place. The wave-length shift to a shorter  $\lambda_{max}$  could occur due to either the solvent used in the measurement or the effect of an auxochrome of the ligand. However in this study, as the solvent used for all measurements was the same which is methanol, the shift in the  $\lambda_{max}$  that occurred must be due to the auxochrome effect. In case of compound **1** and **2**, there is an oxide group which has electron drawing effect bigger in compound **2** than that of chloride group in **1**. As a result, the electron transition in **2** is hard to occur, thus,  $\lambda_{max}$  measured was getting shorter in compound **2** than in compound **1** [19-22]. Similar results are also observed for other changes as shown in Table 3. For instance, in compound **3**, the electron drawing effect of  $3-C_6H_4(NO_2)COOH$  is less than chloride in **1**, so the electron transition in this molecule will be easier (the energy required is less), thus producing longer  $\lambda_{max}$ , 291.3 nm.

|   |       | $\lambda_{max}$ (nm) | )                           |
|---|-------|----------------------|-----------------------------|
| Compound                                    | π- π* | n-π                  | Benzene ring secondary band |
| $[(n-C_4H_9)_2SnCl_2](1)$                   | 210.7 | -                    | -                           |
| $[(n-C_4H_9)_2SnO](2)$                      | 202.9 | -                    | -                           |
| $[(n-C_4H_9)_2Sn(3-OOCC_6H_4(NO_2))_2]$ (3) | 203.5 | 291.8                | -                           |
| $[(C_6H_5)_2Sn(3-OOCC_6H_4(NO_2)_2](6)$     | 201.3 | 299.9                | 407.6                       |
| $[(C_6H_5)_3Sn(3-OOCC_6H_4(NO_2)]$ (9))     | 204.5 | 304.0                | 410.2                       |

Table 3. The  $\lambda_{max}$  of the UV-Vis spectra of the organotin(IV) 3-nitrobenzoate compounds

Table 4. <sup>1</sup>H and <sup>13</sup>C spectra of the compounds synthesized

| Compounds   | H in butyl or<br>phenyl (ppm) | H in<br>benzoate<br>(ppm) | C in butyl, phenyl and<br>benzoate<br>(ppm) |
|---|-------------------------------|---------------------------|---|
|   | Ηα & Ηβ:1.37-                 |                           | Cα: 21.2; Cβ: 26.6; Cγ: 25.9;               |
| $[(n-C_4H_9)_2Sn(3-$  | 1.62 (m); Hy:                 | 7.34-7.86                 | Сб: 14.2; С1: 174.9; С2:                    |
| $OOCC_6H_4(NO_2))_2](3)$  | 1.30 (m); Hδ:                 | (m)                       | 139.3; C3 & C7: 129.7; C4                   |
|   | 0.93 (t)                      |                           | & C6: 128.5; C5: 125.2                      |
|   | H2 & H6 7.59                  |                           | C1-6 (phen): 131.7-126.9;                   |
| $I(C, \mathbf{H}) = S_{\mathbf{T}}(2, \mathbf{OOCC}, \mathbf{H}, (\mathbf{NO})) I(C)$ | (d, 4H); H3 &                 | 7.82 - 7.93               | C7: 175.7; C8: 139.5; C9 &                  |
| $[(C_6H_5)_2SI(5-OOCC_6H_4(NO_2)_2](0)$   | H5 7.48 (t, 4H);              | (m)                       | C13: 130.2; C10 & C12:                      |
|   | H4: 7.36 (t, 2H)              |                           | 129.3; C11: 128.5                           |
|   |                               |                           |   |

| IOP Conf. Series: Journal of Physics: Conf. Series <b>1338</b> (2019) 012012 | doi:10.1088/1742-6596/1338/1/012012 |
|--|-------------------------------------|
|--|-------------------------------------|

|  | H2 & H6 7.58     |             | C1-6 (phen): 131.1-126.2;  |
|--|------------------|-------------|----------------------------|
| $[(C, \mathbf{H}) \ S_{\mathbf{p}}(2, \mathbf{OOCC}, \mathbf{H}, (\mathbf{NO}))] \ (0))$ | (d, 6H); H3 &    | 7.81 - 7.92 | C7: 175.3; C8: 139.2; C9 & |
| $[(C_6H_5)_3SII(3-OOCC_6H_4(INO_2)] (9)]$  | H5 7.46 (t, 6H); | (d)         | C13: 130.3; C10 & C12:     |
|  | H4: 7.03 (t, 3H) |             | 128.6; C11: 128.2          |

The results of <sup>1</sup>H and <sup>13</sup>C NMR for all compounds synthesized are tabulated in Table 4. A number of signals in the spectra recorded have been characterized carefully. The chemical shift ( $\delta$ ) of butyl protons attached to the tin metal appeared in the range of 0.93 ppm for H $\delta$  up to 1.37-1.62 ppm for H $\alpha$  and H $\beta$ , and the carbons of butyl ligands are observed at position comparable and in agreement with other similar compounds reported previously [9-12,14-16, 19, 21-23]. The chemical shift of phenyl protons attached to tin metal appeared in the range of 7.36 – 7.58 ppm, while the carbon of carboxyl group of all compounds as expected appeared in the region of 176 ppm [13,19,21-23]. The carbon atoms of the phenyl ligand as also expected appeared in  $\delta$  of 131 – 126 ppm, while the carbons in the nitrobenzoate derivatives appeared in  $\delta$  range of 140 – 130 ppm close to the reported values of similar compounds [9-12,14-16, 19, 21-23].

In our previous study on the antifungal and anticancer activity of the compounds reported here [9-12,14-16], it has been shown that optimal activity of the antifungal and anticancer has been associated with the number of carbon atoms of the ligand present in the organotin(IV) used [9-12,14-16], where in general, the derivative of triphenyltin(IV) carboxylate which contain 18 carbon atoms has the highest activity [6-9,12,13,24]. The same phenomena interestingly was also observed in this study.

As shown in Table 5, the derivatives of triphenyltin(IV) compounds showed the highest antimalarial activity in the series, and the diphenyltin(IV) compounds are stronger in inhibiting than those of dibutyltin(IV) compounds, similar to those previously reported [9-12,14-16]. Thus the number of carbon atoms present as well as the type of the ligands has significant effect on the antimalarial activity of the organotin(IV) compounds tested [24].

| Compounds                                   | IC <sub>50</sub> (µg/mL) |
|---|--------------------------|
| Chloroquine                                 | 2.0 x 10 <sup>-3</sup>   |
| $[(n-C_4H_9)_2Sn(3-OOCC_6H_4(NO_2))_2]$ (3) | 8.1 x 10 <sup>-2</sup>   |
| $[(C_6H_5)_2Sn(3-OOCC_6H_4(NO_2)_2]$ (6)    | 5.2 x 10 <sup>-2</sup>   |
| $[(C_6H_5)_3Sn(3-OOCC_6H_4(NO_2)] (9))$     | 7.2 x 10 <sup>-3</sup>   |

| Table | 5. | The | $IC_{50}$ | of | the | com | pound | l tested |
|-------|----|-----|-----------|----|-----|-----|-------|----------|
|-------|----|-----|-----------|----|-----|-----|-------|----------|

The organotin(IV) 3-nitrobenzoate compounds synthesized have been found exhibited much higher antimalarial activity compared to those of the ligands, starting materials and intermediate products. In this respect, our results are consistent with a well-known fact that many biologically active compounds become more active upon complexation than in their uncomplexed forms [25]. Crowe stated the actual biological activity of diorganotin compounds of the type RR'SnXY (R and R' = alkyl or aryl; X and Y= anions) was determined solely by the RR'Sn<sup>2+</sup> moiety [26].

#### 4. Conclusion

The organotin(IV) 3-nitrobenzoate compounds were successfully prepared and it is quite clear from the discussion above that they have shown some promising result to be used as antimalarial drug. The fact that triphenyltin(IV) 3-nitrobenzoate derivatives have shown the highest antimalarial activity

was in line with other data relating to the number of carbon atom present in the compound. To find the best explanation for this phenomenom, more studies will be carried out to .

#### Acknowledgments

The authors are grateful to Institute of Research and Community Services, Universitas Lampung that provided fund for this project to be undertaken through International Research Collaboration 2018 and Penelitian Berbasis Kompetensi (Competency Research Grant Scheme) 2018 from Directorate of Research and Community Services, The Ministry of Research, Technology and Higher Education, Indonesia with contract number 384/UN26.21/PN/2018, February 12, 2018 and Basic Research (Penelitian Dasar) 2019 with contract number 857/UN26.21/PN/2019, April 8, 2019.

#### REFERENCES

- [1] Tiekink ERT 1991 Structural Chemistry of Organotin Carboxylates: a Review of the Crystallographic Literature. *App. Organomet. Chem.*, **5**, 1-30.
- [2] Shahid K, Ali S, Shahzadi S and Akhtar Z 2003 Organotin(IV) Complexes on Aniline Derivaties Part-II-Synthesis and Spectroscopic Characterization of Organotin(IV) Derivatives of 2-[4-Bromoanailine)carboxyl]benzoic Acid. *Turk. J. Chem.*, 27, 209-215.
- [3] Pellerito L and Nagy L 2002 Organotin (IV)<sup>n+</sup> Complexes Formed with Biologically Active Ligands: Equilibrium and Structural Studies and Some Biological Aspect. *Coor. Chem. Rev.* 224 pp 111–150
- [4] Gleeson B, Claffey J, Ertler D, Hogan M, Müller-Bunz H, Paradisi F, Wallis D and Tacke M 2008 Novel organotin antibacterial and anticancer drugs. *Polyhedron*, **27**, 3619
- [5] Rehman W, Badshah A, Khan S and Tuyet LTA 2009 Synthesis, characterization, antimicrobial and antitumor screening of some diorganotin(IV) complexes of 2-[(9H-Purin-6-ylimino)]phenol. *Eur. J. Med. Chem.*, 44, 3981
- [6] Hadi S, Rilyanti M 2010 Synthesis and *in vitro* anticancer activity of some organotin(IV) benzoate compounds. *Orient. J. Chem.*, **26**, 775
- [7] Hadi S, Rilyanti M and Suharso 2012 In Vitro Activity and Comparative Studies Of Some Organotin(IV) Benzoate Derivatives Against Leukemia Cancer Cell: L-1210. Indo. J. Chem., 12, 172
- [8] Hadi S, Irawan B and Efri 2008 The Antifungal Activity Test of Some Organotin(IV) Carboxylates. J. Appl. Sci. Res., 4, 1521
- [9] Hadi S, Rilyanti M and Nurhasanah 2009 Comparative Study on the Antifungal Activity of Some Di- and Tributyltin(IV) Carboxylate Compounds. *Modern Appl. Sci.*, **3** (2), 12-17.
- [10] Singh R, Chaudary P and Khausik NK 2010 A Review: Organotin Compounds in Corrosion Inhibition. *Rev. Inorg. Chem.*, **30**, 275 294.
- [11] Kurniasiah H, Nurissalam M, Iswantoro B, Afriyani H, Qudus HI and Hadi S 2015, The Synthesis, Characterization and Comparative Anticorrosion Study of Some Organotin(IV) 4-Chlorobenzoates. *Orient. J. Chem.*, **31**, 2377-2383.
- [12] Hadi S, Afriyani H, Anggraini WD, Qudus HI and Suhartati T 2015 The Synthesis and Potency Study of Some Dibutyltin(IV) Dinitrobenzoate Compounds as Corrosion Inhibitor for Mild Steel HRP in DMSO-HCl Solution. *Asian J. Chem.*, 27, 1509-1512.
- [13] Hadi S, Noviany and Rilyanti M 2018 In vitro antimalarial activity of some organotin(IV)2nitrobenzoate compounds against *Plasmodium falciparum*. *Maced. J. Chem. Chem Eng.* 37, 191
- [14] Hansch C and Verma RP 2009 Larvicidal activities of some organotin compounds on mosquito larvae: A QSAR study. *Eur. J. Med. Chem.*, **44**, 260.

- [15] WHO 2011 Roll Back Malaria program. Available at. http://www.who.int/malaria/en/. 2011. [Verified 20 April 2011]
- [16] James AA 1992 Mosquito molecular genetics: the hands that feed bite back. *Science*, 257, 37-38.
- [17] Harijanto PN 2006 Perubahan Radikal dalam pengobatan Malaria di Indonesia (Radical change on the malaria treatment in Indonesia), *Cermin Dunia Kedokteran*,**152**, 30-36 (in Indonesian)
- [18] Szorcsik A, Nagy L, Gadja-Schrantz K, Pellerito L, Nagy E and Edelmann ET 2002 Structural studies on organotin(IV) complexes formed with ligands containing {S, N, O} donor atoms. *J.Radioanal.Nucl.Chem.* 252, 523-530.
- [19] Nath M, Yadav R, Gielen M, Dalil H, de Vos D and Eng G 1997 Synthesis, characteristic spectral studies and *in vitro* antimicrobial and antitumour activities of organotin(IV) complexes of Schiff bases derived from amino-acids. *Appl. Organometal. Chem.*, **11**, 727-736.
- [20] Sudjadi, The Structure Determination of Organic Compounds. Ghalia Publishers, 1985. (In Indonesian).
- [21] Rilyanti M and Hadi S 2011 Synthesis, characterization and thermal stability of complex *cis*-[Co(bipy)<sub>2</sub>(CN)<sub>2</sub>] and its interaction with NO<sub>2</sub> gas. *Russ. J. Inorg. Chem.*, **56**, 418-421.
- [22] Hadi S, Appleton TG and Ayoko GA 2003 Reactions of *fac*-[PtMe<sub>2</sub>(OMe)(H<sub>2</sub>O)<sub>3</sub>]<sup>+</sup> with halide ions: effect of halide trans effect on methoxide hydrolysis. *Inorg. Chim. Acta.*, **352**, 201-207.
- [23] Hadi S and Appleton TG 2010 Reactions of cisplatin hydrolytes, *cis*-[Pt(<sup>15</sup>NH<sub>3</sub>)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>]<sup>2+</sup>, with N-acetyl-L-cysteine. *Russ. J. Inorg. Chem.*, **55**, 223-228.
- [24] Chohan ZH and Rauf A 1996 Some Biologically Active Mixed Ligand Complexes of Co(II), Cu(II) and Ni(II) with ONO, NNO and SNO Donor Nicotinoylhydrazine-Derived Ligands. *Synth. React.Inorg.Met.-Org. Chem.*, **26**, 591-604.
- [25] Gershon H 1974 Antifungal Activity of Bischelates of 5-, 7-, and 5,7-halogenated 8-quinols with copper(II). Determination of the long and short access of the pores in the fungal spore wall. J. Med. Chem., 17, 824-827.
- [26] Crowe AJ 1989 The antitumour activity of tin compounds; in Gielen M., Ed. *Metal-Based Drugs*. **1**, 103-149.