

Antidiabetic and Antioxidant Activities of Bay, Pandan, Citrus Leaves and their Combination in Vitro

By Samsu Udayana Nurdin; Devi Sabarina; Subeki; Sussi Astuti

Antidiabetic and Antioxidant Activities of Bay, Pandan, Citrus Leaves and their Combination in Vitro

Samsu U. Nurdin*, Devi Sabarina, Subeki and Sussi Astuti

Department of Agricultural Product Technology, Agriculture Faculty,
Lampung University, Jl. Sumantri Brojonegoro, Bandar Lampung, Indonesia.

<http://dx.doi.org/10.13005/bpj/1707>

(Received: 08 March 2019; accepted: 03 May 2019)

The study aimed to evaluate the effects of bay (B), pandan (P), citrus leaves (C) and their combinations against starch hydrolysis enzymes (α -glucosidase and α -amylase enzymes) and antioxidant activity and to examine the role of polyphenol compounds in enzyme inhibition and antioxidant activity. Three single leaves extracts and five of their combinations were applied to inhibit α -glucosidase hydrolyzing *p*-nitrophenyl- α -D-glucopyranoside or α -amylase hydrolyzing starch solution as well as to scavenge free radicals. The leaf extracts and their combination showed inhibition activities against α -glucosidase and α -amylase enzymes with range of inhibition activities were between 17.63% to 26.04% and 20.14% to 35.30% respectively. There is no significant difference among the extracts in modulation of α -glucosidase activity, but each extract exhibited different effect on α -amylase or antioxidant activities. Mixing P with B and C increases the inhibitory activity of the extract against α -amylase as seen that percent of inhibition of BPC is significantly higher than P, even though their total phenolic content was not different. The synergism or antagonism effect was not observed when the extracts were combined as the enzyme inhibition or antioxidant activities are not dependent on the proportion of the extract in the mixtures. The role of polyphenol compounds on inhibition of the starch digestion enzymes and on antioxidant activity was not observed. Further study is required to fully elucidate the effect of the leaf or their combinations on diabetic animal models or diabetic patients.

Keywords: Bay leaf, pandan leaf, citrus leaf, antidiabetic, antioxidant, phenolic compound.

Rice is staple food of most Asia countries including Indonesia. As rice is good source of starch, therefore, consumption of rice is suggested as risk factor of diabetes mellitus^{1 and 2}. Reducing of starch digestibility of the rice is one of promising strategies to reduce hyperglycemic effect of the rice³.

Starch, a polysaccharide composed of alpha 1,4-linked glucose units (amylose) and alpha 1,4-1,6-linked branched structure (amylopectin), is cleaved in the duodenal cavity involved several hydrolytic enzymes such as pancreatic alpha-

amylase and brush border glycosidase⁴. Inhibition of these enzymes is not only considered as a strategy to reduce the digestibility of the starch but also a treatment of carbohydrate intake disorder, such as diabetes and obesity^{5 and 6}. Plants are an important source of phytochemical compounds those have inhibition activities against the enzymes, therefore, they have potentiality for therapeutic drug or functional food for the diseases⁷.

Rice is prepared by cooking (steaming or boiling) rice soaked in water. Addition of aromatic or flavouring ingredients such as Indonesian bay

(*Eugenia polyantha* Wight), pandan (*Pandanus amaryllifolius* Roxb.) and citrus (*Citrus hystrix*) leaves is a common practice in Indonesian rice cooking. Whether the leaves have beneficial effects on rice starch digestibility when they are mixed with rice remain unexplored. However, the therapeutic benefits of these ingredients for diabetes have been reported.

Aqueous extracts of the bay leaves improve glucose and insulin metabolism in in vitro model⁸. Consumption 1 to 3 g a day¹⁷ ground bay leaves by type 2 diabetes patients reduced serum glucose with significant decreases ranging from 21 to 26% after 30 d and improved lipid profile¹⁶ of the subjects⁹. Moreover, methanolic extract of bay leaf displayed scavenging activity against superoxide and hydroxyl radicals in a concentration-dependent manner¹⁰.

Pandan leaf is a tropical plant which is used mostly as a flavoring agent for certain rice and bread²⁸. Water extract of pandan leaves reduced blood glucose level as well as improvement the insulin resistance of obese mice¹². In healthy subjects²⁷ drinking pandan leaf tea effectively decreased postprandial blood sugar through inhibition of α -glucosidase enzyme and induction of insulin production in pancreatic cell¹³.

Citrus leaf is an aromatic Asian leaf most often used in Indonesian recipes including cooked rice recipes¹⁴. The leaf extracts exhibited anticancer activity through reduction of cancer cell line viability¹⁵. Fresh juice from Citrus fruits contains phenolics, tannins and flavonoids and exhibited anti-alpha amylase and alpha-glucosidase in vitro¹⁶.

Mechanism of the protective effect of dietary antioxidants has been hypothesized through inhibition of oxidation chain reactions¹⁷. Thus, consumption plant foods rich in antioxidant compounds could reduce incidence of chronic diseases, such as diabetes³³ through down regulation of oxidative stress¹⁷⁻²⁰. In this study, we aimed to evaluate the effects of bay, pandan, citrus leaves¹⁴ their combinations against starch hydrolysing enzymes (α -glucosidase and α -amylase enzymes) and antioxidant activity. Additionally, due to the leaves are rich in polyphenol^{16, 21, 22}, the role of polyphenol compounds in enzyme inhibition and antioxidative effect was also elucidated.

26

MATERIALS AND METHODS

Plant material

Bay, pandan and citrus leaves were collected from local market in Bandar Lampung, Indonesia. The leaves⁴⁶, immediately, after collection were thoroughly washed with water and dried in oven at 60°C. Dried leaves were powdered using grinder to produce coarsely powder.

Preparation of extracts

The dried leaf powder⁴⁵ of bay, pandan, citrus or their combination (10 g) were boiled in 100 mL water for 20 minutes. The extract was filtered (extract 1), and the residue was reboiled in 100 mL for 20 minutes, then filtered to get extract 2. Extract 1 and extract 2 were then mixed and considered as 100% extract. Proportion of each type of leaves in the combination was 50.0 % when the combination contained two types of plants, and 33.3 % when the combination contained three types of plants (Table 1).

Assay of α -glucosidase inhibitory activity

The slightly modified method⁴⁴ described by Rao *et al.*²³ was applied to measure the effect of the leaf extracts on α -glucosidase activity using α -glucosidase crude enzyme (Shandong Longda Bio-Products Co., Ltd.). The substrate solution p-nitrophenyl glucopyranoside (pNPG) (Sigma Aldrich, Switzerland) was prepared in aquades (0.03012 g/100mL). Briefly, sample of 200 μ L leaf extracts were⁵¹ incubated with 2 mL of α -glucosidase crude enzyme for 10 min at 37°C. The reaction was initiated by adding⁵² 1 mL of pNPG substrate and⁵ incubated at 37°C for 30min. The reaction was stopped by adding 2 mL of 2 % Na_2CO_3 (Merck, Germany). The α -glucosidase activity was determined by measuring the yellow-colored paranitrophenol released from pNPG at⁶⁵ 5 nm (Thermo Scientific Genesys 20, USA). Percentage inhibition is calculated as %Inhibition = $[(\text{Abs}_{\text{control}} - \text{Abs}_{\text{extract}}) / \text{Abs}_{\text{control}}] \times 100$.

Assay of α -amylase inhibitory activity

α -amylase inhibitory activity assay was performed at 37°C using α -amylase crude enzyme (Shandong Longda Bio-Products Co., Ltd.). A⁴³ mixture containing 1 ml α -amylase crude enzyme, 0.1 ml phosphate buffer 0.1 M and 0.2 extract (A) or water (B) was incubated for 10 min at 37°C.

Then, 3 ml of 4% starch solution (wheat starch) was added to the mixture and incubated for 60 min at 37°C²⁴. Reducing sugar released from the starch hydrolysis was measured using DNS method. α -amylase inhibitory activity was calculated by the following formula:

$$\alpha\text{-amylase inhibitory activity} = \frac{(B-A)}{B} \times 100\%$$

Where A and B represent sugar concentrations in the reaction mixture with and without an addition of leaf extract, respectively.

Antioxidant activity measurement

The antioxidant activity assay of the extracts was performed according to protocol describe by Xu *et al.*²⁵ using DPPH (2,2-Diphenyl-1-Picrylhydrazyl) (Merck, Germany) radical scavenging activity method¹⁸. Leaf extract (0.25 mL) was mixed with 2 mL DPPH solution (3.3 mg of DPPH in 100 mL methanol) and 8 mL 11 mol(JT Baker), vortexes gently, then incubated at room temperature for 30 min in the dark, and the absorbance (A1) was measured at 517 nm (Thermo Scientific Genesys 20, USA). The absorbance (A0) control sample (distilled water instead of leaf extract) is also recorded at the same wavelength. Radical scavenging activity (%) was calculated by using the formula = [(A0 - A1)/A0] × 100, where A0 was the absorbance of control sample and A1 was the absorbance of extracts.

Total phenolic analysis

Total phenolic content of the extract was measured using the Folin-Ciocalteu (Merck, Germany) reagent²⁶ with slight modification. 0.2 mL of leaf extract was mixed with 0.2 mL of aquades and 0.2 mL of Folin-Ciocalteu reagent (1 N). Then

4 mL of sodium carbonate (Merck, Germany) solution (2%) was added and then allowed to stand for 30 min in the dark for incubation. The absorbance was measured at 760 nm in a spectrophotometer (Thermo Scientific Genesys 20, USA). A standar curve was prepared using Gallic acid (Tokyo Chemical Industry Co., Ltd) (0.00-0.01 mg/mL). The total phenolic contents were expressed in terms of gallic acid equivalents (mg of per mL extract).

Statistical analysis

Results are expressed as the mean of 3 replicates. Statistical analysis was carried out with a statistical program (SPSS version 18. One way-Anova with Fisher test was used. Results were considered significant if p<0,05.

RESULTS AND DISCUSSION

Alpha glucosidase enzyme is located in the brush border of the small intestine and is involved in starch digestion to release monosaccharide. Inhibition of the enzyme leads to retardation of starch digestion in small intestine. Study of α -glucosidase inhibitor activity of extracts of bay, pandan, citrus leaves or their combination might contribute to the understanding of their potentiality for diabetic management. Commercially, inhibitor of this enzyme is available for glucose-lowering medication for diabetic patients²⁷.

The α -glucosidase inhibitory activity of the extracts of single or mixture of leaves is not depend on the type of extracts (Fig 1). Percent of inhibition of B, P and C or their combinations against α -glucosidase was around 32%. Synergism or antagonism effect was not observed when the

Table 1. Proportion of leaves in dried leaf combination

Treatments	Proportion of each leaf in combinations (%)		
	Bay	Pandan	Citrus
BPC	33.3	33.3	33.3
BC	50.0	0.0	50.0
PC	0.0	50.0	50.0
BP	50.0	50.0	0.0
B	100.0	0.0	0.0
P	0.0	100.0	0.0
C	0.0	0.0	100

Table 2. Correlation coefficients between total phenolic with antiglucosidase, antiamylase and antioxidant activities of the extracts

Activities	Cell Contents	Total Phenol
Antiglucosidase	Pearson correlation	0,104
	P-Value	0,654
Antiamylase	Pearson correlation	-0,112
	P-Value	0,630
Antioxidant	Pearson correlation	-0,527
	P-Value	0,014

*Correlation is considered significant when p <0.05

extracts were combined as the inhibition activity is not changed when these leaves were mixed.

It has been reported that the phenolic compounds of plant extracts have an ability to inhibit α -glucosidase^{28 and 29}. B, P and C extracts were suggested containing different type polyphenol compounds with different affectivity against α -glucosidase. Although the total amount of the phenolic compounds in Indonesian bay leaves

is the highest among other tested leaves (Fig.2), its inhibitory activity against α -glucosidase is similar as others (Fig. 1). Our results are in line with the previously reported results whereas not all phenolic compounds or fractions in plant extracts showed similar inhibition activity against α -glucosidase enzyme^{30 and 31}. The phenolic compounds of *Artemisia* species extracts showed inhibitory activity against α -glucosidase enzyme




Fig. 1. Effect of leaf extracts on α -glucosidase inhibitory activities. Each value represents a mean \pm SEM (n = 3); BPC, BC, PC and BP were combination B, P and C, B and C, P and C, B and P respectively. B, P and C were Bay, Pandan, and Citrus leaves extracts, respectively. The bars represent the mean of three replicates. Data points denoted by different superscripts (letters on the bar) differ significantly with $p < 0,05$




Fig. 2. Total phenolic content of leaves extracts, single or combination. BPC, BC, PC and BP were combination B, P and C, B and C, P and C, B and P respectively. B, P and C were Bay, Pandan, and Citrus leaves extracts, respectively. The bars represent the mean of three replicates. Data points denoted by different superscripts (letters on the bar) differ significantly with $p < 0,05$. Total phenol content of each combination was calculated based on content of single extracts

($IC_{50} = 214.42-754.12 \text{ } \mu\text{g/mL}$) but not all individual phenolic compounds in the extract exhibited high inhibition. Extract of *Artemisia* containing high caffeoylquinic acids was the most pronounced³¹. The ethyl acetate extract of *Clinofolium taxifolium* (Kunth) Govaerts (Lamiaceae) showed stronger inhibitory activity against α -glucosidase than the methanolic and the hexanic extracts where ursolic acid contained in the three extracts was the

individual phenolic compound that showed a strong inhibitory activity³². However, similar inhibition of 2 different fractions ($P < 0.05$) against α -glucosidase also has been observed for fraction of methanol (rich in phenolic and flavonoid compounds) and ethyl acetate (rich in proanthocyanidins) from *Ehretia cymosa* Thonn³⁰. Therefore, it is suggested that phenolic compounds contained by B extract are less effective in inhibiting of α -glucosidase than




Fig. 3. Effect of plant extracts on α -amylase inhibitory activities. Each value represents a mean \pm SEM ($n = 3$); BPC, BC, PC and BP were combination B, P and C, B and C, P and C, B and P respectively. B, P and C were Bay, Pandan, and Citrus leaves extracts, respectively. The bars represent the mean of three replicates. Data points denoted by different superscripts (letters on the bar) differ significantly with $p < 0.05$.




Fig. 4. Effect of plant extracts on antioxidant activities. Each value represents a mean \pm SEM ($n = 3$); BPC, BC, PC and BP were combination B, P and C, B and C, P and C, B and P respectively. B, P and C were Bay, Pandan, and Citrus leaves extracts, respectively. The bars represent the mean of three replicates. Data points denoted by different superscripts (letters on the bar) differ significantly with $p < 0.05$.

the phenolic compounds in P or C extract, but their total phenolic concentration have no correlation with anti- α -glucosidase activity ($p = 0.654$) (Table 2).

Alpha amylase is a carbohydrate hydrolyzing enzyme secreted by salivary glands and pancreas that can hydrolyze starch into oligosaccharide and simple sugars. Inhibition of this enzymes inhibits starch digestion and reduce the rate of glucose absorption in small intestine. Our result shows that extracts of B and C exhibit α -amylase inhibition higher than P but the difference is statistically not significant (Fig. 3). As the total phenolic concentration of the three extracts was difference (Fig 2), it is suggested that the effect of the extract on α -amylase activity merely depend on phenolic composition than on the total phenolic concentration. Phenolic compounds extracted from different plant species or cultivar has different composition and anti α -amylase activity³³. Less than 15% of 126 extracts gained from 17 plants posed inhibition activity against α -amylase with varying degree, whereas 3 of them, those contain different compounds, had inhibition level more than 50%³⁴. Chemical analysis of 2 Oat varieties (Amlal and F11-5) revealed the phenolic composition of their extracts was different, where IC_{50} for amylase inhibition of F11-5 was higher than Amlal, 1027.14 μ g/mL and 723.91 μ g/mL respectively³³.

Inhibition pattern of the extracts (single or mixture) against α -amylase enzyme (Fig 3) was not similar as the pattern of the extracts against α -glucosidase (Fig 1). There is no significantly differ among the extracts in modulation of α -glucosidase activity, but each extract exhibited different effect on α -amylase activity. Mixing P with B and C increases the inhibitory activity of the extract against α -amylase as seen that percent of inhibition of BPC is significantly higher than P, eventhough their total phenolic content was not different. P and BPC extracts reduce the α -amylase activity by 16 % and 26 %, respectively. The increasing of inhibition activity of P due to mixing with B and C (BPC) was not due to increasing of total phenolic concentration (Fig 2) as no correlation between its phenolic concentration and inhibition activity was observed ($p=0.630$; Table 2). It is suggested that synergism effect was occurred when P was mixed with B and C as shown that mixture of plant extracts show superior

effect when compared to single extract at the equivalent concentration³⁵. Previously Lau *et al.*³⁶ identified a synergism effect of a mixture of extracts of *Astragalus membranaceus* and *Mannia glutinosa* roots in wound-healing of a diabetic foot ulcer animal model.

It has been shown that potency of 50 μ B-glucosidase and 50 μ B-amylase inhibition of plant extracts are related to the presence of phenolic compounds^{28,33,37} those have antioxidant activity³³. Therefore, in the current study we investigated the antioxidant powers of the extracts with DPPH radical scavenging assay. Figure 4 shown a variation in DPPH radical scavenging activities of the extracts ranging from 49% to around 77 %. The C extract exhibited a stronger DPPH scavenging ability than the P extract, though the difference is not statistically significant (76% and 74 %, respectively). On the contrary the B extract had the weakest scavenging activity (49%). A Mixture of the C and B extracts was proved to be more effective in the enhancement of the antioxidant activity of the B extract than any other combinations with the scavenging activity of 77% (Fig. 4).

The antioxidant activity patterns of extract combinations does not depend on the antioxidant activity of the single extract or their total phenolic content. For example, the total phenolic compound of the C extract was lower than of the P extract. However, both extracts exhibited a similar antioxidant activity. Additionally, the BC extract mixture showed higher antioxidant activity than the BP extract mixture (21.68 and 19.18 %, respectively), though the total phenolic compound of the BP mixture is higher than the BC mixture (7.57 and 5.90 mg/dL respectively). Therefore, accumulation rather than synergism effect on antioxidant activity was detected when the extracts were mixed. Negative correlation (-0.527) between total phenol concentration of the extracts with antioxidant activity was observed ($p = 0.014$), in which plant extracts with higher concentration of total phenol have lower antioxidant activity (Fig. 2 and Fig. 4). Similar findings have been reported showing that total phenolic compounds in the extracts of ginger, curcuma, cinnamon³⁸ and Korean propolis³⁹ have negative correlation with their antioxidant activity. Antioxidant activity of the plant extracts may not always positively

correspond to the total phenol concentration, but may be determined by the composition of phenolic compounds³⁹.

Furthermore, the molecular interactions between major phenolic compounds in plant extracts determine the antioxidant capacity of the extracts⁴⁰. A mixture of plant extracts with high total phenol compounds will have low antioxidant capacity when major phenolic compounds in the extracts have antagonistic interactions. A mixture of chlorogenic acid and caffeic acid is an example of this description whereas a combination of gallic acid and caffeic acid that showed synergistic interaction, has high antioxidant capacity⁴¹. Therefore, antioxidant activity of mixture of phenolic-rich plant extracts may not always the sum of antioxidant activity of individual extract or individual phenolic compound present in the extracts⁴⁰.

CONCLUSION

The leaf **49** acts and their combination showed inhibition activities against α -glucosidase and α -amylase enzymes and scavenging activity against free radicals. The synergism or antagonism effect was not observed when the extracts were combined as the enzyme inhibition and scavenging activities are not depend on the proportion of the extract in the mixtures. Additionally, the role of polyphenol compounds on inhibition of the starch digestion enzymes or scavenging of free radicals was not observed. Further study is required to fully elucidate the effect of the leaf or their combinations on diabetic animal models or diabetic patients.

AKNOWLEDGEMENT

This work was funded by research grant to Research Centre for Nutrition, Health and Herbal University of Lampung 2015/2016 from Research and Community Bureau, University of Lampung, Indonesia. We would also like to thank Nurlinawati PhD for reviewing the draft of this paper.

REFERENCES

- Sowmya N, Lakshmi priya N, Arumugam K, Venkatachalam S, Vijayalakshmi P, Ruchi V, et al. Comparison of dietary profile of a rural south Indian population with the current dietary recommendations for prevention of non-communicable diseases (CURES 147). *Indian J Med Res* 2006; 144(1):112-119.
- Golozar A, Khalili D, Etemadi A, Poustchi H, Fazeltabar A, Hosseini F, et al. White rice intake and incidence of type-2 diabetes: analysis of two prospective cohort studies from Iran. *BMC Public Health* 2017; 17(1):133.
- Ajala O, English P, Pinkney J. Systematic review and meta-analysis of different dietary approaches to the management of type 2 diabetes. *Am J Clin Nutr* 2013; 97(3):505-16.
- Gray GM. Starch digestion and absorption in nonruminants. *J Nutr* 1992. 122(1):172-7
- Sales PM, Souza PM, Simeoni LA, Silveira D. α -Amylase inhibitors: a review of raw material and isolated compounds from plant source. *J Pharm Pharm Sci* 2012; 15(1):141-83.
- Liu T, Yip YM, Song L, Feng S, Liu Y, Lai F, et al. Inhibiting enzymatic starch digestion by the phenolic compound diboside A: A mechanistic and *in silico* study. *Food Research International* 2013; 54(1):595-600.
- Governa P, Bainsi G, Borgonetti V, Cettolin G, Giachetti D, Magnano AR, et al. Phytotherapy in the Management of Diabetes: A Review. *Molecules* 2018; 23(1). pii: E105..
- Broadhurst CL, Polansky MM, Anderson RA. Insulin-like biological activity of culinary and medicinal plant aqueous extracts in vitro. *J Agric Food Chem* 2000. 48(3):849-52.
- Khan A, Zaman G, Anderson RA. Bay leaves improve glucose and lipid profile of people with type 2 diabetes. *J Clin Biochem Nutr* 2009; 44(1):52-6.
- Devi SL, Kannappan S, Anuradha CV. Evaluation of in vitro antioxidant activity of Indian bay leaf, *Cinnamomum tamala* (Buch. -Ham.) T. Nees & Eberm using rat brain synaptosomes as model system. *Indian J Exp Biol* 2007; 45(9):778-84.
- Faras AF, Wadkar SS, Ghosh JS. Effect of leaf extract of *Pandanus amaryllifolius* (Roxb.) on growth of *Escherichia coli* and *Micrococcus* (*Staphylococcus*) *aureus*. *International Food Research Journal* 2014; 21(1): 421-423.
- Saenthaweesuk S, Naowaboot J, Somparn N. *Pandanus amaryllifolius* leaf extract increases insulin sensitivity in high-fat diet-induced obese mice. *Asian Pacific Journal of Tropical Biomedicine* 2016; 6(10): 866-871.
- Chiabchalard A, Nooron N. Antihyperglycemic effects of *Pandanus amaryllifolius* Roxb. leaf extract. *Pharmacogn Mag* 2015; 11(41):117-22.
- Yel. Tips Memasak Nasi Lebih Enak dan Wangi Secara Alami. 2014. <https://www.vemale.com/tips-dapur/52809-tips-memasak-nasi-lebih-enak>

- dan-wangi-secara-alami.html. Downloaded 10th August 2017.
15. Tunjung WAS, Cinatl Jr J, Michaelis M, Smales CM. Anti-Cancer Effect of Kaffir Lime (*Citrus hystrix* DC) Leaf Extract in Cervical Cancer and Neuroblastoma Cell Lines. *Procedia Chemistry*2015; 14:465-468.
 16. Abirami A, Nagarani G, Siddhuraju P. In Vitro Antioxidant, Anti-Diabetic, cholinesterase and tyrosinase inhibitory potential of fresh juice from *Citrus hystrix* and *C. maxima* fruits. *Food Science and Human Wellness*2014; 3(1):16-25.
 17. Dal S, Sigrist S. The Protective Effect of Antioxidants Consumption on Diabetes and Vascular Complications. *Diseases*2016;4(3). pii: E24. doi: 10.3390/diseases4030024
 18. Lin D, Xiao M, Zhao J, Li Z, Xing B, Li X, et al. An Overview of Plant Phenolic Compounds and Their Importance in Human Nutrition and Management of Type 2 Diabetes. *Molecules*2016;21(10). pii: E1374.
 19. Nazarian-Samani Z, Sewell RDE, Lorigooini Z, Rafieian-Kopaei M. Medicinal Plants with Multiple Effects on Diabetes Mellitus and Its Complications: a Systematic Review. *Curr Diab Rep*2018; 18(10):72.
 20. Al-Waili N, Al-Waili H, Al-Waili T, Salom K. Natural antioxidants in the treatment and prevention of diabetic nephropathy; a potential approach that warrants clinical trials. *Redox Rep*2017; 22(3):99-118.
 21. Lelono RA, Tachibana S, Itoh K. In vitro antioxidative activities and polyphenol content of *Eugenia polyantha* Wight grown in Indonesia. *Pak J Biol Sci*2009; 12(24):1564-70.
 22. Ghasemzadeh A, Jaafar HZ. Profiling of phenolic compounds and their antioxidant and anticancer activities in pandan (*Pandanus amaryllifolius* Roxb.) extracts from different locations of Malaysia. *BMC Complement Altern Med* 2013;13:341.
 23. Rao RR, Tiwari AK, Reddy PP, Babu KS, Ali AZ, Madhusudana K, et al. New furanoflavonoids, intestinal α -glucosidase inhibitory and free radical [DPPH] scavenging, activity from antihyperglycemic root extract of *Derris indica* (Lam). *Bioorg Med Chem*2009; 17:5170–5175.
 24. Hasenah, A, Houghton PJ, Soumyanath A. α -amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*. *Journal of ethnopharmacology*2006. 107(3):449–455
 25. Xu JG, Hu QP, Liu Y. Antioxidant and DNA-protective activities of chlorogenic acid isomers. *Journal of Agricultural and Food Chemistry*2012; 60: 11625–11630.
 26. Ismail, J., Runtuwene, M.R.J., Fatimah, F. Penentuan total fenolik dan uji aktivitas antioksidan pada biji dan kulit buah pinang Yaki (*Areca vestiaria* Giseke). *Jurnal Ilmiah Sains*2012; 12(2):84-88.
 27. Kerru N, Singh-Pillay A, Awolade P, Singh P. Current anti-diabetic agents and their molecular targets: A review. *Eur J Med Chem*2018; 25;152:436-488.
 28. Kalita D, Holm DG, LaBarbera DV, Petrash JM, Jayanty SS. Inhibition of α -glucosidase, α -amylase, and aldose reductase by potato polyphenolic compounds. *PLoS One*2018. 25;13(1):e0191025.
 29. Bothon FTD, Debiton E, Avlessi F, Forestler C, Teulade JC, Sohounhloue DKC. In vitro biological effects of two anti-diabetic medicinal plants used in Benin as folk medicine. *BMC Complement Altern Med*2013; 13:51..
 30. Ogundajo A, Ashafa AT. Phytochemical Compositions and *In vitro* Assessments of Antioxidant and Antidiabetic Potentials of Fractions from *Ehretia cymosa* Thonn. *Pharmacogn Mag*2017;13(Suppl 3):S470-S480.
 31. Olennikov DN, Chirikova NK, Kashchenko NI, Nikolaev VM, Kim SW, Vennos C. Bioactive Phenolics of the Genus *Artemisia* (Asteraceae): HPLC-DAD-ESI-TQ-MS/MS Profile of the Siberian Species and Their Inhibitory Potential Against α -Amylase and α -Glucosidase. *Front Pharmacol*2018;9:756.
 32. Morocho V, Valle A, Garcia J, Gilardoni G, Cartuche L, Suárez AI. α -Glucosidase Inhibition and Antibacterial Activity of Secondary Metabolites from the Ecuadorian Species *Clinopodium taxifolium* (Kunth) Govaerts. *Molecules*2018.23(1). pii: E146.
 33. Marmouzi I, Karym EM, Saidi N, Meddah B, Kharbach M, Masrar A, et al. In Vitro and In Vivo Antioxidant and Anti-Hyperglycemic Activities of Moroccan Oat Cultivars. *Antioxidants (Basel)*2017; 6(4). pii: E102.
 34. Sudha P, Zinjarde SS, Bhargava SY, Kumar AR. Potent α -amylase inhibitory activity of Indian Ayurvedic medicinal plants. *BMC Complementary and Alternative Medicine*2011; 11:5.
 35. Che CT, Wang ZJ, Chow MSS, Lam CWK. Herb-Herb Combination for Therapeutic Enhancement and Advancement: Theory, Practice and Future Perspectives. *Molecules*; 18, 5125-5141 (2013).
 36. Lau KM, Lai KK, Liu CL, Tam JC, To MH, Kwok HF, et al. Synergistic interaction between *Astragalus Radix* and *Rehmannia Radix* in a Chinese herbal formula to promote diabetic wound healing. *J Ethnopharmacol*; 141(1): 250-

- 6 (2012).
37. Ombra MN, d'Acierno A, Nazzaro F, Spigno P, Riccardi R, Zaccardelli M, et al. Alpha-amylase, α-glucosidase and lipase inhibiting activities of polyphenol-rich extracts from six common bean cultivars of Southern Italy, before and after cooking. *Int J Food Sci Nutr*, **16**: 1-11 (2018).
38. Nurdin, S.U., Sukohar, A., Ramadhani, O.S. Antiglucosidase and Antiooxidant Activities of Ginger, Cinnamon, Turmeric and Their Combination. *Internasional Journal of Pharmacy & Pharmaceutical Research*; **10**(1): 296-306 (2017).
39. Wang X, Sankarapandian K, Cheng Y, Woo SO, Kwon HW, Perumalsamy H, et al. Relationship between total phenolic contents and biological properties of propolis from 20 different regions in South Korea. *BMC Complement Altern Med*; **16**: 65 (2016).
40. Palafox-Carlos H, Gil-Chávez J, Sotelo-Mundo RR, Namiesnik J, Gorinstein S, González-Aguilar GA. Antioxidant Interactions between Major Phenolic Compounds Found in 'Ataulfo' Mango Pulp: Chlorogenic, Gallic, Protocatechuic and Vanillic Acids. *Molecules*; **17**(11): 12657-12664 (2012).
41. Hajimehdipoor H, Shahrestani R, Shekarchi M. Investigating the synergistic antioxidant effects of some flavonoid and phenolic compounds. *Research Journal of Pharmacognosy (RJP)*; **1**(3): 35-40 (2014).

Antidiabetic and Antioxidant Activities of Bay, Pandan, Citrus Leaves and their Combination in Vitro

ORIGINALITY REPORT

20%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.mdpi.com Internet	86 words — 2%
2	bmccomplementaltermmed.biomedcentral.com Internet	49 words — 1%
3	www.ncbi.nlm.nih.gov Internet	28 words — 1%
4	mdpi.com Internet	22 words — 1%
5	Kazeem, Mutiu Idowu, and Theophilus Clavell Davies. "Hypoglycaemic potential of leaf extracts of <i>Lonchocarpus cyanescens</i> (Schum. and Thonn) Benth", <i>Transactions of the Royal Society of South Africa</i> , 2015. Crossref	22 words — 1%
6	Paola dos Santos da Rocha, Ana Paula de Araújo Boleti, Maria do Carmo Vieira, Carlos Alexandre Carollo et al. "Microbiological quality, chemical profile as well as antioxidant and antidiabetic activities of <i>Schinus terebinthifolius</i> Raddi", <i>Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology</i> , 2019 Crossref	19 words — 1%
7	www.science.gov Internet	19 words — 1%
8	Manconi, Maria, Maria Letizia Manca, Francesca Marongiu, Carla Caddeo, Ines Castangia, Giacomo	18 words — 1%

Luigi Petretto, Giorgio Pintore, Giorgia Sarais, Guy D'hallewin, Marco Zaru, Gianluigi Bacchetta, and Anna Maria Fadda. "Chemical characterization of Citrus limon var. pompia and incorporation in phospholipid vesicles for skin delivery", International Journal of Pharmaceutics, 2016.

Crossref

9 www.lebanoncardiology.com 18 words — 1%

Internet

10 ejournal2.undip.ac.id 17 words — < 1%

Internet

11 Maria Manconi, Maria Letizia Manca, Francesca Marongiu, Carla Caddeo et al. "Chemical characterization of Citrus limon var. pompia and incorporation in phospholipid vesicles for skin delivery", International Journal of Pharmaceutics, 2016

Crossref

12 Suktilang Majaw, Soonil Challam, Donkumar Syiem. "Effect of Potentilla Fulgens L. on Selected Enzyme Activities and Altered Tissue Morphology in Diabetic Mice", Journal of Morphological Sciences, 2018

Crossref

13 pubs.acs.org 15 words — < 1%

Internet

14 Park, Mi-Sung, Ya Xin Zhu, Hyun-Ock Pae, and Seong Hoon Park. "In Vitro and In Vivo α -Glucosidase and α -Amylase Inhibitory Effects of the Water Extract of Leaves of Pepper (*Capcicum Annuum* L. Cultivar Dangjo) and the Active Constituent Luteolin 7-O-Glucoside : Hypoglycemic Activity of Pepper Leaves", Journal of Food Biochemistry, 2016.

Crossref

15 Zhang, Liang Liang, Sai Sai Feng, and Jian Guo Xu. "Profile of Phytochemicals and Antioxidant Activities of Different Solvent Extracts of Cumin Seeds", Applied Mechanics and Materials, 2014.

Crossref

16	www.zhion.com Internet	14 words — < 1%
17	www.greenmedinfo.com Internet	14 words — < 1%
18	Rashmi S. Tupe, Neena M. Sankhe, Shamim A. Shaikh, Devyani V. Phatak, Juhi U. Parikh, Amrita A. Khaire, Nisha G. Kemse. "Aqueous extract of some indigenous medicinal plants inhibits glycation at multiple stages and protects erythrocytes from oxidative damage—an in vitro study", <i>Journal of Food Science and Technology</i> , 2013 Crossref	13 words — < 1%
19	docplayer.net Internet	13 words — < 1%
20	innovareacademics.in Internet	13 words — < 1%
21	ijbms.mums.ac.ir Internet	12 words — < 1%
22	Podsedek, Anna, Iwona Majewska, Małgorzata Redzyna, Dorota Sosnowska, and Maria Koziółkiewicz. "In Vitro Inhibitory Effect on Digestive Enzymes and Antioxidant Potential of Commonly Consumed Fruits", <i>Journal of Agricultural and Food Chemistry</i> Crossref	12 words — < 1%
23	repositorio.ufes.br Internet	11 words — < 1%
24	BahaaAldeen, Al-Trad (Prof.Dr. Gäbel Gotthold and Universität Leipzig, Veterinärmedizinische Fakultät). "Effects of Increasing Intravenous Glucose Infusions on Lactation Performance, Metabolic Profiles, and Metabolic Gene Expression in Dairy Cows", <i>Universitätsbibliothek Leipzig</i> , 2010. Publications	11 words — < 1%
25	journals.plos.org Internet	10 words — < 1%

-
- 26 Pancasakti Kusumaningrum, Hermin, Budiharjo Anto, Suprihadi Agung et al. "The characterization of Citrus sp. from Parang Island Karimunjawa based on morphological, DNA barcoding and nutritional analysis", International Journal of Genetics and Molecular Biology, 2018
Crossref 10 words — < 1%
-
- 27 Zedong Jiang, Gang Yu, Yan Liang, Tianyuan Song, Yanbing Zhu, Hui Ni, Kenichi Yamaguchi, Tatsuya Oda. " Inhibitory effects of a sulfated polysaccharide isolated from edible red alga on α -amylase and α -glucosidase ", Bioscience, Biotechnology, and Biochemistry, 2019
Crossref 10 words — < 1%
-
- 28 doaj.org
Internet 10 words — < 1%
-
- 29 M Hamburger, V Butterweck. "Abstracts of the 65th Annual Meeting of the Society for Medicinal Plant and Natural Product Research (GA)", Georg Thieme Verlag KG, 2017
Crossref 9 words — < 1%
-
- 30 Palafox-Carlos, H., E.M. Yahia, and G.A. González-Aguilar. "Identification and quantification of major phenolic compounds from mango (*Mangifera indica*, cv. Ataulfo) fruit by HPLC-DAD-MS/MS-ESI and their individual contribution to the antioxidant activity during ripening", Food Chemistry, 2012.
Crossref 9 words — < 1%
-
- 31 www.hal.inserm.fr
Internet 9 words — < 1%
-
- 32 Shruti Sancheti, Sandesh Sancheti, Mayur Bafna, Sung-Yum Seo. "2,4,6-Trihydroxybenzaldehyde as a potent antidiabetic agent alleviates postprandial hyperglycemia in normal and diabetic rats", Medicinal Chemistry Research, 2010
Crossref 9 words — < 1%
-
- 33 www.ecco-org.eu
Internet

9 words — < 1%

34 files.aiscience.org
Internet

8 words — < 1%


35 Didem Şöhretoğlu, Yasin Genç, Ü. Şebnem Harput, Suna Sabuncuoğlu, Michal Šoral, Gülin Rendad, Tibor Liptaj. "Phytochemical Content, Antioxidant and Cytotoxic Activities of ", Natural Product Communications, 2016
Crossref

36 Amit D. Kandhare, Pinaki Ghosh, Subhash L. Bodhankar. "Naringin, a flavanone glycoside, promotes angiogenesis and inhibits endothelial apoptosis through modulation of inflammatory and growth factor expression in diabetic foot ulcer in rats", Chemico-Biological Interactions, 2014
Crossref

37 Nele Gheldof, Nicki J. Engeseth. "Antioxidant Capacity of Honeys from Various Floral Sources Based on the Determination of Oxygen Radical Absorbance Capacity and Inhibition of in Vitro Lipoprotein Oxidation in Human Serum Samples", Journal of Agricultural and Food Chemistry, 2002
Crossref

38 www.pharmascope.org
Internet

8 words — < 1%

39 Xu, Donglan, Qingchuan Wang, Wenqin Zhang, Bing Hu, Li Zhou, Xiaoxiong Zeng, and Yi Sun. "Inhibitory Activities of Caffeoylquinic Acid Derivatives from Ilex Kudingcha C.J. Tseng on -Glucosidase from Saccharomyces cerevisiae", Journal of Agricultural and Food Chemistry
Crossref

40 Jianmin Man, Yang Yang, Changquan Zhang, Xinghua Zhou, Ying Dong, Fengmin Zhang, Qiaoquan Liu, Cunxu Wei. "Structural Changes of High-Amylose Rice Starch Residues following in Vitro and in Vivo Digestion", Journal of Agricultural and Food Chemistry, 2012

8 words — < 1%

-
- 41 theses.ulaval.ca 8 words — < 1%
Internet
-
- 42 www.ukaazpublications.com 8 words — < 1%
Internet
-
- 43 www.cambridge.org 8 words — < 1%
Internet
-
- 44 www.tandfonline.com 8 words — < 1%
Internet
-
- 45 Gokhan Zengin, Abdurrahman Aktumsek, Ramazan Ceylan, Sengul Uysal, Andrei Mocan, Gokalp Ozmen Guler, M. Fawzi Mahomoodally, Jasmina Glamočlija, Ana Ćirić, Marina Soković. "Shedding light on the biological and chemical fingerprints of three Achillea species (*A. biebersteinii*, *A. millefolium* and *A. teretifolia*)", *Food Funct.*, 2017 8 words — < 1%
Crossref
-
- 46 "Sustainable Technologies for the Management of Agricultural Wastes", Springer Nature, 2018 8 words — < 1%
Crossref
-
- 47 link.springer.com 8 words — < 1%
Internet
-
- 48 Jianbo Xiao, Guoyin Kai, Koichiro Yamamoto, Xiaoqing Chen. "Advance in Dietary Polyphenols as α -Glucosidases Inhibitors: A Review on Structure-Activity Relationship Aspect", *Critical Reviews in Food Science and Nutrition*, 2013 7 words — < 1%
Crossref
-
- 49 Mohammed A. Ibrahim, June C. Serem, Megan J. Bester, Albert W. Neitz, Anabella R.M. Gaspar. "Structure - Function Analysis of Peptide Analogs of SQSPA with Respect to α -glucosidase and α -amylase Inhibition", *Protein & Peptide Letters*, 2019 6 words — < 1%
Crossref
-

50 Hassan Fahmi Ismail, Zanariah Hashim, Wong Tet Soon, Nur Syukriah Ab Rahman, Ain Nabihah Zainudin, Fadzilah Adibah Abdul Majid. "Comparative study of herbal plants on the phenolic and flavonoid content, antioxidant activities and toxicity on cells and zebrafish embryo", Journal of Traditional and Complementary Medicine, 2017
Crossref 6 words — < 1%

51 Gangadhar Yamanappa Meti, Ravindra Ramappa Kamble, Dhareesh Bhimaraya Biradar, Sheetal Babu Margankop. "Synthesis of biphenyl derivatives as ACE and α -amylase inhibitors", Medicinal Chemistry Research, 2013
Crossref 6 words — < 1%

52 www.jipbs.com
Internet 5 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF