

THE TOXICITY OF PURIFIED ISOLATE
OF POLAR EXTRACT POWDER LEAFS
GLIRICIDIA MACULATA HBR. TO
CACAO MEALYBUG (PLANOCOCCUS
MINOR MASKELL)

By Ratih Andriyani



ISBN: 978-602-0860-13-8

Proceedings of 3rd International Wildlife Symposium October 18-20, 2016

*“Conserving Sumatran Wildlife Heritage
for Sustainable Livelihood”*



Institute for Research and Community Service
University of Lampung

3rd INTERNATIONAL WILDLIFE SYMPOSIUM



"Conserving Sumatran Wildlife Heritage for Sustainable Livelihood"

PROCEEDING

ISBN: 978-602-0860-13-8

Organized by:



1
RESEARCH AND DEVELOPMENT CENTER OF ENVIRONMENT
INSTITUTE FOR RESEARCH AND COMMUNITY SERVICE
UNIVERSITY OF LAMPUNG

2016

PROCEEDING IWS 2016

Person in charge:

Warsono, Ph.D.

Steering Committee:

Dr. Hartoyo, M.Si.

Organizing Committee:

Dr. Erdi Suroso, M.T.A.

Editors:

Dr. Endang Nurcahyani, M.Si.

Dr. Ir. Sumaryo Gs, M.Si.

Published by:

1

Research and Development Center of Environment

Institute for Research and Community Service

12

University of Lampung

Jl. Sumantri Brojonegoro No. 1, Bandar Lampung 35145

Phone: +62-721-705173, Fax. +621-721-773798

E-mail: lppm@kpa.unila.ac.id

ISBN: 978-602-0860-13-8

3

All right reserved (including those of translation into other languages).
No part of this book may be reproduced in any form – by photoprinting,
microfilm, or any other means – nor transmitted or translated into a
machine language without written permission from the publishers.
Registered names, trademarks, etc. Used in this book, even when not
specially marked as such, are nor to be considered unprojected by law.

WELCOMING SPEECH FROM CHAIR PERSON OF THE ORGANIZING COMMITTEE

Distinguished guests,

- Minister of Environment and Forestry Republic of Indonesia, Dr. Siti Nurbaya or representing, ⁹
- Rector University of Lampung, Prof. Dr. Ir. Hasriadi Mat Akin, M.P.
- Honorable Keynote Speaker, Invited Speakers, participants, sponshorships, ladies and gentlemen

Assalamu'alaikum warohmatullohi wabarokatuh.
May God bless all of us.
Tabik pun.

It gives me great pleasure to extend to you all a very warm welcome to the 3rd International Wildlife Symposium (IWS 2016), here in Bandar Lampung.

Ladies and gentlemen, it is gratifying to note that symposium is designed to improve awareness on wildlife conservation and sustainability in order to improve the welfare of society. To increase the consciousness and understanding on the potensial, economic value, and sustainable management of tropical wildlife through bioengineering application and to strengthen international scientific network of biological and related scientiests to share and exchange progress in various fields of wildlife research.

No matter how much we can do by ourselves on the institutional and national level, it is never enough. International level of collaboration work would be the best answer. Therefore I wish that this event which is attended by distinguished speaker and attendants from Malaysia, India, US, and Indonesia, would be a great opportunity for us to establish scientific collaboration between scientist internationally.

Hereby, on the behalf of Organizing Committee I acknowledge Dr. Siti Nurbaya, Minister of Environment and Forestry Republic of Indonesia or representing, and also to Mrs. Siti Nur Hidayati, Ph.D. (Middle Tennessee State University), as a keynote speaker, and also to the following invited speakers, Dr. Ashley Brooks (WWF Tigers Alive Initiativ), Dr. Barney Long (Global Wildlife Conservation), and drh. Dedi Candra (Way Kambas National Park) for willingness to share their valuable knowledge and scientific information.

To make this symposium happen, I would like to gratefully acknowledge to the valuable contributions from personal and institutional sponshorships including University of Lampung, Doctor Coffee, Aska Jaya, PT. Nestle Indonesia, Levi's Indonesia, and Rumah Kolaborasi (Ru-Ko). In particular, thanks a lot to the World Wide Fund (WWF) for supporting the financing of this symposium.

¹⁴
I would like also to take this opportunity to express my sincere thanks to the Head and Secretary of Research Institution and Community Service University of Lampung, for giving us opportunity and support to organize this symposium. Heartfelt thank is delivered to

steering committee, academic reviewers, organizing committee, for all participation and hard works. All of them have been working since the beginning of the planning stage and they are still here today for all of us.

Despite our best efforts, it is inevitable that there is a lack in organizing this symposium and I proudly apologize to all invited speakers, oral and poster presenters, attendants, donators, and committee members.

Finally, I would like to offer my best wishes for a highly enjoyable, succesful, productive and fruitful symposium.

Thank you so much.

Dr. Erdi Suroso
Chair Person of the Organizing Committee

OPENING REMARKS FROM THE HEAD OF RESEARCH INSTITUTION AND COMMUNITY SERVICE, UNIVERSITY OF LAMPUNG

Distinguished guests

- Minister of Environment and Forestry Republic of Indonesia, Dr. Siti Nurbaya or representing, 9
- Rector University of Lampung, Prof. Dr. Ir. Hasriadi Mat Akin, M.P.
- Honorable Keynote Speaker, Invited Speakers, participants, sponshorships, ladies and gentlemen

Assalamu'alaikum warohmatullohi wabarokatuh.
May God give us health and happiness.
Tabik pun.

It is my great pleasure to welcome all speakers and participants to the 3rd International Wildlife Symposium 2016 (IWS-2016) held in Meeting Room 2nd floor Rektorat University of Lampung, Bandar Lampung, Indonesia. I recognize that this symposium is principally designed to enhance and strengthen the contribution of researchers to the wildlife conservation. The theme of this event is "*Conserving Sumatran Wildlife Heritage for Sustainable Livelihood*". Therefore, I wish that this event will be a great opportunity and wonderful venue to lay down a cooperative framework and to internationally establish scientific collaboration among scientiests.

Hereby, I appreciatively acknowledge Dr. Siti Nurbaya, Minister of Environment and Forestry Republic of Indonesia, and also to Mrs. Siti Nur Hidayati, Ph.D. (Middle Tennessee State University), as a keynote speaker, and also to the following invited speakers, Dr. Ashley Brooks (WWF Tigers Alive Iniativ), Dr. Barney Long (Global Wildlife Conservation), and drh. Dedi Candra (Way Kambas National Park) for delivering their valuable scientific information.

My appreciation also goes to the Steering Committee, Academic Reviewers, and the Organizing Committee that spend almost their valuable time to review, manage and organize this symposium effectively. I also would like to gretefully acknowledge to the valuable contributions from personal and institutional sponshorship and funding to make this program happen.

Finally, I wish you all best wishes to have meaningfull and useful symposium. Thank you.

Wassalamu'alaikum warohmatullohi wabarokatuh.

Warsono, Ph.D.

Head of Research Institutions and Community Service

**KEYNOTE SPEAKER
MINISTER OF ENVIRONMENT AND FORESTRY
REPUBLIC OF INDONESIA**

**AT 3rd INTERNATIONAL WILDLIFE SYMPOSIUM
Bandar Lampung, 18 October 2016**

Distinguished Participants,
Ladies and Gentlemen

Assalamu'alaikum wr.wb

Good morning and May God bless us.

It is my great honor and pleasure to attend to this event and deliver my speak. Let me express my appreciation to University of Lampung in collaboration with WWF Indonesia for organizing this symposium. Hopefully from this symposium which brings together scientific community and field experts, where we share knowledge, experience and concern, can enhance and synergize our efforts to cope issues in various aspects.

Ladies and gentlemen,

Indonesia should be proud to be a country which has rich biodiversity, reported it reaches 47.910 species, that makes Indonesia is well known as mega biodiversity country. At the same time, Indonesia is responsible to promote sustainable use of biodiversity to improve society and country's well-being.

This responsibility will be challenging for Indonesia, considering our country as one of the hot spot of biodiversity loss. The threat mainly coming from habitat loss caused by encroachment, forest fragmentation and forest fires as well as coming from illegal logging, and trade.

Plants and wildlife are one of supporting elements for human life, which their existence hold important and irreplaceable roles. Having that said, I would like to take this opportunity to encourage all of us to protect and conserve the sustainability of wildlife and plants as heritage for the next generation and sustainable livelihood.

Ladies and gentlemen,

Strategy and policy of Indonesian Government to secure the biodiversity are directed into three focuses namely protection, preservation and sustainable use of ecosystem, species and genetic resources. National regulation has been enacted, to mention some, including UU 5/ 1990, UU 41/ 1999, UU 32/ 2009, PP 7/ 1999, PP 8/ 1999 as well as strategic and action plan of several endangered and umbrella species such as sumatran tiger, orangutan, rhino, sumatran elephant, javan eagle, tapir, proboscis monkey (*bekantan*), maleo, banteng and babyrousa.

Our global commitment on biodiversity conservation also reflected on ratification of several conventions such as Convention on Biodiversity (CBD) through UU 5/1994, Ramsar through Keppres No 48/1991, UNFCCC through UU No. 6/1994 and protocols such as Kyoto Protocol through UU No. 17/ 2004, Nagoya Protocol through UU 11/ 2013 and Cartagena Protocol through UU 2/ 2014.

We also aware that in this global era, efforts on the conservation of biodiversity always in the spotlight of international attention. Threats on the biodiversity, for example illegal killing of endangered species have and will always be connected with the issues of deforestation and habitat destruction and these will be the entry points for discrediting and black-campaigning against Indonesia, that in turn can be impacted to Indonesia's products in the global market. Thus saving the biodiversity requires active participation from all stakeholders including private sector.

Biodiversity including ecosystem and genetic resources can and should be utilized in a sustainable manner for human welfare such as for source of food, clothes, medicine, water, energy, and oxygen, for controlling climate and disease, ecosystem balance as well as for leisure.

Sustainable use of biodiversity, for example as I mention before for source of food has been a main discussion in many forums. With the growth of human population, it is a must to conduct study and formulate strategy to maintain our natural resources that can be utilized not only for our generation but also for our kids' generation and further. With this regard, Indonesia as mega-biodiversity country plays an important role as source of germplasm which may contain useful substance for human health or important for bioprospecting to increase country's revenue.

To illustrate, global trade on medicinal herbs reach approximately US\$ 60 billion/year. While protection of coral to support genetic resources for research on medicine can provide revenue US\$ 55-1.110 per ha/year in South East Asia (source: CBD). Indonesia revenue on export from traditional medicine (jamu) reach US\$ 113 million, while for domestic reach US\$ 100 million (source: BPOM 2007).

Ladies and gentlemen,

Despite having all of potency and opportunity, there are also threats and challenges facing our existing biodiversity:

Globally, these includes pressure from human population growth which require demand for land, food, energy and clean water; climate change; and increasing demand of genetic resources for food and energy.

Nationally, these includes illegal logging, forest fire, encroachment, illegal trade, declining of wildlife population, loss of habitat, invasive alien species, as well as low resources capacity and quantity (human and fund) and lack of integrated database.

Hence enormous efforts have been taken by Government Indonesia namely:

- a. public awareness and campaign by involving religious leader and other groups for promoting religious and local wisdoms as well as local engagement.
- b. to restore and protect the population at local level in their habitat to prevent further damage to the population.
- c. Strengthen coordination among government institutions and networking with CSO's.

Currently Ministry of Environment and Forestry Indonesia with relevant law enforcement institutions (Police, General Attorney, Financial Transaction Reports and Analysis Center, and Financial Services Authority) have commitment to support "multi door law enforcement". By implementing multi door law enforcement initiative including applying of

corruption and money laundry act in line with environmental, conservation and forestry act is expected that it could strengthen deterrent effects.

23

Ladies and gentlemen,

As we are all aware, illegal activity related to environmental and forestry including wildlife illegal trafficking is now even more sophisticated, organized and transnational crime, which involves a large network of actors that make up its own chain. We cannot stop it alone. Therefore collaborative among relevant actor/ stakeholder is badly needed to tackle these issues in effectively.

We believe that global collaboration through bilateral, regional and multilateral cooperation can increase the effectiveness in combating illegal activities such as wildlife trafficking. Hence, Indonesia has involved in global collaboration such as:

- a. Bilateral cooperation with Vietnam and USA
- b. Regional cooperation in the framework of the ASEAN-WEN (Wildlife Enforcement Network)
- c. London Conference, Kasane Conference, The Hague Conference on wildlife
- d. Multilateral Cooperation: Interpol, CITES, Global Wildlife Programme
- e. We also have cooperation with International and National NGO concern in combating wildlife crime, such WWF, WCS, etc.

In this moment, I would also inform that we are in the progress renewing our conservation act in order to increase effectiveness of conservation efforts including wildlife law enforcement.

Distinguish ladies and gentlemen,

To develop strategy on biodiversity conservation and sustainable use require strong research and scientific evidence which will be needed to convince government agency and related stakeholder to be aware and act on the right approach. Thus, active participation of scientific community in communicating their knowledge should be appreciated and facilitated such as through this event.

To conclude, the efforts for the conservation of biodiversity required the involvement of us all, not just the governmental institutions only, but also private sectors, NGOs, civil societies and scientific community. I sincerely hope that this symposium can provide a media for us all to share knowledge, experience and concern, and to synergize all efforts.

Wasalamualaikum Warahmatullohi Wabarohkatuh

Bandar Lampung, 18 October 2016

Minister of Environment and Forestry,

Dr. Siti Nurbaya

LIST OF CONTENTS

	Pages
WELCOMING SPEECH FROM CHAIR PERSON OF THE ORGANIZING COMMITTEE	iii
OPENING REMARKS FROM THE HEAD OF RESEARCH INSTITUTION AND COMMUNITY SERVICE, UNIVERSITY OF LAMPUNG	v
KEYNOTE SPEAKER: MINISTER OF ENVIRONMENT AND FORESTRY REPUBLIC OF INDONESIA	vi
SAFE SYSTEMS: HWC Safe Systems Approach and the HWC Rapid Assessment tool (Ashley Brooks, Ph.D.)	x
PROMOTING MULTI-STAKEHOLDER INTERNATIONAL COLLABORATIONS FOR ENDANGERED SPECIES RECOVERY (Barney Long)	xiv
INTEGRATING PLANTS INTO WILDLIFE CONSERVATION PROGRAMS (Siti Nur Hidayati, Ph.D.)	xvii
1. PREVENTION MODELS TOWARDS HUMAN - TIGER CONFLICT (HTC) IN BUKIT BARISAN SELATAN NATIONAL PARK (BBSNP), LAMPUNG (Firdaus Rahman Affandi, Tugiyono, G. Nugroho Susanto, Elly Lestari Rustiaty) ...	1 -- 10
2. IMPACT OF ANIMAL HOUSING TOWARDS WORMS INFECTION IN LOCAL BEEF CATTLE FARMS IN DUKUHBADAG VILLAGE, CIBINGBING, KUNINGAN, WEST JAVA, INDONESIA: AN ANALYSIS (Retno Widyani, Fitri Dian Perwitasari, Mus Nilamcaya, Ida Herawati)	11 -- 17
3. ESTABLISHING BASELINE DATA ON FISHERMAN AND FISH CAUGHT ON THE SERKAP RIVER, KAMPAR PENINSULA, RIAU (Sidiq Purwanto)	18--24
4. WALKING THROUGH CONVERSION: A MONITORING OF ELEPHANT MOVEMENT IN DEGRADED FOREST OF TESSO NILO LANDSCAPE (Febri Anggriawan Widodo, Wishnu Sukmantoro, Heri Irawan, Eka Septayuda, Yansen Gultom, Samsuardi, Sunarto, Nurchalis Fadhlil)	25--29
5. EVALUATING THE INTERVENTION METHODS TO REDUCE HUMAN-ELEPHANT CONFLICT AROUND WAY KAMBAS NATIONAL PARK (Sugiyo, Ardiantiono, Agus Santo, William Marthy, Fahrul Amama)	30--36
6. JAVAN RHINO (<i>RHINOCEROS SONDAICUS</i>), BANTENG (<i>BOS JAVANICUS</i>) & OTHER MAMMALS COEXISTENCE IN UJUNG KULON NATIONAL PARK: SPATIAL AND TEMPORAL OVERLAP (Mahmud R, Rahmaningsih MD, Sunarto, Daryan, Firdaus AY, Muhtarom A, Setiawan R)	37--49
7. FILLING THE KNOWLEDGE GAP ON THE ENDANGERED ASIAN TAPIRS IN SOUTHERN PART OF TROPICAL RAINFOREST HERITAGE OF SUMATRA (Ardiantiono, Fahrudin Surahmat, Tri Sugiharti, Wulan Pusparini)	50--57
8. PEKON MUARA TEMBULIH, NGAMBUR, PESISIR BARAT: PRELIMINARY STUDY ON THE CHARACTERISTICS OF TURTLE HABITAT (Brina Wanda Pratiwi, Sugeng P. Harianto, Elly Lestari Rustiati)	58--65
9. SUMATRAN ELEPHANT (<i>ELEPHAS MAXIMUS SUMATRANUS</i> T) FOOD COMPOSITION AND ITS PREFERENCE IN TESSO NILO NATIONAL PARK (Defri Yoza and Yuliantony)	66--77
10. DIVERSITY AND ABUNDANCE OF AVIAN COMMUNITY AT COASTAL LAGOONS IN BUKIT BARISAN SELATAN NATIONAL PARK, INDONESIA: WHY WATERBIRD IS LACKING? (Ani Mardiasuti, Yeni A. Mulyani, Lina K. Dewi)	78--85

11.	HUMAN ELEPHANT CONFLICT STUDY BASED ON THE COMMUNITY INFORMATION IN RIAU – INDONESIA (Wishnu Sukmantoro, Yansen Gultom, Heri Irawan)	86--90
12.	STUDY ON HEALTH CARE MANAGEMENT SYSTEM OF CAPTIVE SUMATRAN ELEPHANT (<i>Elephas maximus sumateranus</i>) IN Prof. Dr. Ir. M. RUBINI ATMAWIDJAJA ELEPHANT HOSPITAL, WAY KAMBAS NATIONAL PARK (Firda Nur Islami, Dedi Candra, Diah Esti A, Priyambodo)	91--93
13.	A PRELIMINARY STUDY ON POPULATION ESTIMATION TECHNIQUE OF SIAMANG (<i>Sympalangus syndactylus</i>) IN WAY CANGUK RESEARCH STATION, BUKIT BARISAN SELATAN NATIONAL PARK (Nafila Izazaya Idrus, Ryan Setiono, Fahrudin Surahmat)	94--98
14.	HELMINTHES PARASITIC PARAMPHISTOMUM SP INFECTION ON THE SUMATRAN ELEPHANTS IN ELEPHANT TRAINING CENTER WAY KAMBAS NATIONAL PARK LAMPUNG (Dedi Candra, Diah Esti, Elisabeth Devi, Catur Marsudi)	99--101
15.	TRAPPING FRUIT EATING BATS IN WAY CANGUK RESEARCH STATION, BUKIT BARISAN SELATAN NATIONAL PARK: MIST NET VS HARP TRAP (M. Khairul Ikhwan, Eka S. Ariyanti, Fahrudin Surahman, Janjiyanto)	102--105
16.	RESCUE SUMATRAN ELEPHANT BABY WITHOUT TRUNK IN WAY KAMBAS NATIONAL PARK LAMPUNG (Elisabeth Devi K, Dedi Candra, Diah Esti Anggraini, Nazarudin, Mahfud Handoko)	106--108
17.	THE TABANID FLY BIODIVERSITY AND ITS POTENCY AS TRANSMISSION VECTOR OF TRYPANOSOMIASIS TO THE JAVAN RHINO POPULATION WITHIN THE UJUNG KULON NATIONAL PARK (Gita Alvernita, Kurnia O. Khairani, Dariyan, Dyah Lukitaningsih, Supriyono, Dedy S. Pahlawan, Zaenal Gesit Kalbuadi, Upik Kesumawati Hadi)	109--113
18.	ELEPHANT ENDOTHELIOTROPIC HERPESVIRUS (EEHV) MANAGEMENT IN ELEPHANT CONSERVATION CENTER WAY KAMBAS NATIONAL PARK LAMPUNG (Diah Esti, Dedi Candra, Anhar Lubis, M. Wahyu, Elisabeth Devi)	114--116
19.	AN EXPERT SYSTEM TO DIAGNOSE CHICKEN DISEASES WITH CERTAINTY FACTOR BASED ON ANDROID (Aristoteles, Kusuma Adhianto, Puja Putri A)	117--126
20.	COMPARISON EFFECTIVENESS OF ANTIOXIDANT ACTIVITY EXTRACT HERBAL MIXTURE OF SOURSOP LEAF (<i>Annona muricata</i>), BAY LEAF (<i>Syzygium polyanthum</i>) AND PEGAGAN LEAF (<i>Centella asiatica</i>) (Khairun Nisa Berawi, Liana Shidarti, Samsu U. Nurdin)	127--132
21.	THE UTILIZATION OF ISOLATE <i>Bacillus thuringiensis</i> GRAYAK LARVAE PEST (<i>Spodoptera litura</i> Fab.) ON CABBAGE (<i>Brassica oleraceae</i> var. capitata Linn.) (Wibowo Nugroho Jati, Felicia Zahida, Sara Puspareni Prayitno)	133--137
22.	LEG AMPUTATION OF TIMOR DEER (Hastono, S.D)	138--140
23.	IDENTIFICATION OF THE SUMATRAN RHINO FOOD PLANTS IN WAY KAMBAS NATIONAL PARK LAMPUNG (Dedi Candra, Sumadi Hasmaran, Lamijo, Supriyono)	141--146
24.	SURVEILLANCE ANTHRAX (<i>Bacillus anthracis</i>) IN SURROUNDING WAY KAMBAS NATIONAL PARK LAMPUNG INDONESIA (Dedi Candra, Arie Khoiriyah, Diah Esti Anggraini, Joko Siswanto)	147--151
25.	GENOMIC DNA ISOLATION OF GAJAH SUMATERA (<i>Elephas maximus sumatrensis</i>) IN ELEPHANT TRAINING CENTER, WAY KAMBAS NATIONAL PARK, EAST LAMPUNG (Elly L. Rustiati, Priyambodo)	152--155
26.	INDUCE RESISTANCE OF <i>SPATHOGLOTTIS PLICATA</i> BL. TOWARD TO	156--158

22

FUSARIUM OXYSPORUM (Endang Nurcahyani, Rochmah Agustrina, Erdi Suroso)

27. THE EFFECTS OF A HEXANE FRACTION OF RED BETEL LEAF (*Piper cricutum*) ON LEARNING AND MEMORY IN MICE (Pratika Viogenta, Lilik Koernia Wahidah, Yudha Erlangga) 159--163
28. THE LOCAL KNOWLEDGE OF COASTAL ETHNIC COMMUNITIES OF PLANTS THAT EFFICACIOUS AS MEDICINE IN 5 DISTRICTS OF SOUTH LAMPUNG REGENCY (Arum Asterini, Yulianty, Tundjung Tripeni Handayani) .. 164--169
29. PHYTOTELMATA SPECIES AND ITS DISTRIBUTION IN SOUTH PRINGSEWU, LAMPUNG (Putri Minggar Oktaviani, Emantis Rosa, Yulianty) ... 170--174
30. THETOXICITY OF PURIFIED ISOLATE OF POLAR EXTRACT POWDER LEAFS *GLIRICIDIA MACULATA* HBR. TO CACAO MEALYBUG (*PLANOCOCCUS MINOR* MASKELL) (Ratih Andriyani, Nismah Nukmal, Emantis Rosa) 175--181
31. SOCIAL BEHAVIOR OF SPOTTED DEER (*Axis axis*) IN GUNUNG MADU PLANTATIONS INC.SANCTUARY LAMPUNG TENGAH LAMPUNG PROVINCE INDONESIA (Rita Gusmalinda, Bainah Sari Dewi, Niskan Walid Masruri) 182--188
32. THE COMPARISON OF TOXICITY PURIFIED ISOLATE OF WATER AND METHANOL EXTRACTS OF PAWDER LEAF *GLIRICIDIA MACULATA* ON MORTALITY SOURSOP MEALYBUG *PSEUDOCOCCUS CRYPTUS* (Fahrul Aksah, Nismah Nukmal, Emantis Rosa) 189--196
33. DEVELOPMENT OF BOTANICAL INSECTICIDE FROM FLAVONOID OF COMPOUND LEAF EXTRACT *GLIRICIDIA MACULATA* TO CONTROL COFFEE MEALYBUG *PLANACOCCUS CITRI* (Apriliyani, Nismah Nukmal, Emantis Rosa) 197--204

1
**THE TOXICITY OF PURIFIED ISOLATE OF POLAR EXTRACT
 POWDER LEAFS *GLIRICIDIA MACULATA* HBR. TO CACAO
 MEALYBUG (*PLANOCOCCUS MINOR* MASKELL)**

Ratih Andriyani¹, Nismah Nukmal^{2,3} and Emantis Rosa²

¹) Biology Magister Student ²) Biology Department, ³) Corresponding Author
 e-mail: nismah.nukmal@fmipa.unila.ac.id

ABSTRACT

One contributing factor in decreasing productivity of cocoa in the last years is due to pest attack. Cacao mealybug (*P. minor*) attack the young cacao fruits, by sucking them until dry and die. Therefore it should be controlled. Alternatives pest control of the insecticide has been widely searched. *G. maculata* leaves consist of rich flavonoid that potential as botanical insecticide. In order to get the purified isolate of polar extract powder leaf *G. Maculata* that named (PIGR), and test its toxicity to cacao mealybug (*P. minor*), the powder leaf of *G. maculata* were extracted by using various organic solvents (n-hexane, dichloromethane, methanol and water). A set of laboratory experiment was conducted to test the toxicity by bioassay, and to know the type and structure of PIGR by spectroscopic analysis. Five different concentrations (0 %, 0,015 %, 0,030 %, 0,045 % and 0,060 %) of PIGR with each of 3 replications were tested to cacao mealybug mortality. Mortality observed at 12, 24, 48 and 72 hours after treatment. Probit analysis was conducted to obtain LC₅₀. The result indicated the PIGR was toxic to cacao mealybug (*P. minor*) with LC₅₀ 72 hours metanol extract 0,054% and water extract 0,047%. Therefore water extract more toxic than metanol extract. The toxic compound of methanol extract and water extract *G. maculata* is flavon with the structural frame is 2-phenyl-1,4-benzopyron.

Keywords : polar extract, powder leaf, cacao mealybug, flavon.

1. INTRODUCTION

Cocoa (*Theobroma cacao* L) is one of the important tree crops in Indonesia, which has many beneficial value. Production of cocoa in Indonesia is often decreased. The cause of the decline in the production of cocoa beans is due to pest attack mealybugs (Wijaya, 2007). Mealybug (*Planococcus minor* Maskell) was sucking immature fruit causing dry and die (Sumarno, 2015). The use of synthetic insecticides are not right will bring bad effects, more harm than benefit resulting. Leaves gamal (*Gliricidia maculata* Hbr.) Has the active ingredient coumarin which are insecticides, rodenticides and bactericide that can be used as environmentally friendly insecticide plant (Kementerian Pertanian Ditjen Peternakan dan Keswan, 2009). *Gliricidia* leaves contain toxins that dikumerol main compound, a compound capable of binding vitamin K to clot blood. Dikumerol represents the conversion of coumarin due to bacterial activity when it ferments. Coumarin is suspected flavonoid compounds can irritate the skin and can inhibit the transport of amino acids leucine (Nukmal et al, 2010)

The results of the study Nukmal et al., (2009 and 2010) also proved that extracts polar (water and ethanol) *Gliricidia* leaves can cause 100% mortality in the pest imago boils dadap (*Quadrastichus erythrinae*) after 72 hours of treatment at the laboratory scale.

From the results of studies have been done to prove that the *G. maculata* leaf powder extract has potential as botanical insecticide. Purposes this study to know the toxicity, the type and structure of the *gliricidia* (*G. maculata*) pure isolates powder leaf are effective in control (*P. minor*) to the cocoa (*T. cacao*).

2. RESEARCH METHODS

A. Tools and Materials

The tools used this research are Machete, sacks, grinding machines, scales, glass jars, filter paper, Rotary evaporation, separating funnel, Freezedryer, UV light, aluminum foil, erlenmeyer flask, test tubes, spatulas, analytical balance, oven, beaker, beakers, pipettes, funnel, hot plate, electric heating, capillary pipette. The tools used to prepare the test insects is a plastic knife, jars, gauze pads, rubber bands, brushes and pins.

The materials are Gliricidia leaves, solvent n-hexane, dichloromethane (DCM), methanol, distilled, KK Amberlite XAD-4, Plate TLC cellulose, H₂SO₄ as an ingredient identifier solution, and HCl is used to adjust the pH at the time of fractionation. CeSO₄ visualization solvent, AlCl₃, H₃BO₃ and NaOH. HCl, NaCl and ethyl acetate to extract water hydrolysis, H₂SO₄ as an ingredient in a solution of identifiers when TLC. Cocoa mealybugs (*P. minor*), cocoa fruits.

B. Isolation and Purification of Flavonoid Compounds Group Methanol extracts

500 gr of Gliricidia powder leaves macerated using a solvent Hexana 1.500 ml, DCM solvent 1000 ml, methanol 1.200 ml and 1.200 ml water. Total filtrate of methanol was evaporated and then recrystallization method using freezedryer for 72 hours. Furthermore, to the purification of the methanol extract conducted by fractionation using column chromatography (CC) Amberlite XAD-4. Amberlite XAD-4 5 gr was added to column chromatography. Fractions that have been obtained are evaporated and TLC analyzed to obtain the active fraction rich in flavonoids.

Water filtrate was concentrated Gliricidia leaf powder by recrystallization method using freezedryer for 72 hours. After that extract water hydrolysed. The results of hydrolysis showed two-phase extraction, the water phase and ethyl acetate phase. Furthermore, the results of hydrolysis was monitored by TLC.

Bioassays were carried out is the test of mortality with residual effect (residual effect) Bioassay made by soaking the test medium with 5 level of concentration levels (0%, 0.015%, 0.030%, 0.045% and 0.060%) for 10 minutes, 10 test insects tail (*P. minor*) females who already acclimatized for 1 day before the treatment is placed on the test medium. Observations of insect mortality trial conducted at 12, 24, 48 and 72 hours after treatment. The percentage of deaths for each extract will be analyzed with probit analysis EXE program, test ANOVA and continued with Tukey's test was used to determine an effective solution as botanical insecticide to determine the relationship with the concentration of insect death. Test solution is said to be effective if the solution is giving the LC₅₀ value $\leq 5\%$ (Prijono, 2005). These trials were conducted each with 3 replications. The data were then analyzed using probit analysis to determine LC₅₀ values. Determination Pure Active Compounds Structure using spectroscopic methods.

3. RESULTS AND DISCUSSION

A. Isolation and Purification of Flavonoid Compounds

The resulting methanol extract as much as 53 gr. The resulting crude extract freezedryer process is as much as 20 grams in the form of pasta that is solid green.

Results fractionation methanol crude extract obtained 44 fractions. Of the 44 fractions only 4 fractions which showed spots on the chromatogram as an indication of flavonoid compound that is the fraction 1, fraction 2, fraction 26 and fraction 27. Fraction 1 and 2 has a R_f value of 0,700. Fraction 26 and 27 have a value of R_f 0,875. R_f equal value can be said that the compounds identified as having the same or similar characteristics (Khopkar, 1990).

TLC analysis results after the evaporated fraction flavonoid addressing their spots on the chromatogram that the fraction 1, fraction 2, fraction 26 and fraction 27 with a value of R_f 0,725

whereas other fractions did not indicate the presence of flavonoid compounds. The same RF value is the active fraction methanol and allegedly have the same classes of compounds.

Extract the water produced as many as 10 grams of crude extract water in the form of a brown paste. Pure water extract obtained by hydrolysis with the aim to break bonds glucosides in the extract. Hydrolysis form crystalline precipitate 2 grams, water phase of 20 ml and ethyl acetate phase as much as 15 ml. Based on the analysis chromatogram TLC Rf value of the water phase amounted to 0,625 while on crystal or precipitation does not show stains on the TLC plate as an indication of flavonoid compounds therefore used during the bioassay is to extract the water phase.

B. Mortality *Planococcus minor* Maskell

Table 1. The average percentage mealybug pest mortality (*Planococcus minor* Maskell) treatment with the methanol extract and water extract leaf *G. maculata* with different concentration and time observation.

Time observation after treatment (hours)	Concentration (%)	The death of insects (%)	
		Methanol Extract	Water Extract
12	0	0,00	0,00
	0,015	3,33	0,00
	0,030	0,00	3,33
	0,045	3,33	3,33
	0,060	0,00	0,00
24	0	0,00	0,00
	0,015	3,33	3,33
	0,030	3,33	10,00
	0,045	6,67	10,00
	0,060	6,67	10,00
48	0	0,00	0,00
	0,015	13,33	13,33
	0,030	23,33	20,00
	0,045	13,33	26,67
	0,060	23,33	26,67
72	0	0,00	0,00
	0,015	33,33	30,00
	0,030	33,33	40,00
	0,045	36,67	46,67
	0,060	53,33	56,67

At 12 hours after treatment methanol extract and water extract can kill insects already test (*P. minor*) as much as 3.33%. But death occurs on lower concentration than the high concentration. This could be due to several factors. According Raini (2007) there are several factors that influence the toxicity of a compound of one of them is endurance test animals. Possibilities that occur in test animals (*P. minor*) were treated extract higher concentrations have more endurance stronger. The higher the concentration of the extract is treated persentation higher death rate insects. When compared with the methanol extract, extract the water was shut off more test insects between 3,34% - 13, 34%. This tends to occur due to the influence of several factors among which the dose of the extract, endurance test animals, and exposure times. Low dose will give the effect of compound toxicity is low. While high doses at the time of initial exposure will force the body to continuously defend themselves from substances that are toxic, but exposure times would make these toxic substances accumulate in the body resulting in chronic poisoning and death (Raini, 2007).

Results of analysis of variance showed a significant difference among the treatments. The average mortality mealybug by treatment concentration and time showed a significantly difference. While the

average mortality mealybug if seen from a comparison between extracts, concentration and extract, and extract the time, the concentration, time, and the extract did not show significant differences.

The results Tukey's at the level α 5% average mortality pest infestation white treatment of methanol extract and water extract at 72 hours after treatment showed that the methanol extract and extract water with a concentration of 0% was significantly different from the concentration of 0,015%, 0,030%, 0,045 and 0,060%. 0,015% concentration was significantly different from the concentration of 0,060%, but not significantly different from the concentration of 0,030% and 0,045%. The concentration of 0,030% and 0,045% is not significantly different from the concentration of 0,060%. The difference is not noticeable due to the difference in mortality rates between white lice are very small concentrations. There are real differences between the control and treatment groups showed that the concentration significantly affect mortality mealybug.

The results Tukey's at the level α 5% of the average number of deaths pest mealybugs were treated extracts of methanol and water extract of leaves of *G. maculata* at a concentration of 0,060%, ie death mealybugs were treated with methanol extract and extract water at 12 hours after treatment was not significantly different with 24 hours after treatment but significantly different with 48 and 72 hours after treatment. This suggests that the effect on the time of death mealybug pests. The longer the period of observation after treatment also increase the mortality rate of mealybugs.

Statistically methanol extract and water extract did not show significant differences but when seen from the effectiveness of the extract showed significant differences. It can be seen Table 2 and Table 3.

Table 2. Value LC_{50} results probit analysis of methanol extract and water at 12-72 hours after treatment.

Time (Hours)	LC_{50} value (%)		Difference (%)
	Methanol Extract	Water Extract	
12	-	-	-
24	-	-	-
48	0,104	0,082	0,022
72	0,054	0,047	0,007

LC_{50} value of the water extract of between 0,007 to 0,022 lower than the methanol extract of the same treatment period. This indicates that the water extract is more effective than the methanol extract to kill 50% of test animals.

Table 3. Value LT_{50} probit analysis results of methanol and water extracts at different concentrations

Concentration (%)	LT_{50} value (hours)		Difference (hours)
	Methanol Extract	Water Extract	
0,015	89,567	87,777	1,790
0,030	81,707	82,399	0,692
0,045	87,676	73,535	14,141
0,060	68,785	66,248	2,537

LT_{50} values were also lower water extract between 0,692- 14,41 hours than methanol extract. This indicates that the water extract is more effective than the methanol extract with the same concentration takes longer than methanol extracts even at concentrations of 0,045% difference very long time is 14,141 hours, it indicates the water extract is more effective than the methanol extract.

The ability of the power to kill the methanol extract and water extract of leaves of *G. maculata* caused their compounds which are toxic secondary metabolites. One of them is of secondary metabolites flavonoid. The content of flavonoid compounds in leaf extracts gliricidia seen from the TLC analysis using several solvents visualization (Table 4).

Table 4. Rf value on the analysis of secondary metabolites (flavonoids) methanol extract and water extract of leafs *G. maculata* with TLC methods using several solvents and solvent developers visualization DCM and methanol (4: 1)

Extract	Solvent Visualization				Average value of Rf
	CeSO ₄	AlCl ₃	NaOH	H ₃ BO ₃	
Methanol Extract	0,727	0,750	0,675	0,850	0,750
Water Extract	0,625	0,828	0,750	0,625	0,707

Rf Value flavonoid compounds using some solvent visualization shows varying values, the value of Rf methanol extract using solvents visualization CeSO₄ and H₃BO₃ higher than the value of the Rf in the water extract but the value of Rf methanol extract using solvents visualization AlCl₃ and NaOH lower than the value of Rf on the extract water. This shows that the solvent visualization has the ability to identify different as the principle of KLT according Soebagio (2002) in which the adsorption is absorption at the surface, while the partition is the spread, or the ability of a substance or substances present in the solution to separate and move upwards depending on plate TLC and the solvent used. From the results of the average value of Rf obtained by the use of various solvents visualization of the average value of Rf extract water smaller than the average value of Rf in methanol extract. According to Yazid (2005) the higher the Rf value obtained, the lower the level of the polarity of a substance that, because the concept of the higher polarity of a substance, then a stationary phase which is polar compounds will bind to each other and form a very strong bond so that the distance spot or stains on the TLC plate gets smaller and the lower Rf value. Thus the degree of polarity of the water extract is higher than the methanol extract.

C. Types and Chemical Structure of Extract Leafs *G. maculata*

Based on the results of spectroscopic analysis of the methanol extract of leaves of *G. maculata* has absorbance value of 0,585 -0,849. The presence of flavonoid compounds can be seen from the maximum wavelength of between 230,00 295,00 nm- nm (Neldawati et al, 2013). The methanol extracts were analyzed to give two maximum absorption is from 310,00 to 350,00 nm and 250,00 to 280,00 nm (Figure 1).




Figure 1. The spectrum of the methanol extract powder leafs *G. maculata*

The results of spectroscopic analysis of aqueous extracts were analyzed giving the maximum absorption is 250,00-280,00nm with the absorbance value of 0,736 (Figure 2).

According Neldawati et al, (2013) from 310,00 to 350,00 nm wavelength and 250,00-280,00nm included into the class of flavonoids types of flavonoids. Thus both polar leaf powder extract containing flavonoids gliricidia kind flavones. Demand Tapas et al, (2008) the characteristics of flavonoids is almost the same as flavonoid compounds that have the basic structural framework C₆-C₃-C₆ consisting of two C₆ aromatic rings (A and B) and heterocyclic ring (C) (Figure 3).

Flavones are flavonoid class consists of the structural frame 2-phenyl-1,4-benzopiron. In chemistry, a phenyl or phenyl ring are one functional group in a chemical formula. In this group, six carbon atoms arranged in a cyclic ring structure. This ring is very stable, and is part of a group of aromatic compounds. The phenyl ring is hydrophobic (water repellent) and aromatic hydrocarbons. These

groups can be found in many organic compounds. This ring is estimated to be derived from benzene, (C₆H₆). Phenyl bergugus simplest compound is phenol, C₆H₅OH. While benzopiren is an organic compound with the formula C₂₀H₁₂ polycyclic aromatic hydrocarbons .Benzopiren itself is a chemical that is toxic (Marais, et al., 2006; Kumar and Pandey, 2013).




Figure 2. The spectrum of water extract leafs *G. maculata*




Figure 3. The structure of the flavonoids
(Source: Tapas et al, 2008)

4. CONCLUSION

Based on the research results obtained the following conclusions:

1. Methanol extract and water extract of powder leafs Gamal (*Gliricidia maculata* Hbr.) Has the power insecticides against mealybugs (*Planococcus minor* Maskell) on the cocoa plant (*Theobroma cacao* L).
2. Methanol extract the and water extract of powder leafs gamal of containing flavonoids, with type flavone that the structural frame 2-phenyl-1,4-benzopyron.

REFERENCES

- 2 Kementerian Pertanian, Ditjen Peternakan dan Keswan. 2009. *Keunggulan Gamal Sebagai Pakan Ternak*. BPTU Sembawa. Sumatera Selatan.
- 15 Khopkar, S. M. 1990. *Konsep Dasar Kimia Analitik*. UI Press. Jakarta.
- 10 Kumar, S., dan Pandey, A. K. 2013. *Chemistry and Biological Activities of flavonoids*. *The Scientific Word Journal*. Vol.2013. Article ID 162750, 16 pages. India.
- Marais, J.P.J., Vours, B. D., Dixon, R.A., Ferreira, D. 2006. *The Stereochemistry of Flavonoids*. Springer Science. ISBN-10 0-387-28821. United Dtates of America. America

- 5 Neldawati, Ratnawulan, dan Gusnedi. 2013. Analisis Nilai Absorbansi Dalam Penentuan Kadar Flavonoid Untuk Berbagai Jenis Daun Tanaman Obat. *Pillar Of Physics, Vol. 2. Oktober 2013*, 76-83
- 2 Nukmal, N., Widiastuti, E.L., dan Sumiyani, E. 2009. Uji Efikasi Ekstrak Daun Gamal (*Gliricidia maculata*) Terhadap Imago Hama Bisul Dadap (*Quadrastichus erythrinae*). *Prosiding Seminar Nasional XX dan Kongres Biologi Indonesia XIV*. Malang 24 -25 Juli 2009.
- Nukmal, N., Utami, N., dan Suprpto. 2010. Skrining Potensi Daun Gamal (*Gliricidia maculata* Hbr.) Sebagai Insektisida Nabati. *Laporan Penelitian Hibah Strategi Unila*. Universitas Lampung. 2010.
- 2 Prijono, D. 2005. Pemanfaatan dan Pengembangan Pestisida Nabati. *Makalah Seminar Ilmiah*. Jurusan Proteksi Tanaman, Fakultas Pertanian, Universitas Lampung. 3 Agustus 2005.
- Raini, M. 2007. Toksikologi Pestisida dan Penanganan Akibat Keracunan Pestisida. *Media Litbang kesehatan Vol. XVII. No. 3*. Departemen Kesehatan. Jakarta.
- 16 Soebagio. 2002. *Kimia Analitik*. Universitas Negeri Makassar Fakultas MIPA. Makassar.
- 21 Sumarno, E. 2015. *Jenis - Jenis Serangga Hama Berdasarkan Tingkat Kerusakan Yang DiTimbulkan*. Universitas Halu Oleo. Kendari.
- 4 Tapas, AR., Sakarkar, DM., Kakde RB., 2008. *Flavonoids as Nutraceuticals*. *Tropical Journal of Pharmaceutical Research*(3): 1089-1099. Faculty of Pharmacy, University of Benin-Nigeria.
- 2 Wijaya, S. Y. 2007. *Kolonisasi Semut Hitam (*Dolichoderus thoracicus smith*) Pada Tanaman Kakao (*Theobroma cacao L.*) Dengan Pemberian Pakan Alternatif*. Skripsi. Universitas Sebelas Maret. Surakarta.
- 19 Yazid, Estien. 2005. *Kimia Fisika untuk Paramedis*. Andi: Yogyakarta.

THE TOXICITY OF PURIFIED ISOLATE OF POLAR EXTRACT POWDER LEAFS GLIRICIDIA MACULATA HBR. TO CACAO MEALYBUG (PLANOCOCCUS MINOR MASKELL)

ORIGINALITY REPORT

11%

SIMILARITY INDEX

PRIMARY SOURCES

1	repository.lppm.unila.ac.id Internet	267 words — 4%
2	digilib.unila.ac.id Internet	133 words — 2%
3	khoathuongmaidulich.edu.vn Internet	60 words — 1%
4	docobook.com Internet	21 words — < 1%
5	www.coursehero.com Internet	21 words — < 1%
6	Laetitia de Jong, Nicolas Pech, Gisela de Aragão Umbuzeiro, Xavier Moreau. "Multi-scale biomarker evaluation of the toxicity of a commercial azo dye (Disperse Red 1) in an animal model, the freshwater cnidarian <i>Hydra attenuata</i> ", <i>Water Research</i> , 2016 Crossref	21 words — < 1%
7	jurnal.fp.unila.ac.id Internet	20 words — < 1%
8	www.nparks.gov.sg Internet	20 words — < 1%

9	www.unila.ac.id Internet	18 words — < 1%
10	Kuo-Kuei Huang, Meng-Nan Lin, Yi-Ling Hsu, I-Huang Lu, I-Hong Pan, Jia-Ling Yang. "Fruit Extract Ameliorates Experimental Autoimmune Encephalomyelitis through the Regulation of Th1/Th17 Cells ", Evidence-Based Complementary and Alternative Medicine, 2019 Crossref	16 words — < 1%
11	www.bis.org Internet	15 words — < 1%
12	isomass.unila.ac.id Internet	15 words — < 1%
13	Liu, Jinxin, Jingyi Hou, Chao Jiang, Geng Li, Heng Lu, Fanyun Meng, and Linchun Shi. "Deep Sequencing of the Scutellaria baicalensis Georgi Transcriptome Reveals Flavonoid Biosynthetic Profiling and Organ-Specific Gene Expression", PLoS ONE, 2015. Crossref	14 words — < 1%
14	developing8.org Internet	13 words — < 1%
15	vdocuments.mx Internet	11 words — < 1%
16	www.scribd.com Internet	10 words — < 1%
17	Yanbo Huang. "Early Detection of Soybean Plant Injury from Glyphosate by Measuring Chlorophyll Reflectance and Fluorescence", Journal of Agricultural Science, 03/31/2012 Crossref	10 words — < 1%
18	library.ukdw.ac.id Internet	10 words — < 1%

19	hudhanuer10.blogspot.com Internet	9 words — < 1%
20	Chih-Min Liu, Shu-Sen Chang, Shih-Cheng Liao, Tzung-Jeng Hwang, Ming-Hsien Shieh, Shi-Kai Liu, Wei J. Chen, Hai-Gwo Hwu. "Absent response to niacin skin patch is specific to schizophrenia and independent of smoking", <i>Psychiatry Research</i> , 2007 Crossref	9 words — < 1%
21	edisumarnoblog.blogspot.com Internet	8 words — < 1%
22	www.sciencepublishinggroup.com Internet	8 words — < 1%
23	www.bi.go.id Internet	8 words — < 1%

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE MATCHES OFF