

# The Moment Generating Function of the FourParameter Generalized F Distribution and Related Generalized Distributions

*By* Dian Kurniasari

## The Moment Generating Function of the Four-Parameter Generalized F Distribution and Related Generalized Distributions

**Warsono<sup>1</sup>, Dian Kurniasari<sup>1</sup>, Widiarti<sup>1</sup>, Mustofa Usman<sup>1</sup> and Faiz A. M. Elfaki<sup>2</sup>**

<sup>1</sup>*Department of Mathematics, University of <sup>13</sup>mpung, Indonesia.*

<sup>2</sup>*Department of Mathematics, Statistics and Physics, College of Arts and Sciences,  
 Qatar University, P. O. Box 2713 Doha, Qatar.*

### Abstract

7

The main objective of this article is to show the derivation of the moment generating function of the four-parameter of generalized (G4F) F distribution. Through parameterization of its moment generating function, the behavior in relation to several well-known generalized distributions is presented. By utilizing MacLaurin series expansion and Stirling formula, it is shown that with parameterization of its moment generating function, the generalized F distribution might have special relationship to several well-known generalized distributions, such as generalized beta of the second kind (GB2), generalized log-logistic (G4LL), and generalized gamma (G3G) distributions.

11

**Keywords:** Moment generating function; Generalized beta of the second kind distribution; Generalized log-logistic distribution; Generalized gamma; MacLaurin series; Stirling formula

### 1. INTRODUCTION

As stated by several authors, the generalized F is one of the most well-known generalized distributions in probability modeling. In fitting survival data of carcinoma patients, Ciampi, et al. (1986) discussed the properties and maximum likelihood

5

inference of the family of the generalized F distribution. Peng, et al. (1998) investigated the application of the generalized F distribution to a mixture model from lymphoma patients. Kalbfleisch and Prentice (2002) discussed the use of the generalized F distribution.

The generalized F distribution is parameterization of other well-known generalized distributions, such as the generalized beta and generalized log-logistic distributions (Pham-Ghia and Duong 1989, Singh 1989). According to Ciampi, et al. (1986) when a random variable X has four-parameter generalized F distribution with degrees of freedom  $2m_1 + 2m_2$ , as denoted by  $X \sim G4F(\mu, \sigma, m_1, m_2)$  or  $G4F_3(m_1, m_2)$ , the corresponding probability density function (PDF) of the  $G4F(\mu, \sigma, m_1, m_2)$  can be written in the following form;

$$f_{G4F}(x) = \frac{e^{-\frac{\mu x}{\sigma}} x^{\left(\frac{m_1}{\sigma}\right)-1} \left(\frac{35}{m_2}\right)^{m_1} \left[1 + \left(\frac{m_1}{m_2}\right)(e^{-\mu} x)^{\frac{1}{\sigma}}\right]^{-(m_1+m_2)}}{\sigma \frac{\Gamma(m_1) \cdot \Gamma(m_2)}{\Gamma(m_1 + m_2)}} ; x \geq 0 \quad (1)$$

where  $\mu > 0$ ,  $\sigma > 0$ ,  $m_1 > 0$ ,  $m_2 > 0$ , and  $\Gamma$  is gamma function.

47

It can be shown that if we let  $\mu = \ln b + (1/a) \ln(m_1/m_2)$ ,  $\sigma = 1/a$ , and  $B(m_1, m_2)$  that is beta function, then the PDF of the G4F distribution becomes

$$f_{GB2}(x) = \frac{a}{xB(m_1, m_2)} \left[ \left( \frac{x}{b} \right)^a \right]^{m_1} \left[ 1 + \left( \frac{x}{b} \right)^a \right]^{m_1+m_2} \quad (34)$$

which is the PDF of the GB2 distribution as deliberated by McDonald (1984), and McDonald and Richards (1987).

33

If we let  $\mu = -\beta - \ln(m_1/m_2)(1/\alpha)$  and  $\sigma = 1/\alpha$ , where and, then the PDF of the G4F distribution can be written in the form

$$f_{G4LL}(x) = \frac{\alpha}{xB(m_1, m_2)} \left[ \frac{1}{1 + e^{-(\beta + \alpha \ln(x))}} \right]^{m_1} \left[ 1 - \frac{1}{1 + e^{-(\beta + \alpha \ln(x))}} \right]^{m_2}$$

which is the PDF of the G4LL distribution considered by Singh, et al.(1997).

32

Moreover, as noted by several authors (Ciampi, et al. 1986; Peng et al. 1998; 5 albfleisch and Prentice 2002; and Zhou et al. 2011), the generalized F contains several commonly used distributions as special cases or limiting distributions, such as exponential, gamma, lognormal<sup>14</sup> and Weibull distributions. Recently, Cox (2008) has linked the G4F distribution to the three-parameter generalized (G3G) and the three-parameter generalized log-logistic (G3LL) distribution<sup>43</sup>. To link the G4F distribution to other generalized distributions, he has parameterized the probability density function (PDF) of the G4F distribution.<sup>20</sup>

However, in term of a moment generating function (MGF), flexibilities of the G4F distribution and relation to other generalized distributions are apparently not too readily accessible in all of above literatures. Based on the MGF, an alternative method could be generated for computing moments and characterizing distributions from their MGFs. Based on MGFs, Warsono (2009) demonstrated the relationship between GB2 and G3G distributions, and Warsono (2010) linked the G4LL distribution to the GB2 and G3G distribution. In developing other aspects of the G4F distribution, it seems worth<sup>27</sup> while to extend all of above endeavors and to record them in this paper. Therefore, the main objective of this paper is to derive the MGF of the generalized F distribution and relate other generalized distributions, such as generalized beta, generalized log-logistic, and generalized gamma.<sup>24</sup>

7

The rest of the article is outlined as follows. In Section 2, the explicit moment generating function of the G4F distribution is derived. By using parameterization of the moment generating function of the G4F, Section 3 provides discussions of the relation between G4F distribution and G<sup>7</sup>2 and G4LL distributions. Section 4 contains a description of limiting behavior<sup>49</sup> of the moment generating function of the G4F distribution as a general case of the moment generating function of G3G and G3F distributions. Finally, some concluding remarks are noted in section 5.

## 2. MOMENT GENERATING FUNCTION OF THE GENERALIZED F DISTRIBUTION

42

In term of a beta function the pdf of GF distribution in equation (1) can be rewritten as:

$$g(x) = \frac{1}{\alpha \Gamma(m_1, m_2)} \frac{\left( e^{-\frac{\mu}{\sigma}} \left( \frac{m_1}{m_2} \right) x^{\frac{1}{\sigma}} \right)^{m_1}}{\left[ 1 + \left( e^{-\frac{\mu}{\sigma}} \right) x^{\frac{1}{\sigma}} \left( \frac{m_1}{m_2} \right) \right]^{m_1+m_2}} ; \quad x \geq 0 \quad (2)$$

where  $\mu > 0$ ,  $\sigma > 0$ ,  $m_1 > 0$ , and  $\Gamma$  is gamma function.

2

**Theorem 2.1** Let  $X$  be a random variable of the G4F( $\mu, \sigma, m_1, m_2$ ) distribution, then moment generating function (MGF) of  $X$  is given by

$$M_{G4F}(t) = e^{tx}$$

**Proof:**

$$\begin{aligned} M_{G4F}(t) &= E(e^{tx}) = \int_{-\infty}^{\infty} e^{tx} \cdot f(x) dx \\ &= \int_0^{\infty} e^{tx} \left( \frac{1}{\sigma x B(m_1, m_2)} \right) \frac{e^{-\frac{\mu m_1}{\sigma}} \left( \frac{m_1}{m_2} \right)^{m_1} x^{\frac{m_1}{\sigma}}}{\left[ 1 + (e^{-\mu} x)^{\frac{1}{\sigma}} \frac{m_1}{m_2} \right]^{m_1+m_2}} dx \end{aligned}$$

By algebra manipulation we may find the following equation

$$= \frac{1}{\sigma B(m_1, m_2)} \int_0^{\infty} \left( \frac{e^{tx}}{x} \right) \frac{\left[ e^{-\frac{\mu}{\sigma}} \left( \frac{m_1}{m_2} \right)^{\frac{1}{\sigma}} x^{\frac{1}{\sigma}} \right]^{m_1}}{\left[ 1 + e^{-\frac{\mu}{\sigma}} \left( \frac{m_1}{m_2} \right)^{\frac{1}{\sigma}} x^{\frac{1}{\sigma}} \right]^{m_1+m_2}} dx \quad (3)$$

By letting  $y = e^{-\frac{\mu}{\sigma}} \left( \frac{m_1}{m_2} \right)^{\frac{1}{\sigma}} x^{\frac{1}{\sigma}}$  we may rewrite the equation (3) in the following form

$$\begin{aligned} M_{G4F}(t) &= \frac{1}{\sigma B(m_1, m_2)} \int_0^{\infty} \left( \frac{\sigma e^{tx}}{y} \right) \frac{y^{m_1}}{(1+y)^{m_1+m_2}} dy \\ &= \frac{1}{B(m_1, m_2)} \int_0^{\infty} e^{tx} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy \end{aligned} \quad (4)$$

Making use of a well-known property of MacLaurin series of the  $e^{tx}$  function, then equation (4) is given by

$$\begin{aligned} M_{G4F}(t) &= \frac{1}{B(m_1, m_2)} \int_0^{\infty} \left( \sum_{n=0}^{\infty} \frac{(tx)^n}{n!} \right) \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy \\ &= \frac{1}{B(m_1, m_2)} \int_0^{\infty} \left( 1 + tx + \frac{(tx)^2}{2!} + \frac{(tx)^3}{3!} + \dots \right) \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{B(m_1, m_2)} \int_0^\infty \left( \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + tx \frac{6}{(1+y)^{m_1+m_2}} + \frac{(tx)^2}{2!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + \frac{(tx)^3}{3!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + \dots \right) dy \\
&= \frac{31}{B(m_1, m_2)} \left[ \int_0^\infty \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \int_0^\infty tx \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \right. \\
&\quad \left. \int_0^\infty \frac{(tx)^2}{2!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \int_0^\infty \frac{(tx)^3}{3!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \dots \right] \\
&= \frac{1}{B(m_1, m_2)} \left[ \int_0^\infty \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \int_0^\infty y^\sigma \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \right. \\
&\quad \left. \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \right)^2}{2!} \int_0^\infty y^{2\sigma} \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \right)^3}{3!} \int_0^\infty y^{3\sigma} \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \dots \right] \\
&= \frac{1}{B(m_1, m_2)} \left[ \int_0^\infty \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \int_0^\infty \frac{y^{m_1+\sigma-1}}{(1+y)^{m_1+\sigma+m_2-\sigma}} dy + \right. \\
&\quad \left. \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \right)^2}{2!} \int_0^\infty \frac{y^{41_{\sigma-1}}}{(1+y)^{m_1+2\sigma+m_2-2\sigma}} dy + \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \right)^3}{3!} \int_0^\infty \frac{y^{m_1+3\sigma-1}}{(1+y)^{m_1+3\sigma+m_2-3\sigma}} dy + \dots \right] \\
&\stackrel{30}{=} \frac{1}{B(m_1, m_2)} \left[ B(m_1, m_2) + t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \cdot B(m_1 + \sigma, m_2 - \sigma) + \right. \\
&\quad \left. \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \right)^2}{2!} \cdot B(m_1 + 2\sigma, m_2 - 2\sigma) + \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^\sigma \right)^3}{3!} \cdot B(m_1 + 3\sigma, m_2 - 3\sigma) + \dots \right]
\end{aligned}$$

$$\begin{aligned}
& \stackrel{18}{=} 1 + t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^{\sigma} \cdot \frac{B(m_1 + \sigma, m_2 - \sigma)}{B(m_1, m_2)} + \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^{\sigma} \right)^2}{2!} \cdot \frac{B(m_1 + 2\sigma, m_2 - 2\sigma)}{B(m_1, m_2)} + \\
& \quad \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^{\sigma} \right)^3}{3!} \cdot \frac{B(m_1 + 3\sigma, m_2 - 3\sigma)}{B(m_1, m_2)} + \dots \\
& = \sum_{n=0}^{\infty} \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^{\sigma} \right)^n}{n!} \cdot \frac{B(m_1 + n\sigma, m_2 - n\sigma)}{B(m_1, m_2)} \\
& = \sum_{n=0}^{\infty} \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^{\sigma} \right)^n}{n!} \cdot \frac{\frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1 + n\sigma + m_2 - n\sigma)}}{\frac{\Gamma(m_1) \cdot \Gamma(m_2)}{\Gamma(m_1 + m_2)}} \\
& \stackrel{40}{=} \sum_{n=0}^{\infty} \frac{\left( t \left( e^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right) \right)^{\sigma} \right)^n}{n!} \cdot \frac{\frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1 + m_2)}}{\frac{\Gamma(m_1) \cdot \Gamma(m_2)}{\Gamma(m_1 + m_2)}}
\end{aligned}$$

Therefore, the MGF of the G4F distribution is

$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left( te^{\frac{\mu}{\sigma}} \left( \frac{m_2}{m_1} \right)^{\sigma} \right)^n}{n!} \cdot \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)} \quad (5)$$

It is well-known that the Stirling's approximation formula of the gamma function (Spiegel 1968) is given by

$$\Gamma(az + b) \sim \sqrt{2\pi} \cdot e^{-az} (az)^{az+b-\frac{1}{2}}$$

By Stirling's approximation, equation (5) can be expressed as

$$\begin{aligned}
 M_{G4F}(t) &= \sum_{n=0}^{\infty} \frac{\left(te^{\mu}\left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \cdot \frac{\sqrt{2\pi}e^{-m_1}(m_1)^{m_1+n\sigma-\frac{1}{2}} \cdot \sqrt{2\pi}e^{-m_2}(m_2)^{m_2+n\sigma-\frac{1}{2}}}{\sqrt{2\pi}e^{-m_1}(m_1)^{m_1-\frac{1}{2}} \cdot \sqrt{2\pi}e^{-m_2}(m_2)^{m_2-\frac{1}{2}}} \quad (29) \\
 M_{G4F}(t) &= \sum_{n=0}^{\infty} \frac{\left(te^{\mu}\left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \cdot \left(m_1^{\sigma}\right)^n \left(\left(\frac{1}{m_2}\right)^{\sigma}\right)^n \\
 M_{G4F}(t) &= \sum_{n=0}^{\infty} \frac{\left(te^{\mu}\left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \cdot \left(\left(\frac{m_1}{m_2}\right)^{\sigma}\right)^n \quad (6)
 \end{aligned}$$

By MacLaurin series, equation (6) can be written as

$$\begin{aligned}
 M_{G4F}(t) &= e^{te^{\mu}\left(\frac{m_2}{m_1}\right)^{\sigma}\left(\frac{m_1}{m_2}\right)^{\sigma}} \\
 M_{G4F}(t) &= e^{te^{\mu}}
 \end{aligned}$$

## 2.2. Moment Generating Function (MGF) of Three-Parameter G3F Distribution

Based on the gamma distribution (Malik 1967, and Dyer 1982) and generalized beta with three-parameter distribution (Pham-Ghia and Duong 1989) obtained three-parameters of generalized F distribution with the following pdf

$$f_{G3F}(x) = \frac{\alpha^m}{B(m_1, m_2)} \frac{x^{m_1-1}}{[1+\alpha x]^{m_1+m_2}} dx; x > 0, m_1, m_2, \alpha > 0 \quad (39)$$

38

**Theorem 2.2** Let  $X$  be a random variable of the  $G3F(\alpha, m_1, m_2)$  distribution, then moment generating function (MGF) of  $X$  is given by

$$M_{G3F}(t) = \sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^n}{n!} \cdot \frac{\Gamma(m_1+n) \cdot \Gamma(m_2-n)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$

**Proof:**

$$\begin{aligned} M_{G_{3F}}(t) &= E(e^{tx}) = \int_{-\infty}^{\infty} e^{tx} \cdot f(x) dx \\ &= \int_0^{\infty} e^{tx} \left( \frac{\alpha^{m_1}}{B(m_1, m_2)} \right) \frac{x^{m_1-1}}{(1+\alpha x)^{m_1+m_2}} dx \end{aligned}$$

By algebra manipulation we may find the following equation

$$= \frac{\alpha^{m_1}}{B(m_1, m_2)} \int_0^{\infty} (e^{tx}) \frac{x^{m_1-1}}{(1+\alpha x)^{m_1+m_2}} dx \quad (7)$$

By letting  $y = \alpha x$  we may rewrite the equation (7) in the following form

$$\begin{aligned} M_x(t) &= \frac{\alpha^{m_1}}{B(m_1, m_2)} \int_0^{\infty} (e^{tx}) \frac{\left(\frac{y}{\alpha}\right)^{m_1-1}}{(1+y)^{m_1+m_2}} \frac{1}{\alpha} dy \\ &= \frac{1}{B(m_1, m_2)} \int_0^{\infty} (e^{tx}) \frac{(y)^{m_1-1}}{(1+y)^{m_1+m_2}} dy \end{aligned} \quad (8)$$

Making use [15](#) a well-known property of MacLaurin series of the  $e^{tx}$  function, then equation (8) can be written as

$$\begin{aligned} M_x(t) &= \frac{1}{B(m_1, m_2)} \int_0^{\infty} \left( \sum_{n=0}^{\infty} \frac{(tx)^n}{n!} \right) \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy \\ &= \frac{1}{B(m_1, m_2)} \int_0^{\infty} \left( 1 + tx + \frac{(tx)^2}{2!} + \frac{(tx)^3}{3!} + \dots \right) \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy \\ &= \frac{1}{B(m_1, m_2)} \int_0^{\infty} \left( \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + tx \frac{6}{(1+y)^{m_1+m_2}} + \frac{(tx)^2}{2!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + \frac{(tx)^3}{3!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + \dots \right) dy \\ &= \frac{1}{B(m_1, m_2)} \left| \int_0^{\infty} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \int_0^{\infty} tx \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \right. \\ &\quad \left. \int_0^{\infty} \frac{(tx)^2}{2!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \int_0^{\infty} \frac{(tx)^3}{3!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \dots \right| \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{B(m_1, m_2)} \left[ \int_0^{\infty} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left( \frac{1}{\alpha} \right) \int_0^{\infty} y \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \right. \\
&\quad \left. \frac{\left( t \left( \frac{1}{\alpha} \right) \right)^2}{2!} \int_0^{\infty} y^2 \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \frac{\left( t \left( \frac{1}{\alpha} \right) \right)^3}{3!} \int_0^{\infty} y^3 \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \dots \right] \\
&= \frac{1}{B(m_1, m_2)} \left[ \int_0^{\infty} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \left( \frac{t}{\alpha} \right) \int_0^{\infty} \frac{y^{m_1+1-1}}{(1+y)^{m_1+1+m_2-1}} dy + \right. \\
&\quad \left. \frac{\left( \frac{t}{\alpha} \right)^2}{2!} \int_0^{\infty} \frac{y^{m_1+2-1}}{(1+y)^{m_1+2+m_2-2}} dy + \frac{\left( \frac{t}{\alpha} \right)^3}{3!} \int_0^{\infty} \frac{y^{m_1+3-1}}{(1+y)^{m_1+3+m_2-3}} dy + \dots \right] \\
&= \frac{10}{B(m_1, m_2)} \left[ B(m_1, m_2) + \left( \frac{t}{\alpha} \right) \cdot B(m_1+1, m_2-1) + \right. \\
&\quad \left. \frac{\left( \frac{t}{\alpha} \right)^2}{2!} \cdot B(m_1+2, m_2-2) + \frac{\left( \frac{t}{\alpha} \right)^3}{3!} \cdot B(m_1+3, m_2-3) + \dots \right] \\
&= 1 + \frac{t}{\alpha} \cdot \frac{B(m_1+1, m_2-1)}{B(m_1, m_2)} + \frac{\left( \frac{t}{\alpha} \right)^2}{2!} \cdot \frac{B(m_1+2, m_2-2)}{B(m_1, m_2)} + \frac{\left( \frac{t}{\alpha} \right)^3}{3!} \cdot \frac{B(m_1+3, m_2-3)}{B(m_1, m_2)} + \dots \\
&= \sum_{n=0}^{\infty} \frac{\left( \frac{t}{\alpha} \right)^n}{n!} \cdot \frac{B(m_1+n, m_2-n)}{B(m_1, m_2)} \\
&\equiv \sum_{n=0}^{\infty} \frac{\left( \frac{t}{\alpha} \right)^n}{n!} \cdot \frac{\frac{\Gamma(m_1+n) \cdot \Gamma(m_2-n)}{\Gamma(m_1+n+m_2-n)}}{\frac{\Gamma(m_1) \cdot \Gamma(m_2)}{\Gamma(m_1+m_2)}}
\end{aligned}$$

$$\begin{aligned}
 &= \sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^n}{n!} \cdot \frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1 + m_2)} \cdot \frac{\Gamma(m_1 + m_2)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
 &= \sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^n}{n!} \cdot \frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1) \cdot \Gamma(m_2)}
 \end{aligned}$$

Therefore, the MGF of the G3F distribution is

$$M_X(t) = \sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^n}{n!} \cdot \frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1) \cdot \Gamma(m_2)} \quad (9)$$

### 3. THE RELATION OF THE GF DISTRIBUTION WITH GB2 DISTRIBUTION

Warsono (2010) has mathematically derived the MGF of the GB2 distribution using MGF defintion. Based on reparameterization of the MGF of the GLF distribution, use passive sentence- The MGF of the GB2 distribution is provided in this section. The reparameterization proposition is stated and proved.

**Proposition 3.1** Let  $X$  be a random variable having the G4F  $(\mu, \sigma, m_1, m_2)$  moment and

$\mu = \ln b + \frac{1}{a} \ln \left( \frac{m_1}{m_2} \right)$  and  $\sigma = \frac{1}{a}$ , then  $X$  has the GB2( $a, b, m_1, m_2$ ) moment.

**Proof:**

$$\begin{aligned}
 M_{G4F}(t) &= \sum_{n=0}^{\infty} \frac{\left( t e^{\mu} \left( \frac{m_2}{m_1} \right)^{\sigma} \right)^n}{n!} \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
 &= \sum_{n=0}^{\infty} \frac{\left( t e^{\ln b + \frac{1}{a} \ln \left( \frac{m_1}{m_2} \right)} \left( \frac{m_2}{m_1} \right)^{\frac{1}{a}} \right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{a}\right) \cdot \Gamma\left(m_2 - \frac{n}{a}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)}
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \frac{\left( t e^{ln b} e^{\frac{1}{a} \ln \left( \frac{m_1}{m_2} \right)} \left( \frac{m_2}{m_1} \right)^{\frac{1}{a}} \right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{a}\right) \cdot \Gamma\left(m_2 - \frac{n}{a}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&= \sum_{n=0}^{\infty} \frac{\left( tb \left( \frac{m_1}{m_2} \right)^{\frac{1}{a}} \left( \frac{m_2}{m_1} \right)^{\frac{1}{a}} \right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{a}\right) \cdot \Gamma\left(m_2 - \frac{n}{a}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&= \sum_{n=0}^{\infty} \frac{(tb)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{a}\right) \cdot \Gamma\left(m_2 - \frac{n}{a}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)}
\end{aligned}$$

**1**

This is the moment generating function of the GB2 ( $a, b, m_1, m_2$ ) stated by Warsono (2010).

#### 4. THE RELATION OF THE GF DISTRIBUTION WITH GLL DISTRIBUTION

Warsono (2010) derived mathematically the MGF of the GLL distribution using definition of MGF's. Based on reparameterization of the MGF of the GF distribution, the MGF of the GLL distribution is provided in this section.

**2**

**Proposition 4.1** Let  $X$  be a random variable having the GF  $(\mu, \sigma, m_1, m_2)$  moment and

$$\mu = \left( -\beta - \ln \left( \frac{m_2}{m_1} \right) \right) \frac{1}{\alpha} \text{ and } \sigma = \frac{1}{\alpha}, \text{ then } X \text{ has the GLL}(\alpha, \beta, m_1, m_2) \text{ moment.}$$

**Proof:**

$$\begin{aligned}
M_{G4F}(t) &= \sum_{n=0}^{\infty} \frac{\left( t e^{\mu} \left( \frac{m_2}{m_1} \right)^{\sigma} \right)^n}{n!} \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&= \sum_{n=0}^{\infty} \frac{\left( t e^{\left( -\beta - \ln \left( \frac{m_2}{m_1} \right) \right) \frac{1}{\alpha}} \left( \frac{m_2}{m_1} \right)^{\frac{1}{\alpha}} \right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{\alpha}\right) \cdot \Gamma\left(m_2 - \frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)}
\end{aligned}$$

45

$$\begin{aligned}
 & \stackrel{12}{=} \sum_{n=0}^{\infty} \frac{\left( t e^{-\frac{\beta}{\alpha} - \frac{1}{\alpha} \ln\left(\frac{m_2}{m_1}\right)} \left(\frac{m_2}{m_1}\right)^{\frac{1}{\alpha}} \right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{\alpha}\right) \cdot \Gamma\left(m_2 - \frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
 & \stackrel{1}{=} \sum_{n=0}^{\infty} \frac{\left( t e^{-\frac{\beta}{\alpha}} e^{-\frac{1}{\alpha} \ln\left(\frac{m_2}{m_1}\right)} \left(\frac{m_2}{m_1}\right)^{\frac{1}{\alpha}} \right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{\alpha}\right) \cdot \Gamma\left(m_2 - \frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
 & \stackrel{48}{=} \sum_{n=0}^{\infty} \frac{\left( t e^{-\frac{\beta}{\alpha}} \left(\frac{m_2}{m_1}\right)^{\frac{1}{\alpha}} \left(\frac{m_2}{m_1}\right)^{\frac{1}{\alpha}} \right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{\alpha}\right) \cdot \Gamma\left(m_2 - \frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
 & \stackrel{3}{=} \sum_{n=0}^{\infty} \frac{\left( t e^{-\frac{\beta}{\alpha}} \right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{\alpha}\right) \cdot \Gamma\left(m_2 - \frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)}
 \end{aligned}$$

## 5. THE RELATION OF THE G4F DISTRIBUTION WITH G3F DISTRIBUTION

Equation (8) has mathematically derived the MGF of the G3F distribution. Based on reparameterization of the MGF of the GF distribution, the MGF of the G3F distribution is provided in this section.

<sup>2</sup>

**Proposition 5.1** Let  $X$  be a random variable having the G4F  $(\mu, \sigma, m_1, m_2)$  moment and

$$\mu = \left( -\ln\left(\frac{m_2}{m_1}\right) - \ln\alpha \right) \text{ and } \sigma = 1, \text{ then } X \text{ has the G3F}(\alpha, m_1, m_2) \text{ moment.}$$

**Proof:**

$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left( t e^{\mu} \left(\frac{m_2}{m_1}\right)^{\sigma} \right)^n}{n!} \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \frac{\left( t e^{\left( -\ln\left(\frac{m_2}{m_1}\right) - \ln\alpha \right)} \left( \frac{m_2}{m_1} \right)^n \right)^n}{n!} \frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&\stackrel{1}{=} \sum_{n=0}^{\infty} \frac{\left( t e^{-\ln\left(\frac{m_1}{m_2}\right)} e^{-\ln\alpha} \left( \frac{m_2}{m_1} \right)^n \right)^n}{n!} \frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&= \sum_{n=0}^{\infty} \frac{\left( t e^{\frac{\ln\frac{1}{\alpha}}{\alpha} \left( \frac{m_2}{m_1} \right)^{-1}} \left( \frac{m_2}{m_1} \right)^n \right)^n}{n!} \frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&\stackrel{25}{=} \sum_{n=0}^{\infty} \frac{\left( \frac{t}{\alpha} \right)^n}{n!} \frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1) \cdot \Gamma(m_2)}
\end{aligned}$$

1 This is the moment generating function of the G3F( $\alpha$ ,  $m_1$ ,  $m_2$ ) stated in equation (9).

## 6. THE MOMENT OF THE GG AS A LIMITING MOMENT OF THE GF DISTRIBUTIONS

In this section, propositions of limiting moment properties of the GF distribution are stated and proved.

**Proposition 6.1** The GF( $\mu, \sigma, m_1, m_2$ ) distribution converges to the GG distribution as  $m_2$  tends to  $\infty$  and  $\mu = \ln\gamma + \frac{1}{a} \ln(m_1)$ , and  $\sigma = \frac{1}{a}$ .

**Proof:**

$$\lim_{m_2 \rightarrow \infty} M_{G4F}(t) = \lim_{m_2 \rightarrow \infty} \sum_{n=0}^{\infty} \frac{\left( t e^{\mu} \left( \frac{m_2}{m_1} \right)^{\sigma} \right)^n}{n!} \cdot \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$

$$\begin{aligned}
&= \lim_{m_2 \rightarrow \infty} \sum_{n=0}^{\infty} \frac{te^{\ln \gamma + \frac{1}{a} \ln(m_1)} \left( \frac{m_2}{m_1} \right)^{\frac{1}{a}}}{n!} \cdot \frac{\Gamma(m_1 + \frac{n}{a}) \cdot \Gamma(m_2 - \frac{n}{a})}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&\stackrel{37}{=} \lim_{m_2 \rightarrow \infty} \sum_{n=0}^{\infty} \frac{te^{\ln \gamma} e^{\frac{1}{a} \ln(m_1)} \left( \frac{m_2}{m_1} \right)^{\frac{1}{a}}}{n!} \cdot \frac{\Gamma(m_1 + \frac{n}{a}) \cdot \Gamma(m_2 - \frac{n}{a})}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&\stackrel{36}{=} \lim_{m_2 \rightarrow \infty} \sum_{n=0}^{\infty} \frac{t \gamma(m_1)^{\frac{1}{a}} \left( \frac{m_2}{m_1} \right)^{\frac{1}{a}}}{n!} \cdot \frac{\Gamma(m_1 + \frac{n}{a}) \cdot \Gamma(m_2 - \frac{n}{a})}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&= \lim_{m_2 \rightarrow \infty} \sum_{n=0}^{\infty} \frac{\left( t \gamma(m_2)^{\frac{1}{a}} \right)^n}{n!} \cdot \frac{\Gamma(m_1 + \frac{n}{a}) \cdot \Gamma(m_2 - \frac{n}{a})}{\Gamma(m_1) \cdot \Gamma(m_2)} \\
&= \lim_{m_2 \rightarrow \infty} 1 + \lim_{m_2 \rightarrow \infty} \left( t \gamma(m_2)^{\frac{1}{a}} \right) \cdot \frac{\Gamma(m_1 + \frac{1}{a}) \cdot \Gamma(m_2 - \frac{1}{a})}{\Gamma(m_1) \cdot \Gamma(m_2)} + \\
&\quad \lim_{m_2 \rightarrow \infty} \frac{\left( t \gamma(m_2)^{\frac{1}{a}} \right)^2}{2!} \cdot \frac{\Gamma(m_1 + \frac{2}{a}) \cdot \Gamma(m_2 - \frac{2}{a})}{\Gamma(m_1) \cdot \Gamma(m_2)} + \\
&\quad \lim_{m_2 \rightarrow \infty} \frac{\left( t \gamma(m_2)^{\frac{1}{a}} \right)^3}{3!} \cdot \frac{\Gamma(m_1 + \frac{3}{a}) \cdot \Gamma(m_2 - \frac{3}{a})}{\Gamma(m_1) \cdot \Gamma(m_2)} + \dots \\
&= \lim_{m_2 \rightarrow \infty} 1 + (t \gamma) \cdot \frac{\Gamma(m_1 + \frac{1}{a})}{\Gamma(m_1)} \lim_{m_2 \rightarrow \infty} (m_2)^{\frac{1}{a}} \frac{\Gamma(m_2 - \frac{1}{a})}{\Gamma(m_2)} +
\end{aligned}$$

$$\frac{(t\gamma)^2}{2!} \cdot \frac{\Gamma(m_1 + \frac{2}{a})}{\Gamma(m_1)} \lim_{m_2 \rightarrow \infty} \cdot (m_2)^{\frac{2}{a}} \frac{\Gamma(m_2 - \frac{2}{a})}{\Gamma(m_2)} + \\ \frac{(t\gamma)^3}{3!} \cdot \frac{\Gamma(m_1 + \frac{3}{a})}{\Gamma(m_1)} \lim_{m_2 \rightarrow \infty} \cdot (m_2)^{\frac{3}{a}} \frac{\Gamma(m_2 - \frac{3}{a})}{\Gamma(m_2)} + \dots$$

Using Stirling's approximation formula of the gamma function, the limiting moment property of the G4F( $\mu, \sigma, m_1, m_2$ ) distribution can be written as:

$$\lim_{m_2 \rightarrow \infty} M_X(t) GF\left(\mu = \ln \gamma + \frac{1}{a} \ln(m_1), \sigma = \frac{1}{a}, m_1, m_2\right) \\ = \lim_{m_2 \rightarrow \infty} 1 + (t\gamma) \cdot \frac{\Gamma(m_1 + \frac{1}{a})}{\Gamma(m_1)} \lim_{m_2 \rightarrow \infty} (m_2)^{\frac{1}{a}} \cdot \frac{1}{(m_2)^{\frac{1}{a}}} + \frac{(t\gamma)^2}{2!} \cdot \frac{\Gamma(m_1 + \frac{2}{a})}{\Gamma(m_1)} \lim_{m_2 \rightarrow \infty} (m_2)^{\frac{2}{a}} \cdot \frac{1}{(m_2)^{\frac{2}{a}}} + \dots \\ = 1 + (t\gamma) \cdot \frac{\Gamma(m_1 + \frac{1}{a})}{\Gamma(m_1)} + \frac{(t\gamma)^2}{2!} \cdot \frac{\Gamma(m_1 + \frac{2}{a})}{\Gamma(m_1)} + \\ = \sum_{n=0}^{\infty} \frac{(t\gamma)^n}{n!} \cdot \frac{\Gamma(m_1 + \frac{n}{a})}{\Gamma(m_1)}$$

This result is the MGF of the GG stated by Warsono(2010). Thus, the G4F distribution converges to the GG distribution as  $m_2$  tends to  $\infty$  and  $\alpha = a$ , and  $\beta = -a \ln\left(\gamma(m_2)^{\frac{1}{a}}\right)$

## 7. CONCLUSION

The moment of the generalized F distribution is parameterization of the generalized beta of the second kind (GB2) and the generalized log-logistic (GLL). The moment of the generalized gamma (GG) distribution is the limiting moment of (GF) distribution. Moreover, since the moments of the gamma and exponential distributions are special cases of the moment of <sup>16</sup> generalized gamma distribution (Warsono 2009), the moments of both special distributions are also special cases of the moments of the moment of the GF distribution.

**REFERENCES**

- [1]. Ciampi, A., S.A. Hogg, and L. Kates. (1986). Regression analysis of censored survival data with the generalized F family – an alternative to the proportional hazards model. *Statistics in Medicine* Vol 5: 85-96.
- [2]. Cox, C. (2008). The generalized F distribution: An umbrella for parametric survival analysis. *Statistics in Medicine* 27: 4301-4312.
- [3]. Dyer, D. (1982). The convolution of generalized F distribution. *J. Am. Statist. Ass.* 77: 184-189.
- [4]. Kalbfleisch, J. D. and Prentice, R. L. (2002). *The Statistical Analysis of Failure Time Data*, Wiley, New York.
- [5]. Malik, H.J. (1967). Exact distributions of the quotient of independent generalized gamma variables. *Can. math. BUN.* **10**,463465.
- [6]. McDonald, J.B. (1984). Some generalized functions for the size distribution of income. *Econometrica* 52(3): 647-663.
- [7]. McDonald, J.B. and D.O. Richards. (1987). Hazard rates and generalized beta distributions. *IEEE Transactions reliability*, Vol. R-36: 463-466.
- [8]. Pham-Gia, T. And Q.P. Duong. (1989). The generalized beta- and F-distributions in statistical Modelling. *Math. Comput. Modelling*. Vol 12(12): 1613-1625.
- [9]. Peng, Y.P., K.B.G. Dear, and J.W. Denham. (1998). A generalized F mixture model for cure rate estimation. *Statistics in Medicine* 17: 813-830.
- [10]. Singh, K.P. (1989). A generalized log-logistic regression model for survival analysis: hazard rate characteristics. *Biometrie-Praximetrie* 29: 63-74
- [11]. Singh, K.P., Warsono, and A.A. Bartolucci. (1997). Generalized log-logistic model for analysis of environmental pollutant data. Proc. Int. Congress on Modeling and Simulation. Hobart, Tasmania. December 8-11 1997.
- [12]. Spiegel, M.R. (1968). Mathematical Handbook of Formulas and Tables. McGraw-Hill Inc.
- [13]. Warsono. (2009). Moment Properties of the Generalized Gamma Distribution. *Proceedings on Seminar Nasional Sains MIPA dan Aplikasinya*, Bandar Lampung: 157-162.
- [14]. Warsono. (2010). Remarks on Moment Properties of Generalized Distributions. *Proceedings of the Third International Conference on Mathematics and Natural Sciences (ICMNS 2010)*. Faculty of Mathematics and Natural Sciences Institut Teknologi Bandung. p. 1123 -1131.
- [15]. Xiaojing Zhou, X., L.Yan, D.R. Prows , and Runqing Yang. (2011). Generalized F accelerated failure time model for mapping survival trait loci. *Genomics* 97: 379-385.

# The Moment Generating Function of the FourParameter Generalized F Distribution and Related Generalized Distributions

---

ORIGINALITY REPORT

---

24%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

- |   |  |               |
|---|--|---------------|
| 1   | docslide.us<br>Internet  | 60 words — 2% |
| 2   | Stapleton. "Moment Generating Functions and Limit Theory", Models for Probability and Statistical Inference, 12/12/2007<br><small>Crossref</small>   | 54 words — 2% |
| 3   | math.ucsb.edu<br>Internet  | 52 words — 2% |
| 4   | Xu, S.p.. "Completeness of general solutions for three-dimensional transversely isotropic piezoelectricity", International Journal of Solids and Structures, 200809<br><small>Crossref</small> | 37 words — 1% |
| 5   | Yingwei Peng. "A generalized F mixture model for cure rate estimation", Statistics in Medicine, 04/30/1998<br><small>Crossref</small>  | 34 words — 1% |
| 6   | ALESSANDRO FILISETTI. "SYNCHRONIZATION PHENOMENA IN PROTOCELL MODELS", Biophysical Reviews and Letters, 2008<br><small>Crossref</small>  | 28 words — 1% |
| 7   | T. Pham-Gia, Q.P. Duong. "The generalized beta- and F-distributions in statistical modelling", Mathematical and Computer Modelling, 1989<br><small>Crossref</small>                            | 26 words — 1% |
| Huang Wen-Hua. "Doubly Periodic Propagating Wave for (2+1)- |  |               |

- 8 Dimensional Breaking Soliton Equation", Communications in Theoretical Physics, 02/2008 Crossref 22 words — 1%
- 9 www.ripublication.com Internet 22 words — 1%
- 10 etd.lsu.edu Internet 20 words — 1%
- 11 Cummins, J.. "Applications of the GB2 family of distributions in modeling insurance loss processes", Insurance Mathematics and Economics, 199012 Crossref 20 words — 1%
- 12 cseweb.ucsd.edu Internet 19 words — 1%
- 13 www.inderscience.com Internet 19 words — 1%
- 14 www.unc.edu Internet 17 words — 1%
- 15 M Sebawe Abdalla. "Effect of Kerr-like medium on a two-level atom in interaction with bimodal oscillators", Journal of Physics B Atomic Molecular and Optical Physics, 04/14/2006 Crossref 16 words — < 1%
- 16 Xiaojing Zhou, Li Yan, Daniel R. Prows, Runqing Yang. "Generalized F accelerated failure time model for mapping survival trait loci", Genomics, 2011 Crossref 13 words — < 1%
- 17 T. Clarke, S. J. Griffin, J. Ensor. "A polynomial approach to eigenstructure assignment using projection with eigenvalue trade-off", International Journal of Control, 2003 Crossref 13 words — < 1%
- 18 Farham. "The Concept of Event-Driven Time", 13 words — < 1%

- 
- 19 Robert N. Rodriguez. "Frequency Curves, Systems of", Encyclopedia of Statistical Sciences, 13 words — < 1%  
08/15/2006  
Crossref
- 
- 20 Mihajlo C. Stefanovic. "The Joint Probability Density Function of the SSC Combiner Output Signal in the Presence of Nakagami-m Fading", 2008 The Fourth International Conference on Wireless and Mobile Communications, 13 words — < 1%  
07/2008  
Crossref
- 
- 21 Saralees Nadarajah. "Multitude of beta distributions with applications", Statistics, 13 words — < 1%  
2007  
Crossref
- 
- 22 research.bus.wisc.edu 12 words — < 1%  
Internet
- 
- 23 www.dtic.mil 11 words — < 1%  
Internet
- 
- 24 hub.hku.hk 11 words — < 1%  
Internet
- 
- 25 www.mdpi.com 10 words — < 1%  
Internet
- 
- 26 Keisuke Okamura. "Higher loop Bethe ansatz for open spin-chains in AdS/CFT", Journal of High Energy Physics, 10 words — < 1%  
09/29/2006  
Crossref
- 
- 27 K.-H. Wang. "Profit Analysis of the Machine-repair Problem with a Single Service Station Subject to Breakdowns", Journal of the Operational Research Society, 10 words — < 1%  
2017  
Crossref
- 
- 28 Cristiano Calcagno. "Context logic as modal logic", ACM SIGPLAN Notices, 9 words — < 1%  
1/17/2007  
Crossref

- 29 Martin-Herran, G.. "A time-consistent open-loop Stackelberg equilibrium of shelf-space allocation", *Automatica*, 200506  
Crossref 9 words — < 1%
- 30 [www.restode.cfwb.be](http://www.restode.cfwb.be) Internet 9 words — < 1%
- 31 Jung, E.. "Bulk versus brane in the absorption and emission: 5d rotating black hole case", *Nuclear Physics, Section B*, 20051219  
Crossref 9 words — < 1%
- 32 Balakrishnan, N., and Suvra Pal. "An EM algorithm for the estimation of parameters of a flexible cure rate model with generalized gamma lifetime and model discrimination using likelihood- and information-based methods", *Computational Statistics*, 2015.  
Crossref 9 words — < 1%
- 33 [iptek.its.ac.id](http://iptek.its.ac.id) Internet 9 words — < 1%
- 34 [archive.numdam.org](http://archive.numdam.org) Internet 9 words — < 1%
- 35 Rassias, J.M.. "Landau's Type Inequalities", *Journal of Mathematical Analysis and Applications*, 19960815  
Crossref 9 words — < 1%
- 36 Doliwa, A.. "Geometric discretization of the Bianchi system", *Journal of Geometry and Physics*, 200411  
Crossref 8 words — < 1%
- 37 A. Krivov, A. Jurewicz. "The ethereal dust envelopes of the Martian moons", *Planetary and Space Science*, 1998  
Crossref 8 words — < 1%
- 38 [cran.r-project.org](http://cran.r-project.org) Internet 8 words — < 1%

- 39 [www.jvtc.jx.cn](http://www.jvtc.jx.cn)  
Internet 8 words — < 1%
- 40 Michel Enock. "QUANTUM GROUPOIDS OF COMPACT TYPE", Journal of the Institute of Mathematics of Jussieu, 01/2005 8 words — < 1%
- 41 van der Hofstad, R.. "The survival probability for critical spread-out oriented percolation above 4+1 dimensions. II. Expansion", Annales de l'Institut Henri Poincare / Probabilites et statistiques, 200709 8 words — < 1%
- 42 Heng Xing-Chen, Qin Zheng, Tian Lei, Shao Li-Ping. "Research on Structure Learning of Dynamic Bayesian Networks by Particle Swarm Optimization", 2007 IEEE Symposium on Artificial Life, 2007 8 words — < 1%
- 43 Zhenmin Chen. "Exact confidence interval for the shape parameter of a log-logistic distribution", Journal of Statistical Computation and Simulation, 2/1997 8 words — < 1%
- 44 [ccsenet.org](http://ccsenet.org)  
Internet 8 words — < 1%
- 45 Cereceda, J.L.. "Mermin's n-particle Bell inequality and operators' noncommutativity", Physics Letters A, 20010813 7 words — < 1%
- 46 Lou, Y.-Q., and Y. Wu. "Global perturbation configurations in a composite disc system with an isopedic magnetic field", Monthly Notices of the Royal Astronomical Society, 2005. 7 words — < 1%
- 47 Hunter, A.. "Fusion rules for merging uncertain information", Information Fusion, 200603 7 words — < 1%

- 
- 48 N. Balakrishnan, Erhard Cramer. "The Art of Progressive Censoring", Springer Nature, 2014  
Crossref 6 words — < 1%
- 49 Kleiber, C.. "The existence of population inequality measures", Economics Letters, 19971128  
Crossref 6 words — < 1%
- 50 Christopher A. Stone. "Extensional equivalence and singleton types", ACM Transactions on Computational Logic, 10/1/2006  
Crossref 6 words — < 1%
- 

EXCLUDE QUOTES

ON

EXCLUDE

BIBLIOGRAPHY

EXCLUDE MATCHES

OFF