



International Journal of Waste Resources

E-ISSN: 2252-5211

Open Access





 25+ Million Readerbase

Editorial Board


Useful Links

- > [Aims and Scope](#)
- > [Advertising](#)
- > [Citations Report](#)
- > [Indexing and Archiving](#)
- > [Table of Contents](#)
- > [Submit Paper](#)
- > [Track Your Paper](#)
- > [Funded Work](#)

Share This Page



Recommended Conferences

 June 22-23, 2020
Frontiers in Renewable Energy and Environmental Engineering
 New York, USA

 July 17-18, 2020
Current Trends in Recycling and E-Waste Management
 Vienna, Austria

Advertise HERE!



Budiyo
 Editor-in-Chief
 Department of Chemical Engineering
 University of Diponegoro, Indonesia

[Biography](#)

[Research Interest](#)



Elena Cristina Rada
 Editor-in-Chief
 Department of Civil Environmental and Mechanical Engineering
 University of Trento, Italy

[Biography](#)

[Research Interest](#)



Shoichiro Ozaki
 The Institute of Physical and Chemical Research
 Japan

[Biography](#)

[Research Interest](#)



Yan Luo
 Department of Chemical Engineering
 Halliburton West Virginia University, Dallas/Fort Worth Area, USA

[Biography](#)


[Research Interest](#)



Meine Pieter van Dijk
 Professor, UNESCO-IHE Institute for Water Education
 Erasmus University in Rotterdam, Netherlands

[Biography](#)

[Research Interest](#)



Baoyu Gao
 School of Environmental and Chemical Engineering
 East China University of Science and Technology, Shanghai, China

[Biography](#)


[Research Interest](#)

HERE!

25+ Million Website Visitors
9036+ Journal Visitors

Join Us

Journal Flyer



Tweets by @Carolin50507446

International Journal of Was
@Carolin50507446

Solid waste management: The sordid saga

Read more at:
longdom.org/scholarly/soli...

Embed View on Twitter



Baoyu Gao
School of Environmental Science and Engineering
Shandong University, China

[Biography](#)

[Research Interest](#)



Dimitris P Makris
Department of Food Science & Nutrition
University of the Aegean, Greece

[Biography](#)

[Research Interest](#)



Kaimin Shih
Department of Civil Engineering
Hong Kong University, Hong Kong

[Biography](#)

[Research Interest](#)



Kai Qin Xu
Professor, Center for Material Cycles and Waste Management Research, National Institute for Environmental Studies
Japan

[Biography](#)

[Research Interest](#)



Gabriela Ionescu
Department of Civil, Environmental and Mechanical Engineering
University of Trento, Italy

[Biography](#)

[Research Interest](#)



Margarida Marchetto
Department of Sanitary and Environmental Engineering
Federal University of Mato Grosso do Sul, Brazil

[Biography](#)

[Research Interest](#)



Vasudevan Subramanyan
Principal Scientist, Central Electrochemical Research Institute
India

[Biography](#)

[Research Interest](#)



Nigel Freestone
Head of the Department
The University of Northampton, United Kingdom

[Biography](#)

[Research Interest](#)



Tutuk Djoko Kusworo
Department of Chemical Engineering
University of Diponegoro, Indonesia

[Biography](#)

[Research Interest](#)



Moinuddin Sarker
Vice President, Natural State Research
USA

[Biography](#)

[Research Interest](#)



Moinuddin Sarker

Vice President, Natural State Research
USA

[Biography](#)

[Research Interest](#)



Abdullah M Busyairi

Professor, Department of Chemical Engineering
University of Diponegoro, Indonesia

[Biography](#)

[Research Interest](#)

Wenshan Guo

Wenshan Guo

Lecturer, School of Civil and Environmental Engineering
University of Technology Sydney, Australia

[Biography](#)

[Research Interest](#)



Mohd Fadhil Md Din

Associate Professor, Department of Environmental Engineering
Universiti Teknologi Malaysia, Malaysia

[Biography](#)

[Research Interest](#)



Atiq Uz Zaman

Zero Waste SA Research
University of South Australia, Australia

[Biography](#)

[Research Interest](#)



Gurpreet Singh Dhillon

Department of Agricultural, Food and Nutritional Sciences
University of Alberta, Canada

[Biography](#)

[Research Interest](#)



Nurul Fadly bin Habidin

Department of Management and Economics
Universiti Pendidikan Sultan Idris, Malaysia

[Research Interest](#)



I Nyoman Widiasta

Department of Chemical Engineering
Diponegoro University, Indonesia

[Biography](#)

[Research Interest](#)



Irvan Matseh



25+ Million Readerbase

Useful Links

- > Aims and Scope
- > Advertising
- > Citations Report
- > Indexing and Archiving
- > Table of Contents
- > Submit Paper
- > Track Your Paper
- > Funded Work

Share This Page



Recommended Conferences

- June 22-23, 2020
Frontiers in Renewable Energy and Environmental Engineering
New York, USA
- July 17-18, 2020
Current Trends in Recycling and E-Waste Management
Vienna, Austria

Advertise
HERE!

25+ Million Website Visitors

Join Us

Volume 4, Issue 2 (2014)

Editorial

Pages: 1 - 2

Increasing Trend towards Production of High Value Bioproducts from Biomass Feedstocks

Surinder Kaur, Gurpreet Singh Dhillon and Satinder Kaur Brar

DOI:10.4172/2252-5211.1000e105



Share this article



Research Article

Pages: 1 - 6

Comparison of Mercury Emissions in USA and China-The Way of Effective Control of Hg from the Power Plant

Haoyi Chen, Xiaolong WANG, Yu HUANG, Li Zhong, Shisen Xu and Tiancun Xiao

DOI:10.4172/2252-5211.1000138



Share this article



Research Article

Pages: 1 - 5

Estimation of Current and Future Generation of Medical Solid Wastes In Hanoi City, Vietnam

Duc Luong Nguyen, Xuan Thanh Bui and The Hung Nguyen

DOI:10.4172/2252-5211.1000139



Share this article



Editorial

Pages: 1 - 2

Roadmap towards Zero Waste Cities

Atiq Uz Zaman

DOI:10.4172/2252-5211.1000e106



Share this article



Research Article

Pages: 1 - 5

Transforming Municipal Waste into a Valuable Soil Conditioner through Knowledge-Based Resource-Recovery Management

Golabi MH, Kirk Johnson, Takeshi Fujiwara and Eri Ito

DOI:10.4172/2252-5211.1000140



Share this article



Research Article

Pages: 1 - 6

Not Too Little, Not Too Much and Shortcut: A Review on the Effectualness of Per Capita Pollutant Discharge Indicators

Yoshiaki Tsuzuki

DOI:10.4172/2252-5211.1000141



Share this article



Research Article

Pages: 1 - 4

White Shrimp (*Litopenaeus vannamei*) Culture using Heterotrophic Aquaculture System on Nursery Phase

Supono, Johannes Hutabarat, Slamet Budi Prayitno and YS Darmanto

DOI:10.4172/2252-5211.1000142



Share this article



Research Article

Pages: 1 - 4

How Economic Situation Affect the Decisions on the Building Energy Efficiency(BEE) Development in Malaysia



Tweets by @Carolin50507446

Apr 8, 2020



International Journal of Waste Management

@Carolin50507446

#COVID2019 Work Practices for #waste, #recycling Workers.

[Journal Home](#) [Editorial Panel](#)

[resources@esandsciencejournals.com](#) or [online on our website](#)

[Embed](#)

[View on Twitter](#)

Queen K Qian, Abd Ghani Bin Khalid and Edwin HW Chan

DOI:10.4172/2252-5211.1000143



Share this article



Pages: 1 - 8

Research Article

Adsorption Mechanisms and Transport Behavior between Selenate and Selenite on Different Sorbents



Share this article



Pages: 1 - 6

Research Article

Rapid Waste Composition Studies for the Assessment of Solid Waste Management Systems in Developing Countries

Timothy G Townsend and Max J Krause

DOI:10.4172/2252-5211.1000145



Share this article



Pages: 1 - 5

Research Article

Newly-Isolated Laccase High Productivity Streptomyces Sp. Grown In Cedar Powder as the Sole Carbon Source

Akihisa Aoyama, Kazuhiro Yamada, Yoshinobu Suzuki, Yuta Kato, Kazuo Nagai and Ryuichiro Kurane

DOI:10.4172/2252-5211.1000146



Share this article



White Shrimp (*Litopenaeus vannamei*) Culture using Heterotrophic Aquaculture System on Nursery Phase

Supono^{1*}, Johannes Hutabarat², Slamet Budi Prayitno² and YS Darmanto²

¹Department of Aquaculture, Faculty of Agriculture, Lampung University, Indonesia

²Department of Fisheries, Faculty of Fisheries and Marine Science, Diponegoro University, Indonesia

*Corresponding author: Supono, Department of Aquaculture, Faculty of Agriculture, Lampung University, Indonesia, Tel: +628127240191; E-mail: supono_unila@yahoo.com

Received date: January 9, 2014; Accepted date: April 16, 2014; Published date: April 23, 2014

Copyright: ©2014 Supono et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Heterotrophic aquaculture system is an environmental friendly shrimp culture that has a huge potency to improve yields of *Litopenaeus vannamei*. Biofloc grown in a heterotrophic aquaculture system that can be used as an alternative feed for shrimp due to its high nutrition. Biofloc contains bacterial protein and polyhydroxybutyrate that are able to enhance growth. Biofloc also contains bacteria that have peptidoglycan and lipopolysaccharide on their cell walls. The aim of the research was to study the effect of heterotrophic aquaculture system on culturing of *Litopenaeus vannamei* during nursery phase. The experiment was arranged in split plot design in three replicates. The treatments consisted of two factors namely various densities and different aquaculture systems. The aquaculture systems were autotrophic and heterotrophic aquaculture system, while densities were 1,000, 1,500, and 2,000 PLm⁻³. The result showed that there was no significant interaction between densities and aquaculture system toward the growth rate, protein efficiency ratio and yield of *Litopenaeus vannamei*. The heterotrophic aquaculture system was able to increase the yield of *Litopenaeus vannamei* on nursery phase. However heterotrophic aquaculture system did not significantly affect growth rate and protein efficiency ratio of *Litopenaeus vannamei*. While, the density significantly affected survival rate and yield of *Litopenaeus vannamei*.

Keywords: Environmental friendly; *Litopenaeus vannamei*; Heterotrophic; Biofloc

Introduction

The intensive development of the shrimp culture has been accompanied by an enhancement in environmental impact. In the autotrophic aquaculture system, waste of shrimp culture can be a serious problem both in culture pond and in the environment. The effluent as uneaten feed, feces, and excretions is mostly inorganic nitrogen (mobile nitrogen) in form of ammonia and nitrite due to high protein content in feed (30-40%). Nitrogen in feed is only 25% that is recovered in the shrimp on harvest while about 75% is released into the pond ecosystem, mostly as TAN [1]. Ammonia and nitrite in culture pond is toxic to shrimp. Inorganic nitrogen built up in ponds, is controlled by algae and nitrification. Inorganic nitrogen is converted to organic nitrogen to build algae cell. This process is limited by the rate carbon assimilation by algae. Nitrification is a slow process and need a few weeks to complete it [2].

In order to minimize impact of the shrimp culture effluent, it is important to develop shrimp farming with zero water exchange. The effluent produced by shrimp should be recycled in the pond before releasing to the surrounding environment. The method that can be applied to overcome the problem is heterotrophic aquaculture system (biofloc system). A carbon source namely sugar, molasses, and starch is added into culture pond increasing ratio of C:N to immobilize inorganic nitrogen and to stimulate the growth of heterotrophic bacteria to form biofloc.

The shrimp cultivated in heterotrophic system has higher price than that in autotrophic system due to an environmental friendly product.

Heterotrophic system in aquaculture has huge potential to increase growth and survival rate of shrimp. Heterotrophic system is capable to decrease cost production from feed because biofloc can be an alternative feed for shrimp that has high nutrition content [2].

Biofloc dominated by bacteria is high protein content and able to produce polyhydroxybutyrate as reserve energy and carbon [3]. Polyhydroxybutyrate is capable to accelerate the animal growth and to inhibit pathogenic vibrio in intestine tract. The objective of this experiment was to study the effect of heterotrophic system on the performance of on nursery *Litopenaeus vannamei* phase.

Materials and Methods

The experiment was arranged in split plot design with two factors in three replicates. The treatments consisted of aquaculture system (as sub plot) and density (as main plot). The aquaculture systems were heterotrophic system (HS) and autotrophic system (AS) while stocking densities of *Litopenaeus vannamei* were 10, 15, and 20 PL in 10 litres container equivalent to 1,000, 1,500, and 2,000 PLm⁻³.

Eighteen plastic containers were filled with sterile saline water. Water salinity was adjusted to osmolarity of *Litopenaeus vannamei* haemolymph at intermolt phase. Osmolarity of *Litopenaeus vannamei* haemolymph was measured with method conducted by Anggoro and Muryati [4]. Nine containers were used to culture *Litopenaeus vannamei* in heterotrophic system. Shrimp feed and glucose was added to get C:N ratio of 21 [2]. Bacterium of *Bacillus cereus* (10⁶ CFU/ml) was inoculated into media to stimulate heterotrophic system. Each container was equipped by aeration and was installed on the container bottom to maintain dissolved oxygen min. at 4 mg/l and water movement. After 10 days of growing biofloc, *Litopenaeus vannamei*

postlarvae (PL 17) with an average body weight of 11 ± 1.0 mg and average length of 1.36 ± 0.15 cm was stocked into culture media.

Shrimp culture was conducted in 30 days. Shrimp were fed by formulated feed of 38% protein on feeding rate of 5%. Zero water exchange was applied in heterotrophic system. The growth and survival rate of *Litopenaeus vannamei* were calculated with methods conducted by Far et al. [5]. Protein efficiency ratio was analyzed with a method described by Hoffman and Falvo [6]. All data were further analyzed statistically using Two-Way-Anova after testing of normality, homogeneity, and additivity using SPSS statistical software. Statistical significance of differences required that $p < 0.05$.

Results and Discussion

Haemolymph osmolarity of *Litopenaeus vannamei*

Haemolymph osmolarity of PL 11- *Litopenaeus vannamei* on premolt phase was 933,89 mOsm/l H₂O, equivalent to 32‰, meanwhile haemolymph osmolarity of *Litopenaeus vannamei* on intermolt phase was 861,00 mOsm/l H₂O, equivalent to 29,5‰ (Table 1).

Phase	Osmolarity (m Osm/l H ₂ O)	Salinity (‰)
Premolt	1. 933.89	32
	2. 933.90	32
	3. 933.87	32
	Average	32
Intermolt	1. 861.02	29.5
	2. 860.98	29.5
	3. 861.00	29.5
	3. 861.00	29.5

Table 1: Osmolarity haemolymph of *Litopenaeus vannamei*.

According to the data, applied salinity of media was 30‰. Osmolarity of haemolymph on premolt phase and intermolt phase is the best range for growth (4).

Water quality

Dissolved oxygen, total ammonia nitrogen (TAN), and pH of culture media of heterotrophic system and autotrophic system are performed at Figures 1, 2 and 3, respectively. The range of dissolved oxygen (DO) in culture media was 5.0–8.0 mg/l, TAN was 0.01–0.05 mg/l, and pH was 6.8–7.2. The values were in range for growing *Litopenaeus vannamei* [5].

Shrimp Performance

In general, performance of *Litopenaeus vannamei* in heterotrophic aquaculture system was better than that in autotrophic one. Specific Growth Rates (SGR), survival Rate (SR), protein efficiency ratio (PER), and yield of *Litopenaeus vannamei* at harvest are summarized in Table 2.




Figure 1: DO of culture media.




Figure 2: TAN of culture media.




Figure 3: pH of culture media.

According to statistical analytic (ANOVA), there was no significant interaction between densities and aquaculture system to performance of *Litopenaeus vannamei*. The use of heterotrophic system did not enhance the growth rate and protein efficiency ratio (Figures 4 and 5). However survival rate and yield of *Litopenaeus vannamei* significantly increased in heterotrophic system (Figures 6 and 7). Stocking densities significantly affected the survival rate and yield of *Litopenaeus vannamei*.

Density (PL/m ³)	System	SGR (%)	SR (%)	PER	Yield (g/m ³)
1000	AS	12.75 ± 2.08	60.0 ± 10.0	0.83 ± 0.37	256.6 ± 107.8
	HS	13.07 ± 1.27	73.3 ± 5.8	1.08 ± 0.35	329.9 ± 101.9

1500	AS	12.85 2.08	±	60.0 ± 6.7	0.82 ± 0.37	378.1 85.7	±	
	HS	14.04 0.80	±	66.7 ± 6.7	1.25 ± 0.26	571.4 114.5	±	
2000	AS	12.90 2.08	±	41.7 10.4	±	0.55 ± 0.10	344.3 59.2	±
	HS	13.15 2.08	±	50.0 10.0	±	0.73 ± 0.19	450.8 113.6	±

Table 2: Performance of *Litopenaeus vannamei*.




Figure 4: Specific growth rates (SGR) of *Litopenaeus vannamei*.




Figure 5: Protein efficiency ratio (PER) of *Litopenaeus vannamei*.




Figure 6: Survival rate (SR) of *Litopenaeus vannamei*.

Discussion

Ammonia in pond is produced as a major end product of the metabolism due to high content protein of feed and is excreted as ammonia across the gill of shrimp [7]. No water exchange applied in heterotrophic system was able to control TAN in aquaculture system. By adding organic carbon source to the water, it forces the bacteria to immobilize any inorganic nitrogen present in the pond. Inorganic nitrogen was recycled in the culture pond resulting in microbial protein biomass needed for cell growth and multiplication. At high ratio of C:N, heterotrophic bacteria will assimilate ammonium nitrogen directly from water metabolized to cell biomass. The addition of carbon source is most effective method in decreasing inorganic nitrogen mostly TAN [7] and is often more stable and reliable than algal uptake or nitrification [1].




Figure 7: Yields of *Litopenaeus vannamei*.

Specific growth rates and protein efficiency ratio of *Litopenaeus vannamei* in heterotrophic aquaculture system tended to be better than those in autotrophic system. Biofloc, formed in heterotrophic system, can be benefited by shrimp as an alternative feed (Figure 8). Biofloc contains bacterial protein [8] and polyhydroxybutyrate [3] produced by bacteria. Biofloc was consumable due to the appropriate size for shrimp. Polyhydroxybutyrate is the most dominant polymer

and is useful in aquaculture. The advantages of PHB are an energy reserve for fish, digestible in intestine, increasing unsaturated fatty acid, and increasing growth of fish [1]. Biofloc also contains methionine, lysine, vitamins and minerals, especially phosphorus




Figure 8: Biofloc.

Survival rate of *Litopenaeus vannamei* in heterotrophic system experienced enhancement due to biofloc contains bacteria. Bacteria are capable to produce polyhydroxybutyrate. Polyhydroxybutyrate will release 3-hydroxy butyric acid (short chain fatty acid) in the gastro intestinal tract as inhibitor of pathogenic bacteria. According to several researches, PHB is capable to inhibit pathogen in the intestinal tract and to be antimicrobial against *Vibrio*, *E. coli*, and *Salmonella*, to control pathogen of *Vibrio harveyi*, and to enhance survival rate of *Artemia franciscana* larvae [9]. Far et al. [5] investigated that Bacillus is able to increase survival rate of *Litopenaeus vannamei* and to decrease luminous *Vibrio* densities in the pond water. Bacteria also contain peptidoglycan and lipopolysaccharide on their cell wall. Peptidoglycan and lipopolysaccharide are immunostimulant being capable to increase nonspecific immunity of shrimp. The substances influence prophenoloxidase activity and phagocytosis of hyaline cells [10].

The yields of *Litopenaeus vannamei* in heterotrophic system enhanced significantly compared to autotrophic one. The enhancement was affected by growth (Figure 4) and survival rate (Figure 6). According to polynomial orthogonal analysis, optimal density of *Litopenaeus vannamei* in autotrophic system on nursery phase was 1612 PL/m³ yielded 466 g/m³, while in heterotrophic system, optimal density was 1638 yielded 639 g/m³ (Figure 9). Heterotrophic system was capable to increase yield of *Litopenaeus vannamei* of 37% compared to autotrophic system.

Conclusions

Heterotrophic aquaculture system can be an alternative method to culture *Litopenaeus vannamei* in pond. The system is environmental-friendly one overcoming the problem in shrimp culture related to deterioration of environment quality. Pond water quality was controlled resulting in good growth, high survival rate, and increased yield of *Litopenaeus vannamei* on nursery phase. Heterotrophic system aquaculture was capable to increase the yield of *Litopenaeus vannamei* on nursery phase. However heterotrophic system did not significantly affect the growth rate and protein efficiency ratio of *Litopenaeus vannamei*. The density affected the survival rate and yield of *Litopenaeus vannamei*.




Figure 9: Optimal density of *Litopenaeus vannamei* on nursery phase.

References

1. Crab R, Avnimelech Y, Defoirdt T, Bossier P, Verstraete W (2007) Nitrogen removal techniques in aquaculture for a sustainable production. *Aquaculture* 270: 1–14.
2. Avnimelech Y (2009) *Biofloc Technology—A Practical Guide Book*. The World Aquaculture Society Baton Rouge Louisiana United State.
3. De Schryver P, Sinha AK, Kunwar PS, Baruah K, Verstraete W, et al. (2010) Poly-beta-hydroxybutyrate (PHB) increases growth performance and intestinal bacterial range-weighted richness in juvenile European sea bass, *Dicentrarchus labrax*. *Applied Microbiology and Biotechnology* 86: 1535–1541.
4. Anggoro S, Muryati (2006) Osmotic respons of tiger shrimp (*Penaeus monodon* Fab.) juvenile and adult at various level of molting stages and salinity. *Buletin Penelitian dan Pengembangan Industri* 1: 59–63.
5. Far HZ, Saad CRB, Daud HM, Harmin SA, Shakibazadeh S (2009) Effect of *Bacillus subtilis* on the growth and survival rate of shrimp (*Litopenaeus vannamei*). *African Journal of Biotechnology* 8: 3369–3376.
6. Hoffman JR, Falvo MJ (2004) Protein - Which is Best? *J Sports Sci Med* 3: 118–130.
7. Ebeling JM, Michael B, Timmons JJ, Bisogni (2006) Engineering analysis of the stoichiometry of photoautotrophic, autotrophic, and heterotrophic removal of ammonia-nitrogen in aquaculture systems. *Aquaculture* 257: 346–358.
8. Hargreaves JA (2013) *Biofloc Production System for Aquaculture*. Southern Rregional Aquaculture Center Publication No: 4503.
9. Crab R, Lambert A, Defoirdt T, Bossier P, Verstraete W (2010) The application of bioflocs technology to protect brine shrimp (*Artemia franciscana*) from pathogenic *Vibrio harveyi*. *J Appl Microbiol* 109: 1643–1649.
10. Yeh ST, Li CC, Tsui WC, Lin YC, Chen JC (2010) The protective immunity oh white shrimp *Litopenaeus vannamei* that had been immersed in the hot water extract of *Gracilaria tenuistipitata* and subjected to combined stresses of *Vibrio alginolyticus* injection and temperature change. *Fish & Shellfish Immunology* 29: 271–278.