



**FACULTY OF ANIMAL SCIENCE
BOGOR AGRICULTURAL UNIVERSITY**

**THE FIRST INTERNATIONAL SEMINAR ON ANIMAL INDUSTRY
"Sustainable Animal Production for Food Security and Safety"**

23-24 November 2009

IPB International Convention Center, Bogor-Indonesia

**PROGRAM
ABSTRACTS**



BOGOR AGRICULTURAL UNIVERSITY



MINISTRY OF AGRICULTURE




**FACULTY OF ANIMAL SCIENCE
BOGOR AGRICULTURAL UNIVERSITY**

**THE FIRST INTERNATIONAL SEMINAR ON ANIMAL INDUSTRY
"Sustainable Animal Production for Food Security and Safety"**

23-24 November 2009

IPB International Convention Center, Bogor-Indonesia

PROGRAM
ABSTRACTS



BOGOR AGRICULTURAL UNIVERSITY



MINISTRY OF AGRICULTURE

LEMBAR PENGESAHAN

Judul : Ration with Different Dietary Cation Anion Difference to Mineral Status of Blood and Urine Garut Ewes

Penulis Utama : Farida Fathul

Penuls Anggota : Toto Toharmat, Arief Boediono, Idat Galih Permana

Jurusan/Fakultas : Peternakan/Pertanian

Seminar : The First International Seminar on Animal Industry "Sustainable Animal Production for Food Security and Safety" di IPB International Convention Center, Bogor Indonesia, 23-24 November 2009

Mengetahui,
aa. Dekan Fakultas Pertanian
Universitas Lampung
Pembantu Dekan I,

Dr. Ir. Erwanto, M.S.
NIP 19610225 198603 1 004

Bandar Lampung, 30 Maret 2010

Ketua Jurusan Peternakan
Fakultas Pertanian
Universitas Lampung,

Prof. Dr. Ir. Muhtarudin, M.S.
NIP 19610307 198503 1 006

Mengesahkan,
Ketua Lembaga Penelitian
Universitas Lampung

Prof. Dr. John Hendri, M.S.
NIP 19581021 198703 1 000

| | |
|--|--------------------------------|
| KORUMBATASILEMBAKA PENELITIAN UNIVERSITAS LAMPUNG | |
| TGL. | 5 maret 2010 |
| No. INVEN | 161 / H26 / 3 / PC / FP / 2010 |
| JENIS | prosidy |
| PARAF | |



REMARKS FROM THE DEAN OF FACULTY OF ANIMAL SCIENCE

Assalamu'alaikum Warahmatullahi Wabarakatuh

First of all, let us pray to Allah SWT the Almighty for His blessings bestowed to all of us.

As the Dean of Faculty of Animal Science, it is surely a great pleasure for me to welcome all of you on the 1st International Seminar on Animal Industry 2009, entitled "Sustainable Animal Production for Food Security and Safety". This seminar is organized by the Faculty of Animal Science, Bogor Agricultural University.

As one of the faculties of animal science in Indonesia, it is our responsibility to take a real action for developing animal production for food security and safety. In this seminar, we expect to have interesting discussion on current animal research especially on efficiencies and productivities of livestock; perspectives of stakeholders on potencies, prospect and constraints on animal industry; animal biotechnology, animal business in global era; and other relevant topics. We hope that this seminar supplies a scientific recommendation to government and non government institutions on policy development of food security and safety, improvement of international linkage for solving the problems in animal production, strengthen the international collaborative research and the exchange of information.

In this special occasion I would like to express appreciation to Ir. Suswono, MMA., the Minister of Agriculture for support and encouragement. We also extend our gratitude to the Directorate General of Livestock Services, who has given the support to this seminar. I would like to express my appreciation to the invited speakers and other speakers both oral and poster presentation, who are willing to share their experience and vision with us. To the contributors, sponsors and exhibitors I would like to express our great thank to every effort which have been done to make this event successful. Last but not least, please accept my gratitude for the members of steering committee and organizing committee, without their effort and hard work, this meeting will never be carried out. Please, enjoy the seminar and hopefully you will get the benefits of this scientific and professional gathering. Thank you very much.

Wabillahi taufiq wal hidayah,
Wassalamu'alaikum warahmatullahi wabarakatuh

Bogor, November 23rd, 2009
Dean,

Dr. Ir. Luki Abdullah, M.Sc.Agr.



GUIDELINES OF PRESENTATION

Guidelines for Oral Presentation

1. The official language of the 1st ISAI is English.
2. The material of presentation must be submitted to the organizing committee during registration on day 1 in order to arrange it in the computer.
3. A chair person will act as a moderator and has authority to organize the presentation.
4. Each presenter will be allocated 10 minutes for presentation (the moderator may stop presentation if over time), and 5 minutes for discussion.
5. Discussion is not allowed during presentation.
6. After presentation, the moderator will guide the discussion.
7. The presenters must attend the full 2 days program to be eligible as the ISAI Oral Presenter Award nominee.

Guidelines for Poster Presentation

1. Poster must be placed in the allocated space at least in the first day after registration and displayed for two days. Double-sided sticky taped must be provided by the presenters.
2. Poster must be attended by the presenter(s) during the provided time (break time) as scheduled in the program.
3. The presenter must attend the full 2 days program of ISAI seminar to be eligible as the ISAI Poster Award nominee.

SEMINAR PROGRAM

Tentative Schedule

Monday, November 23, 2009

| Time | Ballroom 3 | |
|-------------|---|---|
| | Event | Speaker |
| 08.00-09.00 | Registration | |
| 09.00-10.00 | Opening Ceremony <ul style="list-style-type: none"> Organizing Committee Report Welcome Address from Rector of the Bogor Agricultural Institute Opening and Keynote Speech by Vice Minister of Agriculture | Prof. Dr. Ir. Muladno, MSA. Prof. Dr. Ir. Hery Suhardiyanto, M.Sc. Dr. Ir. Bayu Khrisnamurthi, MSc. |
| 10.00-10.15 | Coffee Break and Poster Session | |
| 10.15-10.45 | Plenary 1: New development in animal production technology for sustainable use of animal genetic resources (AnGR) | Orlando Fernandez, DVM, FPCCP |
| 10.45-11.15 | Plenary 2: Recent feed technology: Role of feed technology to enhance efficiency and productivity of animal | Ahmad Mujahid, PhD |
| 11.15-11.45 | Plenary 3: Anticipating the Outbreak of Zoonotic Infectious Diseases Related to Animal Industry | Dr. Drh. Retno Dewi Bagja |
| 11.45-12.15 | Plenary 4: New Development of Animal Production in Indonesia | Prof. Dr. Ir. Toto Toharmat, M.Agr.Sc. |
| 12.15-13.30 | Lunch | |
| 13.30-14.00 | Poster Session | |

| Time | Room C | Room D |
|-------------|---|---|
| | Genetics, Breeding and Reproduction Moderator: Prof. Dr. Ir. Cece Sumantri, M.Agr.Sc. | Feed and Nutrition Moderator: Dr. Ir. Sumiati, M.Sc. |
| 14.10-14.25 | Anneke Anggraeni: Genetic Polymorphism of the Kappa-Casein Gene in Holstein-Friesian Dairy Cattle in West Java Province | Retno Murwani: Effect of Mungbean as Local Feed Ingredients to Substitute Soybean Meal in the Diet on the Performance of Broilers |

The Effect of Ration with Dietary Cation Anion Difference to Blood Mineral and Urine Status of Garut Ewes

F. Fathul¹⁾, T. Toharmat²⁾, A. Boediono³⁾, and I. G. Permana⁴⁾

¹⁾ Department of Animal Husbandry, Faculty of Agriculture, Lampung University, Bandar Lampung, Indonesia


^{2,4)} Department of Nutrition and Feed Technology, Faculty of Animal Science, Bogor Agricultural University, Indonesia

³⁾ Department of Anatomy, Physiology, and Pharmacology, Faculty of Veterinary, Bogor Agricultural University, Indonesia

Abstract

The objectives of the present experiment were to evaluate the effect of dietary cation anion difference (DCAD) on mineral status of in blood and urine. Rations with DCAD value of -28, -18, 0, +14, and +32 mEq were offered to 15 ewes in a completely randomized block design. The ewes were grouped into ewes previously had twin female offsprings (I); twin male offsprings (II); twin male and female offsprings (III). Each treatment rations with different DCAD had been offered for 3 ewes. On day 21, blood samples were taken anaerobically using heparinized syringes from the coccygeal jugular venipuncture. Each syringe was capped and placed on ice immediately following collection to determine on blood pH and plasma Na, K, Cl, Ca, and P concentrations. The DCAD value had no effect on blood pH and plasma Na, K, Cl, Ca, and P concentrations indicating that there was homeostasis to maintain the physiological status of the body. The DCAD value of -18, 0, +14 and +32 mEq resulted in the normal blood with Na : K ratio closed to 20 : 1. Plasma Cl concentration was associated with plasma Na, but the concentration of Cl was lower than that of Na. The DCAD value significantly influenced P, but had no effect on urinary Na, K, Cl, S, and Ca. The DCAD value of -28 and -18 meq resulted in the low acidity of urine at level of 5.73 ± 0.20 and 5.84 ± 0.27 , respectively. The DCAD value of 0, +14, dan +32 mEq resulted in normal urinary pH.

Keywords: dietary cation-anion difference (DCAD), blood, urin, mineral, ewe



LIST OF ISAI COMMITTEE

| | |
|-----------------------------|--|
| Honorable advisors | : Minister of Agriculture Minister of National Education Minister of Micro, Small and Middle Entrepreneurship Rector of IPB |
| Steering Committee | |
| Chair person | : Dr. Ir. Luki Abdullah, M.Sc.Agr. |
| Secretary | : Prof. Dr. Ir. Muladno, M.S.A. |
| Members | : Prof. Dr. Ir. Wiranda G. Pilliang, M.Sc. Prof. Dr. Ir. Pollung H. Siagian, M.S. Prof. Dr. Ir. Toto Toharmat, M.Sc. Prof. Dr. Ir. Iman Rahayu H.S., M.S. Prof. Dr. Ir. Ronny R. Noor, M.Rur.Sc. Prof. Dr. Ir. Cece Sumantri, M.Agr.Sc. Dr. Ir. Moh. Yamin, M.Agr.Sc. Dr. Ir. Idat G. Permana, M.Sc.Agr |
| Organizing Committee | |
| Chair person | : Prof. Dr. Ir. Muladno, M.S.A. |
| Executive secretaries | : Indah Wijayanti, S.TP., M.Si. (Domestic affairs) Maria Ulfah, S.Pt., M.Sc.Agr. (International affairs) Dr. Rudi Afnan, S.Pt., M.Sc.Agr. (General affairs) |
| Administration | : Nur Rochmah Kumalasari, S.Pt., M.Si. Dewi Murtini, S.Pt. Ani Nurnina, S.E. Nindyar Harumi Sukma, S.Pt. |
| Treasures | : Dr. Ir. Dewi Apri Astuti, M.S. Triyati |
| Scientific Committee | : Prof. Dr. Ir. I Komang Gede Wiryawan, M.Sc. Prof. Dr. Ir. Ronny R. Noor, M.Rur.Sc. Prof. Dr. Ir. Toto Toharmat, M.Sc. Prof. Dr. Ir. Cece Sumantri, M.Agr.Sc. Dr. Ir. Nahrowi Ramli, M.Sc. Dr. Ir. Rudy Priyanto Dr. Ir. Rarah Ratih Adjie Maheswari, D. E. A. Dr. Ir. Sri Mulatsih Sri Suharti, S.Pt., M.Si. Tuti Suryati, S.Pt., M.Si. |

| | |
|---|--|
| Funding | : Dr. Ir. Asnath Maria Fuah, M.S. Dr. Ir. Erika B. Laconi, M.S. Dr. Ir. Panca Dewi MHKS, M.S. |
| Documentation/Publication | : Ir. Asep Tata Permana, M.Sc. Bramada Winar Putra, S.Pt. |
| Promotion/Website Excursion | : Sugeng Triwahyono, Amd. : Dr. Ir. Mohammad Yamin, M.Agr.Sc. Dr. Ir. Henny Nuraini, M.S. |
| Consumption | : Ir. Sri Rahayu, M.S. Ir. Widya Hermana, M.S. Ir. Hotnida CH. Siregar, M.Si. |
| Logistic, Transportation and General | : Dr. Jakaria, S.Pt., M.Si Edit Lesa Aditia, S.Pt. Jaka Saputra Ahmad Fuadi Muhammad Sarwar Khan Yesua Gidsan Tondas Meri Afiza Titis A. P. Apdini Fastasqi Nuniek Setiorini Adriana Herawati Rosalia Praharyanti Erliyani Nurhidayah |

RATION WITH DIFFERENT DIETARY CATION ANION DIFFERENCE TO MINERAL STATUS OF BLOOD AND URINE GARUT EWES

F. Fathul^{*1}, T. Toharmat[†], A. Boediono[‡], and I. G. Permana[†]

^{*}Department of Animal Husbandry, Faculty of Agriculture, Lampung University, Bandar Lampung, Indonesia

[†]Department of Animal Feed and Nutrition, Faculty of Animal Animal Science, Bogor Agricultural University, Indonesia

[‡]Department of Anatomy, Phisiology, and Pharmacology, Faculty of Veterinary, Bogor Agricultural University, Indonesia

1 Corresponding address:

Email : ida_fathul@unila.ac.id or farida_fathul@yahoo.com

Office : Animal Husbandry Department, Agricultural Faculty, Lampung University

Jl Sumantri Brojonegoro 1

Telp 0721-773552

Bandar Lampung. 35135



Abstract

The objectives of the present experiment were to evaluate the effect of dietary cation-anion difference (DCAD) on mineral status in blood and urine. Rations with DCAD value of -28, -18, 0, +14, and +32 mEq were offered to 15 ewes in a randomized complete block design. On day 21, blood samples were taken anaerobically using heparinized syringes from the coccygeal jugular venipuncture. Each syringe was capped and placed on ice immediately following collection to determine on plasma Na, K, Cl, Ca, and P concentrations.

The DCAD values had no effect on plasma Na, K, Cl, Ca, and P concentrations indicating that there was homeostasis to maintain the physiological status of the body. The DCAD value of -18, 0, +14 and +32 mEq resulted in the normal blood with Na : K ratio closed to 20 : 1. Plasma Cl concentration was associated with plasma Na, but the concentration of Cl was lower than that of Na.

The DCAD values significantly influenced P urine, but had no effect on urinary Na, K, Cl, S, and Ca. The DCAD value of -28 and -18 mEq resulted in the low acidity of urine at level of 5.73 ± 0.20 and 5.84 ± 0.27 , respectively. The DCAD value of 0, +14, dan +32 mEq resulted in normal urinary pH.

Rations with DCAD values of -18, 0, +14, and +32 mEq in garut ewes had normal ratio of plasma's $\text{Na}^+ : \text{K}^+$ and were able to perform regulation of minerals control inside their blood to be homeostatis, and some excessive minerals would be secreted through urine. Rations with DCAD values of -28 and -18 mEq in garut ewes had the highest ratio of plasma's $\text{Ca}^{2+} : \text{P}^{2-}$ which was 2.2 : 1.0, so it could be used as an action to prevent milk fever.

Keywords: dietary cation-anion difference, blood, urin, ewe

INTRODUCTION

Consumed ration will affect physiological condition of the livestock. According to Stewart (1983), addition of anions (Cl and S) into the ration would lower the pH of body fluid. Blood condition is the result of acid base balance in body fluid and regulation of nutrition metabolism inside it. Blood is consisted of cells and plasma. Plasma contains water as much as 90 % and anorganic minerals in form of soluted ion as electrolytes, proteins, metabolic waste products, respiration gases and hormones. Concentration of this combined ions is important for maintenance of blood osmotic balance. Acid base balance is highly affected by the function of lungs and kidneys.

Kidneys have vital role as controller of volume and composition of blood chemicals by secreting solution and water selectively. Vital functions of kidneys are done by the filtration of plasma through *glomerulus* followed with reabsorption some amount of solution and water with correct volume along the kidneys tubulus. The excess of solution and water will be secreted out as urine through collector system. Epitelic cells help maintaining constant pH of body fluid by controlling secretion of hydrogen ion. Secretion of acid in urine as the result of potential acid and H^+ formation rate from blood buffer. Acidic urine is also secreting Ca^{2+} .

In this research, the rations experimented with DCAD values of -28, -18, 0, +14, and +32 mEq. The objectives of this research was, to identify the effect of different DCAD to mineral in blood and urine of Garut ewes (*Ovis aries*).

MATERIALS AND METHODS

This experiment was conducted at the Pen Field Laboratorium A of Animal Husbandry Faculty and Integrated Laboratorium of Veterinary Faculty Bogor Agricultural University on January 11th - July 14th 2007.

Experimental Design and Animal Care

Fifteen Garut ewes were 2.50 ± 0.25 years old were assigned randomly to randomized complete block design. The ewes were blocked into groups of 3 according to (I) ewes previously had twin female offsprings; (II) ewes previously

had twin male offsprings; and (III) ewes previously had twin male and female offsprings. Ewes were housed and fed in individual cage. The composition of basal ration contained 89.30 % dry matter, 8.12 % ash, 15.00 % crude protein, 5.12 % ether extract, 14.73 % crude fiber, and 57.03 % nitrogen free extract (Tabel 1). Determination of crude protein ration contents of 15.00 % based on Wodzicka-Tomaszewska et al. (1991), Na mineral of 0.09 - 0.18 %, K of 0.50 - 0.80 % (maximum 3.00 %), Cl had no clause (based on NRC 1985).

Tabel 1. Composition and nutrient content of basal ration base on dry matter

| Feed type | Percent | Proximate analysis (%) | | | | | Minerals (%) | | | |
|---------------|------------|-----------------------------|--------------|-------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Ash | CP | EE | CF | NFE | Na | K | Cl | S |
| Corn forage | 35.0 | 2.81 | 3.24 | 0.43 | 10.18 | 18.34 | 0.020 | 0.305 | 0.011 | 0.070 |
| Rice bran | 6.0 | 0.63 | 0.59 | 0.84 | 0.65 | 3.30 | 0.000 | 0.079 | 0.014 | 0.008 |
| Onggok | 9.5 | 1.83 | 0.26 | 0.13 | 0.83 | 6.45 | 0.007 | 0.017 | 0.002 | 0.003 |
| Corn meal | 18.5 | 0.26 | 1.48 | 0.55 | 0.42 | 15.79 | 0.001 | 0.068 | 0.002 | 0.013 |
| Coconut meal | 7.0 | 0.43 | 1.17 | 1.01 | 1.08 | 3.31 | 0.007 | 0.144 | 0.053 | 0.017 |
| Soybean meal | 22.0 | 1.98 | 8.24 | 0.37 | 1.58 | 9.84 | 0.001 | 0.458 | 0.111 | 0.041 |
| Fish oil | 2.0 | 0.18 | 0.03 | 1.79 | | | 0.003 | 0.006 | 0.000 | 0.000 |
| Total | 100 | 8.12 | 15.00 | 5.12 | 14.73 | 57.03 | 0.039 | 1.077 | 0.192 | 0.152 |
| Description : | | CP = crude protein | | | | | Na = natrium | | | |
| | | EE = ether extract | | | | | K = kalium | | | |
| | | CF = crude fiber | | | | | Cl = chlor | | | |
| | | NFE = nitrogen-free extract | | | | | S = sulfur | | | |

The value of basal rations DCAD was +14 mEq/100 g of DM and treatment rations in this research with five dietary cation anion difference values (DCAD).

1. -28 mEq = basal ration added with 14.375 mEq S and 27.884 mEq Cl
2. -18 mEq = basal ration added with 14.375 mEq S and 17.884 mEq Cl
3. 0 mEq = basal ration added with 14.259 mEq S
4. +14 mEq = basal ration
5. +32 mEq = basal ration added with 10.21 mEq Na and 7.531 mEq K

Method of operating decreasing of DCAD value to 0 mEq/100 g of DM with basal ration was added CaSO_4 (Brataco Chemika, Cikarang, Jakarta). Decreasing DCAD to -28 and -18 mEq/100 g of DM with basal ration were added CaCl_2 , dan CaSO_4 (Brataco Chemika, Cikarang, Jakarta). The value of DCAD increased to +32 by addition of Na_2CO_3 and K_2CO_3 in basal ration (Brataco Chemika, Cikarang, Jakarta). Treatment rations had been offered for three weeks before the samples were collected.

Sample Collection and Analysis

On 21st day after treatment rations were offered, blood samples were taken anaerobically using heparinized syringes (Franklin Lakes NJ USA) from the coccygeal jugular venipuncture. Then, the blood were centrifuged for 15 minutes at 3000 rpm. Afterward, the plasma were taken to be analyzed for its Na, K, and Ca contents by using Automatic Absorbance Spectrofotometer (AAS), while Cl, P, and S by titration.

Sample of urine were collected by using plastic apron in the morning around 07.00 - 08.00 o'clock. The urine were tested for its pH by using pH-meter pocket HANNA, then the urine were analyzed for its Na, K, Ca, Cl, P, and S mineral contents by using the same method with the blood sample.

Statistical Analysis

Data were analyzed with GLM procedure in SAS System for Windows 6.12. Treatments effects were compared using the multiple comparison approach of Duncan Multiple Range Test. Regression analyses were conducted with the Proc REG procedure, whereas correlation coefficients were obtained from the Proc CORR procedure of SAS (Mattjik and Sumertajaya, 2006).

RESULTS AND DISCUSSIONS

Blood Acidity and Blood Plasma Mineral Status

Average data of Na⁺, K⁺, Cl⁻, S²⁻, Ca²⁺, and P²⁻ of garut ewes' blood plasma offered with rations with various DCAD values were presented on Table 2.

Table 2. Average of Na^+ , K^+ , Cl^- , S^{2-} , Ca^{2+} , and P^{2-} of garut ewes' blood plasma offered with different DCAD

| Variables | Dietary cation-anion difference (mEq) | | | | |
|--------------------------------|---------------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| | -28 | -18 | 0 | +14 | +32 |
| Na^+ (ppm) | 17308 \pm 3281 | 18547 \pm 1661 | 18397 \pm 4940 | 16520 \pm 516 | 18397 \pm 1915 |
| K^+ (ppm) | 983 \pm 183 | 926 \pm 174 | 918 \pm 104 | 811 \pm 268 | 945 \pm 55 |
| Cl^- (ppm) | 4580 \pm 646 | 4449 \pm 82 | 4509 \pm 268 | 4307 \pm 102 | 4698 \pm 182 |
| S^{2-} (ppm) | 63 \pm 18 ^a | 35 \pm 2 ^b | 33 \pm 2 ^b | 32 \pm 5 ^b | 29 \pm 3 ^b |
| Ca^{2+} (ppm) | 473 \pm 27 | 471 \pm 22 | 449 \pm 20 | 421 \pm 44 | 426 \pm 43 |
| P^{2-} (ppm) | 211 \pm 114 | 217 \pm 51 | 320 \pm 50 | 331 \pm 51 | 339 \pm 25 |
| $\text{Na}^+:\text{K}^+$ | 18 : 1 | 20 : 1 | 20 : 1 | 20 : 1 | 19 : 1 |
| $\text{Ca}^{2+}:\text{P}^{2-}$ | 2.2 : 1.0 | 2.2 : 1.0 | 1.4 : 1.0 | 1.3 : 1.0 | 1.3 : 1.0 |

description : value with different letter on same row mean different ($P < 0.05$)

The DCAD values had no effect ($P > 0.05$) on plasma's Na^+ . The differences of DCAD values were consumed by garut ewes had no effect on plasma's Na^+ ($P > 0.05$). It meant that, the ewes succeeded in performing homeostatis. Average values of Na^+ of experimented ewes' plasma were ranging 17308 \pm 3281 to 18397 \pm 1915 ppm (Table 2). The amount of plasma's Na^+ was not related to DCAD value ($r = 0.01$). Hu and Murphy (2004) stated that there was no effect of DCAD on Na^+ of blood plasma. However, Roche et al. (2005) reported that the increase of DCAD value, would increase Na^+ of blood plasma. Ewes offered with rations with DCAD value of -28 mEq produced acidic blood (Fathul et al., 2008) so the Na^+ of blood plasma in ewes offered with rations with DCAD value of -28 mEq were relatively lower than those offered with rations with DCAD values of 0, +14, dan +32 mEq.

The DCAD values had no effect ($P > 0.05$) on plasma's K^+ . The differences of DCAD values were consumed by garut ewes had no effect on plasma's K^+ ($P > 0.05$). It meant that, the ewes also succeeded in performing homeostatis. Average values of K^+ of experimented ewes' plasma were ranging 811 \pm 268 to 983 \pm 183 ppm (Table 2). The amount of plasma's K^+ was not related to DCAD value ($r = -0.16$).

At normal condition, extracellular fluid performed balance between Na^+ and K^+ at constanta of 20 : 1 (Georgievskii 1982). The ewes on this research performed homeostatis, the contents of Na^+ and K^+ of blood plasma were not affected by DCAD. But, the body regulated K^+ value inside the plasma to always

lower than Na^+ value. The relatively highest average value of plasma's K^+ (983 ± 183 ppm) was found in ewes offered with rations with DCAD value of -28 mEq. This was because in those ewes there were occurred metabolic acidosis indicated by the decrease of blood's HCO_3^- concentration (-2.53 ± 2.42 mmol/L) (Fathul et al. 2008). Therefore, the decrease of blood's HCO_3^- concentration would be followed by alteration of plasma's K^+ so plasma's K^+ became relatively highest. On Table 2, ewes offered with rations with DCAD values of -28 , -18 , 0 , $+14$, and $+32$ mEq had each ratio plasma's $\text{Na}^+ : \text{K}^+$ of $18 : 1$, $20 : 1$, $20 : 1$, $20 : 1$, and $19 : 1$, respectively. The values ratio plasma's $\text{Na}^+ : \text{K}^+$ in ewes offered with rations with DCAD value of -18 , 0 , $+14$, and $+32$ mEq (approximately $19 : 1 - 20 : 1$) were close to normal because the normal ratio plasma's $\text{Na}^+ : \text{K}^+$ was $20 : 1$. Ewes offered with rations with DCAD values of -28 had ratio plasma's $\text{Na}^+ : \text{K}^+$ of $18 : 1$, it meant that its plasma's Na^+ was lower and its plasma's K^+ was the highest compared with ewes offered with other DCAD values. Odongo et al. (2006) stated that metabolic acidosis was acid-base upset caused by the decrease of blood's $[\text{HCO}_3^-]$ and generally, followed by alteration of $[\text{K}^+]$ to become hyperkalemia (Weiderseiner et al. 2004). Ewes offered with rations with DCAD value of -28 mEq had the lowest blood's HCO_3^- and included in metabolic acidosis (Fathul et al., 2008), so it had relatively the most K^+ content. High content of K^+ in blood was called hyperkalemia. This, maybe because the ewes offered with rations with DCAD value of -28 mEq had very acidic blood pH and there was occurred metabolic acidosis, eventually ratio $\text{Na}^+ : \text{K}^+$ was not at normal condition. Ratio $\text{Na}^+ : \text{K}^+$ had to be performed by the livestock in order homeostatis. Determination of $\text{Na}^+ : \text{K}^+$ homeostasis mechanism inside the body was done by kidneys. Regulation of $\text{Na}^+ : \text{K}^+$ homeostasis was involving corticoid-aldosterone and deoxycorticosterone mineral which acted on K^+ secretion in consequence of reabsorption of Na^+ ion inside the kidneys' ducts. Corticoid mineral was also likely affecting the regulation of membrane permeability and $\text{Na}^+ : \text{K}^+$ pair mechanism (Pratas 1992).

Block (1994) explained that the unbalance of one ion with another, will caused poisoning that produced alkalosis or acidosis. This was likely because the unbalance of HCO_3^- and H^+ variables. If the presence of Na^+ was not enough to

initiate absorption of NaCl (neutral), the excess of HCO_3^- in blood vessels could drive to acidosis condition. Further explained by Horst et al. (1997) that Cl^- was absorbed more than SO_4^{2-} so Cl^- was a stronger aciditive to acidified the blood. Acid-base balance was related to exchange of H^+ ion internal media component which was able to donate or recieve ion. Substances that was able to donate ion was acid, while the one that was able to bind hydrogen was base.

The DCAD values had no effect ($P > 0.05$) on plasma's Cl^- . The differences of DCAD values consumed by garut ewes had no effect on plasma's Cl^- ($P > 0.05$). It meant that, the ewes succeeded in performing homeostatis. Average values of Cl^- of experimented ewes' plasma were ranging 4307 ± 102 to 4698 ± 182 ppm (Table 2). The amount of plasma's Cl^- was not related to DCAD value ($r = 0.07$). The content of plasma's Cl^- was following presence trend of plasma's Na, but the amount of Cl^- was lower than Na and the presence of Na^+ to form NaCl (neutral). Ewes offered with rations with DCAD value of -28 mEq had the lowest plasma's Na so to perform neutralization with Cl^- was relatively fewer than ewes offered with rations with other DCAD values.

The DCAD values had very high effect ($P < 0.01$) on plasma's S^{2-} . Ewes offered with rations with DCAD value of -28 mEq had the highest plasma's S^{2-} ($P < 0.05$) as much as 63.38 ± 17.94 ppm. Ewes offered with rations with DCAD value of -18, 0, +14, dan +32 mEq had no differences on plasma's S^{2-} ($P < 0.05$). The amount of plasma's S^{2-} was highly related to DCAD value ($r = -0.67$).

The DCAD values had no effect ($P > 0.05$) on plasma's Ca^{2+} . The differences of DCAD values consumed by garut ewes had no effect on plasma's Ca^{2+} ($P > 0.05$). It meant that, the ewes also succeeded in performing homeostatis. Average values of Ca^{2+} of experimented ewes' plasma were ranging 421 ± 44 to 473 ± 27 ppm. The relatively highest average value of plasma's Ca^{2+} (473 ± 27 ppm) was found in ewes offered with rations with DCAD value of -28 mEq. This was because the ewes offered with rations with DCAD values of -28 and -18 mEq had very acidic blood. That acidic condition would increase cells of intenstines tissue's sensitivity to paratyroid hormone (PTH) so the absorption of Ca^{2+} on intestines increased. In addition, acidic condition increased synthesis of 1.25 dihydroxyvitamin D3 from 25 hydroxyvitamin D3 by 1α -hydroxylase

enzyme in the kidneys so increasing reabsorption of Ca^{2+} from glomerular filtrate. Therefore, ewes offered with rations with DCAD value of -28 and -18 mEq had more plasma's Ca^{2+} than those offered with other DCAD values. Block (1994) stated that the increase of ration's anion, would increase reabsorption of osteoclastic bones and increase synthesis of 1.25 dihydroxyvitamin D3 regulated by PTH. Parathyroid hormone also regulated reabsorption of Ca^{2+} and HPO_4^{2-} .

Ewes offered with rations with different DCAD values performed ratio $\text{Ca}^{2+} : \text{P}^{2-}$ inside the blood therefore obtained different number of ratio, depended on cells of intestines tissue's sensitivity to PTH, 1.25 dihydroxyvitamin D3, and PTH utilization depended on its blood acidity. Experimented ewes offered with rations with various DCAD values had ratio of plasma's $\text{Ca}^{2+} : \text{P}^{2-}$ ranging from 1.3 : 1.0 to 2.2 : 1.0 (Table 2). Ewes offered with rations with DCAD values of -28 and -18 mEq had the highest ratio of plasma's $\text{Ca}^{2+} : \text{P}^{2-}$ (2.2:1.0) than ewes offered with rations with other DCAD values. Ratio of normal plasma's $\text{Ca}^{2+} : \text{P}^{2-}$ balance was 2 : 1 (Georgievskii et al. 1982). If the ratio of plasma's $\text{Ca}^{2+} : \text{P}^{2-} < 2 : 1$, it was likely that the livestock would have milk fever especially dairy cattle with high dairy production. Therefore, rations supplies at late months pregnancy (dry condition) could act as prevention to milk fever. The amount of plasma's Ca^{2+} was highly related to DCAD value ($r = -0.59$) and less related to plasma's K ($r = 0.47$).

Block (1994) explained that low DCAD value could reduce hypokalemia peripartum by the increase of Ca^{2+} ion in blood and urine. Addition of Ca^{2+} in blood was caused by the decreasing DCAD value (low) which causing Ca^{2+} had homeostatis by increasing absorption on intestines so that also increasing the secretion (Schonewille et al. 1994, Roche et al. 2003b). Findings of this research were appropriate with ideas of Moore et al. (2000), Roche et al. (2003), and Charbonneau et al. (2006) who stated that the decrease of DCAD value would cause increase of blood Ca^{2+} . Relation between plasma's Ca^{2+} and plasma's K was explained by Yingst et al. (2001) that K^+ ion increased the pump of Na^+ in order to increase the concentration of Ca^{2+} so that free Ca^{2+} in blood was increased by the pump of Na^+ in some cells. In the other part, polarization affected

the decrease of K^+ concentration in order to perform K^+ balance (Quinn et al., 1987).

The DCAD values had no effect ($P > 0.05$) on plasma's P^{2-} . The differences of DCAD values consumed by garut ewes had no effect on plasma's P^{2-} ($P > 0.05$). It meant that, the ewes were trying to perform homeostatis. Average values of P^{2-} of experimented ewes' plasma were ranging 211 ± 44 to 473 ± 27 ppm. The amount of plasma's P^{2-} was quite highly related to DCAD value ($r = 0.67$) but not related to blood pH ($r = 0.17$).

In this research, the order of plasma's mineral from the most to the less were Na^+ which then followed by Cl^- , K^+ , Ca^{2+} , P^{2-} , and S^{2-} . Isnaeni (2006) stated that in extracellular fluid or blood plasma the order of mineral from the most to the less were Na^+ , Cl^- , K^+ , Ca^{2+} , P^{2-} , and Mg^{2+} .

Urinary Mineral Status

The DCAD values had high effect ($P < 0.01$) on urinary pH. Ewes offered with rations with DCAD values of -28 and -18 mEq had the lowest urinary pH ($P < 0.05$) each of 5.73 ± 0.20 and 5.84 ± 0.27 , because in rations with DCAD values of -28 and -18 mEq there were addition of $CaCl_2$ and $CaSO_4$ anionic minerals.

Table 3. Average values of pH, Na^+ , K^+ , Cl^- , S^{2-} , Ca^{2+} , and P^{2-} in urine of garut ewes offered with different DCAD

| Variables | Dietary cation-anion difference (mEq) | | | | |
|-----------------|---------------------------------------|-------------------|-------------------|-------------------|-------------------|
| | -28 | -18 | 0 | +14 | +32 |
| pH | 5.73 ± 0.20^b | 5.84 ± 0.27^b | 7.60 ± 0.51^a | 7.51 ± 0.78^a | 8.28 ± 0.33^a |
| Na^+ (ppm) | 68 ± 10 | 94 ± 43 | 121 ± 82 | 460 ± 661 | 907 ± 734 |
| K^+ (ppm) | 27397 ± 8162 | 33039 ± 6704 | 21258 ± 8874 | 36697 ± 16258 | 39895 ± 12109 |
| Cl^- (ppm) | 2374 ± 2528 | 2870 ± 3014 | 1400 ± 1329 | 2192 ± 778 | 3072 ± 990 |
| S^{2-} (ppm) | 977 ± 456 | 1274 ± 676 | 630 ± 574 | 117 ± 117 | 66 ± 23 |
| Ca^{2+} (ppm) | 2295 ± 1733 | 2119 ± 1951 | 224 ± 323 | 272 ± 118 | 17 ± 27 |
| P^{2-} (ppm) | 120 ± 67^b | 95 ± 30^b | 96 ± 36^b | 203 ± 93^b | 407 ± 191^a |

Description : values with different letters on same row mean different ($P < 0.05$)

It known that body of livestock perform homeostatis so the excess of Cl^- and S^- were secreted outside the body. Secretion of S^- dan Cl^- excess were through urine so the urine would become more acidic because Cl^- dan S^- were acidic. Value of urinary pH was the picture of cation-anion of consumed rations. If the livestock

consumed excessive anion, its urine will be acid. Otherwise, if the Na^+ was consumed excessively, urinary pH will become base. On Table 6, in rations with DCAD values of -28 and -18 mEq there were many addition of acidic anionic salts, while in rations with DCAD value of +32 mEq there were many addition of base cationic salts. So ewes offered with rations with DCAD values of -28 and -18 mEq had the lowest urinary pH than those offered with other DCAD values, but ewes offered with rations with DCAD values of +32 mEq had the highest urinary pH. This had been explained by Chan et al. (2006) that the decrease of urinary pH was reflection of the effect from anion contained in the rations. Urinary pH values caused by consuming rations with DCAD value of -28 and -18 mEq were 5.73 ± 0.20 dan 5.84 ± 0.27 , respectively; those urinary pH were acid because the urinary pH were <6.0 . Ewes offered with rations with DCAD values of 0, +14, and +32 mEq had urinary pH of 7.60 ± 0.51 , 7.51 ± 0.78 , and 8.28 ± 0.33 , respectively; those urinary pH were normal because the urinary pH were between 7.50 – 8.50. Pratas (1992) stated that acidic urinary pH was <7.50 ; normal urinary pH was between 7.50 – 8.50; base urinary pH was >8.50 . Moore et al. (2000) reported that if urinary pH was lower than 6.0, than the rations offered contained excessive anionic salts. Based on those facts above, the addition of anionic salts into the rations with DCAD values of -28 dan -18 were excessive. Low urinary pH showed blood pH was also low (Vagnoni dan Oetzel, 1998). This matched with this research, ewes offered with rations with DCAD value of -28 mEq had very acidic blood as well as the urinary pH. The urinary pH was highly related to DCAD value ($r = 0.89$). Roche et al. (2002) reported that the increase of K^+ concentration in rations would increase urinary pH. The increase of K^+ concentration in rations meant that there was an increase of DCAD value. This was similar to the findings by West et al. (1992), Moore et al. (2000), Riond (2001), Dersjant-Li et al. (2002), Roche et al. 2003, Borucki Castro et al. (2004), Hu and Murphy (2004), Roche et al. (2005), Apper-Bossard et al. (2006), Charbonneau (2006), and Kienzle et al. (2006); they stated that the increase of DCAD value would also increase urinary pH.

Body would always balancing its body fluids, in this case, the one that took role was kidneys. The excess of anion or cation carried by rations would be

regulated by kidneys to be secreted through urine. Renal tubular cells responded directly to the changes in blood pH and intracellular pH. Kalium ion was moving from cells into the blood by releasing H^+ . The body cells pumped the excess of H^+ ion into the urine. Ion of H^+ caused the decrease of pH. In this research, changes in urinary pH were matched with its rations' cation-anion balance. Escobasa et al. (1984) reported that the increase of Cl^- consumption on cattle would decrease its urinary pH.

The DCAD values had no effect ($P > 0.05$) on urinary Na^+ . The average values of experimented ewes' urinary Na^+ were ranging from 68 ± 10 to 907 ± 734 ppm (Table 3). Although the urinary Na^+ was not affected by DCAD value, but it close related ($r = 0.66$) with DCAD value. In Rations with DCAD value of +32 mEq there was addition of Na_2CO_3 salt, but there were no differences in plasma's Na^+ and urinary Na^+ ($P > 0.05$) and normal urinary pH, so it meant that the addition of that Na_2CO_3 was not excessive.

The DCAD values had no effect ($P > 0.05$) on urinary K^+ . The average values of experimented ewes' urinary K^+ were ranging from 21258 ± 8874 to 39895 ± 12109 ppm (Table 3). Urinary K^+ content was not too related to DCAD ($r = 0.35$). In Rations with DCAD value of +32 mEq there was addition of K_2CO_3 salt, but there were no differences in plasma's K^+ and urinary K^+ ($P > 0.05$) and normal urinary pH, so it meant that the addition of that K_2CO_3 was not excessive.

The DCAD values had no effect ($P > 0.05$) on urinary Cl^- . The average values of experimented ewes' urinary Cl^- were ranging from 1400 ± 1329 to 3072 ± 990 ppm (Table 3). Urinary Cl^- content was not related to DCAD ($r = 0.06$), Cl^- consumption ($r = 0.04$), and blood Cl^- ($r = 0.11$).

The DCAD values had no effect ($P > 0.05$) on urinary S^{2-} . The average values of experimented ewes' urinary S^{2-} were ranging from 66 ± 23 to 977 ± 456 ppm (Table 3). Urinary Cl^- content was highly related to DCAD ($r = -0.72$).

The DCAD values had no effect ($P > 0.05$) on urinary Ca^{2+} . The average values of experimented ewes' urinary Ca^{2+} were ranging from 17 ± 27 to 2295 ± 1733 ppm (Table 3). Urinary Ca^{2+} content was highly related to DCAD ($r = -0.65$). Roche (1999) stated that there was an increase in absorption of Ca^{2+} and secretion of Ca^{2+} in dairy cattle, if anionic salts were added in its DCAD rations.

The DCAD values had no effect ($P > 0.03$) on urinary P^{2-} . The lowest urinary P^{2-} contents ($P < 0.05$) were found in ewes offered with rations with DCAD values of -28, -18, 0, and +14 mEq each of 120 ± 67 , 95 ± 30 , 96 ± 36 , and 203 ± 93 ppm, respectively, while the highest value ($P < 0.05$) was found in ewes offered with rations with DCAD values of +32 mEq, which was 407 ± 191 ppm. This was because urinary P^{2-} was highly related ($r = 0.82$) to DCAD value. Secretion through urine was the main homeostatis in the regulation of Na^+ , K^+ , and Cl^- (Maltz and Silanikove 1996) in order to maintain constant $Na^+ : K^+$ ratio in extracellular fluids. On Table 2 appeared that ratio plasma's $Na^+ : K^+$ was close to normal balance (as much as 20 : 1) in rations with DCAD values of -18, 0, +14, and +32 mEq. It meant that the ewes consuming rations with DCAD values of -18, 0, +14, and +32 mEq were able to perform regulation of minerals control inside thier blood to be homeostatis, and some excessive minerals would be secreted through urine. Bannink et al. (1999) and Nennich et al. (2006) stated that that urine secretion was directly related with consumption of Na^+ , K^+ , and N. Maltz and Silanikove (1996) explained that urine secretion was the main method of homeostatis regulation. Furthermore, according to Price and Wilson (1995), minerals filtrated by kidneys were Na^+ , K^+ , Ca^{2+} , Mg^{2+} , and Cl^- .

CONCLUSION

Rations with DCAD values of -18, 0, +14, and +32 mEq in garut ewes resulting normal ratio of plasma's $Na^+ : K^+$ and were able to perform regulation of minerals control inside their blood to be homeostatis, and some excessive minerals would be secreted through urine. Rations with DCAD values of -28 and -18 mEq in garut ewes resulting highest ratio of plasma's $Ca^{2+} : P^{2-}$ which was 2.2 : 1.0, so it could be used as an action to prevent milk fever.

REFERENCES

- Apper-Bossard, E., J. L. Peyraud, P. Faverdin, and F. Meschy. 2006. Changing dietary cation-anion difference for dairy cows fed with two contrasting levels of concentrate in diets. *J. Dairy Sci.* 89:749–760.
- Bannink, A., H. Valk, and A.M. Van Vuuren. 1999. Intake and excretion of sodium, potassium, and nitrogen and the effects on urine production by lactating dairy cows. *J. Dairy Sci.* 82: 1008–1018.
- Block, E. 1994. Manipulation of dietary cation-anion difference on nutritionally related production diseases, productivity, and metabolic responses of dairy cows. *J. Dairy Sci.* 77:1437–1450.
- Borucki-Castro, S. I., L. E. Phillip, V. Girard, and A. Tremblay. 2004. Altering dietary cation-anion difference in lactating dairy cows to reduce phosphorus excretion to the environment. *J. Dairy Sci.* 87:1751–1757.
- Charbonneau, E., D. Pellerin, and G. R. Oetzel. 2006. Impact of lowering dietary cation-anion difference in nonlactating dairy cows: a meta-analysis. *J. Dairy Sci.* 89:537–548.
- Desjant-Li Y., M. W. A. Verstegen, A. Jansman, H. Schulzes, J. W. Schrama, and J. A. Verreth. 2002. Changes in oxygen content and acid-base balance in arterial and portal blood in response to the dietary electrolyte balance in pigs during a 9-h period after a meal. *J. Anim. Sci.* 80:1233–1239.
- Escobosa, A., C. E. Coppock, L. D. Rowe Jr, W. L. Jenkins, and C. E. Gates. 1984. Effects of dietary sodium bicarbonate and calcium chloride on physiological responses of lactating dairy cows in hot weather. *J. Dairy Sci.* 67: 574–584.
- Fatmali, F., T. Toharmat, I. G. Permana, & A. Boediono. 2008. Keasaman Cairan Tubuh dan Ratio Kelamin Anak Domba Garut (*Ovis aries*) yang Diberi Kation-Anion Ransum yang Berbeda. *Media Peternakan* 31 (2) : 87 – 98.
- Georgievskii, V. I., B. N. Annenkov, and V. T. Samokhin. 1982. *Mineral Nutrition of Animals*. London, Boston: Butterworths.
- Hurst, R. L., J. P. Goff, T. A. Reinhardt, and D. R. Buxton. 1997. Strategies for preventing milk fever in dairy cattle. *J. Dairy Sci.* 80 :1269 – 1280.
- Hu, W., and M. R. Murphy. 2004. Dietary cation-anion difference effects on performance and acid-base status of lactating dairy cows: a meta-analysis. *J. Dairy Sci.* 87:2222–2229.
- Ismaeni, W. 2006. *Fisiologi Hewan*. Yogyakarta : Penerbit Kanisius.
- Kienzle E., K. Stumer, D. Ranz, and M. Clauss. 2006. A high roughage / concentrate ratio decreases the effect of ammonium chloride on acid-base balance in horses. *J. Nutr.* 136:2048S– 2049S.
- Maltz, E. and N. Silanikove. 1996. Kidney function and nitrogen balance of high yielding dairy cows at the onset of lactation. *J. Dairy Sci.* 79:1621–1626.
- Manjik, A. A. and I. M. Sumertajaya. 2006. *Perancangan Percobaan dengan Aplikasi SAS dan MINITAB*. Bogor: IPB Press.

- Moore, S.J., M.J. VanderHaar, B.K. Sharma, T.E. Pilbeam, D.K. Beede, H.F. Bucholtz, J.S. Liesman, R.L. Horst & J.P. Goff. 2000. Effects of altering dietary cation-anion difference on calcium and energy metabolism in peripartum cow. *J Dairy Sci* 83 : 2095-2104.
- Nennich, T. D., J. H. Harrison, L. M. VanWieringen, N. R. St- Pierre, R. L. Kincaid, M. A. Wattiaux, D. L. Davidson, and E. Block. 2006. Prediction and evaluation of urine and urinary nitrogen and mineral excretion from dairy cattle. *J Dairy Sci* 89: 353 –364.
- [NRC] National Research Council. 1985. Nutrient Requirements of Sheep. 6th Ed. Washington DC: National Academy Press.
- Odongo, N. E., O. AlZahal, M. I. Lindinger, T. F. Duffied, E.V. Valdes, S. P. Terrell, and B. W. McBride. 2006. Effects of mild heat stress and grain challenge on acid-base balance and rumen tissue histology in lambs. *J. Anim. Sci.* 84:447–455.
- Pratas, R. G. 1992. Influence of sodium, potassium, and chloride on the mineral metabolism, acid-base status and salivation in ruminants [disertation]. Denmark: Graduate School at Department of Animal Science and Animal Health Division of Animal Nutrition. The Royal Veterinary and Agricultural University Copenhagen.
- Price, S. A. and L. M. Wilson. 1995. Patofisiologi: Konsep Klinis Proses-proses Penyakit. Alih bahasa Anugerah P. Editor Wijaya C. Jakarta: Penerbit Buku Kedokteran EGC.
- Quinn, S. J., M. C. Cornwall, and G. H. Williams. 1987. Electrophysiological responses to angiotensin II of isolated rat adrenal glomerosa cells. *Endocrinology* 120:1581–1586.
- Rioud, J. L. 2001. Animal nutrition and acid-base balance. *Eur. J. Nutr.* 40:245–254.
- Roche, J. R. 1999. Dietary cation-anion difference for grass-fed dairy cows. [disertation]. Univ. College, Dublin.
- Roche, J. R., D. Dalley, P. Moate, C. Grainger, M. Rath & F.O. Mara. 2003. Dietary cation-anion difference and the health and production of pasture-fed dairy cows. 2. Dairy cows in early lactation. *J Dairy Sci* 86:979–987.
- Roche, J. R., J. Morton, and E. S. Kolver. 2002. Sulfur and chlorine play a non-acid base role in periparturient calcium homeostasis. *J. Dairy Sci.* 85:3444–3453.
- Roche, J. R., S. Petch, and J. K. Kay. 2005. Manipulating the dietary cation-anion difference via drenching to early-lactation dairy cows grazing pasture. *J. Dairy Sci.* 88: 264 –276.
- Schonewille, J. T., A. T. Van't Klooster, A. Dirksdagwe, and A. Bayen. 1994. Stimulatory effect of an anion (chloride)-rich ration on apparent calcium absorption in dairy cows. *Livest Prod. Sci.* 40:233–240.
- Stewart, P. A. 1983. Modern quantitative acid-base chemistry. *Can. J. Physiol. Pharmacol.* 61:1444–1461.
- Vagnoni, D. B. and G. R. Oetzel. 1998. Effects of dietary cation-anion difference on the acid-base status of dry cows. *J. Dairy Sci.* 81:1643–1652.

- Weidenseiner, J. M. *et al.* 2004. Acute metabolik acidosis: characterization and diagnosis of the disorder and the plasma potassium response. *J Am Soc Nephrol* 15:1589-1796.
- West, J. W., K. D. Haydon, B. G. Mullinix, and T. G. Sandifer. 1992. Dietary cation-anion and cation source effects on production and acid-base status of head stressed cows. *J. Dairy Sci.* 75:2776-2786.
- Wodnicka-Tomaszewska, M., I. K. Utama, I.G. Putu, and T. D. Chaniago. 1991. Reproduksi, Tingkah laku, dan Produksi Ternak di Indonesia. Diterbitkan atas kerjasama Direktorat Jenderal pendidikan Tinggi dan International Development Program of Australian Universities and Colleges. Jakarta: Gramedia Pustaka Utama.
- Yingst, D. R., J. Davis, and R. Schiebinger. 2001. Effects of extracellular calcium and potassium on the sodium pump of rat adrenal glomerulosa cells. *Am. J. Physiol. Cell Physiol.* 280:C119-C125.



**FACULTY OF ANIMAL SCIENCE
BOGOR AGRICULTURAL UNIVERSITY
INDONESIA**

Certificate

is presented to

Dr. Farida Fathul

for active contribution as
Poster Presenter

**in The First International Seminar on Animal Industry
Bogor, November 23-24, 2009**



Dr. Ir. Luki Abdullah, M.Sc.Agr.
Dean of Faculty of Animal Science