## AGRIVITA JOURNAL OF AGRICULTURAL SCIENCE ACCREDITED BY DIRECTORATE GENERAL OF HIGHER EDUCATION THE MINISTRY OF EDUCATION AND CULTURE, REPUBLIC OF INDONESIA No. 81/DIKTI/Kep/2011

**FEBRUARI-2015** 

VOLUME 37

NUMBER: 1

ISSN NO. 0126- 0537

TABLE OF CONTENTS	
Isolation and Characteristic of Nitrogen-Fixing Bacteria and Phosphate-Solubilizing Bacteria from Soil High in Mercury in Tailings and Compost Areas of Artisanal Gold Mine Yulia Nuraini, Novi Arfarita, and Bambang Siswanto	1
Progress from Selection of Some Maize Cultivars' Response to Drought in the Derived Savanna of Nigeria O.J. Olawuyi, O.B. Bello, C.V. Ntube, and A.O. Akanmu	8
An Effective Method for DNA Extraction of Mature Leaf of Sapodilla ( <i>Manilkara zapota</i> (L.) van Royen) Vega Kartika Sari, and Rudi Hari Murti	18
The Growth and Yield of Paddy Ciherang Planted in Dry and Rainy Season and Fertillized With Organic and Inorganic Fertilizers <i>Mudji Santosa and Agus Suryanto</i>	24
An Interaction Model Between Environmental Factors and Black Rice Growth in Irrigated Organic Paddy Field Budiman and Endang Arisoesilaningsih	30
Exogenous Application of Tryptophan and Indole Acetic Acid (IAA) to Induce Root Nodule Formation and Increase Soybean Yield in Acid, Neutral and Alkaline Soil Sudadi and Suryono	37
Estimates of Genetic Variance Component of an Equilibrium Population of Corn Hamirul Hadini, Nasrullah, Taryono, and Panjisakti Basunanda	45
In Vitro Shoot Regeneration of Indonesian Bananas ( <i>Musa</i> Spp.) cv. Ambon Kuning and Raja Buld, Plantlet Acclimatization and Field Performance <i>Yusnita, Ekawati Danial, and Dwi Hapsoro</i>	51
The Relationship Between Altitudes and the Contents of Protein, Carbohydrates, Lipids of Pumpkin (Cucurbita moschata) Suranto Tiiptowibisono, Tedianto, Edi Purwanto, and Prabang Sety	59
Hypovirulent Isolates of Fusarium Collected from Chili Crops in Boyolali Regency, Central Java, Indonesia Supyani and Sri Widadi	67
Pod Number and Photosynthesis as Physiological Selection Criteria in Soybean (Gycine max L. Mernin) Breeding for High Yield S.M. Sitompul, Dian Indra Sari, Erna Krisnawati, Rachmad Harri Mulia, and M. Taufiq	75
Study of Wood Sawdust with Addition of Plantation Wastes as a Growth Medium on Yields and Quality of White Oyster Mushroom Imam Mudakir and Utami Sri Hastuti	89
	and the second sec

ISSN: 0126-0537 (Printed) E-ISSN: 2302 -6766 (Electronic)

# AGRIVITA

Accredited by Decree of The Directorate General of Higher Education The Ministry of Education and Culture, Republic of Indonesia No: 81/DIKTI/Kep/2011 Indexed in Elsevier Product (Scopus), Proquest, Crossref, International Scientific Indexing (ISI), Directory of Open Access Journal (DOAJ), Google Scholar, DRJI (Directory of Research Journal Indexing) and Global Impact Factor (GIF)

Agrivita is Journal of Agricultural Science (AJAS) publishes articles in plant science such as agronomy, horticulture, plant breeding, soil-plant sciences and pest or disease-plant sciences and has been not published at least five years old. Published by Faculty of Agriculture University of Brawijaya Indonesia in collaboration with Indonesian Agronomy Association (PERAGI).

## Editor in Chief Kuswanto

Editoral Board Moch. Dawam Maghfoer Anton Muhibuddin Budi Prasetya

## **Managing Editor**

Silvia Santi Wahyuni

Banking

Andik Yudianto, Bank Negara Indonesia (BNI Bank) Branch of Malang, Acc No.0254013689 Swift Code : BNINIDJA

## **EDITORIAL ADDRESS**

Faculty of Agriculture University of Brawijaya 5<sup>th</sup> Floor Agriculture Building JI. Veteran Malang 65145 East Java Indonesia Phone/Fax :+62-341– 575743 E-mail : <u>agrivita@ub.ac.id</u>, <u>agrivitafaperta@yahoo.com</u>, <u>agrivitafaperta@gmail.com</u> Website: http://www.agrivita.ub.ac.id

## **Publication Schedule**

AGRIVITA, journal of agricultural science publish three times in a year (February, June and October)

## **Manuscript Submission**

Manuscript should be written in MS words format or other software program. Figure, illustration and picture included in the manuscript file. Send the manuscript directly to http://www.agrivita.ub.ac.id

## **Manuscript Publishing**

Feasible manuscript was determined by Editorial Board after obtaining recommendations from peer reviewer. Manuscript revision is author responsibility and manuscripts that are not feasible will be returned to the author.

## International Editorial Board

Prof. Hunsa Punnapayak (Microbiology - University of Chulalongkorn Thailand)

B. Mohan Kumar, Ph.D, FNASc, FNAAS, FNIE (Agroforestry - University of Agricultural Kerala India)

Dr. Chris Anderson (Bio-mining Remediation - University of Massey New Zealand)

Sivananda Tirumalaraju, Ph.D. (Plant Genetic and Plant Breeding - University of South Dakota State, United State of America)

> Dr. Jeroen C.J. Groot (Organic Farming - University of Wageningen The Netherlands)

Dr. A.K. Srivastava (Soil Science - National Research Centre for Citrus Indian Council of Agricultural Research India)

Prof. Bambang Guritno (Agricultural Ecology, Cassava, Multiplecropping - University of Brawijaya Indonesia)

> Prof. Wani Hadi Utomo (Land Management - University of BrawijayaIndonesia)

> > Prof. Kurniatun Hairiah (Soil Biology - University of Brawijaya Indonesia)

Prof. Eko Handayanto (Soil organic management - University of Brawijaya Indonesia)

Prof. Ika Rochdjatun Sastrahidayat (Epidemiology - University of Brawijaya Indonesia)

Editorial Board would like to thank to peer reviewer who has been invited by Agrivita Journal of Agricultural Science Volume 37 number 1, 2015

Prof. Xinbin Feng
(Institute of Geochemistry - Chinese Academy of Sciences (IGCAS))

Prof. Pablo Tittonell (Farming Systems Ecology - Wageningen UR Plant Sciences)

Dennis B. Egli, PhD (Plant and Soil Sciences - University of Kentucky, Pennsylvania State)

Wang Pu (College of Agriculture and Biotechnology - China Agricultural University)

> Dr. Buang Abdullah (Genetics Plant Breeding - Indonesian Rice Research Institute)

> Dr. Sujin Patarapuwadol (Department of Plant Pathology - Kasetsart University Thailand)

Dr. Ragapadmi Purnamaningsih (Agricultural biotechnology - Indonesian Biotechnology and Genetic Resources Research Institute) B. Mohan Kumar, PhD (Agronomy, Agroforestry and Climate Change - Indian Council of Agricultural Research (ICAR))

> Dr. Moch. Dawam Maghfoer (Horticulture - University of Brawijaya Indonesia)

Dr. Budi Prasetya (Soil Science - University of Brawijaya Indonesia)

Dr. Ali Jamil (Biology and Soil Fertility - Indonesian Rice Research Institute)

Dr. Darmawan Saptadi (Plant Breeding - University of Brawijaya Indonesia)

Panca Jarot Santoso, M.Sc. (Plant Breeding - Indonesian Tropical Fruits Research Institute)

Prof. Kuswanto (Plant Breeding - University of Brawijaya Indonesia)

Dr. Bambang Tri Rahardjo (Plant Protection - University of Brawijaya Indonesia)

Prof. Estri Laras Arumingtyas (Department of Biology - University of Brawijaya Indonesia)

Hagus Tarno, PhD (Plant Protection - University of Brawijaya Indonesia)

Farzane Porianejad (Ministry of Jihad e-Agriculture - Islamic Republic of Iran)

Dr. Noer Rahmi Ardiarini (Plant Breeding - University of Brawijaya Indonesia)

## IN VITRO SHOOT REGENERATION OF INDONESIAN BANANAS (*Musa* spp.) cv. AMBON KUNING AND RAJA BULU, PLANTLET ACCLIMATIZATIONAND FIELD PERFORMANCE

Yusnita <sup>1\*</sup>), Ekawati Danial <sup>2)</sup> and Dwi Hapsoro <sup>1)</sup>

 <sup>1)</sup> Plant Science Laboratory, Department of Agrotechnology, College of Agriculture, University of Lampung Jl. Sumantri Brojonegoro No. 1 Bandar Lampung, Lampung Indonesia
 <sup>2)</sup> Department of Agrotechnology, College of Agriculture, University of Baturaja, Jl. Ratu Penghulu No.02301 Karang Sari, Baturaja, Kab. Ogan Komering Ulu Indonesia

<sup>\*)</sup> Corresponding author Phone: +62-8128145990 E-mail: yusnita.said@yahoo.com

#### Received: August 21, 2014/ Accepted: January 30,2015

#### ABSTRACT

The aim of this study was to observe responses of two banana cultivars 'Ambon' and 'Raia Bulu' on different BA concentrations and effects of different media and fertilizer on survival and arowth of plantlets. Sterilized explants were cultured on initiation medium (MS with 1mg L BA) for 4 weeks, then subjected to media MS with 2.5, 5.0, or 7.5 mg  $L^{-1}$  BA. Numbers of shoot buds, shoots and propagules were recorded after 4 consecutive passages with 4 weeks intervals. Rooted plantlets were acclimatized in three different media, then treated with or without NPK (32:10:10) fertilizer solution once a week. After 2 months, the survival and growth of plantlets were recorded. Cultures of banana 'Ambon Kuning' showed higher regenerative capacity compared to 'Raja Bulu', producing higher numbers of shoot buds, shoots and propagules. The best medium for propagule proliferation of both banana cultivars was MS+5 mg L<sup>-1</sup> BA, producing 40.7 propagules for 'Ambon Kuning', and 12.3 propagules for 'Raja Bulu' per explant. In all acclimatization media tested, 100% of plantlet survival was achieved. The best plantlet growth was found in sand: compost (1:1,v/v) with application of NPK solutions. The in vitro-derived plants were planted in the field and produced fruits of high quality.

Keywords: acclimatization, benzyladenine, fertilizer, genotypes, media mixtures

#### INTRODUCTION

Banana (*Musa* spp.) is the fourth most important crops in the world, belonging to the Musaceae family. Indonesia is the sixth top

#### Accredited SK No.: 81/DIKTI/Kep/2011

http://dx.doi.org/10.17503/Agrivita-2015-37-1-p051-058

banana producing country, which produced 6.19 millions MT, after India (in millions MT) (24.86), China (10.55), The Philippines (9.22), Ecuador (7.01) and Brazil (6.90) (Maps of World, 2014). In Indonesia, banana ranks first, becoming the most important fruit in term of production and harvested area. Two most popular bananas in Indonesia are 'Ambon Kuning' (AAA group) and 'Raja Bulu' (AAB group). Both cultivars are commonly consumed as fresh fruits or as the raw materials for banana powder. In Indonesia, banana cultivation is conducted mostly in small scales (0.5-5 ha), with minimum inputs, but the farmer involved in this cultivation is very large, i.e., up to 21,482,000 households (Indonesia Ministry of Agriculture, 2012).

Currently, the major constrains for banana production in Indonesia are banana wilt caused by Fusarium oxysporum f.sp. cubense (Foc), bacterial blood disease and moko disease caused by Ralstonia solanacearum. These diseases damage most of banana plantation in Indonesia, causing significant economic losses, i.e., a decline in production by 60-70% (Suyamto et al., 2012). It was also reported that Black Sigatoka and Eumusae leaf spot disease infected most banana orchards in West and North Sumatera and Bengkulu, Indonesia at severity level of 15% to 62.3% (Sahlan and Soemargono, 2011). Monoculture cultivation of banana in a large area has met with difficulty for the availability of healthy and uniform suckers. Plant materials can be prepared from suckers or However, rhizome division (bit). those conventional propagation methods are prone to causing endogenous bacterial or fungal pathogen contamination of the planting materials. Plant tissue culture is a good alternative to suckers in providing disease-free and uniform plant materials of these cultivars in a large number and in a relatively short time. Established micropropagation protocols for these genotypes are needed to overcome shortage of planting materials and to synchronize harvest time for local banana growers.

Success in tissue culture of banana from various genome groups has been reported by several researchers and it was well-documented that the regenerative capacity in vitro is highly genotype specific and greatly influenced by growth regulator amended in shoot-inducing medium (Karim et al., 2009; Sipen and Davey, 2012; Rahman et al., 2013; Igbal et al., 2013). Different genome groups, especially with balbisiana (AAB, ABB or BBB groups) in their ploidy appeared to be more difficult to propagate than all acuminata (AAA) groups (Hirimburegama and Gamage, 1997; Saha-Roy et al., 2010; Sipen and Davey, 2012). Previous procedures in micropropagation of various cultivars of banana have been reported (Kalimuthu et al., 2007; Al-Amin et al., 2009; Ali et al., 2011; Hapsoro et al., 2010); however, the best medium for shoot multiplication of each cultivar was different. The objective of the present study was to investigate the response of banana cultivars 'Ambon Kuning' (AAA group) and 'Raja Bulu' (AAB group) to different benzyladenine (BA) concentrations, to develop a reproducible method of plantlet acclimatization, and to evaluate performance of the tissue-derived plants in the field.

#### MATERIALS AND METHODS

This research was conducted in Plant Tissue Culture Laboratory, Department of Agrotechnology, College of Agriculture, University of Lampung from May 2012 to January 2014. Healthy shoot buds of bananas 'Ambon Kuning' and 'Raja Bulu' were taken from public collections in Lampung Province, Indonesia. Apical buds approximately 3 x 3 x 4 cm with rhizome parts at the base were excised from rhizomes, rinsed under tap water, and soaked in a solution containing 150 mg  $L^{-1}$  ascorbic acid, 50 mg  $L^{-1}$  citric acid and 2 g  $L^{-1}$  of fungicide for 30 minutes before being surface sterilized in laminar air flow cabinet with 2.6% sodium hypochlorite (NaOCI) plus 2 drops of Tween-20 solution for 30 minutes. This was followed by aseptically trimming the buds to smaller size (approximately 1.5 x1.5 x 2 cm) and shaking them in the 0.8% NaOCI solution for 10 minutes, followed by three-time rinsing with sterile distilled water. All explants were incubated on a preconditioning medium (Figure 3a) for 14 days to ensure aseptic condition of explants after sterilization and to induce early growth. The preconditioning medium was MS (Murashige and Skoog, 1962) basal supplemented with 1 mg L<sup>-1</sup>N6salts, benzyladenine (BA) (Phyto Technology Laboratories<sup>®</sup>). Aseptic explants from the preconditioning medium were transferred into MS medium with different concentrations of BA(2.5, 5.0 or 7.5 mg L<sup>-1</sup>). The basal media consisted of MS macro and micro salts supplemented with MS vitamines, 100 mg L<sup>-1</sup> myo-inositol and 3% (w/v) sucrose. The pH of medium was adjusted to 5.8 prior to being solidified with 0.8% (w/v) agar (Swallow Globe, Indonesia) and autoclaving. Subcultures into the same growth regulatorcontaining media were done every 4 weeks. All cultures were incubated in a culture room at 26 ± 2<sup>°</sup>C under continuous light from cool-white fluorescent lamps (Phillips) at 1500 lux, and 60% relative humidity.

The experiment was set up in a completely randomized design with three replicates. Each experimental unit consisted of four culture bottles each of which contained one explant. After 20 weeks of culture, the effects of different medium on axillary buds and shoot proliferation were observed by recording the percentage of explants forming multiple shoots, average number of shoot buds ( $\leq 0.5$  cm), shoots (> 0.5 cm) and propagules (shoot buds + shoots) per explant. All data were subjected to analysis of variance (ANOVA) followed by mean separation test using least significant difference (LSD) at  $\alpha 0.05$ .

Shoot clusters of 'Ambon Kuning' in roliferation medium (Figure 1c) were separated into individual shoot and transferred to MS + 1 mg L<sup>-1</sup> BA for 4 weeks for shoot elongation and rooting. Cultures with rooted shoots of 6-7cm in length (Figure 1b) were taken out from culture room and hardened off at room temperature with diffused sunshine for 1 week before being used for subsequent acclimatization experiment. The healthy rooted plantlets were selected for uniformity in size and were washed under tap water from the sticking media, then planted in plastic pots containing one of three different potting-mix media, namely: M1:rice husk:compost (1:1;v/v), M2:rice husk:sand (1:1;v/v), M3:rice

husk:compost:sand (1:1:1;v/v). The plantlets were covered with transparent polyethylene bags for a few (3-5) days to retain moisture and were gradually acclimatized to a shaded greenhouse.

After 2 weeks in potting media, the plantlets were treated with or without fertilizer NPK (32-10-10) solution at 2 g  $L^{-1}$  weekly, 10 ml per pot. Data of the percentage of acclimatized plantlets, plant height, number of leaves, root length, and plant fresh weight were recorded at 10 weeks after acclimatization. This experiment was also set up in a completely randomized design with three replicates, each of which consisted of 10 plantlets.

Acclimatized plantlets were then transferred to bigger polyethylene bags (5 kg capacity) with soil-mix media consisting of top soil: compost (2:1,v/v). Some of these planting materials were planted in the field, maintained and grown until fruiting, and harvested after 12 months in the field.

### **RESULTS AND DISCUSSION**

Since the early culture establishment, there was a big challenge with 'Raja Bulu' (AAB group)

due to high rate of contamination and blackening of explants. In contrast, establishment of 'Ambon Kuning' (AAA group) cultures was much easier with lighter blackening. In fact, our preliminary research showed that all 'Raja Bulu' explants cultured in MS medium without growth regulator underwent severe blackening and eventually died. Therefore, to avoid the dead of 'Raja Bulu' explants, the MS without growth regulator as the control treatment was not assigned in this experiment.

In the first two passages of cultures, most 'Raja Bulu' explants produced severe black exudation compared to light blackening of 'Ambon Kuning' explants (Figure 1a, 1b). The severe blackening exudation in 'Raja Bulu' cultures remained high for up to three passages of cultures, and this apparently played an important role in inhibition of shoot bud initiation. Blackening or browning of explants in tissue culture is widely known as the oxidation of phenolic compounds resulting in the formation of quinines which are highly reactive and toxic to plant tissue (Titov *et al.*, 2006; Ko *et al.*, 2009; Ahmad *et al.*, 2013).



Figure 1. (a) 'Ambon Kuning' explants underwent less blackening; (b) severe blackening occurred in 'Raja Bulu' explants in the second passage; (c) Cultures of 'Ambon Kuning' with normal shoots in MS + 5 mg L<sup>-1</sup> BA; (d) A few shoots and many buds that failed to elongate in 'Raja Bulu' culture at the fourth passage

One way to control blackening of *in vitro*cultured banana explants was by treating the explants with 1.2 g L<sup>-1</sup> ascorbic acid, followed by addition of 100 mg L<sup>-1</sup> ascorbic acid in the medium (Ngomuo *et al.*, 2014). Treatments of explants with potassium citrate and citrate (K-C:C) as an antioxidant during explants preparation were also reported to overcome browning in plantain cultures (Onuoha *et al.*, 2011). In this experiment, explants were soaked for 30 minutes in 150 mg L<sup>-1</sup> ascorbic acid and 50 mg L<sup>-1</sup> citric acid prior to sterilization, and it appeared that this treatment was effective for 'Ambon Kuning', but less effective for 'Raja Bulu'.

The results of this experiment showed that numbers of shoot buds, shoots and propagules (buds + shoots) per explant after 16 weeks in 'Ambon Kuning' cultures in each concentration of BA being tested were much higher than those in 'Raja Bulu' cultures (Table 1). The blackening of 'Raja Bulu' cultures in BA-containing media seemed to decrease along with further subcultures to fresh culture media. As seen in Table 1, all (100%) of 'Ambon Kuning' explants produced multiple shoots on media with various concentrations of BA tested (2.5, 5.0, and 7.5 mg L<sup>-1</sup>), whereas the 'Raja Bulu' explants produced 100% multiple shoots only on media with 5 mg  $L^{-1}$ or 7.5 mg L<sup>1</sup> BA. Only 50% of 'Raja Bulu' explants produced multiple shoots on media with 2.5 mg  $L^{-1}$ BA. These data showed significant genotype effects on regenerative capacity of 'Ambon Kuning' vs 'Raja Bulu' cultivars. The number of shoots and propagules were significantly higher in 'Ambon Kuning' compared to 'Raja Bulu'.

Explants from both cultivars produced maximum number of shoots and propagules (25.6

shoots and 40.7 propagules per explants in 'Ambon Kuning', and 1.9 shoots and 12.3 propagules per explant for' Raja Bulu') in MS amended with 5 mg L<sup>-1</sup> BA. Almost similar result was reported by Saha-Roy et al. (2010), that the highest shoot proliferation among different BA concentrations tested for 'Malbhog' (AAB group) was BA 5 mg  $L^{-1}$  + 1 mg  $L^{-1}$  IAA. However, these results were different from other results reported previously. The highest shoot proliferation of 'BARI-1' banana was obtained in MS + BA 7.5 mg  $L^{-1}$  + NAA 0.5 mg  $L^{-1}$  (Al-Amin *et al.*, 2009), whereas the best medium for maximum multiplication of 'Agnishwar' banana was MS + BA 4 mg L<sup>-1</sup> (Rahman et al., 2013). Hapsoro et al. (2010) reported that the bud regenerative capacity of 'Tanduk' (horn type AAB group) was higher than that of 'Ambon Kuning' (AAA group), and the maximum shoots of 'Tanduk' was obtained in MS + 2 mg L<sup>-1</sup> BA, while that of 'Ambon Kuning' was obtained in MS + 5 mg  $L^{-1}$  BA. In addition, it was shown from the figures with respect of the proportion of shoot buds over the propagules produced, that the propagules of 'Raja Bulu' were predominated by shoot buds that failed to elongate into shoots, whereas 'Ambon Kuning' produced more normal shoots. The average length of shoots of 'Ambon Kuning' was higher than that of 'Raja Bulu', especially in lower (2.5-5.0 mg  $L^{-1}$ ) BA concentrations. Therefore, the subsequent acclimatization experiment was conducted only with 'Ambon Kuning' plantlets. The culture performance of 'Ambon Kuning' vs 'Raja Bulu' in media with 5 mg L<sup>-1</sup> BA is presented in Figure 1c and 1d, respectively.

Banana cultivars	BA concentrations (mg L⁻¹)	% of explants forming multiple shoots	Mean number of shoot buds per explant	Mean number of shoots per explant	Mean number of propagules per explant	Mean length of shoots (cm)
'Ambon	2.5	100	4.70 c	7.70 b	12.30 c	5.10 a
Kuning'	5.0	100	15.10 ab	25.60 a	40.70 a	4.70 a
(AAA)	7.5	100	18.90 a	11.90 b	30.80 b	2.60 b
'Raja	2.5	50	5.30 c	1.10 c	6.40 c	2.80 b
Bulu'	5.0	100	10.40 bc	1.90 c	12.30 c	2.70 b
(AAB)	7.5	100	5.80 c	1.30 c	7.10 c	2.10 b
LSD 0.05		-	7.75	4.77	7.19	1.46

Table 1. Effects of benzyladenine (BA) concentrations on bud, shoot and propagule proliferation of banana 'Ambon Kuning' and 'Raja Bulu' after 20 weeks of culture

Remarks: Means in the same column followed by different letters differ significantly at P ≤ 0.05 according to LSD multiple range test. All values are an average of 12 explants

These findings and the supportive literatures discussed above confirmed that responses of in vitro shoot regeneration in banana were highly influenced by genotypes as well as cytokinin concentrations. Hirimburegama and Gamage (1997) reported that bananas having genome with balbisiana (AAB, ABB or BBB groups) have poorer regenerative capacity than those of the pure acuminata (AAA group) as shown by lower axillary shoot numbers or propagule multiplication. Sipen and Davey (2012) also found that for banana 'Nangka' (AAA group), the maximum number of shoots was obtained in MS medium with 5 mg L<sup>-1</sup> BA, whereas for 'Mas' (AA) and 'Berangan' (AAA) with 6 mg L<sup>-1</sup> BA and for 'Awak' (AAB) with 7 mg  $L^{-1}$  BA. In their experiment, it was found that at the same concentration of growth regulator, namely MS + 6 mg L<sup>-1</sup> BA, the shoot number of 'Awak' (AAB) was less than those obtained in 'Nangka' (AAA), 'Mas' (AA) and 'Berangan' (AAA). Apparently, this was due to the stronger apical dominance in the AAB type compared to the AAA group as suggested by Ortiz and Vuylsteke (1994) and discussed by Saha-Roy et al. (2010). The fact that the horn type banana 'Tanduk" (AAB group) showed higher shoot bud regeneration capacity than 'Ambon Kuning' (AAA group) found by Hapsoro et al. (2010) suggests that difficulties in shoot bud

multiplication might not be solely influenced by the involvement of balbisiana (B group) in the genome. Rather, it might also be influenced by the variation of ability to re-generate among the balbisiana types. Furthermore, the difficulties to regenerate buds, shoots or propagules seem to be closely related with the blackening of explants caused by exudation of phenolic compounds.

For acclimatization experiment, 'Ambon Kuning' shoots were cultured individually in MS containing 1 mg L<sup>-1</sup> BA and 3% sucrose for further growth and rooting. After 6 weeks, when the shoots reach 6-7 cm in length all of the shoots showed satisfactory root formation (Figure 3b). These cultures of rooted shoots were transferred to a room with diffused sunshine at room temperature  $(28^{0}-29^{0}C)$  for 1 week to harden-off the shoots. After this period of hardening-off, healthy plantlets with relatively uniform size and roots were obtained and could be used for the subsequent acclimatization experiment (Figure 3c).

The results showed that after 8 weeks in *ex vitro* condition, all plantlets (100%) from all treatments were fully acclimatized (Figure 2) and ready to be transplanted to bigger polyethylene bags for full exposure to full sunshine. It was observed that both the media composition and fertilizer treatments affected growth of the plantlets.



Figure 2. Eight week-old plantlets acclimatized under different media mixture and applicant of NPK solution. M1: rice husk:compost (1:1,v/v), M2: sand:compost (1:1;v/v), M3: rice husk:compost:sand (1:1:1;v/v); P0: without NPK fertilizer, P1: with application of NPK fertilizer solution



Figure 3. (a) Primary explants; (b) micro shoots of banana 'Ambon Kuning' on medium for shoot elongation and rooting, i.e MS with 1.0 mg L<sup>-1</sup> BA; (c) plantlet acclimatization and (d) three-month-old plantlets ready to be transferred to the field



Figure 4. (a) Field performance of *in vitro*-derived banana cv. Ambon Kuning, 10 months, (b,c) 11 months in the field, (d) fruits at harvest time, and (e) ripe 'Ambon Kuning' fruits

In general, application of NPK (32:10:10) solution in acclimatized banana plantlets increased growth as shown by the increase in plant height and fresh weight. In addition, acclimatization media consisting a mixture of sand : compost (1:1,v/v) proved to be the best media for banana acclimatization since it produced the highest value of plant height, number of leaves and fresh weight, in both fertilized and without fertilizer treatment (Table 2). This finding was consistent with the previous

results reported in acclimatization of *Sansevieria trifasciata* cv. Lorentii and cv. Hahnii that were successfully *ex-vitro* rooted and acclimatized with 95.9%-100% survival in sand:compost (1:1,v/v) (Yusnita *et al*, 2011). Figure 2 shows performance of acclimatized plantlets in different media and fertilizer treatment after 8 weeks in *ex vitro* condition, while Figure 3d shows performance of tissue cultured plants in polyethylene bags under full sunshine after 12 weeks.

Fertilizer NPK (32:10:10)2 g L <sup>-1</sup>	Media mixture	Plant height (cm)	Number of leaves	Length of roots (cm)	Plant fresh weight (g)
Without Fertilizer	M1=Rice Husk Charcoal : Compost (1:1.v/v)	31.00 d	3.00 b	24.10 a	10.88 c
	M2=Sand : Compost (1:1,v/v)	40.00 b	3.20 b	25.90 a	15.45 bc
	M3=Rice Husk Charcoal: Sand:Compost (1:1:1,v/v)	33.80 c	3.10 b	24.00 a	11.27 c
With Fertilizer	M1=Rice Husk Charcoal : Compost (1:1.v/v)	41.40 b	3.30 b	20.60 a	21.30 a
	M2=Sand : Compost (1:1,v/v)	45.70 a	3.70 a	22.30 a	25.63 a
	M3=Rice Husk Charcoal: Sand:Compost (1:1:1,v/v)	44.60 a	3.30 b	19.70 a	20.85 ab
LSD 0.05		2.45	0.35	ns	5.69

Table 2. Effects of media mixture and NPK fertilizer solution on growth of banana cv. 'Ambon Kuning' plantlets after 8 weeks in *ex vitro* condition

Remarks: Means in the same column followed by different letters differ significantly at P ≤ 0.05 according to LSD multiple range test. All values are an average of 30 plantlets

When these *in vitro* derived 'Ambon Kuning' plants were planted in the field and maintained under farmer procedures of banana cultivation for 12-13 months or until harvesting time, vigorous growth of plants with 8-16 suckers were observed among tissue cultured banana plants in the field as shown in Figure 4a-c. Most plants produced flowers at 8-8.5 months after planting in the field and produced fruit bunches with 8-12 hands per bunch and 12-17 fingers/hands. The green and yellow fruits of 'Ambon Kuning' (Figure 4d, 4e) were of high quality.

#### CONCLUSIONS AND SUGGESTIONS

Banana 'Ambon Kuning' showed higher shoot and propagule multiplication capacity than 'Raja Bulu'. The best shoot and propagule proliferation medium for both banana cultivars were MS + 5 mg L<sup>-1</sup> BA. All acclimatization media tested caused 100% plantlet survival, and the mixture of sand:compost (1:1;v/v) was the best medium for plant growth. The *in vitro*-derived plants were planted in the field and produced fruits of high quality. Further research is needed to overcome severe blackening in early passages of 'Raja Bulu' cultures, and to increase its multiplication rate using stronger type (s) of cytokinin (s).

#### ACKNOWLEDGMENTS

The authors would like to thank The Directorate General of Higher Education, Ministry of National Education for funding this research through the scheme of The University Priority Research Grant in 2013.

#### REFERENCES

- Ahmad, I., T. Hussain, I. Ashraf, M. Nafees, Maryam, M. Rafay and M. Iqbal. 2013. Lethal effect of secondary metabolites on plant tissue culture. American-Eurasian J. Agric and Environ. Sci. 13(4): 539-547.
- Al-Amin, M.D., M.R. Karim, M.R. Amin, S. Rahman and A.N.M. Mamun. 2009. In vitro micropropagation of banana (*Musa* spp.). Bangladesh J. Agril. Res. 34(4): 645-659.
- Ali, A., A. Sajid, N.H. Naveed., A. Majid., A. Saleem, U.A. Khan, F.I. Jafery and S. Naz. 2011. Initiaton, proliferation and development of micro-propagation system for mass scale production of banana through meristem culture. Afr. J. Biotech. 10(70): 15731-15738.
- Hapsoro, D., M.I. Alisan, T. Ismaryati and Yusnita. 2010. Effects of benzyladenine on *in vitro* shoot multiplication of banana (*Musa paradisiaca* Linn.) cv. Ambon Kuning and Tanduk. Proceed. Int. Sem.

Hort. to Support Food Security. June 22-23, 2010 in Bandar Lampung, Indonesia.

- Hirimburegama, K. and N. Gamage. 1997. Cultivar specificity with respect to *in vitro* micropropagation of *Musa* spp. (banana and plaintain). Horti. Sci. 75: 205-211.
- Indonesia Ministry of Agriculture. Directorate General of Fruit Cultivation and Post Harvest Handling. 2012. Procedures of Banana Post Harvest Handling. Jakarta. p 93.
- Iqbal, M.M., A. Muhammad, I. Hussain and H. Bilal. 2013. Optimization of *in vitro* micropropagation protocol for banana (*Musa Sapientum* L.) under different hormonal concentrations and growth media. Intl. J Agric. Innov. Res. 2(1): 23-27.
- Kalimuthu, K., M. Saravanakumar and R. Senthilkumar. 2007. In vitro propagation of *Musa sapientum* L. (Cavendis Dwarf). Afric. J. Biotech. 6(9): 1106-1109.
- Karim, M.R., M.A. Malek, S.Rahman, M. Al-Amin and M.R. Amin. 2009. *In vitro* propagation of banana. Bangladesh J. Agric. Res. 34(2): 269-278.
- Ko, W.H., C.C. Su, C.I. Chen and C.P. Chao. 2009. Control of lethal browning of tissue culture plantlets of Cavendish banana cv. Formasana with ascorbic acid. Plant Cell Tiss. Organ Cult. 96: 137-141.
- Maps of Worlds. 2014. Top ten banana producing countries. Available at <u>www.mapsofworld.com/world-top-ten/ ba</u> <u>nana-producing-countries.html</u>. Accessed on January 18, 2015.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol. Plant. 15: 473-497.
- Ngomuo, M., E. Mneney and P. Ndakidemi. 2014. Control of browning by using ascorbic acid on shoot tip cultures of local *Musa* spp. (banana) cv. Mzuzu in Tanzania. Afric. J. Biotech. 13 (16): 1721-1725.
- Onuoha, I.C., C.J. Eze and C.I.N. Unamba. 2011. In vitro prevention of browning in plantain culture. Online J. Biol. Sci. 11 (1): 13-17.
- Ortiz, R. and D.R. Vuylsteke. 1994. Genetics of apical dominance in plantain (*Musa* spp.,

AAB group) and improvement of suckering behaviour. J. Amer Soc. Hort. Sci. 119: 1050-1053.

- Rahman, S., N. Biswas, M.M. Hassan, M.G. Ahmed, A.N.K. Mamun, M.R. Islam, M. Moniruzzaman and M.E. Haque. 2013. Micropropagation of banana (*Musa* sp.) cv. Agnishwar by *in vitro* shoot tip culture. Intl. Res. J. Biotech. 4(4): 83-88.
- Saha-Roy O., P. Bantawa, S.K. Gosh, J.A. Texeira da Silva, P. DebGhosh and T.K. Mondal. 2010. Micropropagation and field performance of 'Malbhog' (*Musa paradesiaca*, AAB group): A popular banana cultivar with high keeping quality of North East India. Tree and Forestry Sci. Biotech 4. (Special Issue 1): 52-58.
- Sahlan and A. Soemargono. 2011. Distribution and incidence leaf disease of banana in several banana production centers in North Sumatera, West Sumatera, Bengkulu and West Java. Agrivita J. Agric. Sci. 33(2): 174-181.
- Sipen, P. and M.R. Davey. 2012. Effects of N6benzylaminopurine and indoleacetic acid on *in vitro* shoot multiplication, nodulelike meristem proliferation and plant regeneration of Malaysian bananas. Trop. Life Sci. Res. 23(2): 67-80.
- Suyamto, I. Djatnika and A. Sutanto. 2012. Banana RandD in Indonesia: Updates and highlights. Advancing banana and 13: 81-88.
- Titov, S.S.K., A. Bhowmik, A. Mandal, M.S. Alam and S.N. Uddin. 2006. Control of phenolic compound secretion and effects of growth regulators for organ formation from *Musa* spp. cv. Kanthali floral bud explants. Am. J. Biochem Biotech. 2: 97-104.
- Yusnita, W. Pungkastani and D. Hapsoro. 2011. In vitro organogenesis of two Sansevieria cultivars on different concentrations of benzyladenine (BA). Agrivita J. Agric. Sci. 33(2): 147-153.