

Reprinted from the

**Advances in Differential Equations
and Control Processes**
Volume 19, Number 1, 2018, pp 37-48

**THE INTEGRAL NORMAL FORM OF A
THREE-DIMENSIONAL TRAVELING WAVE
SOLUTION MAPPING DERIVED FROM
GENERALIZED Δ -M-KdV EQUATION**

by

L. Zakaria, Notiragayu,

A. Nuryaman and S. Suharsono



Pushpa Publishing House

Vijaya Nilwas, 198 Mumfordganj

Allahabad 211002, INDIA

<http://pphmj.com/journals/adecp.htm>

de@pphmj.com & arun@pphmj.com



Information for Authors

Aims and Scope: The *Advances in Differential Equations and Control Processes* is a peer-reviewed international journal, which publishes original research papers and critical survey articles in all aspects (theory and applications) of Differential Equations and Control Processes. The *ADECP* is published in four issues per volume annually appearing in February, May, August and November.

Abstracting, Indexing and Reviews: The Emerging Sources Citation Index (ESCI), CrossRef DOIs databases, AMS Digital Mathematics Registry, ProQuest, EBSCOhost, Zentralblatt MATH, IndexCopernicus, Ulrichsweb, Google Scholar, Academic Keys, Excellence in Research for Australia (ERA), Global Impact Factor: 0.861, J-Gate, Scilit.

Submission of Manuscripts: Authors may submit their papers for consideration in the *ADECP* by the following modes:

1. **Online submission:** Please visit journal's homepage at <http://www.pphmj.com/journals/adecp.htm>
2. **Electronically:** At the e-mail address: de@pphmj.com

An effort is made to publish a paper duly recommended by a referee within a period of three months. One set of galley proofs of a paper will be sent to the author submitting the paper, unless requested otherwise, without the original manuscript, for corrections.

Abstract and References: Authors are requested to provide an abstract of not more than 250 words and latest Mathematics Subject Classification. Statements of Lemmas, Propositions and Theorems should be set in *italics* and references should be arranged in alphabetical order by the surname of the first author.

Page Charges and Reprints: Authors are requested to arrange page charges of their papers @ USD 25.00 per page from their institutions/research grants, if any. However, for authors in India this charge is Rs. 800.00 per page. Twenty-five reprints in print version and a copy in soft version are provided to the corresponding author ex-gratis. Additional sets of reprints may be ordered at the time of proof correction.

Copyright: It is assumed that the submitted manuscript has not been published and will not be simultaneously submitted or published elsewhere. By submitting a manuscript, the authors agree that the copyright for their articles is transferred to the Pushpa Publishing House, Allahabad, India, if and when, the paper is accepted for publication.

Subscription Information for 2018

Institutional Price for all countries except India

Print Subscription	€ 345.00	US\$ 465.00
--------------------	----------	-------------

Price in Indian Rs. (For Indian Institutions in India only)

Print Subscription	Rs. 6000.00
--------------------	-------------

The subscription year runs from January 1, 2018 through December 31, 2018.

Information: The journals published by the "Pushpa Publishing House" are solely distributed by the "Vijaya Books and Journals Distributors".

Contact Person: Subscription Manager, Vijaya Books and Journals Distributors, Vijaya Niwas, 198 Mumfordganj, Allahabad 211002, India; sub@pphmj.com; arun@pphmj.com

THE INTEGRAL NORMAL FORM OF A THREE-DIMENSIONAL TRAVELING WAVE SOLUTION MAPPING DERIVED FROM GENERALIZED $\Delta\Delta$ -mKdV EQUATION

L. Zakaria, Notiragayu, A. Nuryaman and S. Subarsono

Mathematics Department
Lampung University
Indonesia

Abstract

The integrability of traveling wave solution mappings can be obtained as reductions of the discrete generalized $\Delta\Delta$ modified Korteweg-de Vries ($\Delta\Delta$ -mKdV) equation. The properties of the integrable discrete dynamical system can be examined through the level set of integral function. In this paper, we show that the integral of a three-dimensional traveling wave solution mapping derived from generalized $\Delta\Delta$ -mKdV equation can be made in the normal form.

1. Introduction

Discrete integrable systems can be classified into two main classes: integrable partial difference equations and integrable ordinary difference equations. Integrable ordinary difference equations are equivalent to integrable mappings. By imposing a periodicity condition, not only integrable lattice equations can be reduced to ordinary difference equations

Received: November 9, 2017; Accepted: February 5, 2018

2010 Mathematics Subject Classification: 37J10, 37J35, 39A10, 70K43.

Keywords and phrases: traveling wave solution, generalized $\Delta\Delta$ -mKdV, integral.

(or mappings/maps) [3, 6, 2, 1] but also there is a connection between the two classes so that many integrable maps can be obtained from integrable partial difference equations [5, 4, 6]. In this paper, we study the integrability of the discrete generalized $\Delta\Delta$ modified Korteweg-de Vries ($\Delta\Delta$ -mKdV) equation.

Consider the standard $\Delta\Delta$ -mKdV equation on the 2D lattice (\mathbb{Z}^2) defined as [6]

$$q(V_{l,m+1}V_{l+1,m+1} - V_{l,m}V_{l+1,m}) = p(V_{l+1,m}V_{l+1,m+1} - V_{l,m}V_{l,m+1}), \quad (1)$$

where the V fields are defined on the lattice sides of the $l, m \in \mathbb{Z}$, which are two discrete variables. Suppose $\xi_{l,m}(k)$ denotes a vector containing a wave function that depends on a spectral parameter k . The above equation can be derived through the following mappings:

$$\begin{aligned} \xi_{l+1,m}(k) &= \frac{1}{p-k} M_{l,m}^{hor} \xi_{l,m}(k), \\ \xi_{l,m+1}(k) &= \frac{1}{q-k} M_{l,m}^{vert} \xi_{l,m}(k) \end{aligned}$$

with

$$M_{l,m}^{hor} = \begin{pmatrix} p & -V_{l+1,m} \\ -\left(\frac{k^2}{V_{l,m}}\right) & p\left(\frac{V_{l+1,m}}{V_{l,m}}\right) \end{pmatrix}$$

and

$$M_{l,m}^{vert} = \begin{pmatrix} q & -V_{l,m+1} \\ -\frac{k^2}{V_{l,m}} & q\frac{V_{l,m+1}}{V_{l,m}} \end{pmatrix}$$

which are Lax pair matrices. These mappings are well defined when the

following condition is satisfied:

$$(M_{l+1,m}^{vert} M_{l,m}^{hor} - M_{l,m+1}^{hor} M_{l,m}^{vert}) \xi_{l,m} = 0 \quad (2)$$

for all $(l, m) \in \mathbb{Z}^2$. This condition is known as *compatibility condition*.

Tuwankotta and Quispel [7] have made the derived discrete dynamic system through the effort of multiplying the parameters on the Lax matrix system. It is intended that the system's integrity properties are always maintained in examining more dynamics occurring from a discrete dynamic system (see [7] and [8]). By the same procedure, the following shows the attempt to elaborate (1).

Consider the following Lax pair matrices:

$$P_{l,m}^{hor} = \begin{pmatrix} \alpha_1 p & -\alpha_2 V_{l+1,m} \\ -\alpha_3 \left(\frac{k^2}{V_{l,m}}\right) & \alpha_4 p \left(\frac{V_{l+1,m}}{V_{l,m}}\right) \end{pmatrix}$$

and

$$P_{l,m}^{vert} = \begin{pmatrix} \beta_1 q & -\beta_2 V_{l,m+1} \\ -\beta_3 \frac{k^2}{V_{l,m}} & \beta_4 q \frac{V_{l,m+1}}{V_{l,m}} \end{pmatrix}. \quad (3)$$

Based on *compatibility condition*, the following four nonlinear equations will be obtained:

$$\begin{aligned}
 & \left\{ \begin{aligned}
 & k^2(\alpha_3\beta_2 - \alpha_2\beta_3)V_{1+l,1+m} = 0 \\
 & -k^2(\alpha_3\beta_2 - \alpha_2\beta_3) = 0 \\
 & p\alpha_1\beta_2V_{l,m}V_{l,1+m} - q\alpha_2\beta_1V_{l,m}V_{1+l,m} \\
 & + q\alpha_2\beta_4V_{l,1+m}V_{1+l,1+m} - p\alpha_4\beta_2V_{1+l,m}V_{1+l,1+m} = 0 \\
 & -k^2(p\alpha_1\beta_3V_{l,m}V_{l,1+m} - q\alpha_3\beta_1V_{l,m}V_{1+l,m}) \\
 & -k^2(p(q\alpha_3\beta_4V_{l,1+m}V_{1+l,1+m}) \\
 & -k^2(p\alpha_4\beta_3V_{1+l,m}V_{1+l,1+m})) = 0.
 \end{aligned} \right. \quad (4)
 \end{aligned}$$

To be consistent with each other, the parameters α_j and β_j with $j = 1, 2, 3, 4$ in equation (4) must be consistent.

As a result, the first two equations are obtained

$$\alpha_3\beta_2 - \alpha_2\beta_3 = 0. \quad (5)$$

In addition, from the last two equations in (4) obtained:

$$q(\alpha_3\beta_2 - \alpha_2\beta_3)(\beta_1V_{l,m}V_{1+l,m} - \beta_4V_{l,1+m}V_{1+l,m}) = 0. \quad (6)$$

From the relationship (5), equation (6) becomes consistent when $\alpha_2 = \alpha_3$ and $\beta_2 = \beta_3$. Thus, the system with the Lax matrix (3) will be consistent if $\alpha_2 = \alpha_3$ and $\beta_2 = \beta_3$. For example,

$$(\alpha, \beta) = (\alpha_1, \alpha_2, \alpha_2, \alpha_4, \beta_1, \beta_2, \beta_2, \beta_4).$$

As a result, the generalized Lax matrices for the system (1) can be written as

$$P_{l,m}^{hor} = \begin{pmatrix} \alpha_1 p & -\alpha_2 V_{1+l,m} \\ -\alpha_2 \left(\frac{k^2}{V_{l,m}} \right) & \alpha_4 p \left(\frac{V_{1+l,m}}{V_{l,m}} \right) \end{pmatrix}$$

and

$$P_{l,m}^{vert} = \begin{pmatrix} \beta_1 q & -\beta_2 V_{l,m+1} \\ -\beta_2 \frac{k^2}{V_{l,m}} & \beta_4 q \frac{V_{l,m+1}}{V_{l,m}} \end{pmatrix}.$$

By substituting $P_{l,m}^{hor}$ and $P_{l,m}^{vert}$ into compatibility condition (2), we get the form of mappings derived from the generalized equation $\Delta\Delta$ -mKdV which is a part of the four-parameter family mapping:

$$0_1 V_{l,m} V_{l,m+1} - 0_2 V_{1+l,m} V_{1+l,m+1} - 0_3 V_{l,m} V_{1+l,m} + 0_4 V_{l,m+1} V_{1+l,m+1} = 0 \quad (7)$$

with $\theta_1 = \alpha_1\beta_2\rho$, $\theta_2 = \alpha_4\beta_2\rho$, $\theta_3 = \alpha_2\beta_1q$ and $\theta_4 = \alpha_2\beta_4q$.

2. Main Result

A. The traveling wave solution of the generalized $\Delta\Delta$ -mKdV equation

In 2016, Zakaria and Tuwankotta described a procedure to derive a two-dimensional traveling wave solution mapping of $\Delta\Delta$ -sine Gordon equation [8]. Using the same procedure, a two-dimensional traveling wave solution mapping of $\Delta\Delta$ -mKdV equation (7) can be obtained.

Consider the following discrete traveling wave solution relationship:

$$V_{l,m} = V_n, \text{ where } n = z_1 l + z_2 m$$

with z_1 and z_2 being relatively prime integers. Substituting these terms into (7), we obtain

$$0_1 V_n V_{n+z_1} - 0_2 V_{n+z_1} V_{n+z_1+z_2} - 0_3 V_n V_{n+z_1} + 0_4 V_{n+z_2} V_{n+z_1+z_2} = 0. \quad (8)$$

Equation (8) is a form of the traveling wave solution derived from the generalized $\Delta\Delta$ -mKdV equation. For the fixed values of z_1 and z_2 , equation

We call equation (10) as *three-dimensional mapping* derived from generalized $\Delta\Delta$ -mKdV equation.

Consider equation (10). Then we denoted ζ_n as the sequence in \mathbb{R}^2 which is defined as

$$\zeta_n = \begin{pmatrix} V_{n+2} \\ V_{n+1} \\ V_{n+1} \\ V_n \end{pmatrix}$$

and by θ a parameter vector in $\mathbb{R}^4 : (\theta_1, \theta_2, \theta_3, \theta_4)$.

Therefore, we can express the mapping as the two-dimensional mapping, i.e.:

$$\zeta_{n+1} = g_0(\zeta_n), \tag{11}$$

where

$$g_0 : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ (W_1, W_0) \mapsto \begin{pmatrix} -1 & \mathbb{R}^2 \\ W_1 W_0 & (\theta_3 W_1 - \theta_1) \\ & (\theta_2 W_1 - \theta_4) \end{pmatrix} W_1$$

with $W_0 = \frac{V_{n+1}}{V_n}$ and $W_1 = \frac{V_{n+2}}{V_{n+1}}$.

$\zeta_{n+1} = g_0(\zeta_n)$ has an integral. In other words, there is a function $S : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $S(\zeta_{n+1}) = S(\zeta_n)$ for all natural number n . The procedure to obtain the integrals in more detail is described in [6].

For the mapping (10), the integral can be explicitly computed from the monodromy matrix below:

$$\begin{pmatrix} \beta_1 q & -\beta_2 V_2 \\ -\beta_2 k^2 & \beta_4 q V_2 \\ -V_0 & V_0 \end{pmatrix}^{-1}$$

(8) is a mapping of $\mathbb{R}^{z_1+z_2} \rightarrow \mathbb{R}^{z_1+z_2}$. It can be checked that equation (8) is invariant for a transformation $z_1 \rightarrow -z_1, \theta_1 \rightarrow -\theta_2$, and $z_1 \leftrightarrow z_2$. In addition, it also fulfills the periodic nature, namely $(i+z_2, j-z_1)$.

Equation (8) is equivalent to the following mapping:

$$V'_{z_1+z_2-1} = \frac{V_0(\theta_3 V_{z_1} - \theta_1 V_{z_2})}{(\theta_4 V_{z_2} - \theta_2 V_{z_1})} \tag{9}$$

$$V'_{z_1+z_2-2} = V_{z_1+z_2-1}$$

$$\vdots$$

$$V'_1 = V_2$$

$$V'_0 = V_1.$$

Note that the mapping in [6] can be obtained from (9) by setting $\theta_1 = \theta_2 = p$ and $\theta_3 = \theta_4 = q$.

B. The integral normal form for a mapping derived from the generalized $\Delta\Delta$ -mKdV equation: three-dimensional mapping case

Let us take $z_1 = 1$ and $z_2 = 2$. So from (9), the third order of difference equation will be obtained as follows:

$$\theta_1 V_n V_{n+2} - \theta_2 V_{n+1} V_{n+3} - \theta_3 V_n V_{n+1} + \theta_4 V_{n+2} V_{n+3} = 0$$

which is equivalent to a three-dimensional mapping:

$$V''_{n+2} = \frac{V_n(\theta_3 V_{n+2} - \theta_1 V_{n+1})}{(\theta_4 V_{n+1} - \theta_2 V_{n+2})}, \tag{10}$$

$$V'_{n+1} = V_{n+2},$$

$$V'_n = V_{n+1}.$$

$$\begin{pmatrix} \alpha_1 p & -\alpha_2 V_2 \\ -\alpha_2 k^2 & \alpha_4 p V_2 \\ \alpha_1 p & -\alpha_2 V_1 \\ \alpha_4 p V_1 & -\alpha_4 p V_0 \end{pmatrix} \begin{pmatrix} \alpha_1 p & -\alpha_2 V_1 \\ -\alpha_2 k^2 & \alpha_4 p V_1 \\ \alpha_1 p & -\alpha_2 V_0 \\ \alpha_4 p V_0 & -\alpha_4 p V_1 \end{pmatrix}. \quad (12)$$

The trace of the monodromy matrix in (12) is

$$\begin{aligned} S(W_0, W_1) &= p^2 q (\alpha_4^2 \beta_1 + \alpha_1^2 \beta_4) \\ &\quad - k^2 \left\{ \theta_1 \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + \theta_2 (W_0 + W_1) \right\} \\ &\quad - k^2 \left\{ \theta_3 \left(\frac{1}{W_0 W_1} \right) - \theta_4 (W_0 W_1) \right\}. \end{aligned} \quad (13)$$

For all $n \in \mathbb{N}$, solution ζ_n from system (11) is a level set of $S(W_0, W_1)$.

C. The normalization processing for the integral

Assume that $\theta_2 \neq 0$. Then the integral (13) can be written as

$$\tilde{S}(W_0, W_1) = \mu_1 \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) - \mu_2 (W_0 W_1) - \mu_3 (W_0 W_1) [0.5cm] \quad (14)$$

with $\mu_1 = \frac{\theta_1}{\theta_2}$, $\mu_2 = \frac{\theta_3}{\theta_2}$, $\mu_3 = \frac{\theta_4}{\theta_2}$, and

$$\tilde{S}(W_0, W_1) = \frac{S(W_0, W_1) - p^2 q (\alpha_4^2 \beta_1 + \alpha_1^2 \beta_4)}{k^2}.$$

Condition: $\mu_2 = \frac{1}{\delta^3} > 0$ where $\delta > 0$.

Let $\mu_2 = \frac{1}{\delta^3} > 0$ and $\delta > 0$. Then choose scaling $W_0 \mapsto \frac{W_0}{\delta}$ and $W_1 \mapsto$

$\frac{W_1}{\delta}$. If $\mu_3 = \delta$, $\mu_3 = -\delta$, and $\mu_3 = 0$, then based on (14), we have,

respectively:

$$S_1(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) - \left(\frac{1}{W_0 W_1} \right) - (W_0 W_1), \quad (15)$$

$$S_2(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) - \left(\frac{1}{W_0 W_1} \right) + (W_0 W_1), \quad (16)$$

$$S_3(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) - \left(\frac{1}{W_0 W_1} \right). \quad (17)$$

Condition: $\mu_2 = -\frac{1}{\delta^3} < 0$ and $\delta > 0$.

Let $\mu_2 = -\frac{1}{\delta^3} < 0$ and $\delta > 0$. Then choose scaling $W_0 \mapsto \frac{W_0}{\delta}$ and

$W_1 \mapsto \frac{W_1}{\delta}$. If $\mu_3 = -\delta$, $\mu_3 = \delta$, and $\mu_3 = 0$, then based on (14), we have, respectively:

$$S_4(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) + \left(\frac{1}{W_0 W_1} \right) - (W_0 W_1), \quad (18)$$

$$S_5(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) + \left(\frac{1}{W_0 W_1} \right) + (W_0 W_1), \quad (19)$$

$$S_6(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) + \left(\frac{1}{W_0 W_1} \right). \quad (20)$$

Condition: $\mu_2 = 0$.

$$S_7(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) - (W_0 W_1), \quad (21)$$

$$S_8(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1) + (W_0 W_1), \quad (22)$$

$$S_9(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + (W_0 + W_1). \quad (23)$$

Condition: $\theta_2 = 0$ and $\theta_4 \neq 0$.

If $\theta_2 = 0$ and $\theta_4 \neq 0$, then the integral (14) can be written as:

$$\hat{S}(W_0, W_1) = \mu_4 \left(\frac{1}{W_0} + \frac{1}{W_1} \right) - \mu_5 \left(\frac{1}{W_0 W_1} \right) - (W_0 W_1), \quad (24)$$

where $\mu_4 = \frac{\theta_1}{\theta_4}$, and $\mu_5 = \frac{\theta_3}{\theta_4}$.

Let $\mu_5 = \frac{1}{\delta_4}$ and $\delta \neq 0$. Then choose scaling $W_0 \mapsto \frac{W_0}{\delta}$ and $W_1 \mapsto \frac{W_1}{\delta}$.

Therefore, we have the following integrals:

$$S_{10}(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) - \left(\frac{1}{W_0 W_1} \right) - (W_0 W_1), \quad (25)$$

$$S_{11}(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) + \left(\frac{1}{W_0 W_1} \right) - (W_0 W_1), \quad (26)$$

$$S_{12}(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) - (W_0 W_1). \quad (27)$$

Condition: $\theta_2 = \theta_4 = 0$.

For the case of $\theta_2 = \theta_4 = 0$, we have two normal forms of integrals normal. Firstly, if $\theta_3 = \frac{1}{\delta^4}$ and $\delta \neq 0$, then the integral (14) can be written as:

$$S_{13}(W_0, W_1) = \mu \left(\frac{1}{W_0} + \frac{1}{W_1} \right) - \left(\frac{1}{W_0 W_1} \right). \quad (28)$$

Secondly, if $\theta_3 = 0$, we have other normal integral function, i.e.

$$S_{14}(W_0, W_1) = \left(\frac{1}{W_0} + \frac{1}{W_1} \right). \quad (29)$$

Note that the level sets of the integral $S(W_0, W_1)$ in (13) for all values of the parameters are completely determined by the level sets of $S_j(W_0, W_1)$, where $j = 1, 2, \dots, 14$.

3. Concluding Remark

Based on the above discussion, we conclude that the integral in three-dimensional maps derived from a $\Delta\Delta$ -mKdV equation for all values of the parameters is completely determined by fourteen integral forms. After some derivations, three-dimensional traveling wave solution mapping is expressed into two-dimensional mapping, having solution on the level sets of fourteen integral functions $S_j(W_0, W_1)$, where $j = 1, 2, \dots, 14$.

Acknowledgments

The authors would like to thank The Directorate of Research and Community Services at Kemenristek Dikti RI for funding this research through DIPA-BLU Research Grants 2017 of Lampung University and also thanks to the Head of Institute of Research and Community Services of Lampung University who supported this research. L. Zakaria acknowledges fruitful interactions with J. M. Tuwankotta in the early stages of this research.

References

- [1] P. H. van der Kamp and G. R. W. Quispel, The staircase method: integrals for periodic reductions of integrable lattice equations, *J. Phys. A: Math. Theor.* 43 (2010), 465207.
- [2] P. H. van der Kamp, Initial value problems for lattice equations, *J. Phys. A: Math. Theor.* 42 (2009), 404019.
- [3] H. W. Capel, F. W. Nijhoff and V. G. Papageorgiou, Complete integrability of Lagrangian mappings and lattices of KdV type, *Phys. Lett. A* 155 (1991), 37787.
- [4] V. G. Papageorgiou, F. W. Nijhoff and H. W. Capel, Integrable mappings and nonlinear integrable lattice equations, *Phys. Lett. A* 147 (1990), 10614.

Advances in Differential Equations and Control Processes

Editorial Board

Editor-in-Chief

K. K. Azad, India

Associate Editors

Jinde Cao, China

Weizhong Dai, USA

Zaihui Gan, China

Fatma Karakoc, Turkey

Michael A. Karls, USA

Christian Klingenberg, Germany

Nikolai Kosmatov, USA

Yansheng Liu, China

Ming Mei, Canada

Angelo B. Mingarelli, Canada

Yusuf Pandir, Turkey

Senjo Shimizu, Japan

Mingxin Wang, China

Zhitao Zhang, China

48 L. Zakaria, Notiragayu, A. Nuryaman and S. Suharsono

- [5] P. H. van der Kamp, O. Rojas and G. R. W. Quispel, Closed-form expressions for integrals of MKdV and sine-Gordon maps, *J. Phys. A: Math. Theor.* 40 (2007), 1278912798. doi:10.1088/1751-8113/40/42/S21.
- [6] G. R. W. Quispel, H. W. Capel, V. G. Papageorgiou and F. W. Nijhoff, Integrable mappings derived from soliton equations, *Physica A* 173 (1991), 243-266.
- [7] J. M. Tuwankotta and G. R. W. Quispel, Dynamics of 2-dimensional maps derived from a discrete sine-gordon equations, Unpublished, 2012.
- [8] L. Zakaria and J. M. Tuwankotta, Dynamics and bifurcations in a two-dimensional map derived from a generalized $\Delta\Delta$ -sine-Gordon equation, *Far East J. Dyn. Syst.* 28(3) (2016), 165-194.

L. Zakaria: lazakaria.1969@fmipa.unila.ac.id

Notiragayu: notiragayu@fmipa.unila.ac.id

A. Nuryaman: aangnuryaman@gmail.com

Suharsono Suharsono: suharsono.1962@fmipa.unila.ac.id