

Sperms quality of mice effectively decreased by essential oil extracted from plant tuber of nutgrass (*Cyperus rotundus*)

Hendri Busman*, Sutyarso, Mohammad Kanedi, Salman Farisi, Dzul Fitria Mumtazah

Department of Biology, Faculty of Mathematics and Sciences, University of Lampung, Bandar Lampung, Indonesia.

Correspondence: Hendri Busman, Department of Biology, Faculty of Mathematics and Sciences, University of Lampung, Bandar Lampung, Indonesia.

Email: hendri_busman@yahoo.com

ABSTRACT

Tuber extract of rumput teki (*Cyperus rotundus*), the nutgrass, has been revealed to induce apoptosis in leukemia cell lines. This effect is probably related to the essential oil content of this plant. This research aimed to determine the effect of nutgrass essential oil on the sperm quality of mice. By using a completely randomized design (CRD), test mice (n = 24) were grouped into four. Group 1 received only distilled water (control). Groups 2, 3, and 4 consecutively were treated with the essential oil at the dose of 9, 18, and 36 mg/30gBW for 35 days. By the end of treatment, the total number of sperm of mice in Groups 1 (control), 2, 3, and 4 consecutively were 51, 25.16, 16.66, and 11.83 million/ml. The sperm motility was 75.5%, 55.66%, 32.66%, and 23.33% respectively. The results revealed that the essential oil from nutgrass caused a decrease in the number and motility of sperm in mice in a dose-dependent manner. It is suggested that the essential oil extracted from the tuber of *Cyperus rotundus* is the potential to be used as antifertility.

Keywords: essential oil, rumput teki, nutgrass, *Cyperus rotundus*, sperm quality

Introduction

The plants have many applications due to their different components^[1-3]. The active component of a chemical compound contained in purple nutsedge tuber is sesquiterpenes. Sesquiterpenes in the purple nutsedge tuber include α -cyperone, β -selinene, cyperene, cyperotundone, patchoulone, sugeonol, kobusone, and isokobusone. In addition to these active components, the purple nutsedge tuber also contains several chemicals, namely alkaloids, flavonoids, tannins, starches, glycosides, and saponins^[4, 5]. Various chemicals contained in the purple nutsedge tuber can be used as an antibacterial, menstrual, and laxative medicine contraception. Contraception that affects the reproductive process, in this case, is inhibiting ovulation of the menstrual cycle in humans and the estrous cycle in mice (*Mus musculus* L.), inhibiting sperm penetration, inhibiting embryo fertilization and

implantation so that the process of pregnancy is difficult to occur and if it occurs then the possibility of the fetus will experience disability^[6].

To diagnose infertility in men can be determined through the measurement of concentration, motility, and morphology of sperm. A decrease in the concentration and motility of sperm, associated with the disability of the secretion function of Sertoli cells and Leydig cells, causes imperfections in the process of spermatogenesis and sperm maturation in the epididymis, resulting in decreased sperm motility, increased abnormalities and morphology of sperm^[7]. The number of spermatogenic cells is the number of spermatogonia cells, primary spermatocytes, secondary spermatocytes, spermatids and sperm which are located in the seminiferous tubules that indicate the process of spermatogenesis that occurs in the testes. The number of spermiogenesis cells is the number of spermatid and sperm cells. The time required for the formation of spermatogonia is 3 days, primary spermatocytes for 16 days, secondary spermatocytes 26 days, spermatids 36 days, and sperm 49 days^[8]. Sperm that form in the testes are channeled into the epididymis to undergo maturation.

Sperm motility is often used as an indicator of sperm fertility. Motility testing is performed to determine the movement of sperm. However, the movement of sperm is also influenced by the integrity of the morphological structure of sperm. The percentage of motility is the percentage of sperm that moves

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: Hendri Busman, Sutyarso, Mohammad Kanedi, Salman Farisi, Dzul Fitria Mumtazah. Sperms quality of mice effectively decreased by essential oil extracted from plant tuber of nutgrass (*Cyperus rotundus*). J Adv Pharm Edu Res 2020;10(2):1-6. Source of Support: Nil, Conflict of Interest: None declared.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

progressively forward^[9]. Cell plasma membrane that is still intact will affect organelles in cells^[10, 11]. This causes sperm to move progressively and stay be alive (viable) so that they are able to fertilize^[10, 11].

Disruption of spermatogenesis through cytotoxic testicular mechanisms, causes a low number of sperm, an increase in the number of abnormal sperm, and a decrease in sperm motility^[12]. Decreased sperm motility with incubation length is thought to be due to sperm running out of energy. Energy is needed by sperm to actively move and is obtained from the decomposition of ATP into ADP and AMP in the mitochondria that are inside the tail of sperm^[13, 14]. If ATP and ADP are depleted, the contraction of fibrils from the tail of the sperm will stop so that the sperm do not move. To maintain the continuity of sperm motility, ATP and ADP must be formed again in the form of alternating reactions^[15]. Provision of essential oils of purple nutsedge grass tubers in experimental animals, it is necessary to know whether essential oils of purple nutsedge tubers can reduce the number and motility of sperm of male mice (*Mus musculus* L.).

Materials and Methods

Making essential oils of nutsedge tubers

The study began with identifying and determining purple nutsedge plants based on observations of plant morphological characteristics such as flowers, leaves, stems, roots, and tubers at the Botanical Laboratory of the Department FMIPA University of Lampung. The tubers were cleaned with water, dried at room temperature for about one week, and cut to small sizes. A total of 1000g of tubers of purple nutsedge were mixed with d distilled water (2/3 the contents of the flask) and heated at a temperature of 100°C for 3 hours until a mixture of oil and water was obtained (until drops of essential oil were not formed). Next, the oil mixture was evaporated in vacuo at a low temperature until essential oil was obtained. Finally, an essential oil that was still mixed with a little water was separated by adding MgSO₄·7H₂O until saturated. The essential oil obtained was used as a sample in the next research process.

Determination of the dose of the essential oil of nutsedge tuber.

The experimental animals used were 24 male mice (*Mus musculus* L.) with an average body weight of 25-35g. The essential oils were administered orally to the animals with the following dosages:

1. Group 1 (Control): mice treated with 0.3 ml of aquabides over 35 days.
2. Group 2: mice were treated with essential oils at a dose of 9 mg/30 g BW in 0.3 ml of aquabides for 35 days.
3. Group 3: mice were treated with essential oils at a dose of 18 mg/30 g BW in 0.3 ml of aquabides for 35 days.

4. Group 4: mice were treated by essential oil with a dose of 36 mg/30 g BW in 0.3 ml of aquabides for 35 days.

Treating experimental animals with essential oils of purple nutsedge tubers

Experimental animals were weighed first, kept in controlled laboratory conditions by feeding them *ad libitum*. Treating with essential oils of nutsedge tubers was done orally using a special syringe whose edges were blunted and given a small rubber pipe. The essential oils were administered to experimental animals by adjusting the volume of suspension with body weight. The treatment was given for 35 days based on the spermatogenic cycle of mice that lasted for 35 days.

Calculation of the number of sperm.

The number of sperm was calculated using the Improved Neubauer booth (hemocytometer). Suspension of sperm was diluted in NaCl solution 0.9%, dropped onto a counting chamber (hemocytometer), and then covered with a cover glass. The number of sperm was counted under a light microscope at 400 times magnification. After knowing the number of sperm, the concentration of sperm was calculated using the following formula:

$$\text{Concentration (million/ml)} = \text{Dilution factor} \times \text{amount in 5 boxes} \times 0.05 \times 10^6.$$

Calculation of sperm motility

After finishing treatment, mice were killed and dissected. Sperm were taken from the epididymis. One of the epididymis was placed on a watch glass that contained 0.9% NaCl, then the organs were chopped into small pieces. One drop of sperm suspension in 0.9% NaCl was dropped on a hemocytometer, covered with a glass cover, and observed under a microscope at 400 times magnification. The percentage of the number of motile sperms was determined by dividing the number of moving sperms by the number of observed sperms and multiplied by 100%.

Statistical Analysis.

Data were analyzed using one way ANOVA and LSD in post hoc test, both ANOVA analysis and LSD test were using $\alpha \leq 0.05$. as a significant difference criteria.

Result and Discussion

The number of sperms

The results of the effect of tuber essential oils on the number of sperms of mice can be seen in Table 1.

Table 1. Sperm counts of mice after treating with essential oils extracted from tubers of nutgrass

Treatments	Sperm counts (million/ml) (mean ± SD)
Control	51.00 ± 8.02 ^a

9 mg / 30g BW	25.16 ± 2.85 ^b
18 mg / 30g BW	16.66 ± 2.33 ^c
36 mg / 30g BW	11.83 ± 2.32 ^c

Values followed by the same superscript were not statistically different at $\alpha = 0.05$.

Sperm Motility

The results of the effect of purple nutsedge tuber essential oils on the motility of sperm of male mice can be seen in Table 2.

Table 2. Sperm motility of mice after treating with essential oils extracted from tubers of nutgrass

Treatments	Sperm motility (%) (mean ± SD)
Control	75.50 ± 9.91 ^a
9 mg/30g BW	55.66 ± 10.42 ^b
18 mg/30g BW	32.66 ± 16.83 ^c
36 mg/30g BW	23.33 ± 11.82 ^c

Values followed by the same superscript were not statistically different at $\alpha = 0.05$.

The results of the effect of purple nutsedge tuber essential oils on the number of sperms of male mice revealed that there are differences in the average number of sperms between the control group and the treatment group. The control group had the highest average number of sperms, while the treatment group, especially T3, had the lowest number of sperms. The results showed that the number of sperms (million/ml) in male mice that were orally administered with 0.3 ml/day essential oils of purple nutsedge tubers for 35 days was decreased compared to the control group. In control mice, the average number of sperms was 51.00 million cells/ml; this number is the normal amount. In T1, the average normal number of sperms was 25.16 million cells/ml, whereas, in T2 and T3, the average number of sperms was less than 20 million per ml of cement which was 16.66 and 11.83 million cells/ml and includes into oligozoospermy light (less normal).

The process of sperm formation or spermatogenesis in male animals occurs when puberty has begun. During this time, the hypothalamus hormones and the pituitary glands actively control reproductive hormones. This causes the testicles to enlarge and develop due to seminiferous tubular activity, and Leydig cells begin to produce the hormone testosterone. Spermatogenesis occurs in the testicular seminiferous tubules and involves spermatogenic cells, Sertoli cells, pituitary hormones, and testosterone.

The results showed that the essential oil of purple nutsedge tubers decreased the number of sperms. This effect causes the number of sperms to be lower with increasing doses of essential oil treatment. The lower number of sperm is thought to occur because the active ingredients of essential oils affect the components involved in spermatogenesis. It is well known that Sertoli cells are responsible for the maturation of spermatogenic cells in the seminiferous tubules, such as for the supply of nutrients and hormones for the maturation of sperms. The

presence of a rough endoplasmic reticulum, a number of mitochondria and Golgi bodies in the basal cytoplasm of Sertoli cells characterizes Sertoli cells for protein metabolic activity, such as Androgen Binding Protein (ABP) and steroid biosynthesis^[16].

No increase in the number of sperms at the reversibility stage was observed; it was decreased or was less when compared to the number of sperms taken at the end of the treatment. It is estimated that this happens in the following ways: For normal spermatogenesis after stopping the treatment for more than 40 days. This is consistent with the results of Hess and Chen (1992), which showed that 90 days after the treatment was stopped, the sperm count was lower than the sperm count in the control group mice^[17].

It can be further explained that, testosterone is needed to begin the process of the first meiosis, namely the formation of primary spermatocyte cells into secondary spermatocyte cells. Al-Makhzoumi (2008) states that testosterone plays a role in the division of the first meiotic prophase that causes a decrease in the number of secondary spermatocytes^[18]. This is also supported by Johnson's statement (2018) that spermatocytes are very sensitive to external influences and tend to suffer damage after the first prophylactic meiosis, especially at the pakiten stage, ie when crossing between homologous chromosomes^[19]. At this stage, the nucleus and cytoplasm grow into the largest cells among the spermatogenic cell layers, when spermatocytes suffer damage such as tubular atrophy, tubular necrosis, and loss of intermedia cells. Cummins et al., (1994) stated that this causes spermatogenic cells to degenerate and phagocytosis by Sertoli cells so that the number of spermatocytes is reduced^[20]. Decreasing the number of spermatocytes causes the number of spermatids to also decrease because spermatocytes that have second meiosis will become spermatids. According to Holdcraft and Braun (2004), the testosterone hormone will maintain all stages of the development of spermatids^[21]. A decrease in the hormone testosterone results in the release of spermatids from the Sertoli cells into the seminiferous tubule lumen which causes the failure of spermiogenesis^[22] or it is suspected that because the number of spermatids decreases it will also decrease the number of sperms produced, because spermatids will experience spermiogenesis to become sperm^[23].

The average percentage of motile sperms of male mice was decreased (in the treatments with essential oils of purple nutsedge tubers with a dose of 9 mg/30gBW, 18 mg/30gBW, and 36 mg/30gBW they were 55.66%, 32.66%, and 23.33% respectively). While the percentage of sperm motility of male mice that were not given essential oils of purple nutsedge tubers (control) was 75.50%. After statistical tests, the results revealed that the control treatments were significantly different from groups 2, 3, and 4 (<0.05). Treatment of 9 mg/30g BW, was significantly different from 18 mg/30gBW, and 36 mg/30gBW (<0.05). Whereas 18 mg/30gBW treatment was not significantly different from the 36 mg/30gBW treatment ($P = 0.213$).

Sperm motility can occur if sperm have mitochondria that function well to produce mobile energy through oxidative

phosphorylation^[24]. It is suspected that the active compounds contained in the purple nutsedge can inhibit the function of mitochondria in producing energy so that the percentage of sperm motility after administration of essential oils of the purple nutsedge tubers decreases. The oxidative phosphorylation process requires the ATP synthase enzyme. It is suspected that the active compounds contained in the purple nutsedge can inhibit the activity of the ATP synthase enzyme so that the available energy cannot be used and causes the sperm to be immobile.

Sperm motility shows the movement of sperm by the presence of flagella that gets energy from the neck of the sperm which is rich in mitochondria. The results of research on the motility of sperm show that papaya seed extract greatly influences motility. Sperms that are released from the testes into the epididymis are not mature because they are not yet motile and cannot be used to fertilize the ovum. The maturation process of sperm occurs in the epididymis.

According to Lohiya *et al.* (2005), decreased sperm motility is not caused by the stage of maturation in the epididymis, but has already occurred while still in the testis^[25]. Sperm motility can occur if sperm have mitochondria that function well to produce mobile energy through oxidative phosphorylation. Januskauskas and Zillinskas, (2002) added that the thing that plays a role in determining sperm motility is the stage of spermiogenesis^[24]. If during the metamorphosis of spermatids, sperms become abnormal (such as decondensation in the nucleus), the formation of acrosomes and mitochondria will affect the success of motility in the epididymis.

Sperm can become motile because of the energy from the neck that is distributed to the tail. This part causes sperm to move forward. So, the main key to the movement of sperm is the production of energy by the mitochondria in the neck of the sperm. It has been suggested that the essential oil of the tubers causes organelle abnormalities in the neck of the sperm, namely vacuolization of the mitochondria and structural abnormalities in the form of a bent neck^[26]. Thus, the mitochondrial function in producing energy is not optimal and ultimately affects the motility of sperm. Lohiya *et al.* (2005) stated that this affects directly the mechanism of sperm formation, while the effect on the epididymis is indirect. Furthermore, it is proved by the correlation between the number and motility of sperm. The number of sperm in the treatment group was less when compared to the number of sperm in the control group. Thus, the essential oil of the tubers affects spermatogenesis at the stage of the metamorphosis of spermatids into sperm. If the spermatozoa produced are abnormal in their formation, of course, this will also affect the maturation of the epididymis.

Menezo *et al.* (2010) reported that sperm motility can occur due to contractions of fibrils in the tail of sperm^[27]. This can happen if sperms have energy in the form of ATP. ATP is produced from simple sugar metabolism through respiration. Respiration can occur if sperm get nutritional intake from outside the cell. Spermatozoa motility occurs because of the movement of the flagellum which consists of microtubules. Flagel motion is the sliding motion between doublets mediated

by dynein. Dynein is a protein that has a group that is responsible for ATPase which contributes to the hydrolysis of ATP. Dynein performs a movement cycle because of the ATP produced by the mitochondria. ATP is activated by the ATPase enzyme to release the first phosphate bonds to form ADP and inorganic phosphate by releasing energy for fibril contraction. If the supply of phosphate, sperm contraction of the fibrils will stop and motility also stops.

Conclusion

The essential oil extracted from the tuber of nutgrass (*Cyperus rotundus*) has revealed to reduce the parameters of sperm quality in male mice in a dose-dependent manner. It can be concluded, therefore, that the essential oil extracted from the tuber of *Cyperus rotundus* is the potential to be used as an anti-fertility agent.

Acknowledgment

The authors are very grateful to Ms. Anis Karimah for her assistance in preparing and keeping samples of the essential oil obtained from tuber extract of rumpu tekki.

References

1. Pucot J R, Manting M M E, Demayo C G. Ethnobotanical Plants used by Selected Indigenous Peoples of Mindanao, the Philippines as Cancer Therapeutics. *Pharmacophores*. 2019; 10(3): 61-69.
2. Kanjekar A P. On Anti-Diabetic Potential of Phytonanoparticles Comparison with Hormonal Therapy and Medicinal Plants. *Int. J. Pharm. Phytopharm. Res.* 2019; 9(1): 103-111.
3. Niazi M, Yari F, Shakarami A. A Review of Medicinal Herbs in the Lamiaceae Family Used to Treat Arterial Hypertension. *Entomol. Appl. Sci. Lett.* 2019; 6(1): 22-27.
4. Kilani S, Abdelwahed A, Ammar RB, Heyder N, Ghedira K, Chraief I, Hammami M, Chekir-Ghedira. Chemical Composition, Antibacterial and Antimutagenic Activities of Essential oil from (Tunisian) *Cyperusrotundus*. *J of Essential Oil Research*, 2005; 17(6):695-700.
5. Lawal, O. A., Adebola, O. Chemical Composition Of The Essential Oils Of *CyperusRotundus* L. From South Africa. *Journal Molecules*, 2009; 14: 2909-2917.
6. Krussel JS, Bielfeld P, Polan ML, Simon C. Regulation of Embryonic Implantation. *European Journal of Obstetrics, Gynecology, and Reproductive Biology*, 2003; 110: S2–S9.
7. Somwanshi, S.D., Madole, M.B., Bikkad, M.D., Bhavthakar, S.S., G.B.S. Ajay. Effect of Cigarette Smoking on Sperm Count and Sperm Motility. *Journal of Medical Education & Research*. 2012; 1: 30-38.

8. Rasgele, P.G. Abnormal Sperm Morphology in Mouse Germ Cell After Short-Term Exposures to Acetamiprid, Propineb, and their Mixture. *Journal of Toksikologi*. 2014; 65: 47-56.
9. Turman EJ, Rich TD. Reproductive Tract Anatomy and Physiology of the Bull. Extension Beef Cattle Resource Committee. Beef Cattle Handbook, 2010.
10. Johnson LA, Weitze KF, Fiser P, Maxwell WMC. Storage of Boar Cement. *Anim Reprod Sci*. 2000; 62 : 143-172.
11. Kaeoket K, Chanapiwat P, Tummaruk P, Techakumphu M, Kunavongkrit A. A Preliminary Study on Using Autologous and Heterologous Boar Sperm Supernatant From Freezing Processes as Post-Thawing Solution: Its Effect on Sperm Motility. *Trop Anim Health Prod*. 2011; 43: 1049-1055.
12. De Lamirande, E., Gagnon, C. Reactive Oxygen Species and Human Sperma. I. Effects on the Motility of Intact sperm and On Sperm Axonemes. *Journal of Andrology*. 1992; 13: 368-378.
13. Hafez, E.S.E. Reproduction in Farm Animals. 6th Ed. Lea and Febiger. Philadelphia, 1993.
14. King, C.J. Reproduction in Domesticated Animals. Elsevier Science Publisher. New York, 1993.
15. Mathews, C.K., Van Holde, K. E. Biochemistry. The Benjamin-Cummings Publishing Company, Inc. California. New York, 1996.
16. Pudney J. Fine Structural Change in Sertoli and Leydig Cells During The Reproductive Cycle of The Ground Squirrel, *Citellus lateralis*. *J. Reprod Fertil*, 1986; 77: 37 - 49
17. Hess RA, Dan Chen PP. Computer of Germ Cells in The Cycle of The Seminiferous Epithelium and Prediction of Change in the The cycle Duration in Animals Commonly Used in Reproductive Biology and Toxicology. *J Andrology*, 1992; 13: 185 - 90.
18. Al-Makhzoomi, A., Lundeheim, N., Haard, M., Rodriguez-Martinez, H. Sperm Morphology and Fertility of Progeny-Tested AI Dairy Bulls in Sweden. *Journal of Theriogenology*. 2008; 70: 682-691.
19. Johnson, M.H. Essential Reproduction. 8th Edition. Wiley Blackwell. Oxford London, 2018.
20. Cummins, J.M., Jequier, A.M., Kan, R. Molecular Biology of the Human Male Infertility: Links With Aging, Mitochondrial Genetics and Oxidative Stress. *Molecular Reproduction and Development*. 1994; 37: 345-362.
21. Holdcraft, R.W., Braun, R.E. Hormonal Regulation of Spermatogenesis. *International Journal of Andrology*. 2004; Vol. 27: 335-342.
22. Cyrus, A., Kabir, A., Goodarzi, D., Moghimi, M. The Effect of Adjuvant Vitamin C After Varicocele Surgery on Sperm Quality and Quantity in Infertile Men: a Double Blind placebo Controlled Clinical Trial. *International Brazil journal Urology*. 2015; 41: 230-8
23. Dada, R., Gupta, N.P., Kucheria, M. Deterioration of Sperm Morphology in Men Exposed to High Temperature. *J. Anat Soc India*. 2001; 50: 107-111
24. Januskauskas A., Zillinskas, H. Bull Semen Evaluation Post-thaw and Relation of Semen Characteristic to Bull's Fertility. *Veterinarijair Zootechnika*. 2002; 17: 1392-2130.
25. Lohiya NK, Mishra PK, Pathak N, Manivannan B, Bhande SS, Panneerdoss S, Sriram S. Efficacy trial on the purified compounds of the seeds of *Carica papaya* for male contraception in albino rat. *Reproductive Toxicology*. 2005; 20(1):135-48.
26. Lohiya NK, Pathak N, Mishra PK, Manivannan B. Reversible contraception with chloroform extract of *Carica papaya* Linn. seeds in male rabbits. *Reproductive toxicology*. 1999 Jan 1; 13(1):59-66.
27. Ménéz Y, Dale B, Cohen M. DNA damage and repair in human oocytes and embryos: a review. *Zygote*. 2010 Nov; 18(4):357-65.