

# ON SOME PETERSEN GRAPHS HAVING LOCATING CHROMATIC NUMBER FOUR OR FIVE

*By Asmiati Asmiati*



## ON SOME PETERSEN GRAPHS HAVING LOCATING CHROMATIC NUMBER FOUR OR FIVE

Asmiati<sup>1</sup>, Wamiliana<sup>1</sup>, Devriyadi<sup>1,2</sup> and Lyra Yulianti<sup>3</sup>

<sup>1</sup>Mathematics Department

Faculty of Mathematics and Natural Sciences

Lampung University

Jl. Brodjonegoro No. 1, Gedung Meneng

Bandar Lampung 35145, Indonesia

e-mail: [asmiati308@yahoo.com](mailto:asmiati308@yahoo.com)

<sup>2</sup>SMKN 1, Mesuji Timur, Lampung, Indonesia

<sup>3</sup>Mathematics Department

Faculty of Mathematics and Natural Sciences

Andalas University

Kampus UNAND Limau Manis

Padang 25163, Indonesia

### Abstract

In this paper, we determine some Petersen graphs having locating chromatic number four or five. Moreover, we give some conjecture to characterize some Petersen graphs having locating chromatic number four.

**17**

Received: March 7, 2017; Revised: April 27, 2017; Accepted: June 12, 2017

2010 Mathematics Subject Classification: 05C12, 05C15.

Keywords and phrases: locating chromatic number, Petersen **3** graph.

## 1. Introduction

The concept of locating chromatic number of a graph is the combination between the partition dimension [5] and vertex coloring of a graph. The following definition of locating chromatic number of a graph is taken from [6].

Let  $G = (V, E)$  be a connected graph. Let  $c$  be a proper coloring of  $G$  using the colors  $1, 2, \dots, k$  for some positive integer  $k$ . Let  $\Pi = \{C_1, C_2, \dots, C_k\}$  be a partition of  $V(G)$ , where  $C_i$  is a set of vertices colored  $i$ , for  $i \in [1, k]$ . The color code  $c_{\Pi}(v)$  of vertex  $v$  in  $G$  is the ordered  $k$ -tuple  $(d(v, C_1), \dots, d(v, C_k))$ , where  $d(v, C_i) = \min\{d(v, x) \mid x \in C_i\}$  for  $i \in [1, k]$ . For every pair of vertices  $u$  and  $v$ , if  $c_{\Pi}(u) \neq c_{\Pi}(v)$ , then  $c$  is called a *locating coloring* of  $G$  using  $k$  colors. The locating chromatic number, denoted by  $\chi_L(G)$ , is the smallest  $k$  such that  $G$  has a locating coloring.

The following theorem taken from [6] is a fundamental theorem about the locating chromatic number of a graph. The neighborhood of a vertex  $u$  in a connected graph  $G$ , denoted by  $N(u)$ , is the set of vertices adjacent to  $u$ .

**Theorem 1.1** [6]. Let  $c$  be a locating coloring in a connected graph  $G$ . If  $s$  and  $t$  are distinct vertices of  $G$  such that  $d(s, u) = d(t, u)$  for all  $u \in V(G) - \{s, t\}$ , then  $c(s) \neq c(t)$ . In particular, if  $s$  and  $t$  are non-adjacent vertices of  $G$  such that  $N(s) = N(t)$ , then  $c(s) \neq c(t)$ .

**Theorem 1.2** [6]. The locating chromatic number of a cycle  $C_n$  is 3 for odd  $n$  and 4 for otherwise.

Determining the locating chromatic number of a graph is an interesting topic because there is no algorithm that can be used to determine the locating chromatic number for any graph. Chartrand et al. [6] discussed the locating chromatic number for some classes of graphs such as paths, cycles, bipartite graphs, caterpillars, double stars, and certain trees. Moreover, Asmiati et al.

[1, 2] determined the locating chromatic number for amalgamation of stars and firecracker graphs.

In 2003, Chartrand et al. [6] characterized a graph having locating chromatic number  $(n - 1)$  or  $(n - 2)$ . Asmiati and Baskoro [3] characterized a graph containing cycles with locating chromatic number three. Next, in 2013, Baskoro and Asmiati [4] characterized the trees with locating chromatic number three. In this paper, we determine the locating chromatic number of some Petersen graphs having locating chromatic number four or five.

The following definition of a Petersen graph is taken from [7]. Let  $\{u_1, u_2, \dots, u_n\}$  be some vertices on the outer cycle and  $\{v_1, v_2, \dots, v_n\}$  be some vertices on the inner cycle, for  $n \geq 3$ . The Petersen graph, denoted by  $P_{n,k}$ ,  $n \geq 3$ ,  $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$ ,  $1 \leq i \leq n$  is a graph that has  $2n$  vertices  $\{u_i\} \cup \{v_i\}$ , and edges  $\{u_i u_{i+1}\} \cup \{v_i v_{i+k}\} \cup \{u_i v_i\}$ .

In this paper, we determine some Petersen graphs having locating chromatic number four or five. Moreover, we give some conjecture to characterize some Petersen graphs having locating chromatic number four.

20

## 2. Main Results

In this section, we determine Petersen graph  $P_{n,1}$ ,  $n \geq 3$  having locating chromatic number four or five. Besides, we discuss Petersen graphs  $P_{4,2}$  and  $P_{5,2}$  having locating chromatic number 4 or 5, respectively.

**Theorem 2.1.** *The locating chromatic number of Petersen graph  $P_{n,1}$  is 4 for odd  $n \geq 3$  or 5 for even  $n \geq 4$ .*

**Proof.** To prove this theorem, we consider the following cases:

**Case 1.**  $\chi_L(P_{n,1}) = 4$ , for odd  $n \geq 3$ .

First, we determine the lower bound of  $\chi_L(P_{n,1})$  for odd  $n \geq 3$ . Because the Petersen graph  $P_{n,1}$  contains some even cycles, then by Theorem 1.2,  $\chi_L(P_{n,1}) \geq 4$ . Thus,  $\chi_L(P_{n,1}) \geq 4$  for odd  $n \geq 3$ .

Let  $c$  be a coloring of Petersen graph  $P_{n,1}$ , for odd  $n \geq 3$ . We make the partition of the vertices of  $V(P_{n,1})$ :  $C_1 = \{u_1\}$ ;  $C_2 = \{u_{2j}, v_{2j-1}\}$ ;  $C_3 = \{u_{2j+1}, v_{2j}\}$ ;  $C_4 = \{v_n\}$ , for  $j > 0$ . Therefore, the color codes of all the vertices of  $G$  are:

$$(a) \quad c_{\Pi}(u_1) = (0, 1, 1, 2).$$

$$(b) \quad C_2 = \{u_{2j}, v_{2j-1}\}. \text{ Let } u_i, \quad 2 \leq i \leq n-1; i = 2j; 1 \leq j \leq \left\lfloor \frac{n}{2} \right\rfloor \text{ and } v_i, \quad 1 \leq i \leq n-2; i = 2j-1; 1 \leq j \leq \left\lfloor \frac{n}{2} \right\rfloor.$$

For  $i < \left\lfloor \frac{n}{2} \right\rfloor$ , we define

$$c_{\Pi}(u_i) = (i-1, 0, 1, i+1),$$

$$c_{\Pi}(v_i) = (i, 0, 1, i).$$

For  $i = \left\lfloor \frac{n}{2} \right\rfloor$ , we define

$$c_{\Pi}(u_i) = c_{\Pi}(u_{n-2j+1}) = (i-1, 0, 1, 2j),$$

$$c_{\Pi}(v_i) = c_{\Pi}(v_{n-2j}) = (i, 0, 1, 2j).$$

For  $i > \left\lfloor \frac{n}{2} \right\rfloor$ ,

$$c_{\Pi}(u_i) = c_{\Pi}(u_{n-2j+1}) = (2j, 0, 1, 2j),$$

$$c_{\Pi}(v_i) = c_{\Pi}(v_{n-2j}) = (2j+2, 0, 1, 2j).$$

(c)  $C_3 = \{u_{2j+1}, v_{2j}\}$ . Let  $u_i, 3 \leq i \leq n; i = 2j + 1; 1 \leq j \leq \lfloor \frac{n}{2} \rfloor$  and  $v_i, 2 \leq i \leq n - 1; i = 2j; 1 \leq j \leq \lfloor \frac{n}{2} \rfloor$ .

For  $i < \lfloor \frac{3}{2} \rfloor$ ,

$$c_{\Pi}(u_i) = (i - 1, 1, 0, i + 1),$$

$$c_{\Pi}(v_i) = (i, 1, 0, i).$$

For  $i = \lfloor \frac{n}{2} \rfloor$ ,

$$c_{\Pi}(u_i) = c_{\Pi}(u_{n-2j+2}) = (i - 1, 0, 1, 2j - 1),$$

$$c_{\Pi}(v_i) = c_{\Pi}(v_{n-2j+1}) = (i, 1, 0, 2j - 1).$$

For  $i > \lfloor \frac{n}{2} \rfloor$ ,

$$c_{\Pi}(u_i) = c_{\Pi}(u_{n-2j+2}) = (2j - 1, 1, 0, 2j - 1),$$

$$c_{\Pi}(v_i) = c_{\Pi}(v_{n-2j+1}) = (2j + 1, 1, 0, 2j - 1).$$

(d)  $c_{\Pi}(v_n) = (2, 1, 1, 0)$ .

Since the color codes of each vertex in  $V(P_{n,1})$  is different,  $c$  is a locating coloring of  $P_{n,1}$  for odd  $n \geq 3$ . Therefore,  $\chi_L(P_{n,1}) \leq 4$ .

**Case 2.**  $\chi_L(P_{n,1}) = 5$ , for even  $n \geq 4$ .

The Petersen graph  $P_{n,1}$ , for even  $n \geq 4$ , contains some even cycles. Then, by Theorem 1.2,  $\chi_L(P_{n,1}) \geq 4$ . Suppose that  $c$  is a locating coloring of  $P_{n,1}$ , for even  $n \geq 4$ . Let  $C_1 = \{u_1\}$ ;  $C_2 = \{u_{2i}, v_{2i-1}\}$ ;  $C_3 = \{u_{2i+1}, v_{2i}\}$  and  $C_4 = \{u_n\}$  for  $i > 0$ . Then there are two vertices with the same color

codes,  $c_{\Pi}(u_2) = c_{\Pi}(v_1)$ , a contradiction. Thus, we need at least five colors.

Therefore,  $\chi_L(P_{n,1}) \geq 5$  for even  $n \geq 4$ .

Next, we determine the upper bound of  $\chi_L(P_{n,1})$  for even  $n \geq 4$ . We assign the vertices like this:

$$C_1 = \{u_1\}; C_2 = \{u_{2j}, v_{2j-1}\}; C_3 = \{u_{2j+1}, v_{2j}\}; C_4 = \{u_n\}; C_5 = \{v_n\}.$$

So, we have the color codes of vertices of  $G$  as follows:

$$(a) c_{\Pi}(u_1) = (0, 1, 2, 1, 2).$$

$$(b) C_2 = \{u_{2j}, v_{2j-1}\}. \text{ Let } u_i, 1 \leq i \leq n-2; i = 2j; 1 \leq j \leq \frac{n}{2}-1 \text{ and } v_i, 1 \leq i \leq n-1; i = 2j-1; 1 \leq j \leq \frac{n}{2}.$$

$$\text{For } i \leq \left\lceil \frac{n}{2} \right\rceil,$$

$$c_{\Pi}(u_i) = (i-1, 0, 1, i, i+1),$$

$$c_{\Pi}(v_i) = (i, 0, 1, i+1, i).$$

$$\text{For } i > \left\lceil \frac{n}{2} \right\rceil,$$

$$c_{\Pi}(u_i) = c_{\Pi}(u_{n-2j}) = (2j+1, 0, 1, 2j, 2j+1),$$

$$c_{\Pi}(v_i) = c_{\Pi}(v_{n-2j+1}) = (2j+1, 0, 1, 2j, 2j-1).$$

$$(c) C_3 = \{u_{2j+1}, v_{2j}\}. \text{ Let } u_i, 1 \leq i \leq n-1; i = 2j+1; 1 \leq j \leq \frac{n}{2}-1 \text{ and } v_i, 1 \leq i \leq n-2; i = 2j; 1 \leq j \leq \frac{n}{2}-1.$$

$$\text{For } i \leq \left\lceil \frac{n}{2} \right\rceil,$$

$$c_{\Pi}(u_i) = (i-1, 1, 0, i, i+1),$$

$$c_{\Pi}(v_i) = (i, 1, 0, i + 1, i).$$

For  $i > \left\lfloor \frac{n}{2} \right\rfloor$ ,

$$c_{\Pi}(u_i) = c_{\Pi}(u_{n-2j+1}) = (2j, 1, 0, 2j - 1, 2j),$$

$$c_{\Pi}(v_i) = c_{\Pi}(v_{n-2j}) = (2j + 2, 1, 0, 2j + 1, 2j).$$

(d)  $C_4 = \{u_n\}$ , then  $c_{\Pi}(u_n) = (1, 2, 1, 0, 1)$ .

(e)  $c_{\Pi}(v_n) = (2, 1, 2, 1, 0)$ .

Since all the vertices have different color codes,  $c$  is a locating coloring of Petersen graphs  $P_{n,1}$  for even  $n \geq 4$ . Therefore, we have  $\chi_L(P_{n,1}) = 5$ , for even  $n \geq 4$ . □




Figure 1. A minimum locating coloring of  $P_{7,1}$ .




Figure 2. A minimum locating coloring of  $P_{8,1}$ .



**Theorem 2.2.**  $\chi_L(P_{4,2}) = 4$ .

**Proof.** Since Petersen graph  $P_{4,2}$  containing some even cycles, by Theorem 2.1,  $\chi_L(P_{4,2}) \geq 4$ . Next, we determine the upper bound of  $\chi_L(P_{4,2})$ . Let  $c$  be a coloring of Petersen graph  $P_{4,2}$  using 4 colors, by assigning the colors like this:  $c(u_1) = 1$ ,  $c(u_2) = 2$ ,  $c(u_3) = 3$ ,  $c(u_4) = 4$ ,  $c(v_1) = 2$ ,  $c(v_2) = 4$ ,  $c(v_3) = 1$ ,  $c(v_4) = 3$ . The color codes of the vertices:

$$c_\pi(u_1) = \{0, 1, 2, 1\}; c_\pi(u_2) = \{1, 0, 1, 1\}; c_\pi(u_3) = \{1, 1, 0, 1\};$$

$$c_\pi(u_4) = \{1, 2, 1, 0\}; c_\pi(v_1) = \{1, 0, 2, 2\}; c_\pi(v_2) = \{2, 1, 1, 0\};$$

$$c_\pi(v_3) = \{0, 1, 1, 2\}; c_\pi(v_4) = \{2, 2, 0, 1\}.$$

Since all the vertices have distinct color codes,  $c$  is a locating coloring of  $P_{4,2}$ , so  $\chi_L(P_{4,2}) \leq 4$ .  $\square$

**Theorem 2.3.**  $\chi_L(P_{5,2}) = 5$ .

**Proof.** Since Petersen graph  $P_{5,2}$  containing some even cycles, by Theorem 2.1,  $\chi_L(P_{5,2}) \geq 4$ . Suppose that the locating chromatic number of  $P_{5,2}$  is 4. Without loss of generality, we assign  $c(u_i) = 1$ ;  $c(u_i) = 2$  for  $i = 2, 4$ ;  $c(u_i) = 3$  for  $i = 3, 5$  and  $c(v_1) = 2$ . Then  $c(v_3) = 1$  or 4. If  $c(v_3) = 1$ , then  $c(v_5) = 2$  or 4. Set  $c(v_2) = 4$  and  $c(v_4) = 3$ , then  $c_\pi(u_3) = c_\pi(u_5)$ , a contradiction. So, we need at least 5 colors. As a result,  $\chi_L(P_{5,2}) \geq 5$ .

Next, we determine the upper bound of  $\chi_L(P_{5,2})$ . Let  $c$  be a coloring of vertices of  $P_{5,2}$  such that the color classes are as follows:

$$C_1 = \{u_1\}; C_2 = \{u_2, u_4, v_1, v_5\}; C_3 = \{u_3, u_5, v_2\};$$

$$C_4 = \{v_3\}; C_5 = \{v_4\}.$$

So, we have the color codes of vertices are:

$$c_{\Pi}(u_1) = (0, 1, 1, 2, 2); c_{\Pi}(u_2) = (1, 0, 1, 2, 2); c_{\Pi}(u_4) = (2, 0, 1, 2, 1);$$

$$c_{\Pi}(v_1) = (1, 0, 2, 1, 1); c_{\Pi}(v_5) = (2, 0, 1, 1, 2); c_{\Pi}(u_3) = (2, 1, 0, 1, 2);$$

$$c_{\Pi}(u_5) = (1, 1, 0, 2, 2); c_{\Pi}(v_2) = (2, 1, 0, 2, 1); c_{\Pi}(v_3) = (2, 1, 1, 0, 2);$$

$$c_{\Pi}(v_4) = (2, 1, 1, 2, 0).$$

Since the color codes of all the vertices in  $P_{5,2}$  are different,  $c$  is a locating coloring. So,  $\chi_L(P_{5,2}) \leq 5$ .  $\square$

Next, we give some conjecture related to the locating chromatic number of Petersen graph.

**Conjecture.** The locating chromatic number of Petersen graph  $P$  is four if and only if  $P = P_{n,1}$ , for odd  $n \geq 3$  or  $P_{4,2}$ .

### 3. Conclusion

Based on the results, locating chromatic number of Petersen graph  $P_{n,1}$  is 4 for odd  $n \geq 3$  and 5 for even  $n \geq 4$ ;  $\chi_L(P_{4,2}) = 4$ ;  $\chi_L(P_{5,2}) = 5$ . Then we have a conjecture to characterize Petersen graph having locating chromatic number four.

### Acknowledgment

This research is supported by Fundamental Research Grant in 2017, Ministry of Research, Technology and Higher Education, Indonesia.

The authors gratefully acknowledge the constructive criticisms of the reviewers which have greatly improved the clarity and presentation of the paper.

**References**

- [1] Asmiati, H. Assiyatun and E. T. Baskoro, Locating-chromatic number of amalgamation of stars, *ITB J. Sci.* 43A (2011), 1-8.
- [2] Asmiati, E. T. Baskoro, H. Assiyatun, D. Suprijanto, R. Simanjuntak and S. Uttunggadewa, Locating-chromatic number of firecracker graphs, *Far East J. Math. Sci. (FJMS)* 63(1) (2012), 11-23.
- [3] Asmiati and E. T. Baskoro, Characterizing of graphs containing cycle with locating-chromatic number three, *AIP Conf. Proc.*, 1450, 2012, pp. 351-357.
- [4] E. T. Baskoro and Asmiati, Characterizing all trees with locating-chromatic number 3, *Electronic J. Graph Theory Appl.* 1(2) (2013), 109-117.
- [5] G. Chartrand, E. Salehi and P. Zhang, On the partition dimension of graph, *Congr. Numer.* 130 (1998), 157-168.
- [6] G. Chartrand, D. Erwin, M. A. Henning, P. J. Slater and P. Zhang, Graph of order  $n$  with locating-chromatic number  $n - 1$ , *Discrete Math.* 269 (2003), 65-79.
- [7] D. A. Holton and J. Sheehan, *The Petersen Graph*, Cambridge University Press, Cambridge, 1993.

# ON SOME PETERSEN GRAPHS HAVING LOCATING CHROMATIC NUMBER FOUR OR FIVE

ORIGINALITY REPORT

24%

SIMILARITY INDEX

## PRIMARY SOURCES

- 1 Gary Chartrand, David Erwin, Michael A. Henning, Peter J. Slater, Ping Zhang. "Graphs of order  $n$  with locating-chromatic number  $n-1$ ", Discrete Mathematics, 2003 116 words — 5%

Crossref
- 2 Ali Behtoei, Behnaz Omoomi. "On the locating chromatic number of Kneser graphs", Discrete Applied Mathematics, 2011 61 words — 3%

Crossref
- 3 Mohan Chris Monica, Samivel Santhakumar. "Partition dimension of rooted product graphs", Discrete Applied Mathematics, 2019 41 words — 2%

Crossref
- 4 H. L. FU, C. A. RODGER. "4-Cycle Group-Divisible Designs with Two Associate Classes", Combinatorics, Probability and Computing, 2001 36 words — 2%

Crossref
- 5 Michitaka Furuya, Naoki Matsumoto. "Upper bounds on the locating chromatic number of trees", Discrete Applied Mathematics, 2019 30 words — 1%

Crossref
- 6 arxiv.org 22 words — 1%

Internet
- 7 Tamer Elgohary, Moussa S. Elbisy, Amir M. Mobasher, Hassan Salah. "DEEP WAVE HEIGHT PREDICTION FOR ALEXANDRIA SEA REGION BY USING NONLINEAR REGRESSION METHOD COMPARED TO SUPPORT VECTOR

- 
- 8 Yunjiang Lou, Wei Gong, Zexiang Li, Jianjun Zhang, Guilin Yang. "Natural frequency based optimal design of a two-link flexible manipulator", 2009 IEEE International Conference on Robotics and Automation, 2009  
18 words — 1%  
Crossref
- 
- 9 Haeder Y. Althoby, Mohamed Didi Biha, André Sesboüé. "Exact and heuristic methods for the vertex separator problem", Computers & Industrial Engineering, 2020  
17 words — 1%  
Crossref
- 
- 10 Debi Oktia Haryeni, Edy Tri Baskoro. "Partition Dimension of Some Classes of Homogenous Disconnected Graphs", Procedia Computer Science, 2015  
17 words — 1%  
Crossref
- 
- 11 Cyriac Grigorious, Paul Manuel, Mirka Miller, Bharati Rajan, Sudeep Stephen. "On the metric dimension of circulant and Harary graphs", Applied Mathematics and Computation, 2014  
15 words — 1%  
Crossref
- 
- 12 [www.combinatorics.cn](http://www.combinatorics.cn)  
Internet  
12 words — 1%
- 
- 13 Clint Dawson, Robert Kirby. "High Resolution Schemes for Conservation Laws with Locally Varying Time Steps", SIAM Journal on Scientific Computing, 2001  
11 words — 1%  
Crossref
- 
- 14 [tesisexarxa.net](http://tesisexarxa.net)  
Internet  
10 words — < 1%
- 
- 15 "DESIGNED STRUCTURE AND MAGNETIC CHARACTERISTIC STUDIES OF MAGNETIC IRON OXIDE (Fe<sub>3</sub>O<sub>4</sub>) NANOPARTICLES COATED BY POLYVINYL ALCOHOL AND POLYVINYL ALCOHOL-LINKED WITH GLUTARALDEHYDE", Rasayan Journal of Chemistry, 2017  
9 words — < 1%  
Crossref

- 
- 16 [www.emis.de](http://www.emis.de)  
Internet 9 words — < 1%
- 
- 17 [research.aalto.fi](http://research.aalto.fi)  
Internet 9 words — < 1%
- 
- 18 [ajc.maths.uq.edu.au](http://ajc.maths.uq.edu.au)  
Internet 8 words — < 1%
- 
- 19 J. M.-S. Lubuma, S. Nicaise. "Edge behaviour of the solution of the Stokes problem with applications to the finite-element method", Proceedings of the Royal Society of Edinburgh: Section A Mathematics, 2007  
Crossref 8 words — < 1%
- 
- 20 [www.discuss.wmie.uz.zgora.pl](http://www.discuss.wmie.uz.zgora.pl)  
Internet 8 words — < 1%
- 
- 21 De Ming Li. "The Star Chromatic Numbers of Some Planar Graphs Derived from Wheels", Acta Mathematica Sinica, English Series, 2002  
Crossref 7 words — < 1%
- 
- 22 László Neumann, Rafael García, János Basa, Ramón Hegedüs. "Acquisition and visualization techniques for narrow spectral color imaging", Journal of the Optical Society of America A, 2013  
Crossref 7 words — < 1%
- 
- 23 Grigorious, Cyriac, Paul Manuel, Mirka Miller, Bharati Rajan, and Sudeep Stephen. "On the metric dimension of circulant and Harary graphs", Applied Mathematics and Computation, 2014.  
Crossref 7 words — < 1%
- 
- 24 Gunaydin, M.. "Erratum to: "Unitary supermultiplets of  $OSp(8^*|4)$  and the AdS<sup>7</sup>/CFT<sup>6</sup> duality" [Nucl. Phys. B 578 (2000) 405]", Nuclear Physics, Section B, 20041004  
Crossref 7 words — < 1%
- 
- 25 Israel, D.. "D-branes in little string theory", Nuclear Physics,

Section B, 20050822

Crossref

7 words — < 1%

26

Ames, G.. "Comparison morphisms and the Hochschild cohomology ring of truncated quiver algebras", Journal of Algebra, 20090901

Crossref

6 words — < 1%

EXCLUDE QUOTES ON

EXCLUDE MATCHES OFF

EXCLUDE BIBLIOGRAPHY ON