

The Locating-Chromatic Number for Certain Operation of Generalized Petersen Graphs $\square A \square A (\square A, \square A)$

By A. Irawan

PAPER · OPEN ACCESS

The Locating-Chromatic Number for Certain Operation of Generalized Petersen Graphs $sP(4,2)$

14

To cite this article: A Irawan *et al* 2019 *J. Phys.: Conf. Ser.* **1338** 012033

11

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices
to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of
every title for free.

The Locating-Chromatic Number for Certain Operation of Generalized Petersen Graphs $sP(4, 2)$

A Irawan^{1,2,a}, Asmiati^{3,b}, Suharsono^{3,c}, and K Muludi^{4,d}

¹ Posgraduate student, Faculty of Mathematics and Natural Sciences Lampung University, Jl. Sumantri Brodjonegoro No.1, Bandar Lampung, Indonesia.

² School of Information and Computer Management (STMIK) Pringsewu Lampung. Jl. Wisma Rini No.09, Pringsewu, Lampung, Indonesia

³ Departement Mathematics, Faculty of Mathematics and Natural Sciences Lampung University. Jl. Sumantri Brodjonegoro No.1, Bandar Lampung, Indonesia

⁴ Computer Sciences, Faculty of Mathematics and Natural Sciences Lampung University. Jl. Sumantri Brodjonegoro No.1, Bandar Lampung, Indonesia

^aagusirawan814@gmail.com; ^basmiati.1976@fmipa.unila.ac.id;

^csuharsono.1962@fmipa.unila.ac.id; ^dkurnia.muludi@fmipa.unila.ac.id

Abstract. The locating-chromatic number of a graph combined two graph concept, coloring vertices and partition dimension of a graph. The locating-chromatic number, denoted by $\chi_L(G)$, is the smallest k such that G has a locating k -coloring. In this paper, we discuss the locating-chromatic number for certain operation of generalized Petersen graphs $sP(4,2)$.

4 Introduction

Chartrand et al. [1] in 2002 introduced the locating-chromatic number of a graph, with derived two graph concept, coloring vertices and partition dimension of a graph. Let $G = (V, E)$ be a connected graph and c be a proper k -coloring of G with color $1, 2, \dots, k$. Let $\Pi = \{C_1, C_2, \dots, C_k\}$ be a partition of $V(G)$ which is induced by coloring c . The color code $c_\Pi(v)$ of v is the ordered k -tuple $(d(v, C_1), d(v, C_2), \dots, d(v, C_k))$ where $d(v, C_i) = \min \{d(v, x) | x \in C_i\}$ for any i . If all distinct vertices of G have distinct color codes, then c is called k -locating coloring of G . The locating-chromatic number, denoted by $\chi_L(G)$, is the smallest k such that G has a locating k -coloring.

In 2003, Chartrand et al. [2] succeeded in constructing $n \geq 5$ tree graphs with locating-chromatic numbers ranging from 3 to n , except $(n - 1)$. Behtoe and Omoomi [3] found locating-chromatic numbers on the Kneser graph. Furthermore, Baskoro and Purwasih [4] found the locating chromatic number for corona product of graphs. Next, Asmiati [5] determined the locating chromatic number of banana tree graph and Asmiati et al. [6] for amalgamation of stars graphs. Asmiati et al. [7] also found the locating chromatic number of firecracker graphs and Syofyan et al. [8] for lobster graph.

Specially for non-homogenous tree graph in 2014, Asmiati [9] determined the locating-chromatic number of non-homogeneous amalgamation of stars, then Asmiati [10] for caterpillar graphs and non-homogenous firecracker graphs. In 2017, Asmiati et al. [11] determined some generalized Petersen graphs $P(n, 1)$ having locating-chromatic number 4 for odd $n \geq 3$ or 5 for even $n \geq 4$.

The generalized Petersen graph $P(n, k)$, $n \geq 3$ and $1 \leq k \leq \lfloor (n-1)/2 \rfloor$, consists of an outer n -cycle u_1, u_2, \dots, u_n , a set of n spokes u_i, v_i , $1 \leq i \leq n$, and n edges v_i, v_{i+k} , $1 \leq i \leq n$, with indices taken modulo n . The generalized Petersen graph was introduced by Watkins in [12].

To define the generalized Petersen graph $sP(4,2)$, suppose there are s generalized Petersen graph $P(4,2)$. Some vertices on the outer cycle u_i , $i = 1,2,3,4$ for the generalized Petersen graph t^{th} , $t = 1,2, \dots, s$, $s \geq 1$ denoted by u_i^t , while some vertices on the inner cycle v_i , $i = 1,2,3,4$ for the generalized Petersen graph t^{th} , $t = 1,2, \dots, s$, $s \geq 1$ denoted by v_i^t . Generalized Petersen graph $sP(4,2)$ obtained from $s \geq 1$ graph $P(4,2)$, which every vertices on the outer cycle u_i^t , $i \in [1,4]$, $t \in [1, s]$ connected by a path $(u_i^t u_i^{t+1})$ $t = 1,2, \dots, s-1$, $s \geq 2$.

Some researchers have determined the locating-chromatic number for certain operation. Behtoei and Omoomi [13] obtained locating-chromatic number from the grid, cartesian multiplication for trajectories and complete graphs, and cartesian multiplication of two complete graphs. Furthermore Behtoei and Omoomi [14] determined the locating-chromatic number of the fan graph, wheel and friendship graph for join multiplication of two graphs. Asmiati [15] found locating-chromatic number for certain operation of tree. In this paper, we discuss the locating-chromatic number for certain operation of generalized Petersen graphs $sP(4,2)$.

The following theorem is basic to determine the locating chromatic number of a graph. The set of neighbours of a vertex s in G , denoted by $N(s)$.

Theorem 1.1. Chartrand et al. [1] *Let c be a locating coloring in a connected graph G . If r and s are distinct vertices of G such that $d(r,w) = d(s,w)$ for all $w \in V(G) - \{r,s\}$, then $c(r) \neq c(s)$. In particular, if x and y are non-adjacent vertices of G such that $N(x) \neq N(y)$, then $c(x) \neq c(y)$.*

Theorem 1.2. Chartrand et al. [1] *The locating chromatic number of a cycle C_n , is 3 for odd n and 4 for otherwise.*

2. Results and Discussion

In this section we will discuss the locating chromatic number of $sP(4,2)$.

Theorem 2.1. The locating chromatic number of generalized Petersen graph $sP(4,2)$ is 5 for $s \geq 2$.

Proof : First, we determine lower bound of $\chi_L(sP(4,2))$ for $s \geq 2$. Because generalized Petersen graph $P(4,2)$, for $s \geq 2$, contains some even cycles. Then by Theorem 2, $\chi_L(sP(4,2)) \geq 4$. Next, we will show that $\chi_L(sP(4,2)) \geq 5$, for $s \geq 2$. For a contradiction, suppose that c is 4-locating coloring on $sP(4,2)$ for $s \geq 2$. Consider $c(u_i^1) = i$, $i = 1,2,3,4$ and $c(v_j^1) = j$, $j = 1,2,3,4$ such that $c(u_i^1) \neq c(v_j^1)$ for $c(u_i^1)$ adjacent to $c(v_j^1)$. Observe that if we assign color 4 for any vertices in u_i^2 or v_i^2 , then we have two vertices which have color codes. Therefore, c is not locating 4-coloring on $sP(4,2)$. As the result $\chi_L(sP(4,2)) \geq 5$ for $s \geq 2$.

Next, we determine the upper bound of $\chi_L(sP(4,2))$ for $s \geq 2$. Let c be a coloring of generalized Petersen graph $sP(4,2)$ for $s \geq 2$. We make the partition of the vertices of $V(sP(4,2))$:

$$\begin{aligned} C_1 &= \{u_1^t | \text{for odd } s\} \cup \{u_2^t, v_4^t | \text{for even } s\} \\ C_2 &= \{u_2^t, u_4^t | \text{for odd } s\} \cup \{u_3^t, v_1^t | \text{for even } s\} \\ C_3 &= \{u_3^t, v_1^t, v_2^t | \text{for odd } s\} \cup \{u_4^t, v_2^t, v_3^t | \text{for even } s\} \\ C_4 &= \{v_3^t | \text{for odd } s\} \cup \{u_1^t | \text{untuk } s \text{ genap}\} \cup \{v_4^t | \text{for odd } s \geq 3\} \\ C_5 &= \{v_4^t\} \end{aligned}$$

Therefore the color codes of all the vertices of G are :

(a) $C_1 = \{u_1^t | \text{for odd } s\} \cup \{u_2^t, v_4^t | \text{for even } s\}$

For odd s , the color codes of $sP(4,2)$ are:

$$c_{\Pi}(u_1^t) = \begin{cases} 0 & , \text{ for } 1^{st} \text{ component} \\ 1 & , \text{ for } 2^{nd}, 3^{rd} \text{ and } 4^{th} \text{ component} \\ s+1 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

For even s , the color codes of $sP(4,2)$ are:

$$c_{\Pi}(u_2^t) = \begin{cases} 0 & , \text{ for } 1^{st} \text{ component} \\ 1 & , \text{ for } 2^{nd}, 3^{rd} \text{ and } 4^{th} \text{ component} \\ s+1 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

$$c_{\Pi}(v_4^t) = \begin{cases} 0 & , \text{ for } 1^{st} \text{ component} \\ 2 & , \text{ for } 2^{nd} \text{ and } 4^{th} \text{ component} \\ 1 & , \text{ for } 3^{rd} \text{ component} \\ s+1 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

(b) $C_2 = \{u_2^t, u_4^t | \text{for odd } s\} \cup \{u_3^t, v_1^t | \text{for even } s\}$

For odd s the color codes of $sP(4,2)$ are:

$$c_{\Pi}(u_2^t) = \begin{cases} 1 & , \text{ for } 1^{st} \text{ and } 3^{rd} \text{ component} \\ 0 & , \text{ for } 2^{nd} \text{ component} \\ 4 & , \text{ for } 4^{th} \text{ component} \\ s+1 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

$$c_{\Pi}(u_4^t) = \begin{cases} 1 & , \text{ for } 1^{st}, 3^{rd} \text{ and } 5^{th} \text{ component} \\ 0 & , \text{ for } 2^{nd} \text{ component} \\ 2 & , \text{ for } 4^{th} \text{ component} \end{cases}$$

For odd $s \geq 3$, the color codes of $sP(4,2)$ are:

$$c_{\Pi}(u_4^t) = \begin{cases} 1 & , \text{ for } 1^{st}, 3^{rd} \text{ and } 4^{th} \text{ component} \\ 0 & , \text{ for } 2^{nd} \text{ component} \\ s & , \text{ for } 5^{th} \text{ component} \end{cases}$$

For even s , the color codes of $sP(4,2)$ are:

$$c_{\Pi}(u_3^t) = \begin{cases} 1 & , \text{ for } 1^{st} \text{ and } 3^{rd} \text{ component} \\ 0 & , \text{ for } 2^{nd} \text{ component} \\ 2 & , \text{ for } 4^{th} \text{ component} \\ s+1 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

$$c_{\Pi}(v_1^t) = \begin{cases} 2 & , \text{ for } 1^{st} \text{ component} \\ 0 & , \text{ for } 2^{nd} \text{ component} \\ 1 & , \text{ for } 3^{rd} \text{ and } 4^{th} \text{ component} \\ s+2 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

(c) $C_3 = \{u_3^t, v_1^t, v_2^t | \text{for odd } s\} \cup \{u_4^t, v_2^t, v_3^t | \text{for even } s\}$.

For odd s , the color codes of $sP(4,2)$ are:

$$c_{\Pi}(u_3^t) = \begin{cases} 2 & , \text{ for } 1^{st} \text{ component} \\ 1 & , \text{ for } 2^{nd} \text{ and } 4^{th} \text{ component} \\ 0 & , \text{ for } 3^{rd} \text{ component} \\ s+1 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

$$c_{\Pi}(v_1^t) = \begin{cases} 1 & , \text{ for } 1^{st} \text{ and } 4^{th} \text{ component} \\ 2 & , \text{ for } 2^{nd} \text{ component} \\ 0 & , \text{ for } 3^{rd} \text{ component} \\ s+2 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

$$c_{\Pi}(v_2^1) = \begin{cases} 2 & , \text{ for } 1^{st} \text{ component} \\ 1 & , \text{ for } 2^{nd} \text{ and } 5^{th} \text{ component} \\ 0 & , \text{ for } 3^{rd} \text{ component} \\ 3 & , \text{ for } 4^{th} \text{ component} \end{cases}$$

For odd $s \geq 3$ the color codes of $sP(4,2)$ are:

$$c_{\Pi}(v_2^t) = \begin{cases} 2 & , \text{ for } 1^{st} \text{ and } 4^{th} \text{ component} \\ 1 & , \text{ for } 2^{nd} \text{ component} \\ 0 & , \text{ for } 3^{rd} \text{ component} \\ s + 2 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

For even s the color codes of $sP(4,2)$ are:

$$c_{\Pi}(u_4^t) = \begin{cases} 1 & , \text{ for } 1^{st}, 2^{nd} \text{ and } 4^{th} \text{ component} \\ 0 & , \text{ for } 3^{rd} \text{ component} \\ s & , \text{ for } 5^{th} \text{ component} \end{cases}$$

$$c_{\Pi}(v_2^t) = \begin{cases} 1 & , \text{ for } 1^{st} \text{ component} \\ 2 & , \text{ for } 2^{nd} \text{ and } 4^{th} \text{ component} \\ 0 & , \text{ for } 3^{rd} \text{ component} \\ s + 2 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

$$c_{\Pi}(v_3^t) = \begin{cases} 2 & , \text{ for } 1^{st} \text{ and } 4^{th} \text{ component} \\ 1 & , \text{ for } 2^{nd} \text{ component} \\ 0 & , \text{ for } 3^{rd} \text{ component} \\ s + 2 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

(d) $C_4 = \{v_3^t | \text{for odd } s\} \cup \{u_1^t | \text{for even } s\} \cup \{v_4^t | \text{for odd } s \geq 3\}$

For odd s the color codes of $sP(4,2)$ are:

$$c_{\Pi}(v_3^t) = \begin{cases} 2 & , \text{ for } 1^{st} \text{ and } 2^{nd} \text{ component} \\ 1 & , \text{ for } 3^{rd} \text{ component} \\ 0 & , \text{ for } 4^{th} \text{ component} \\ s + 2 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

For odd $s \geq 3$ the color codes of $sP(4,2)$ are:

$$c_{\Pi}(v_4^t) = \begin{cases} 2 & , \text{ for } 1^{st} \text{ component} \\ 1 & , \text{ for } 2^{nd} \text{ and } 3^{rd} \text{ component} \\ 0 & , \text{ for } 4^{th} \text{ component} \\ s + 1 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

For even s the color codes of $sP(4,2)$ are:

$$c_{\Pi}(v_4^t) = \begin{cases} 1 & , \text{ for } 1^{st}, 2^{nd} \text{ and } 3^{rd} \text{ component} \\ 0 & , \text{ for } 4^{th} \text{ component} \\ s + 1 & , \text{ for } 5^{th} \text{ component} \end{cases}$$


(e) $C_5 = \{v_4^1\}$

$$c_{\Pi}(v_4^1) = \begin{cases} 2 & , \text{ for } 1^{st} \text{ component} \\ 1 & , \text{ for } 2^{nd} \text{ and } 3^{rd} \text{ component} \\ 3 & , \text{ for } 4^{th} \text{ component} \\ 0 & , \text{ for } 5^{th} \text{ component} \end{cases}$$

8

Since all the vertices have different color codes, c is a locating coloring of generalized Petersen graphs $sP(4,2)$, so $\chi_L(sP(4,2)) = 5$, for even $s \geq 2$. □

In figure 1 is illustrated a locating coloring of generalized Petersen graphs $4P(4,2)$ with the locating chromatic number 5.



6

Figure 1. A minimum locating coloring of $4P(4,2)$

3. Conclusion

Based on the results, locating chromatic number of generalized Petersen graph $sP(4,2)$ is 5 for $s \geq 2$.

References

- [1] Chartrand G, Erwin D, Henning M, Slater P and Zhang P 2002 The locating-chromatic number of a graph *Bull. Inst. Combin. Appl.* **36** pp 89–101
- [2] Chartrand G, Erwin D, Henning M, Slater P, and Zhang P 2003 Graph of order n with locating-chromatic number $n-1$ *Discrete Math* **269** 1-3 pp 65–79
- [3] Behtoei A and Omoomi B 2011 On the locating chromatic number of kneser graphs *Discrete Applied Mathematics* **159** pp 2214–2221
- [4] Baskoro E T and Purwasih I 2012 The locating-chromatic number for corona product of graphs *Southeast-Asian J. Of Sciences* **1** pp 124–134
- [5] Asmiati 2017 Locating chromatic number of banana tree *International Mathematical Forum* **12** (1) pp 39–45
- [6] Asmiati, Assiyatun H and Baskoro E T 2011 Locating-chromatic number of amalgamation of stars *ITB J.of Sci.* **43** 1 pp 1–8

- [7] Asmiati, Baskoro E T, Assiyatun H, Suprijanto D, Simanjuntak R, and Uttunggadewa S 2012 The Locating-chromatic number of firecracker graphs *Far East Journal of Mathematical Sciences* **63** 1 pp 11 – 23
- [8] Syofyan D K, Baskoro E T, Assiyatun H 2013 The locating-chromatic number of homogeneous lobsters *AKCE Int. J. Graphs Comb.* **10** 3 pp 215 – 252
- [9] Asmiati 2014 The locating-chromatic number of non-homogeneous amalgamation of stars *Far East Journal of Mathematical Sciences* **93** 1 pp 89 – 96
- [10] Asmiati 2016 On the locating-chromatic numbers of non-homogeneous caterpillars and firecrackers graphs *Far East Journal of Mathematical Sciences* **100** 8 pp 1305 – 1316
- [11] Asmiati, Wammiliana, Devriyadi and Yulianti L 2017 On some Petersen graphs having locating chromatic number four or five *Far East Journal of Mathematical Sciences* **102** 4 pp 769 – 778
- [12] Watkins M E 1969 A theorem on tait colorings with application to the generalized Petersen graphs *Journal of Combinatorial Theory* **6** pp 152-164

The Locating-Chromatic Number for Certain Operation of Generalized Petersen Graphs $A_n(A_m)$ (A_n, A_m)

ORIGINALITY REPORT

20%

SIMILARITY INDEX

PRIMARY SOURCES

- 1** Manal Ghanem, Hasan Al-Ezeh, Ala'a Dabbour. "Locating Chromatic Number of Powers of Paths and Cycles", Symmetry, 2019
Crossref 67 words — 3%
- 2** fib-indonesia.org
Internet 50 words — 3%
- 3** Welyyanti, Des, Edy Tri Baskoro, Rinovia Simanjuntak, and Saladin Uttunggadewa. "On Locating-chromatic Number for Graphs with Dominant Vertices", Procedia Computer Science, 2015.
Crossref 37 words — 2%
- 4** www.m-hikari.com
Internet 32 words — 2%
- 5** www.ijc.or.id
Internet 21 words — 1%
- 6** Purwasih, Ira Apni, Edy Tri Baskoro, Hilda Assiyatun, and Djoko Suprijanto. "The Bounds on the Locating-Chromatic Number for a Subdivision of a Graph on One Edge", Procedia Computer Science, 2015.
Crossref 21 words — 1%
- 7** export.arxiv.org
Internet 20 words — 1%
- 8** Syofyan, Dian Kastika, Edy Tri Baskoro, and Hilda Assiyatun. "The Locating-Chromatic Number of Binary 19 words — 1%

-
- 9 S. C. LÓPEZ, F. A. MUNTANER-BATLE, M. RIUS-FONT. "BI-MAGIC AND OTHER GENERALIZATIONS OF SUPER EDGE-MAGIC LABELINGS", Bulletin of the Australian Mathematical Society, 2011 18 words — 1%
Crossref
-
- 10 Martin Bača, Mirka Miller, Joe Ryan, Andrea Semaničová-Feňovčíková. "Magic and Antimagic Graphs", Springer Science and Business Media LLC, 2019 17 words — 1%
Crossref
-
- 11 re.public.polimi.it 16 words — 1%
Internet
-
- 12 Asmiati. "Locating chromatic number of banana tree", International Mathematical Forum, 2017 13 words — 1%
Crossref
-
- 13 id.scribd.com 11 words — 1%
Internet
-
- 14 A M Purnamasari, C E A Pah, M D I Yoga, A S Girsang, S M Isa. "Business Intelligent in an E-Commerce Industry", IOP Conference Series: Materials Science and Engineering, 2019 10 words — 1%
Crossref
-
- 15 Communications in Computer and Information Science, 2011. 9 words — < 1%
Crossref
-
- 16 perso.telecom-paristech.fr 9 words — < 1%
Internet
-
- 17 digilib.unila.ac.id 9 words — < 1%
Internet
-
- 18 Ali Behtoei, Behnaz Omoomi. "On the locating chromatic number of Kneser graphs", Discrete Applied Mathematics, 2011 8 words — < 1%

19 Changping Wang. "Lower negative decision number in a graph", Journal of Applied Mathematics and Computing, 08/12/2009 6 words — < 1%
Crossref

20 Minhong Wang, William K. Cheung, Jiming Liu, Xiaofeng Xie, Zongwei Luo. "Chapter 19 E-Service/Process Composition Through Multi-agent Constraint Management", Springer Science and Business Media LLC, 2006 6 words — < 1%
Crossref

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF