



## Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Lilis Hermida  
Assignment title: pdf files  
Submission title: Effect of Slow Release Nitrogen (SR...  
File name: lilis-rugayahFINAL\_PROCEEDINGS...  
File size: 1.27M  
Page count: 9  
Word count: 4,201  
Character count: 20,406  
Submission date: 23-Jan-2021 05:33AM (UTC+0700)  
Submission ID: 1492512896



INTERNATIONAL SYMPOSIA  
ON HORTICULTURE 458  
Empowering Horticulture and Horticulture Science  
Advancing Horticulture and Horticulture Science

---

**Effect of Slow Release Nitrogen (SRN) Addition to NPK Fertilization on Cauliflower Growth**

*RUGA YAH I, HERMIDA Lilis<sup>1</sup>, GINTING Yohannes Cahya<sup>1</sup>, AGUSTIAN Joni<sup>2</sup>, and SURYANI Afrida<sup>1</sup>*

<sup>1</sup> Department of Agrotechnology, Faculty of Agriculture, University of Lampung  
<sup>2</sup> Department of Chemical Engineering, Faculty of Engineering, University of Lampung  
Prof. Dr. Soemanto Brjongsoro No. 1 Bandar Lampung 35145  
Email: rugayah\_lilis@yahoo.co.id

**Abstract.** Urea is fertilizer nitrogen source that is high water solubility, so its not efficient. In this study, three kinds of slow release nitrogen (SRN) fertilizers were evaluated for cauliflower growth. These fertilizer is the standard fertilization, while single N is addition. Hence, in the discussion should focus on the effect of such nitrogen addition. The experiment was carried out in a greenhouse, using randomized block design in 5x2 factorial scheme with three replicates. Treatments corresponded to five kinds of nitrogen sources (a : without fertilizer (A0), Urea (A1), SRN-Bentahite (A2), SRN-BBA (A3) and SRN-Mesopore (A4) and two treatments of NPK without NPK (B0) and NPK addition (B1). The dosage of fertilizer is equivalent to Urea 200 kg/ha-1. Data obtained were identified using analysis of variance and followed by Orthogonal Contrast at 5% level. It was found that the use of fertilizer nitrogen significantly increased the growth of cauliflower compared to without fertilizer. There is any difference between urea and SRN application on the number of leaves and cord weight. The growth of cauliflower fertilized by the SRN's with the NPK addition was better than that without NPK, especially the number of leaves, leaf length, leaf width, root dry weight, cord diameter, and cord weight. The results also showed that the effects of SRN types were no significant differences on the cauliflower growth but on the basis of variable observed, SRN-Mesopore agronomically was potential to be developed compared to SRN-Bentahite and SRN-BBA.

**Keywords:** nitrogen, addition of NPK, fertilizer, cauliflower

**1. Introduction**

Cauliflower (*Brassica oleracea* var. *Botrytis* L.) is one of the horticultural products that has complete nutritional. Nutrient content in 100 grams of cauliflower are calories (25.0 cal), protein (2.1 g), fat (0.3 g), carbohydrates (5.2 g), calcium (41 mg), phosphorus (42, 0 mg), iron (1.1 mg), vitamin A (23 RE), vitamin B2 (0.12 mg), vitamin C (62 mg), niacin (0.7 mg) [1]. These nutrients are needed by human body to meet the needs of a healthy life. In addition cauliflower also has the benefits of being an antioxidant, anticancer, reducing estrogen levels, neutralizing carcinogens, and improving bowel movements.

Based on data from the Central Statistics Agency (BPS), the average production of flower cabbage in Indonesia in 2011- 2013 were 12.02 tons/ha-1, 11.53 tons/ha-1, and 12.18 tons ha-1 respectively, but in 2014 the production decreased to 12.08 tons. ha-1 [2] due to less effective of fertilization. Fertilization will be effective if soil and fertilizer conditions support each other. The most common soil conditions in Indonesia are hard soil conditions, high acidity (average pH <4.50), high saturation of Al, low organic matter [3]. This soil condition is included in the Unisol soil type, which is poor in nutrients and difficult to be fertilized due to intensive alkaline washing. Fertilizers that are easily absorbed by plants are fertilizers that are not volatile, washed, or bound by other substances.

# Effect of Slow Release Nitrogen (SRN) Addition to NPK Fertilization on Cauliflower Growth

*by* Lilis Hermida

---

**Submission date:** 23-Jan-2021 05:33AM (UTC+0700)

**Submission ID:** 1492512896

**File name:** lilis-rugayahFINAL\_PROCEEDINGS\_OF\_ISH\_2018\_edit\_4\_turnitin.pdf (1.27M)

**Word count:** 4201

**Character count:** 20406

## Effect of Slow Release Nitrogen (SRN) Addition to NPK Fertilization on Cauliflower Growth

<sup>1,2</sup> GAYAH<sup>1</sup>, HERMIDA Lilis<sup>2</sup>, GINTING Johannes Cahya<sup>1</sup>, AGUSTIAN Joni<sup>2</sup>, and SURYANI Afrida<sup>1</sup>

<sup>1</sup> Department of Agrotechnology, Faculty of Agriculture, University of Lampung

<sup>2</sup> Department of Chemical Engineering, Faculty of Engineering, University of Lampung

Prof. Dr. Soemantri Brojonegoro No.1 Bandar Lampung 35145

Email: [ruguayah\\_unila@yahoo.co.id](mailto:ruguayah_unila@yahoo.co.id)

**Abstract.** Urea is fertilizer nitrogen source that is high water solubility, so its not efficient. In this study, three kinds of slow release nitrogen (SRN) fertilizers were evaluated for cauliflower growth. These fertilizer is the standard fertilization, while single <sup>6</sup> is addition. Hence, in the discussion should focus on the effect of such nitrogen addition. The experiment was carried out in a greenhouse, using randomized block design in 5x2 factorial scheme with three replicates. Treatments corresponded to five kinds of nitrogen sources i.e : without fertilizer (A0), Urea (A1), SRN-Bentonite (A2), SRN-BBA (A3) and SRN-Mesopore (A4) and two treatments of NPK: without NPK (B0) and NPK addition (B1). The dosage of fertilizer is equivalent to Urea 200 kg.ha-1. Data obtained were identified <sup>17</sup> using analysis of variance and followed by Orthogonal Contrast at 5% level. It was found that the use of fertilizer nitrogen significantly increased the growth of cauliflower compared to without fertilizer. There is any difference between urea and SRN application on the number of leaves and curd weight. The growth of cauliflower fertilized by the SRN's with the NPK addition was better than that without NPK, especially the number of leaves, leaf length, leaf width, root dry weight, curd diameter, and curd weight. The results also showed that the effects of SRN- types were no significant differences on the cauliflower growth but on the basis of variable observed, SRN-Mesopore agronomically was potential to be developed compared to SRN-Bentonite and SRN-BBA.

<sup>2</sup>  
**Keywords:** nitrogen, addition of NPK, fertilizer, cauliflower

### 1. Introduction

Cauliflower (*Brassica oleraceae* var. *Botrytis* L.) is one of the horticultural products that has complete nutritional. Nutrient content in 100 grams of cauliflower are calories (25.0 cal), protein (2.1 g), fat (0.3 g), carbohydrates (5.2 g), calcium (41 mg), phosphorus (42, 0 mg), iron (1.1 mg), vitamin A (23 RE), vitamin B2 (0.12 mg), vitamin C (82 mg), niacin (0.7 mg) [1]. These nutrients are needed by human body to meet the needs of a healthy life. In addition cauliflower also has the benefits of being an antioxidant, anticancer, reducing estrogen levels, neutralizing carcinogens, and improving bowel movements.

Based on data from the Central Statistics Agency (BPS), the average production of flower cabbage in Indonesia in 2011- 2013 were 12.02 tons.ha-1, 11.53 tons.ha-1, and 12.18 tons ha-1 respectively, but in 2014 the production decreased to 12.08 tons. ha-1 [2] due to less effective of fertilization.. Fertilization will be effective if soil and fertilizer conditions support each other. The most common soil conditions in Indonesia are hard soil conditions, high acidity (average pH <4.50), high saturation of Al, low organic matter [3]. This soil condition is included in the Ultisol soil type, which is poor in nutrients and difficult to be fertilized due to intensive alkaline washing. Fertilizers that are easily absorbed by plants are fertilizers that are not volatile, washed, or bound by other substances.

Fertilization is the addition of nutrients needed by plants because of lack availability in the soil for optimal plant growth. Nutrients that function for tissue formation are nitrogen elements [4]. This nitrogen nutrient is needed in large quantities, especially in vegetable crops. The highest type of fertilizer containing nitrogen is urea and is widely used by farmers. Besides Urea, other N sources fertilizer which also contain elements of P and K are Nitrofoska or NPK compounds. Urea fertilizer has a nitrogen content of 45% which has very hygroscopic so that it is easily soluble and volatile before is absorbed by plants.

Efforts to improve the efficiency of providing nitrogen nutrients in the soil are by modifying the physical and chemical forms of conventional urea fertilizer into urea fertilizer slowly (Slow Release urea = SRU). Slow-release urea fertilizer is a fertilizer with the mechanism of release of nitrogen elements periodically following the pattern of nutrient absorption by plants. The mechanism of urea fertilizer works slowly (Slow release nitrogen = SRN), which is by coating the urea fertilizer with water-insoluble compounds such as sulphur, polyethylene, alkyd resin, polyurethane [5]; [6] or by melting urea and then mixing it bentonite as with a substrate: and polyacrylamides a binder [7]. The release of nutrients is in accordance with the time and amount needed by the plant, as well as maintaining the presence of fertilizer in the soil so that absorption by plants is more optimal. In this study SRN fertilizer types that were utilized were synthesized by melting urea and then mixing it with one of various substrates (bentonite, silica from bagasse bottom ash ((BBA), mesostructuredcellulair form silica) and corn starch as a binder [8]. SRN using bentonitewas named SRN-Bentonite, using silica from bagasse bottom ash (BBA) was named SRN-BBA and using mesostructuredcellulair form silica was named SRN-Mesopore.

## 2. Methodology

This research was carried out in the greenhouse of the Faculty of Agriculture Lampung University in September to November 2017. The treatment was applied in factorial pattern (5x2) in a randomized block design (RBD) with 3 replications. The first factor is the type of source fertilizer N (A): no fertilizer (A0), Urea (A1), SRN- Bentonite (A2), SRN- BBA (A3), and SRN- Mesopori (A4). The second factor is the addition of NPK (B): without NPK (B0) and addition of NPK (B1). The dosage of fertilizer given is equivalent to Urea 200 kg ha<sup>-1</sup> and NPK 600 kg ha<sup>-1</sup>. The application of SRN fertilizer was carried out at the age of one week after planting and NPK fertilizer 10 days after planting each with the amount of 1.6 g Urea, 1.8 g SRN, and 5 g NPK per plant.

The variables observed included: plant height, leaf number, leaf length, leaf width, root dry weight, curd diameter, and curd weight. Data were analyzed by variance, followed by Orthogonal Contrast at 5% level.

## 3. Results And Discussion

### 3.1. Results

The results showed that the various types of nitrogen source fertilizers produced higher growth of cauliflower than without fertilization, which was shown on variables: plant height, leaf number, leaf length, leaf width, root dry weight, curd diameter, and curd weight. Among the types of nitrogen sources fertilizer do not show a significant difference, except for the number of leaves and curd weight of urea fertilizer results are higher than SRN, but if added NPK, SRN fertilizer is the higher yield.

### 3.1.1. Plant height

The results showed that the various types of nitrogen source fertilizers produced higher plant heights than without fertilization, but among the types of fertilizers used did not show differences in both single N fertilizer and added of NPK fertilizer compounds (Table 1).

Table 1. Effect of type N fertilizer and NPK addition on plant height

Comparison	Difference		F <sub>count</sub>
	(cm)	(%)	
P1: No fertilizer vs. Fertilization	40.36-48.90= -8.54	17.46	23.73*
P2: Urea vs SRN	52.01-48.43= 3.58	7.40	0.97 <sup>10</sup>
P3: Bentonit vs BBA and Mesopori	43.22-51.04= -7.81	15.31	1.39 <sup>tn</sup>
P4: BBA vs Mesopori	50.68-51.09= -0.71	1.39	0.08 <sup>tn</sup>
P5: Urea vs Urea +NPK	52.01-47.85= 4.16	8.70	1.46 <sup>tn</sup>
P6: SRN vs SRN+NPK	48.53-48.79= -0.36	0.74	0.29 <sup>tn</sup>
P7: Urea +NPKvs SRN+NPK	47.85-48.79= -0.71	1.45	0.76 <sup>tn</sup>
P8: Bentonit +NPKvs BBA +NPK and Mesopori+NPK	46.14-50.12= -3.97	7.93	1.72 <sup>tn</sup>
P9: BBA +NPK <sup>1</sup> Mesopori+NPK	46.55-53.69= -3.57	6.65	4.36 <sup>tn</sup>

Descript<sup>1</sup>: \* = significantly different at level  $\alpha$  5%

tn = not significantly different at level  $\alpha$  5%

### 3.1.2. Number of leaves

The results showed that the various types of nitrogen source fertilizers increased the number of leaves of the cauliflower compared to no fertilizer. Among the various types of nitrogen source fertilizer, urea fertilizer significantly produces more leaves than SRN fertilizer. Mesopori SRN fertilizer and BBA (Bagasse Bottom Ash) significantly produce more leaves compared to SRN- Bentonite fertilizer. However, there is no difference in the number of leaves between Mesopori fertilizer and BBA (Bagasse Bottom Ash). The type of SRU and addition of NPK significantly produced more leaves than without fertilizer of NPK (Table 2).

Table 2. Effect of type N fertilizer and NPK addition on number of leaves

Comparison	Difference		F <sub>count</sub>
	(strand)	(%)	
P1: No fertilizer vs. Fertilization	14.83-16.76= -1.93	11.63	8.12*
P2: Urea vs SRN	17.83-14.89= 2.94	19.78	15.82*
P3: Bentonit vs BBA and Mesopori	13.50-15.58= -2.08	13.37	7.04*
P4: BBA vs Mesopori	14.83-16.33= -1.50	9.18	2.74 <sup>tn</sup>
P5: Urea vs Urea +NPK	17.83-18.67= -0.83	4.46	0.84 <sup>tn</sup>
P6: SRN vs SRN+NPK	14.83-17.44= -2.56	14.65	23.83*
P7: Urea +NPKvs SRN+NPK	18.67-17.44= 1.22	6.98	2.73 <sup>tn</sup>
P8: Bentonit +NPKvs BBA +NPK and Mesopori+NPK	17.50-17.42= 0.08	0.48	0.01 <sup>tn</sup>
P9: BBA +NPK <sup>1</sup> Mesopori+NPK	17.33-17.50 = -0.17	0.95	0.03 <sup>tn</sup>

Descript<sup>1</sup>: \* = significantly different at level  $\alpha$  5%

tn = not significantly different at level  $\alpha$  5%

### 3.1.3. Leaf length

The results showed that the various types of nitrogen source fertilizer increased the length of the leaves of the cauliflower compared to without fertilizer. SRN fertilizer with the addition of NPK significantly produced a longer leaf length than SRN fertilizer alone (Table 3).

### 3.1.4. Leaf width

The results showed that the various types of nitrogen source fertilizers increased the width of the leaves in the flower beds compared to without fertilizer. Among the various types of nitrogen source fertilizers, fertilizers actually produce more water than fertilizer SRN. SRN fertilizer with the addition of NPK significantly produced a wider yield than SRU fertilizer alone (Table 4).

### 3.1.5. Root dry weight

The results showed that the various types of nitrogen source fertilizers produced the dry weight of roots of heavier cabbage plants compared to without fertilizer. Among the administration of various types of nitrogen source fertilizers, SRU with the addition of NPK produces heavy dry weight compared to SRN fertilizer alone (Table 5).

Table 3. Effect of N source fertilizer and NPK addition on leaf length

Comparison	Difference		F <sub>count</sub>
	(cm)	(%)	
P1: No fertilizer vs. Fertilization	12.38-18.63=-7.53	39.33	45.22*
P2: Urea vs SRN	20.00-18.17=1.83	7.35	2.21 <sup>tn</sup>
P3: Bentonit vs BBA and Mesopori	17.04-18.74=-1.70	9.34	1.70 <sup>tn</sup>
P4: BBA vs Mesopori	18.19-19.29=-1.11	5.63	0.54 <sup>tn</sup>
P5: Urea vs Urea +NPK	20.00-20.77=-0.77	3.92	0.26 <sup>tn</sup>
P6: SRN vs SRN+NPK	18.17-21.26=-3.09	14.62	12.68*
P7: Urea +NPKvs SRN+NPK	20.77-21.26=-0.49	2.32	0.16 <sup>tn</sup>
P8: Bentonit +NPKvs BBA +NPK and Mesopori+NPK	19.71-22.07=-2.36	10.97	3.21 <sup>tn</sup>
P9: BBA +NPK vs Mesopori+NPK	21.26-22.04=-0.75	3.40	0.25 <sup>tn</sup>

Descript<sup>1</sup>: \* = significantly different at level  $\alpha$  5%

tn = not significantly different at level  $\alpha$  5%

Table 4. Effect of N source fertilizer and NPK addition on leaf width

Comparison	Difference		F <sub>count</sub>
	(cm)	(%)	
P1: No fertilizer vs. Fertilization	9.21-21.75=-3.55	27.80	58.18*
P2: Urea vs SRN	13.79-12.43=1.36	10.98	10.34*
P3: Bentonit vs BBA and Mesopori	12.35-11.21=-1.15	9.57	3.64 <sup>tn</sup>
P4: BBA vs Mesopori	11.92-12.79=-0.88	6.81	1.59 <sup>tn</sup>
P5: Urea vs Urea +NPK	13.79-13.92=-0.13	0.97	0.03 <sup>tn</sup>
P6: SRN vs SRN+NPK	11.97-13.93=-1.96	14.06	23.95*
P7: Urea +NPKvs SRN+NPK	13.92-13.93=-0.01	0.10	0.00 <sup>tn</sup>
P8: Bentonit +NPKvs BBA +NPK and Mesopori+NPK	13.17-13.93=-0.76	5.48	3.64 <sup>tn</sup>
P9: BBA +NPK vs Mesopori+NPK	14.04-14.58=-0.54	3.78	0.61 <sup>tn</sup>

Descript<sup>1</sup>: \* = significantly different at level  $\alpha$  5%

tn = not significantly different at level  $\alpha$  5%

### 3.1.6. Curd Diameter

The results showed that the various types of nitrogen source fertilizers increased the diameter of the curd of cauliflower compared to no fertilizer. Mesopori SRU fertilizer and BBA significantly produced higher curd diameter than Bentonite SRU fertilizer. SRU fertilizer with the addition of NPK significantly produced higher curd diameter than without fertilizer of NPK (Table 6).



Table 5. Effect of types N fertilizer and NPK addition on the dry weight of oots

Comparison	Difference		F <sub>count</sub>
	(cm)	(%)	
P1: No fertilizer vs. Fertilization	1.54-3.81=-2.27	59.60	21.43*
P2: Urea vs SRN	4.34-3.65=0.69	18.76	2.18 <sup>tn</sup>
P3: Bentonit vs BBA and Mesopori	3.11-3.42=-0.31	9.18	0.55 <sup>tn</sup>
P4: BBA vs Mesopori	3.56-3.61=-0.05	1.31	0.00 <sup>tn</sup>
P5: Urea vs Urea +NPK	4.34-4.73=-0.39	10.01	0.29 <sup>tn</sup>
P6: SRN vs SRN+NPK	3.42-4.40=-0.98	21.78	5.35*
P7: Urea +NPKvs SRN+NPK	4.73-4.40=0.33	7.34	0.30 <sup>tn</sup>
P8: Bentonit +NPKvs BBA +NPK and Mesopori+NPK	4.41-4.53=-0.39	9.49	0.38 <sup>tn</sup>
P9: BBA +NPK <sup>1</sup> Mesopori+NPK	4.13-4.94=-0.81	17.80	1.22 <sup>tn</sup>

Descript <sup>1</sup>: \* = significantly different at level  $\alpha$  5%

tn = not significantly different at level  $\alpha$  5%

Table 6. Effect of types N fertilizer and NPK addition on curd diameter

Comparison	Difference		F <sub>count</sub>
	(cm)	(%)	
P1: No fertilizer vs. Fertilization	7.00-9.05 = -2.05	22.66	12.77*
P2: Urea vs SRN	9.83-8.80 = 1.03	11.69	3.86 <sup>tn</sup>
P3: Bentonit vs BBA and Mesopori	7.22-8.46 = -1.24	14.71	6.35*
P4: BBA vs Mesopori	8.79-9.38 = -0.58	6.26	0.46 <sup>tn</sup>
P5: Urea vs Urea +NPK	9.83-10.50 = - 0.67	7.42	0.61 <sup>tn</sup>
P6: SRN vs SRN+NPK	8.46-9.51 = -1.05	10.79	4.54*
P7: Urea +NPKvs SRN+NPK	10.50-9.51 = 0.99	10.10	1.99 <sup>tn</sup>
P8: Bentonit +NPKvs BBA +NPK and Mesopori+NPK	9.25-9.65 = - 0.40	4.16	0.29 <sup>tn</sup>
P9: BBA +NPK <sup>1</sup> Mesopori+NPK	9.54-9.75 = - 0.21	2.16	0.06 <sup>tn</sup>

Descript <sup>1</sup>: \* = significantly different at level  $\alpha$  5%

tn = not significantly different at level  $\alpha$  5%

### 3.1.7. Curd Weight

The results showed that the various types of nitrogen fertilizer increased the curd weight of the cabbage flower compared to no fertilizer. Among various types of nitrogen source fertilizers, urea actually produces heavier curd weights than SRU fertilizer. The addition of NPK produced heavier curd weights in both urea and SRU fertilizers than compared with no addition of NPK (Table 7).

Table 7. Effect of types N fertilizer and NPK addition on curd weight

Comparison	Difference		F <sub>count</sub>
	(g)	(%)	
P1: No fertilizer vs. Fertilization	46.85-98.82= -51.97	52.59	29.35*
P2: Urea vs SRN	107.45-85.63=21.83	25.49	6.21*
P3: Bentonit vs BBA and Mesopori	62.82-78.35= -15.54	19.83	3.54 <sup>tn</sup>
P4: BBA vs Mesopori	85.58-86.66= - 1.09	1.17	0.01 <sup>tn</sup>
P5: Urea vs Urea +NPK	107.45-139.11= - 31.66	31.91	4.90*
P6: SRN vs SRN+NPK	78.35-120.07=-41.72	33.42	25.54*
P7: Urea +NPKvs SRN+NPK	139.11-120.07=19.04	15.25	2.66 <sup>tn</sup>
P8: Bentonit +NPKvs BBA +NPK and Mesopori+NPK	111.19-120.07=-8.88	7.40	1.16 <sup>tn</sup>
P9: BBA +NPK <sup>1</sup> Mesopori+NPK	127.90-121.12= 6.78	5.44	0.22 <sup>tn</sup>

Descript <sup>1</sup>: \* = significantly different at level  $\alpha$  5%

tn = not significantly different at level  $\alpha$  5%

### 3.2. Discussion

The provision of various types of nitrogen source fertilizer and the addition of NPK serves to provide nutrients for both vegetative and generative plant growth. The availability of nutrients in the soil will affect plant growth. Vegetative growth of plants can be stimulated by applying nitrogen source fertilizer [9]. Nitrogen elements play a role in the formation of chlorophyll, fat, protein, other assimilates needed by plants [10]. Besides N, P and K elements are also needed for flower cabbage plants. According [11] P elements play a role in photosynthesis, respiration, electron transfer, root extension, storage of energy and increase the efficiency of N elemental functions. The element K plays a role in the metabolic process as photosynthetic regulator, transport of nutrients from root to leaf, and translocate assimilate throughout plant parts. Flower cabbage plants require the availability of high amounts of nutrients for metabolic processes during their life cycle, especially the final phase of growth. [12] recommended fertilizer for cauliflower is 300 kg SP36 ha, 225 kg ha KCl, and 200 kg ha urea. The administration of urea fertilizer compared to slow release fertilizer (SRU) shows better growth, especially in the number of leaves and curd weight. This increase in growth is due to the availability of N nutrients from urea available more quickly and at the beginning of the growth of flower cabbage requires large amounts of N elements so that the administration of urea at the age of one week after planting is very helpful in the supply of nitrogen. However, if the source N fertilizer is combined with NPK, the slow release nitrogen fertilizer shows better results than urea fertilizer. The addition of NPK fertilizer causes the availability of balanced nutrients and in sufficient quantities is met for the growth of flower cabbage plants. The N and K fertilizers can increase the activity of PEP carboxylase in leaves, although the role of N in stimulating the activity of photosynthetic enzymes is more dominant than K. P elements also play a role in the formation of ATP energy which will then be used for photosynthetic translocation to the part of the plant organ that needs it. The P element contained in NPK plays an important role in the formation and growth of mustard roots, so that the roots are better able to absorb more water and nutrients and in the end the plant will grow and develop better overall.

The SRU fertilizer and addition of NPK produced better growth than without NPK, especially in the number of leaves, leaf length, leaf width, root dry weight, curd diameter, and curd weight. While the better growth of urea and NPK fertilizers is only seen in the number of leaves and curd weight. This is because plant growth is dominated by vegetative growth of plants such as number of leaves, leaf size and root system development. Increasing the nitrogen dose increases the leaf area index, relative growth rate, and the rate of assimilation. which in turn will increase the resulting photosynthetic to be translocated to the curd so that the weight increases. The addition of NPK, causes the supply of K, which functions in regulating the pattern of photosynthetic partition so that photosynthetic partitions are more directed to the economic part of the curd.

The absorption of N by cauliflower was very low before the leaf formation stage to -4 to 6 and there was a tendency to increase absorption of N at the beginning of curd initiation and then continue to increase until harvest [13]. In general, the SRU fertilizer has the potential to be developed as an alternative source of N fertilizer, although in the cauliflower plants need additional sources of other elements, such as P and K to support plant growth. The added of P and K as basic fertilizers serves to supplement the nutrient requirements other than N from the SRU fertilizer. Another advantage with the use of SRU is that it saves the use of labor for fertilization because of its pellet shape, while the urea fertilizer is prill-shaped.

Cauliflower depends on N fertilizer and excessive application of N rarely has a negative impact. Increasing the number of leaves and the size of the leaves will increase the chlorophyll content in the leaves. This chlorophyll will capture sunlight in photosynthesis. The photosynthesis process will produce photosynthates which are translocated to increase leaf size. The development and increase in leaf size



are influenced by the availability of water and the dissolved nutrients are then transported to the upper part of the plant and some will be used to increase the pressure of leaf cell turgor [14].

Overall this study shows that various sources of nitrogen (N) and adding NPK can increase the growth of cauliflower. N source fertilizers from both urea or SRU fertilizers and addition of NPK produce heavier curd weights compared to without NPK. The use of SRN fertilizer and NPK increased the curd weight with 3 leaves by 41.72 grams or 33.42% compared to without NPK. The urea fertilizer with the addition of NPK increased the weight of curd (31.66 grams or 31.91%) compared without NPK. This shows that the use of SRU fertilizer and addition of NPK has the potential to be developed and used for fertilizing crops, especially Mesopori SRN fertilizer. In connection with the addition of NPK fertilizer in this study, which did not use basic fertilizer, for the next study can be tested the use of SRN which is added basic fertilizer, especially the source elements of P and K.

8

#### 4. Conclusions and Suggestions

##### 4.1. Conclusion

Based on the results of the research that has been done, the following conclusions are obtained:

The provision of various types of N fertilizer resulted in the growth of plant height, leaf number, leaf length, leaf width, root dry weight, curd diameter, and curd weight higher than without fertilization. Addition of NPK increases the growth of cauliflower compared to without NPK, especially in curd diameter and curd weight. The provision of N-source fertilizers from both urea and SRN and addition of NPK further increases the growth and yield of cauliflower compared to without NPK.

##### 4.2. Recommendations

The advice given for the next research is that it is necessary to try the use of SRN fertilizer in the field with the addition of basic fertilizer containing P and K elements because in this study no basic fertilizer was used.

#### 5. References

- [1] Pangkalan Ide. 2011. Health Secret of Broccoli. PT Elex Media Komputindo Gramedia Group. Jakarta. 156–167 p.
- [2] Central Statistics Agency. 2015. Production Data of Cauliflower [internet] Jakarta BAS; [downloaded November 26 2017] is available at: [www.bps.go.id](http://www.bps.go.id).2015
- [3] Prasetyo, B. H. and D. A.Suriadikarta. 2006. Characteristics, Potential, and Technology of Ultisol Soil Management for the Development of Dry Land Agriculture in Indonesia. *Agricultural Research and Development*. 2 (25). 39 p.
- [4] Sutejo, M.M. 2010. *Fertilizers and Fertilization Methods*. Rienekacipta. Jakarta. 8–150 p
- [5] Ali, S., Danafar. F., 2015. Controlled-release fertilizers: Advances and challenges. *Life Sci. J.*,12 (11). [www.lifesciencesite.com/lj/life121115/005\\_17565\\_life\\_121115\\_33\\_45.pdf](http://www.lifesciencesite.com/lj/life121115/005_17565_life_121115_33_45.pdf)
- [6] Salman, O.A., 1989. Polyethylene-coated urea: 1. Improved storage and handling properties. *Ind. Eng. Chem. Res.* 28: 630--632.
- [7] Xiaoyu, N., Yuejin, W., Zhengyan, W., Lin, W., Guannan, Q., Lixiang, Y., 2013. A novel slow release urea fertilizer: Physical and chemical analysis of its structure and study of its release mechanism. *Biosystems Eng.* 115: 274--282.

- [8] Hermida, L. and Agustian, J. 2019- Slow release urea fertilizer synthesized through recrystallization of urea incorporating natural bentonite using various binders. *Environmental Technology & Innovation*. 13: 113—121.
- [9] Istiana, H. 2007. How to apply nitrogen fertilizer and its effects on Madurese tobacco plants. *Agricultural Engineering Bulletin*. 12 (2).
- [10] Marsono. 2002. *Instructions for Using Fertilizer. Self-helpers*. Jakarta. 160 p.
- [11] Winarso, S. 2005. *Soil Fertility: Basic Health and Soil Quality*. Gava Media. Yogyakarta. 269 p.
- [12] Rubatzky V.E and M. Yamaguchi. 1998. *World Vegetables Principles, Production and Nutrition* Volume II. Institute of Technology Bandung. 292 p
- [13] Rahayu, Y.S. and N. Nurlenawati. 2011. Effect of combination of nitrogen cattle fertilizer dose on growth and yield of cauliflower (*Brassica oleracea* var. *Botrytis* L) sub var cauliflower PM 126 F1 cultivars in the lowlands in the dry season. *Plant Production Journal* 5 (3).
- [14] Lakitan, B. 2004. *Introduction of Plant Physiology*. PT Raja Grafindo Persada, Jakarta.


## **6. Acknowledment**

Acknowledgments the authors to the Directorate General of Strengthening Research and Development at the Ministry of Research, Technology and Higher Education through the Directorate of Research and Community Service with a Research Contract Number: 062 / SP2H / LT / DRPM / 2018

**Illustration**



Picture 1. The appearance of plants with types of N fertilizer source and NPK addition



Picture 2. The difference of influence of N fertilizer source and NPK addition to Curd weights

# Effect of Slow Release Nitrogen (SRN) Addition to NPK Fertilization on Cauliflower Growth

## ORIGINALITY REPORT

9%

SIMILARITY INDEX

6%

INTERNET SOURCES

3%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

- |   |  |    |
|---|--|----|
| 1 | <a href="http://idus.us.es">idus.us.es</a><br>Internet Source  | 2% |
| 2 | <a href="http://mafiadoc.com">mafiadoc.com</a><br>Internet Source  | 1% |
| 3 | Submitted to Sriwijaya University<br>Student Paper   | 1% |
| 4 | Lilis Hermida, Joni Agustian. "Slow release urea fertilizer synthesized through recrystallization of urea incorporating natural bentonite using various binders", Environmental Technology & Innovation, 2019<br>Publication                                       | 1% |
| 5 | Peri Hardiansyah, Uswatun Nurjanah, Widodo Widodo. "Growth Response and Yield of Pakcoy (Brassica rapa L.) on Various Concentrations Liquid Organic Fertilizer of Jiringa Hulls [Phithecellobium jiringa (Jack) Prain Ex King]", Akta Agrosia, 2019<br>Publication | 1% |

6

[www.mspp.org.my](http://www.mspp.org.my)

Internet Source

&lt;1%

7

Ishak, Siti Amirah, and Irza Sukmana.  
"Modulation of Fibrin Gel Extracellular Matrix  
Properties by Fibrinogen and Thrombin  
Concentrations for Angiogenesis Assay",  
Advanced Materials Research, 2014.

Publication

&lt;1%

8

[seminar.fmipa.unp.ac.id](http://seminar.fmipa.unp.ac.id)

Internet Source

&lt;1%

9

Submitted to Higher Education Commission  
Pakistan

Student Paper

&lt;1%

10

Submitted to Universitas Brawijaya

Student Paper

&lt;1%

11

[www.gssrr.org](http://www.gssrr.org)

Internet Source

&lt;1%

12

[iopscience.iop.org](http://iopscience.iop.org)

Internet Source

&lt;1%

13

Fageria, . "Functions of Nitrogen in Crop  
Plants", Nitrogen Management in Crop  
Production, 2014.

Publication

&lt;1%

14

E Abdillah, H Supriyo, M Na'iem. "Soil Fertility  
Evaluation for Strategic Direction Jati Plus

&lt;1%

# Perhutani (JPP)", IOP Conference Series: Earth and Environmental Science, 2019

Publication

15

Heru Kuswantoro. "Increasing Grain Size Improves Grain Yield of Acid-Adaptive Soybean Lines in Optimal Soil Condition", Journal of Plant Sciences, 2015

Publication

<1%

16

Rudi Saputra Hutabarat, Edhi Turmudi, Nanik Setyowati. "Effect of Tithonia Compost (Tithonia diversifolia) and Phosphorus On The Growth and Yield of Peanuts", Akta Agrosia, 2019

Publication

<1%

17

[pearl.plymouth.ac.uk](http://pearl.plymouth.ac.uk)

Internet Source

<1%

18

[protan.studentjournal.ub.ac.id](http://protan.studentjournal.ub.ac.id)

Internet Source

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On