

**IN VITRO STUDY: INDUCED  
RESISTANCE OF CASSAVA (MANIHOT  
ESCULENTA CRANTZ.) PLANTLET  
AGAINST FUSARIUM OXYSPORUM  
BASED ON ANALYSIS OF PHENOL  
CONTENT**

*By* Sumardi Sumardi



## IN VITRO STUDY: INDUCED RESISTANCE OF CASSAVA (*MANIHOT ESCULENTA* CRANTZ.) PLANTLET AGAINST *FUSARIUM OXYSPORUM* BASED ON ANALYSIS OF PHENOL CONTENT

Dr. Endang Nurcahyani\*, Dr. Sumardi, Dr. Bambang Irawan, Evi Yunita Sari and Tika Lidia Sari

Dept. of Applied Biology, Faculty of Mathematics and Natural of Science, University of Lampung, Bandar Lampung, Indonesia.

\*Corresponding Author: Dr. Endang Nurcahyani

Dept. of Applied Biology, Faculty of Mathematics and Natural of Science, University of Lampung, Bandar Lampung, Indonesia.

Article Received on 20/12/2018

Article Revised on 11/01/2019

Article Accepted on 01/02/2019

### ABSTRACT

Cassava (*Manihot esculenta* Crantz.) is a tropical and subtropical annual shrub from the Euphorbiaceae tribe. Tuber is widely known as a staple food-producing carbohydrates and leaves as a vegetable. Indonesia is the second largest country producing Cassava after Nigeria with an average of five years' total supply of 9.67 million tons or 10.61% of the total supply of world cassava. Central Cassava land in Indonesia is controlled by Lampung Province with cassava production in Lampung Province reaching 8.33 million tons in 2013. This situation makes Lampung a supplier of one-third of the national Cassava production from national production of 23.92 million tons. However, there are still many production constraints in Cassava cultivation, including Fusarium wilt. This disease is caused by the *Fusarium oxysporum* fungus, which until now still cannot be treated effectively. The use of Cassava cultivars that are resistant to fusarium wilt with high yields is expected to be an important disease control alternative. This study aims to determine and determine the specific expression character of Fo-resistant Cassava plantlets, which are phenol content. The Cassava Plantlet has been initiated and selected in vitro in the Murashige and Skoog (MS) medium with the addition of fusaric acid (FA) at a concentration of 0 ppm (control), 20 ppm, 40 ppm, 60 ppm and 80 ppm. The purpose of this study is to determine the phenol content after being induced by fusaric acid and inoculated by *Fusarium oxysporum*. The results showed that in vitro, the concentration range of fusaric acid tolerant for selection of Cassava plantlets with optimum growth was between 20 ppm - 80 ppm; the higher the concentration of FA, the higher the phenol content of the Cassava plantlets.

**KEYWORDS:** Cassava plantlet; *Fusarium oxysporum*; Induced Resistance; in vitro; the phenol content.

### I. INTRODUCTION

Indonesia is the second largest country producing Cassava after Nigeria with an average of five years' total supply of 9.67 million tons or 10.61% of the total supply of world cassava, followed by Brazil, India and United Republic of Tanzania respectively ranged from 8.67 - 4.96 million tons or 9.52% - 5.44%, the rest contributed under 5.30%.<sup>[1]</sup>

One of the problems encountered in cultivating Cassava is Fusarium wilt caused by the fungus *F. oxysporum* (Fo) and is medium-transmitted.<sup>[2,3]</sup> reported that 50% of Cassava tubers produced and harvested in Nigeria were lost due to disease. The main causes of decay of Cassava include: *Aspergillus flavus*, *Aspergillus niger*, *Botryodiplodia theobromae*, *Collectotrichum* spp, *Geotrichum candidum*, *Penicillium chrysogenum*, *Penicillium digitatum*, *Fusarium oxysporum*.<sup>[4,5,6]</sup> This organism reduces the quantity and quality of the plant tubers.<sup>[7]</sup>

Disease control on Cassava has been carried out using pesticides which often cause pollution to the environment. Chemicals have been shown to help control Cassava disease, especially when the tuber has been attacked by pathogens.<sup>[8]</sup> The main problem with chemicals is that it is a big challenge for ecosystems and also the use of chemicals that often predispose them to resistance. One way to control diseases that are safe for the environment is to use varieties that are resistant to induced resistance.<sup>[9]</sup> The use of superior varieties that are resistant to Fo with high yield power is one of the important disease control alternatives and does not cause negative impacts such as the use of pesticides. The development of the Fo-resistant Cassava cultivar can be carried out, among others, by the in vitro selection method that is culturing explants in the form of tissue or organs in a medium containing selective concentration fusaric acid.<sup>[10]</sup>

Fusaric acid (FA) is a metabolite produced by several fungal species from the genus *Fusarium*. Chemically FA is called 5-n-butylpicolinic acid. This acid can be toxic so that it inhibits growth and culture regeneration,<sup>[11,10]</sup> but at non-toxic concentrations it helps to induce the synthesis of phytoalexin, a form of plant response to inhibit pathogenic activity.<sup>[10]</sup> Some parameters can illustrate the occurrence of plant resistance mechanisms against pathogenic infections such as an increase in phenol compounds, an increase in peroxidase enzymes (including PR-protein groups), and the presence of lignification.<sup>[12,13,14]</sup> Identification of insensitive mutants or variants of FA with in vitro selection was carried out on tomato plants,<sup>[15]</sup> bananas,<sup>[16,17]</sup> gladiolus,<sup>[18]</sup> pineapple,<sup>[19]</sup> *Sphatoglottis plicata*<sup>[20,21]</sup> as well as vanilla.<sup>[22,23,24,25]</sup> The results of the researchers' research showed that somaclonal regeneration of cell mass that is resistant to the toxin was also resistant to pathogens, and this trait was passed on to the next generation. The use of FA in a tolerant concentration so far has never been reported with certainty and precisely in the Induced Resistance Cassava plantlets against *Fo*. Therefore, research on the role of FA as an in vitro inducing resistance is interesting to do.

## II. MATERIAL AND METHODS

Materials research in the form of *Cassava* obtained from previous research. For resistance analysis and phenol content used 70% alcohol, Potato Dextrose Agar (PDA), gallic acid, Folin-Ciocalteu reagent, Na<sub>2</sub>CO<sub>3</sub>, sodium carbonate, 80% ethanol.

Analysis of total phenol compounds using the Singleton & Rossi method.<sup>[26]</sup> Preparation of standard calibration curve for gallic acid phenol is used as a standard solution. Preparation of sample plantlets prepared according to the method of.<sup>[27]</sup> Measurement of total phenol was carried out by taking a supernatant of 0.5 mL into a 100 mL volumetric flask, plus 250  $\mu$ L of Folin-Ciocalteu reagent, after leaving it for 5 minutes then 1 mL Na<sub>2</sub>CO<sub>3</sub> was added. After mixing evenly, it was put into cuvettes with a volume of 5 mL and observed absorption values at a wavelength of 765 nm using a spectrophotometer (*Beckman DU-65*), distilled water was used as a control. Based on the absorption value then determined the content of phenol compounds based on the gallic acid regression equation, namely the relationship between the absorption value of plantlet extract and the series of gallic acid concentrations.

Qualitative information as the result on this research was consisted as narratives descriptive and supported by photographs. After that, data's were statistically analyzed by Completely Randomized Design. As quantitatively data's from each parameter measured, were compiled and statistically analyzed by analysis of variance (ANOVAs). If the result showed a significantly different, then was continued analyzed by using Duncan Multiple Range Test (DMRT) analysis with accuracy 95%.<sup>[28]</sup>

## III. RESULTS AND DISCUSSION

Phenol compounds are the result of plant metabolism that is formed with one of the functions as a system of chemical resistance of plants that can prevent the growth and development of pathogens.<sup>[12]</sup> Before observing the analysis of total phenol compounds in Cassava plantlets for each treatment, a standard curve measurement was performed using gallic acid first. The measurement of this curve is done to estimate the quantity of total phenol with linear regression.<sup>[26]</sup> From the measurement of this standard curve produced a linear regression line equation ( $y = 0.002x - 0.005$ ) and has a high positive correlation value ( $R_2 = 0.996$ ). This shows that the diversity is homogeneous between the concentration of gallic acid and absorbance. Based on the standard gallic acid curve, the total phenol content of each treatment can be searched based on the regression line equation (Table 1).

**Table 1: Total phenol content (%) of control and fusaric acid (0, 20, 40, 60, and 80 ppm).**

Treatment (ppm)	Average total phenol content (%)
0 ppm	5,41 $\pm$ 0,03 <sup>a</sup>
20 ppm	6,35 $\pm$ 0,02 <sup>b</sup>
40 ppm	6,54 $\pm$ 0,03 <sup>b</sup>
60 ppm	6,88 $\pm$ 0,03 <sup>c</sup>
80 ppm	6,92 $\pm$ 0,02 <sup>c</sup>

**Description:** The numbers followed by unequal letters on one column are significantly different based on the Duncan test at 95% confidence degree, after being transformed to  $\sqrt{x}$ . Based on Table 1, it is clear that the increase in total phenol levels from about 5.41% in controls, increased to 6.35% at FA stress 20 ppm, followed by 6.54% at 40 ppm, 6.88 ppm at 60 ppm and 6,92% at 80 ppm. This proves that due to the higher concentration of FA stress, the total phenol content produced will also increase. The increase in the total phenol compound in the Cassava plantlets induced by FA, is another proof of the increase in plant resistance in resisting the rate of *Fo* infection. This is in accordance with the opinion of,<sup>[12]</sup> which states that one of the parameters for increasing plant resistance to pathogens is the increase in phenol compounds. According to,<sup>[10]</sup> FA on non-toxic concentrations ( $10^{-7}$ ) stimulates the formation of H<sub>2</sub>O<sub>2</sub> which is strongly associated with the peroxidase enzyme. Furthermore, this enzyme will oxidize phenol compounds.

The results of this study are similar to the results of a study conducted by,<sup>[29]</sup> patchouli plants infected by endophytic bacteria and nematodes showed that there was an indication of an increase in phenolic content by *Achromobacter xylosoxi*.<sup>[11]</sup> Total phenol levels were also reported by,<sup>[30]</sup> on Chickpea plants infected with *Fusarium oxysporum* f.sp. *ciceris*. From the study it was found that there was an increase in total phenol of around 16-17%.

#### IV. CONCLUSION

Based on the results and discussions that have been described in advance can be concluded as follows. The concentration range of fusaric acid tolerant for selection of cassava plantlets with optimum growth is between 20 ppm - 80 ppm. The higher the concentration of fusaric acid, the higher the total phenol level, the Cassava plantlet with *Fusarium oxysporum*.

#### ACKNOWLEDGEMENT

Thanks the authors to the Institute for Research and Community Service through the BLU fund of University of Lampung, based on the Letter of Assignment of "Penelitian PASCASARJANA" 2018 Number of Contract: 1344/UN26.21/PN/2018/25 June 2018.

#### REFERENCES

1. Anonymous. *Statistik Harga Komoditas Pertanian Tahun 2013*. Pusat Data dan Sistem Informasi Pertanian (Pusdatin). Jakarta, 2013.
2. Arinze, AE. Plant Pathology and Post-harvest Food Loss. *An Inaugural Lecture Series*, 2005; 43: 29-72.
3. Okigbo RN, Putheti R, Achusi CT, Post-harvest deterioration of cassava and its control using extracts of *Azadirachta indica* and *Afromonium meleguata*. *E-J Chem*, 2009; 6(4): 1274-1280.
4. Ogunleye AO, Ayansola OT. Studies of Some Isolated Rot-Causing Mycoflora of Yams (*Dioscorea Spp.*). *American Journal of Microbiology and Biotechnology*, 2014; 1(1): 9-20.
5. Okigbo NR, Enweremadu CE, Agu CK, Irondi RC, Okeke BC. Control of White Yam (*Dioscorea rotundata*) Rot Pathogen using Peel Extract of Water Yam (*Dioscorea alata*). *Advances in Applied Science Research*, 2015; 6(10): 7-13.
6. Gwa IV, Bem AA, Okoro JK. Yams (*Dioscorea rotundata* Poir and *D. Alata* Lam.) Fungi Etiology in Katsina-Ala Local Government Area of Benue State, Nigeria. *Journal of Phytopathology and Plant Health*, 2015; 3: 38-43.
7. Amusa NA, Adegbite AA, Muhammed S, Baiyewu RA. Yam diseases and its management in Nigeria. *African Journal of Biotechnology*, 2003; 2(12): 497-502.
8. Markson, A. A., Amadioha, A. C., Omosun, G., Madunagu, B. E., Udo, S. E., Umana, E. J. Control of *Botryodiplodia theobromae* causing Tissue Rot of White Yam (*Dioscorea rotundata* Poir). *Scholarly Journal of Agricultural Science*, 2012; 2(1): 1-7.
9. Agrios GN. *Plant Pathology*. Fifth Edition. Academic Press. New York, 2005; 922.
10. Bouizgarne, B, Bouteau HEM, Frankart C, Reboutier D, Madiona K, Pennarun AM, Monestiez M, Trouverie J, Amiar Z, Briand J, Brault M, Rona JP, Ouhdouch Y, & Hadrami EI., Early Physiological Responses of *Arabidopsis thaliana* Cells to Fusaric Acid : Toxic and Signalling Effects. *New Phytologist*, 2006; 169: 209 – 218.
11. Landa BB, Cachinero-Diaz JM, Lemanceu P, Jimenez-Diaz RM, & Alabouvette C, Effect of Fusaric Acid and Phytoanticipins on Growth of Rhizobacteria and *Fusarium oxysporum*. *Canadian Journal of Microbiology*, 2002; 48: 971-985.
12. Vidhyasekaran P. *Fungal Pathogenesis in Plants and Crops, Molecular Biology and Host Defense Mechanism*. Marcel Dekker. New York, 1997; 553.
13. Agrawal AA, Tuzun S, & Bent E. *Induced Plant Defenses Against Pathogens and Herbivores, Biochemistry, Ecology, and Agriculture*. APS Press, St. Paul, Minnesota, 1999; 390.
14. Lea P. & Leegood RC, *Plant Biochemistry and Molecular Biology*. 2<sup>nd</sup> ed. John Wiley & Sons Ltd. Chichester, 1999; 364.
15. Toyoda H, Hayashi H, & Yamamoto K. Selection of Resistance Tomato Calli to Fusaric Acid. *Ann. Phytopathol. Soc. Japan*, 1984; 50: 538-540.
16. Morpurgo R, Lopato SV, Afza R, & Novak FJ. Selection parameters for Resistance to *Fusarium oxysporum* f.sp. *cubense* race 1 and 4 on Diploid Banana (*Musa acuminata* Colla). *Euphytica*, 1994; 75: 121-129.
17. Matsumoto K, Barbosa ML, Souza LAC, & Teixeira JB. Race 1 Fusarium Wilt tolerance on banana plants selected by Fusaric Acid. *Euphytica*, 1995; 84: 67-71.
18. Remotti PC, Löffler HJM & Loten-Doting LV. Selection of Cell Lines and Regeneration of Plants Resistance to Fusaric Acid from Gladiolus x grandiflorus cv. 'Peter Pear'. *Euphytica*, 1997; 96: 237-245.
19. Boras O, Santos R, Matos A, Cabral P, & Arzola RS., A First Attempt to Use A Fusarium subglutinans Culture Filtrate For The Selection of Pineapple Cultivar to Fusariose Disease. *Plant Breeding*, 2001; 120(5): 345-438.
20. Nurcahyani E, R. Agustrina, T. T. Handayani. The Protein Profile of the Plantlets of *Spathoglottis plicata* Bl. Induced Resistance to *Fusarium oxysporum*. *Journal of Plant Sciences*, 2016a; 4(5): 102-105.
21. Nurcahyani E, Rochmah Agustrina, Erdi Suroso, Gardis Andari. Analysis of Peroxidase Activity and Total Phenol from *Spathoglottis plicata* Bl Plantlet Toward to *Fusarium oxysporum*. *International Journal of Applied Agricultural Sciences*, 2016b; 2(6): 79-82.
22. Nurcahyani E, Sumardi I, Hadisutrisno B, & Suharyanto E. Suppression of Development of Vanilla Foot Rot Disease (*Fusarium oxysporum* f. sp. *vanillae*) Through In Vitro Fusaric Acid Selection. *Journal of Tropical Plant Pests and Diseases*. Accreditation of DIKTI No.110/DIKTI/Kep/2009. ISSN: 1411-7525, 2012; 12(1): 12-22.
23. Nurcahyani, E., Hadisutrisno, B., Sumardi, I., & Suharyanto, E., Identifikasi galur planlet vanili (*Vanilla planifolia* Andrews) Resisten terhadap infeksi *Fusarium oxysporum* f. sp. *vanillae* hasil

- seleksi *in vitro* dengan asam fusarat. *Prosiding Seminar Nasional: "Pengendalian Penyakit Pada Tanaman Pertanian Ramah Lingkungan"*. Perhimpunan Fitopatologi Indonesia Komda Joglosemar-Fakultas Pertanian UGM. ISBN 978-602-71784-0-3/2014 Hal, 2014; 272-279.
24. Nurchayani E, Sumardi I, Hadisutrisno B and Suharyanto E. DNA Pattern Analysis of *Vanilla planifolia* Andrews Plantlet Which Resistant To *Fusarium oxysporum* f. sp. *vanillae*. *WJPLS*, 2017; (3)4: 27-34.
  25. Nurchayani E, Yulianty, and Suharyanto E, In Vivo Study: Characterization of Mutants *Vanilla planifolia* Andrews Resistant To *Fusarium* Wilt Disease Based On Analysis of the Lignin and the Phenol Content. *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)*, 2018; 11(3): 15-18. ISSN 2319-2380.
  26. Aberouman AA & Deokule SS. Comparison of Phenolic Compounds of Some Edible Plants of Iran and India. *Pakistan Journal of Nutrition*, 2008; 7: 582-585.
  27. Ozygi II, Kahraman MV & Ercan O. Relation Between Explant Age, Total Phenol and Regeneration Response in Tissue Cultured Cotton (*Gossypium hirsutum* L.). *African Journal of Biotechnology*, 2007; 6: 3-8.
  28. Gomez KA, Gomez AA, Statistical Procedures for Agricultural Research (2nd edn). John Wiley and sons, USA, 1984; 680.
  29. Harni R, Supramana, Sinaga MS, Giyanto, & Supriadi. Mekanisme Bakteri Endofit mengendalikan nematoda *Pratylenchus brachyurus* Pada Tanaman Nilam. *Bul. Littro*, 2012; 23(1): 102-114.
  30. Khan IA, Alam SS, Haq A, & Jabbar A. Biochemistry of Resistance in *Chicpea* Against Wilt Disease Caused by *Fusarium oxysporum* f. sp. *ciceris*. *Pak. J. Bot.*, 2005; 37: 97-104.

# IN VITRO STUDY: INDUCED RESISTANCE OF CASSAVA (MANIHOT ESCULENTA CRANTZ.) PLANTLET AGAINST FUSARIUM OXYSPORUM BASED ON ANALYSIS OF PHENOL CONTENT

ORIGINALITY REPORT

# 19%

SIMILARITY INDEX

## PRIMARY SOURCES

1	<a href="http://www.scribd.com">www.scribd.com</a> Internet	103 words — 6%
2	<a href="http://ementoring.suttontrust.com">ementoring.suttontrust.com</a> Internet	44 words — 2%
3	<a href="http://sciencepublishinggroup.com">sciencepublishinggroup.com</a> Internet	34 words — 2%
4	<a href="http://article.jplantsciences.org">article.jplantsciences.org</a> Internet	25 words — 1%
5	<a href="http://vegetable-salad.blogspot.com">vegetable-salad.blogspot.com</a> Internet	23 words — 1%
6	<a href="http://ejournal.unmus.ac.id">ejournal.unmus.ac.id</a> Internet	22 words — 1%
7	<a href="http://eprints.umm.ac.id">eprints.umm.ac.id</a> Internet	12 words — 1%
8	<a href="http://www.imedpub.com">www.imedpub.com</a> Internet	11 words — 1%
9	Moustafa Bani, Nicolas Rispaill, Antonio Evidente, Diego Rubiales, Alessio Cimmino. " Identification of the Main Toxins Isolated from f. sp. Race 2 and Their Relation with Isolates' Pathogenicity ", Journal of Agricultural and Food	10 words — 1%

# Chemistry, 2014

Crossref

---

10	<a href="http://jurnal.polinela.ac.id">jurnal.polinela.ac.id</a> Internet	10 words — 1%
11	<a href="http://fupress.net">fupress.net</a> Internet	9 words — 1%
12	<a href="http://journals.plos.org">journals.plos.org</a> Internet	9 words — 1%
13	<a href="http://juniperpublishers.com">juniperpublishers.com</a> Internet	9 words — 1%
14	<a href="http://article.sciencepublishinggroup.com">article.sciencepublishinggroup.com</a> Internet	8 words — < 1%
15	<a href="http://agrosym.ues.rs.ba">agrosym.ues.rs.ba</a> Internet	8 words — < 1%

---

EXCLUDE QUOTES      ON

EXCLUDE  
BIBLIOGRAPHY      ON

EXCLUDE MATCHES      OFF