

Effectiveness of the combination of organonitrofos and inorganic fertilizers on soil chemical properties and the yields of cucumber (*Cucumis sativus* L.) in Ultisols

By Dermiyati; E Aprilia; K Hendarto; S B Yuwono and Y C Ginting

Effectiveness of the combination of organonitrofos and inorganic fertilizers on soil chemical properties and the yields of cucumber (*Cucumis sativus* L.) in Ultisols

Dermiyati¹, E Aprilia¹, K Hendarto¹, S B Yuwono² and Y C Ginting¹

¹ Department of Agrotechnology, Faculty of Agriculture, University of Lampung, Bandar Lampung, Indonesia

² Department of Forestry, Faculty of Agriculture, University of Lampung, Bandar Lampung, Indonesia

E-mail: dermiyati.1963@fp.unila.ac.id

Abstract. Ultisols has constraints in low organic matter, low pH, low CEC, and low soil nutrients. This study aimed to determine the effect of applying a combination of Organonitrofos and inorganic fertilizers on soil chemical properties and yields of cucumber plants in Ultisols. The study consisted of 11 combinations of Organonitrofos and inorganic fertilizers with 3 replications arranged in a randomized block design. A recommended dose of 100% Organonitrofos fertilizer which was 10,000 kg ha⁻¹ and 100% NPK inorganic fertilizer (Urea = 448 kg ha⁻¹; SP-36 = 413.5 kg ha⁻¹; KCl = 63.3 kg ha⁻¹) were applied. The treatments were various combinations of Organonitrofos and inorganic fertilizers of 0, 25, 50, 75 and 100% from each fertilizer recommendation. The application of a combined Organonitrofos and inorganic fertilizers increased pH and soil available-P, but it did not affect the total-N and organic-C of soil. All combined doses of Organonitrofos and inorganic fertilizer significantly increased all the growth and yields variables of cucumber compared to control (without fertilizer) and application of Organonitrofos fertilizer only. Application of 100% Organonitrofos + 50% NPK was the most economically effective dose because it had the highest RAE value of 101%.

1. Introduction

Cucumber is a fruit vegetable that is consumed by many Indonesians in the form of fresh or processed because it has a fairly good nutritional value. Although the need for cucumbers is continuously increase, but the production of cucumber plants still does not meet the needs of consumers. The production of cucumber has decreased from year to year. The average productivity of cucumber in 2012 and 2013 reached 9.97 t ha⁻¹, while in 2014 it decreased to 9.84 t ha⁻¹ [1]. One reason of the decreased cucumber production might be due to the decreasing harvested area because of land conversion. Cucumber plants are usually cultivated on low-fertility Ultisols. The problems of Ultisols are high soil acidity (average pH <4.50), high Al-saturation, poor macronutrient content especially N, P, K, Ca, and Mg, and low organic matter content [2]. Nutrient contents in Ultisols are generally low due to intensive leaching of alkaline bases, while the low organic matter content is because the decomposition process runs quickly and some of the organic matters are lost through erosion.

One effort to increase the productivity of Ultisols can be done through fertilization. Inorganic fertilizers are easily available and can be directly absorbed by plants, but they can cause physical and biological damage to the soil if they apply in a long term. Therefore, a combination of inorganic fertilizer



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

and organic fertilizer is needed so that soil damage caused by land degradation can be prevented. The combination of NPK fertilizer 30% of the recommended dose and 2 t ha⁻¹ of organic fertilizer increased the yields and NPK uptake of curly chilies [3].

One of the organic fertilizers that can be used is Organonitrofos. It is a solid organic fertilizer that can provide high level of N and P nutrients. It is made from fresh manure, MSG waste, coconut fiber, and enriched with beneficial microbes (N-fixer and P-solubilizer) [4]. To improve the quality of Organonitrofos, reformulation had to be done [5]. Continuous researches regarding Organonitrofos combined with inorganic fertilizer are still done to see its possible role as a substitute for inorganic fertilizers, especially NPK [3, 6, 7]. A partial substitution of recommended chemical fertilizers with Organonitrofos fertilizer by 40% consistently resulted in the highest plant growth performance (stover weight) and maize yields (dry shelled weight and one hundred grain weight) followed by 100% organonitrofos fertilizers only (without chemical fertilizers) [8].

This research was conducted to determine the effect of the combined Organonitrofos and inorganic fertilizers on the changes in the chemical properties of Ultisols and the yields of cucumber plants.

2. Materials and methods

A pot experiment was conducted at the Integrated Field Laboratory and the soil analysis at the Soil Science Laboratory of the Faculty of Agriculture, University of Lampung, Bandar Lampung, Indonesia from July to October 2016. The research consisted of eleven treatments with three replication arranged in a randomized block design (RBD). The treatments were a combination of Organonitrofos (OP) and inorganic fertilizer (NPK) (table 1). The recommended doses of 100% organonitrofos fertilizer (OP) was 10 t ha⁻¹, while 100% inorganic fertilizer (NPK) was Urea= 448 kg ha⁻¹; SP-36= 413.5 kg ha⁻¹; KCl= 63.3 kg ha⁻¹. Organonitrofos is a solid organic fertilizer with the composition of pH (H₂O) 7.0, organic-C 13.89%, total-N 1.08%, total-P 4.96% and total-K 3.03% [9].

Table 1. The combination of Organonitrofos (OP) and inorganic fertilizer (NPK) used in the study.

| Treatment | Fertilizer Combination |
|-----------|--------------------------------|
| P0 | 0% Organonitrofos + 0% NPK |
| P1 | 0% Organonitrofos + 100% NPK |
| P2 | 100% Organonitrofos + 0% NPK |
| P3 | 100% Organonitrofos + 25% NPK |
| P4 | 100% Organonitrofos + 50% NPK |
| P5 | 100% Organonitrofos + 75% NPK |
| P6 | 100% Organonitrofos + 100% NPK |
| P7 | 75% Organonitrofos + 75% NPK |
| P8 | 50% Organonitrofos + 75% NPK |
| P9 | 75% Organonitrofos + 25% NPK |
| P10 | 50% Organonitrofos + 50% NPK |

Data were analyzed by ANOVA (Analysis of Variance). Before testing of ANOVA, homogeneity of variance was tested by the Bartlett test, data additivity was tested by the Tukey test. The difference in the mean value of the treatment was tested by *Duncan's Multiple Range Test* (DMRT) at the 5% level.

The research was started by preparing a polybag with a size of 35 x 40 cm, filled with the planting media consisted of 10 kg Ultisols (oven-dry weight). The soils then mixed with husk charcoal (25 g polybag⁻¹), dolomite (10 g polybag⁻¹) and Organonitrofos (according to each treatment dose). The media then allowed to stand for two weeks before planting. Two cucumber seeds were planted in each polybag. After a week, one of the best emerged plant was selected to be grown. Application of inorganic fertilizers (NPK) was given one week after planting (WAP) by being circled around the seeds. Urea fertilizer was applied twice, namely a half dose together with SP-36 and KCl fertilizers and a half dose when the cucumber plants began flowering. Plant maintenance were carried out by irrigation, replanting, plant

stakes, and controlling pests and diseases. Cucumbers were harvested from 28 to 35 days after planting (DAP). Soil samplings were carried out twice, namely in the time before planting and after harvesting in a composite manner at each replication.

12

3. Results and discussion

3.1. Chemical properties of ultisols before planting and after harvesting

Initial soil analysis shows that Ultisols used in this study had characteristic such as acidic pH and very low nutrient contents of total-N, available-P, organic-C, and cation exchange capacity (CEC) (table 2).

Table 2. Initial soil chemical properties of Ultisols before planting time.

| Soil chemical properties | Value (Criteria) |
|------------------------------------|------------------|
| pH | 4.61 (A) |
| Total-N (%) | 0.08 (VL) |
| Available-P (mg kg ⁻¹) | 3.25 (VL) |
| Organic-C (%) | 0.95 (VL) |
| CEC (cmol kg ⁻¹) | 3.43 (VL) |

Note: The number followed by the letter states A=Acid and VL=Very low.

The initial soil nutrient contents were very low so that the combination of Organonitrofos and inorganic fertilizers was applied to increase the fertility of Ultisols soil. The results of the soil analysis after harvesting of the cucumber is presented in table 3.

There was an increase of soil pH values after harvesting which ranged from 5.72 to 6.60 followed by a change in criteria from being slightly acid to neutral (table 3) compared to the initial soil pH of 4.61 with acid criteria (table 2). Soil pH values affect the nutrient availability in the soil. Cucumber plants can be grown in the soil pH ranged from 5.5 to 7.5. At soil pH less than 5.5, there will be interference with the absorption of nutrients by plant roots so that plant growth is disrupted [10]. The use of dolomite as a basalt treatment in this research was sufficient to increase soil pH and to meet good conditions for growing cucumber plants.

The soil total-N content after harvesting was not significantly increased compared to those before planting for all treatments. These could be assumed that cucumber plants have been using soil N for the growth and due to the occurrence of N evaporation. It was found that the soil total-N content decreased from 0.5% (at an initial stage, before planting) to 0.4% (at harvesting time) by the application of chicken manure enriched with husk ash and Tithonia [11]. Moreover, the decreased of nitrogen levels occurred in almost all treatments except on the treatment of goat manure enriched with natural phosphate, dolomite, and Tithonia extract. Nitrogen uptake by plants used a reduction of nitrogen in the soil.

The soil available-P after harvesting ranged from 11.9 mg kg⁻¹ to 56.7 mg kg⁻¹ (high to very high criteria). It was highly increasing compared to the initial value (before planting) of 3.3 mg kg⁻¹ (very low criteria). This is presumably due to the application of Organonitrofos containing phosphate solubilizing microbes and SP-36 inorganic fertilizer which can contribute to P-nutrient content in Ultisols. The application of SP-36 fertilizer and organic fertilizer as well as the interaction of both treatments have significant effects on the increasing of soil available-P [12].

Furthermore, the increase of soil available-P content might be also due to the addition of soil amendments in the form of dolomite and husk charcoal applied to all treatments including control treatment (P0: 0% Organonitrofos + 0% NPK). The soil available-P increased significantly up to 31.9% by the application of dolomite 1 t ha⁻¹ and increased around 9.6% by the application of biochar 4 t ha⁻¹ [13].

Dolomite contains magnesium (Mg) and calcium (Ca) as soil calcifiers. Applying dolomite in acid soils has a good effect on soil properties. If Mg levels in the soil increased, then the level of N and P in the leaves will also increased. If the soil hydrogen contents decrease, then the soil pH increases [10].

The responses of increasing soil pH because of the supply of Mg and Ca nutrients which can shift the position of H^+ on the surface of the colloid to the soil solution to neutralize the acidity of the soil. Liming could reduce aluminum toxicity, increase the availability of soil P as a result of releasing P from Al-P and Fe-P bonds, increase N fixation, N mineralization, and CEC [10].

Table 3. The soil chemical properties of Ultisols after harvesting of cucumber plant.

| Treatment | Fertilizer Combination (%) | | pH | Total-N (%) | Available-P (mg kg ⁻¹) | Organic-C (%) |
|-----------|-------------------------------|-----|-----------|----------------|---------------------------------------|------------------|
| | OP | NPK | | | | |
| P0 | 0 | 0 | 6.33 (SA) | 0.06 (VL) | 11.9 (H) | 0.96 (VL) |
| P1 | 0 | 100 | 5.72 (SA) | 0.08 (VL) | 48.2 (VH) | 1.13 (L) |
| P2 | 100 | 0 | 6.60 (N) | 0.12 (L) | 50.3 (VH) | 1.34 (L) |
| P3 | 100 | 25 | 6.36 (SA) | 0.12 (L) | 51.0 (VH) | 1.35 (L) |
| P4 | 100 | 50 | 6.21 (SA) | 0.13 (L) | 53.9 (VH) | 1.52 (L) |
| P5 | 100 | 75 | 5.98 (SA) | 0.11 (L) | 55.6 (VH) | 1.25 (L) |
| P6 | 100 | 100 | 5.85 (SA) | 0.11 (L) | 56.7 (VH) | 1.25 (L) |
| P7 | 25 | 75 | 5.91 (SA) | 0.08 (VL) | 50.3 (VH) | 1.08 (L) |
| P8 | 50 | 75 | 5.93 (SA) | 0.11 (L) | 52.9 (VH) | 1.21 (L) |
| P9 | 75 | 75 | 6.06 (SA) | 0.11 (L) | 55.0 (VH) | 1.40 (L) |
| P10 | 50 | 50 | 6.15 (SA) | 0.11 (L) | 51.0 (VH) | 1.20 (L) |

Note: OP = Organonitrofos. The number followed by the letter states SA = Slightly Acid, N = Neutral, VH = Very High, H = High, L = Low, VL = Very Low.

The soil organic-C content after harvesting did not increase in the control treatment, while slightly increase in the other treatments. The provision of organic matter can increase soil organic-C content that can affect soil properties to be better physically, chemically, and biologically. Carbon is an energy source of soil microorganisms, so the presence of soil organic-C will spur microorganism activities thereby increasing the soil decomposition process and also soil reactions that involve microorganisms, such as P-dissolution, and N-fixation [14].

It was reported that soil chemical properties such as pH, organic-C, total-N, and available-P after harvesting were significantly ($P < 0.01$) affected by the application of the different rates of organic and inorganic fertilizers [15]. Moreover, relatively higher pH value, organic-C, and nutrient concentrations for plots were treated with manure and compost [15]. Similarly, it was also reported that the application of animal manure and compost increased soil pH [16].

3.2. Effect of application organonitrofos fertilizer and inorganic fertilizer on the growth and yields of cucumber plants

The application of combined Organonitrofos and inorganic fertilizer had no significant effect on the length of cucumber plants, which is consistent with the previous research by Wijaya *et al.* [17]. It is assumed that the macro and micronutrient content in the soil is sufficient enough for the growth of cucumber plants. Based on the analysis of soil chemical properties after harvesting, there was an increase in soil pH and nutrient content compared to before planting, including the control treatment (P0: 0% Organonitrofos + 0% NPK) because all treatments had been given by dolomite and husk charcoal as basalt soil amendments.

However, based on the results of the DMRT 5% test (table 4), the P4 treatment (100% Organonitrofos + 50% NPK) had the highest plant length and significantly different from the control treatment. The application of the combined Organonitrofos and inorganic fertilizer has a significant effect on the vegetative growth of cucumber plants from the parameters of the number of leaves, stem

diameter, and number of branches of cucumber plants. The P9 treatment (75% Organonitrofos + 75% NPK) had the highest number of leaves, stem diameter, and number of branches, but relatively no different from other fertilizer combination treatments and also not different from the treatment of inorganic fertilizers only (P1). The availability of complete and balanced nutrients that can be absorbed by plants is a factor that determines plant growth and yields [18].

Table 4. Effect of a combined Organonitrofos and inorganic fertilizer on the vegetative growth of cucumber plants.

| Treatment | Plant Length (cm)* | Number of Leaves (strands)* | Rod Diameter (cm)* | Number of Branches* |
|-----------|--------------------|-----------------------------|--------------------|---------------------|
| P0 | 161.0 b | 28.0 d | 0.7 c | 3.0 c |
| P1 | 202.3 ab | 78.7 ab | 1.2 ab | 11.3 a |
| P2 | 195.8 ab | 46.7 c | 1.1 b | 6.0 bc |
| P3 | 196.3 ab | 64.7 b | 1.1 ab | 8.7 ab |
| P4 | 215.0 a | 75.0 b | 1.3 ab | 9.7 a |
| P5 | 197.6 ab | 70.0 b | 1.1 ab | 9.0 ab |
| P6 | 197.9 ab | 69.7 ab | 1.1 ab | 9.3 ab |
| P7 | 200.9 ab | 72.5 ab | 1.2 ab | 9.7 a |
| P8 | 200.5 ab | 67.5 b | 1.1 ab | 10.7 a |
| P9 | 201.9 ab | 80.0 ab | 1.3 a | 12.0 a |
| P10 | 198.2 ab | 88.0 a | 1.1 ab | 9.0 ab |

*The numbers followed by the same letter are not significantly different from the DMRT test at the 5% level.

The P9 treatment (75% Organonitrofos + 75% NPK) had the highest fresh stover weight and dry crop stover weight, but it was not relatively different from the P1, P4, P6, P7, P8, and P10 treatments (table 5). Fresh stover weight is the result of the measurement of the fresh weight of plant biomass as an accumulation of material produced during growth. Giving Organonitrofos fertilizer and inorganic fertilizer with various combinations can provide enough nutrients for the growth of cucumber plants because all macro and micronutrients are provided by both kinds of fertilizers.

From Table 6, it can be seen the shortest flowering age was in the treatments of P3 (100% Organonitrofos + 25% NPK), P4 (100% Organonitrofos + 50% NPK), and P6 (100% Organonitrofos + 100% NPK), which was 25.7 DAP. While the longest flowering age was in the control treatment (P0) and NPK only treatment (P1). Organonitrofos fertilizers can provide high levels of N and P nutrients because it was enriched with beneficial microbes (N-fixer, P-solubilizer, and *Trichoderma* sp.), so it can influence the flower formation process. It was found that the application of nitrogen in the form of organic and inorganic fertilizer has a significant effect on the flowering age of cucumber plants [19]. The availability of nitrogen in a high amount in the soil affects the absorption of phosphorus which plays a role in the process of flower formation. Besides, flowers on cucumber plants appear on the armpits of leaves, so the number of flowers that appear on cucumber plants is influenced by the number of leaves and segments of the cucumber plant itself [20].

Moreover, the P9 treatment (75% Organonitrofos + 75% NPK) can produce the highest number of female and male flowers. In addition to the availability of nutrients, the number of flowers is also influenced by environmental factors, especially sunlight and sunlight intensity. It has been stated that if the length of the sun's irradiation is less than 12 hours every day with low light intensity so that more cucumber plants form female flowers [21]. On the contrary, the high intensity of sunlight makes more cucumber plants from male flowers.

The combination of Organonitrofos and inorganic fertilizers did not have a significant effect on the cucumber harvest age. This is supported by Zulyana [22] where the application of chemical fertilizers and combined by dosages and forms of cow dung did not have a significant effect on the cucumber harvest age. The faster harvest time means that the treatment has a better impact because the fast harvest age shows that the ripening of the fruit is going well, as a result of better nutrient absorptions.

Table 5. Effect of a combined Organonitrofos and inorganic fertilizer on fresh and dry stover weights.

| Treatment | Fresh Stover Weight (g plant ⁻¹)* | Dry Stover Weight (g plant ⁻¹)* |
|-------------------------------------|--|--|
| P0 (0% Organonitrofos + 0% NPK) | 243 d | 29.2 c |
| P1 (0% Organonitrofos + 100% NPK) | 800 ab | 79.7 ab |
| P2 (100% Organonitrofos + 0% NPK) | 297 cd | 37.9 bc |
| P3 (100% Organonitrofos + 25% NPK) | 333 cd | 42.9 bc |
| P4 (100% Organonitrofos + 50% NPK) | 623 abc | 84.1 ab |
| P5 (100% Organonitrofos + 75% NPK) | 497 bcd | 57.8 bc |
| P6 (100% Organonitrofos + 100% NPK) | 513 bcd | 71.0 ab |
| P7 (25% Organonitrofos + 75% NPK) | 747 abc | 87.3 ab |
| P8 (50% Organonitrofos + 75% NPK) | 637 abc | 72.7 ab |
| P9 (75% Organonitrofos + 75% NPK) | 1393 a | 144.9 a |
| P9 (50% Organonitrofos + 50% NPK) | 577 abc | 67.5 ab |

*The numbers followed by the same letter are not significantly different from the DMRT test at the 5% level.

Table 6. Effect of a combination of Organonitrofos fertilizer and inorganic fertilizer on the generative growth of cucumber plants.

| Treatment | Flowering Age (days)* | Amount of Female Flowers* | Number of male flowers* | Early Harvest Age (days)* |
|-----------|--------------------------|------------------------------|----------------------------|------------------------------|
| P0 | 27.7 a | 8.7 c | 25.3 c | 41.0 a |
| P1 | 27.0 ab | 31.0 ab | 40.7 ab | 39.0 ab |
| P2 | 26.3 bc | 21.0 bc | 31.7 cb | 39.0 ab |
| P3 | 25.7 c | 25.7 ab | 43.0 ab | 36.3 b |
| P4 | 25.7 c | 29.7 ab | 46.7 a | 37.0 b |
| P5 | 26.0 bc | 22.7 ab | 45.0 ab | 36.3 b |
| P6 | 25.7 c | 24.0 ab | 41.7 ab | 37.0 b |
| P7 | 26.3 bc | 26.0 ab | 48.3 a | 38.7 ab |
| P8 | 26.0 bc | 29.7 ab | 45.3 ab | 37.7 ab |
| P9 | 26.0 bc | 37.0 a | 57.3 a | 38.0 ab |
| P10 | 26.0 bc | 28.7 ab | 46.0 ab | 36.3 b |

* The numbers followed by the same letter are not significantly different from the DMRT test at the 5% level.

The treatment of inorganic fertilizers without Organonitrofos (P1) gave the highest yields based on the number of fruits per plant (table 7). A small number of female flowers will give a little amount of fruit too, but a large number of female flowers may not necessarily produce a lot of fruit, all depends on the process or obstacles that occur during pollination and the environmental factors during the formation and ripening of fruit [22]. During the development of flowers into fruit, many factors prevent the formation of flowers into fruit, including pests and diseases, flower loss, and pollination.

The combination of Organonitrofos and inorganic fertilizer had a significant effect on the length of the cucumber but did not significantly affect the diameter of the cucumber. The treatment of inorganic

fertilizers without Organonitrofos (P1) fertilizer has the highest length and diameter of cucumber but is relatively not significantly different from the treatment of Organonitrofos fertilizer and its combination with inorganic fertilizer.

The combined treatment of organonitrofos and inorganic fertilizers has a significant effect on the yields of cucumber. The P4 treatment (100% Organonitrofos + 50% NPK) was able to provide the highest cucumber production of 5,732 g plant⁻¹ which was relatively different from the other combination treatments and the treatment of inorganic fertilizers only (P1). The high yields achieved with the best-combined dose namely P4 treatment (100% Organonitrofos + 50% NPK) that have given optimal results based on the description of cucumber varieties of Mercy F1, where these varieties have an average production yield of 3,500 to 5,000 g plant⁻¹.

Table 7. Effect of a combined Organonitrofos and inorganic fertilizer on cucumber yields.

| Treatment | Number of Fruits per Plant | Fruit Length (cm) | Fruit Diameter (cm) | Average Fruit Weight (g) | Fruit Weight per Plant (g plant ⁻¹) |
|-----------|-------------------------------|----------------------|---------------------------|-----------------------------|---|
| P0 | 7.3 c | 20.3 c | 4.4 b | 198.1 d | 14689 c |
| P1 | 23.7 a | 25.1 a | 4.9 a | 287.1 a | 5682 a |
| P2 | 13.0 bc | 22.0 bc | 4.6 ab | 233.1 cd | 3335 bc |
| P3 | 14.3 bc | 23.0 ab | 4.6 ab | 236.8 bcd | 3287 bc |
| P4 | 23.0 a | 24.8 a | 4.8 a | 277.1 ab | 5732 a |
| P5 | 16.0 b | 23.5 ab | 4.7 a | 254.5 abc | 3923 ab |
| P6 | 17.0 ab | 23.6 ab | 4.6 ab | 276.8 ab | 4556 ab |
| P7 | 21.3 ab | 24.9 a | 4.8 a | 280.5 a | 5214 ab |
| P8 | 20.0 ab | 25.1 a | 4.6 ab | 267.0 abc | 4925 ab |
| P9 | 21.0 ab | 25.1 a | 4.7 a | 278.5 ab | 5468 a |
| P10 | 19.0 ab | 25.2 a | 4.8 a | 276.5 ab | 4859 ab |

* The numbers followed by the same letter are not significantly different from the DMRT test at the 5% level.

Table 8. The calculation of Relative Agronomic Effectiveness (RAE) on total biomass.

| Treatment | Super Weight (t ha ⁻¹) | Cucumber Weight (t ha ⁻¹) | RAE of Biomass Total (%) |
|-------------------------------------|--|--|--------------------------------|
| P0 (0% Organonitrofos + 0% NPK) | 0.50 | 25.0 | 0 |
| P1 (0% Organonitrofos + 100% NPK) | 1.35 | 96.6 | 100 |
| P2 (100% Organonitrofos + 0% NPK) | 0.64 | 56.7 | 44 |
| P3 (100% Organonitrofos + 25% NPK) | 0.73 | 55.9 | 43 |
| P4 (100% Organonitrofos + 50% NPK) | 1.43 | 97.4 | 101 |
| P5 (100% Organonitrofos + 75% NPK) | 0.98 | 66.7 | 58 |
| P6 (100% Organonitrofos + 100% NPK) | 1.21 | 77.5 | 73 |
| P7 (25% Organonitrofos + 75% NPK) | 1.48 | 88.6 | 89 |
| P8 (50% Organonitrofos + 75% NPK) | 1.24 | 83.7 | 82 |
| P9 (75% Organonitrofos + 75% NPK) | 2.46 | 93.0 | 97 |
| P10 (50% Organonitrofos + 50% NPK) | 1.15 | 82.6 | 80 |

Note: If the RAE value is $\geq 100\%$, the fertilizer tested is effective compared to the recommended standard fertilizer treatment.

Based on RAE calculations, an agronomically effective fertilizer treatment compared to standard fertilizer treatment, which has an RAE value greater than 100% can be seen in the P4 treatment (100% Organonitrofos + 50% NPK) (table 8). Score of *Relative Agronomic Effectiveness* (RAE) was obtained from the comparison between the increase in yield due to the use of testing fertilizer and the increase in yield on standard fertilizer multiplied by 100%. The P0 treatment (0% Organonitrofos + 0% NPK) as a control and P1 (0% Organonitrofos + 100% NPK) as a standard fertilizer treatment. Based on Table 8, the P4 treatment (100% Organonitrofos + 50% NPK) is the most effective combination because it has the highest RAE value of 101%. The RAE value is relatively not different from the standard P1 fertilizer treatment (0% Organonitrofos + 100% NPK) due to the total biomass in the form of dry stover weight and the total weight of cucumber produced in P1 treatment (0% Organonitrofos + 100% NPK). It is also relatively not different from the treatment of P4 (100% Organonitrofos + 50% NPK). A single dose of 5 t ha⁻¹ Organonitrofos fertilizer is recommended for cucumber farmers because it is the most effective dose agronomically (Relative Agronomic Effectiveness), economically and can create sustainable agriculture [17].

Table 9. Correlation between some soil chemical properties after harvesting and an average weight of cucumber, cucumber yields, and dry stover weight.

| Variable | Equation | r |
|---|----------------------|---------|
| Soil pH with Average Fruit Weight | $y = 0.075x + 0.720$ | 0.71 ** |
| Soil pH with Fruit Yields | $y = 52.59x + 395.6$ | 0.62 * |
| Soil pH with Dry Stover Weights | $y = 1.017x + 7.402$ | 0.50 ns |
| Soil Total-N with Fruit Weight | $y = 0.347x + 0.224$ | 0.27 ns |
| Soil Total-N with Fruit Yields | $y = 339.0x + 39.73$ | 0.32 ns |
| Soil Total-N with Dry Stover Weight | $y = 3.625x + 0.822$ | 0.14 ns |
| Soil Available- P with Average Fruit Weight | $y = 0.001x + 0.180$ | 0.75 ** |
| Soil Available- P with Fruit Yields | $y = 1.291x + 11.97$ | 0.73 ** |
| Soil Available- P with Dry Stover Weights | $y = 0.020x + 0.199$ | 0.48 ns |
| Soil Organic-C with Fruit Average Weight | $y = 0.053x + 0.194$ | 0.30 ns |
| Soil Organic-C with Fruit Yields | $y = 57.95x + 2.699$ | 0.41 ns |
| Soil Organic-C with dry stover weights | $y = 1.197x - 0.293$ | 0.35 ns |

* = significantly different at the 5% level.

** = significantly different at the level of 5% and 1%.

ns = not significantly different at the 5% level.

Based on the results of the correlation analysis (table 9), there is a correlation between soil pH with the average fruit weight and fruit yields. Likewise, there is also a correlation between soil available-P with the average fruit weight and fruit yields. The pH value affects the availability of nutrients in the soil. If the soil pH is around 5.5 (the minimum limit) and 7.5 (the maximum limit), it is still tolerant for cucumber plants to have good growth and produce optimal yields [10]. The content of soil available-P after harvesting has increased from the initial value before planting, this causes the number of fruit produced and cucumber fruit yields also increased. The P element is useful for plants to stimulate growth, especially the roots so that plants can develop properly because they can get optimal nutrients. Also, the P content has functions to accelerate flowering and fruit formation [23], so that a high P content in soil can increase fruit yield of cucumber plants.

Based on the present study it can be stated that Organonitrofos fertilizer at a dose of 25% to 100% and its combination with inorganic fertilizer from 50% to 100% are the best range combination dose for the growth and yields of the cucumber plants that can reach yields as many as 3,923 g plant⁻¹ to 5,732 g

plant⁻¹. The most effective agronomically dose is found in the treatment of P4 (100% Organonitrofos + 50% NPK) because it has an RAE value greater than 100% that is equal to 101% (table 8). It showed that the recommended dose has increased yield compare to the standard fertilizers. Besides, the highest marginal rate of return for tea production on Ultisols was obtained from the application of 50% FYM (farmyard manure) + 50% recommended nitrogen and phosphorus fertilizer [24].

4. **C**onclusions

The combination of organonitrofos and inorganic fertilizers increased soil pH and soil available-P of Ultisols. The improvement of soil pH and soil available-P were also affected by the application of dolomite as a basalt treatment. Moreover, there was a significant correlation between soil pH and soil available-P with average cucumber fruit weights and fruit yields. Furthermore, the combined dose of organonitrofos and inorganic fertilizer significantly increased all growth and yield variables of cucumber plants compared to control (without fertilizer) and Organonitrofos only. The treatment of 100% Organonitrofos and 50% NPK fertilizer were the most economically effective due to the highest RAE value of 101%. The application of Organonitrofos fertilizer could reduce the use of NPK fertilizer by 50%.

Effectiveness of the combination of organonitrofos and inorganic fertilizers on soil chemical properties and the yields of cucumber (*Cucumis sativus* L.) in Ultisols

ORIGINALITY REPORT

13%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

Ryandi Eka Putra, Dermiyati Dermiyati, Nur Afni Afianti, Henrie Buchari. "PENGARUH KOMBINASI PUPUK ORGANONITROFOS DAN PUPUK ANORGANIK TERHADAP SIFAT KIMIA TANAH DAN PRODUKSI TANAMAN JAGUNG (*Zea mays* L.) DI TANAH ULTISOLNATAR PADA MUSIM TANAM KE DUA", Jurnal Agrotek Tropika, 2020
Crossref

55 words — 1%
- 2

gfzpublic.gfz-potsdam.de
Internet

49 words — 1%
- 3

garuda.ristekbrin.go.id
Internet

44 words — 1%
- 4

E Maftu'ah, Y Lestari, E B Pangaribuan, V Mayasari. "Amelioration of actual acid sulfate soils to improve soil chemical properties and rice yields", IOP Conference Series: Earth and Environmental Science, 2021
Crossref

41 words — 1%
- 5

orgprints.org
Internet

41 words — 1%
- 6

A Krismawati, Z Arifin, C Hermanto, C Tafakresnanto. "Application of biochar to improve maize

36 words — 1%

performance on volcanic and sediment soil based parent materials in dry land", IOP Conference Series: Earth and Environmental Science, 2021

Crossref

7

www.scilit.net

Internet

31 words — 1%

8

repository.unib.ac.id

Internet

31 words — 1%

9

Gaby Chintya, Dermiyati Dermiyati, Sarno Sarno, Jamalam Lumbanraja. "PENGARUH KOMBINASI PUPUK ORGANONITROFOS DAN PUPUK ANORGANIK TERHADAP NITROGEN TOTAL SELAMA PERTUMBUHAN JAGUNG MANIS (*Zea Mays saccharata*) DI TANAH ULTISOL", Jurnal Agrotek Tropika, 2019

Crossref

24 words — < 1%

10

worldwidescience.org

Internet

24 words — < 1%

11

W Anggraini, I Fiteriani, Nur Nunu Prihantini, Fri Rahmawati, A Susanti, E Septiyani. "The effect of organic fertilizers and inorganic fertilizer on mustard growth in Bahway village, Balik Bukit district, West Lampung regency", Journal of Physics: Conference Series, 2021

Crossref

20 words — < 1%

12

Septiyana, Husnain, L R Widowati, A F Siregar, A Samsun. "The use of soil ameliorants and fertilizers to increase the yields of rice and maize in ultisols Lampung, Indonesia", IOP Conference Series: Earth and Environmental Science, 2021

Crossref

20 words — < 1%

13 T Rostaman, H Wibowo, Nurjaya. "The effects of nano inorganic fertilizer application on rice (*Oryza sativa* L) productivity", IOP Conference Series: Earth and Environmental Science, 2021

19 words — < 1%

Crossref

14 agrobreedjournal.ir

Internet

18 words — < 1%

15 repository.lppm.unila.ac.id

Internet

18 words — < 1%

16 F. Yu, D. Driss - Ecole, J. Rembur, V. Legué, G. Perbal. "Effect of microgravity on the cell cycle in the lentil root", *Physiologia Plantarum*, 2002

14 words — < 1%

Crossref

17 Margaret H. N. Mollel, Elisabeth M. A. Goyvaerts. "Preliminary examination of factors affecting *Agrobacterium tumefaciens*-mediated transformation of marula, *Sclerocarya birrea* subsp. *caffra* (Anacardiaceae)", *Plant Cell, Tissue and Organ Culture*, 2004

12 words — < 1%

Crossref

18 medcraveonline.com

Internet

12 words — < 1%

19 www.rspo.org

Internet

12 words — < 1%

20 Yu Qin, Wu Ye, Suili Feng. "Scheduling for Wireless Networks Using Fuzzy Logic", 2006 International Conference on Wireless Communications, Networking and Mobile Computing, 2006

11 words — < 1%

Crossref

21 Dermiyati, A Niswati, B P Sanjaya, J Lumbanraja, M A S Arif, P Amalia, S Triyono, S Yusnaini. " Soil fauna population during the maize (.) growth with the addition of organonitrophos, inorganic fertilizer and biochar ", IOP Conference Series: Earth and Environmental Science, 2018
Crossref 11 words — < 1 %

22 repository.ub.ac.id
Internet 10 words — < 1 %

23 Catur Herison, Rustikawati Rustikawati, Hasanudin Hasanudin, Usman K.J. Suharjo, Merakati Handajaningsih, Nursalim Nursalim. "Manually Feed NPK Fertigation Improves Growth and Yield of Chili Pepper Hybrids in Ultisol", Akta Agrosia, 2020
Crossref 9 words — < 1 %

24 media.neliti.com
Internet 9 words — < 1 %

25 www.vulhm.cz
Internet 9 words — < 1 %

26 F. O. OLASANTAN, A. W. SALAU, E. E. ONUH. " INFLUENCE OF CASSAVA () INTERCROP ON GROWTH AND FRUIT YIELDS OF PEPPER (SPP.) IN SOUTH-WESTERN NIGERIA ", Experimental Agriculture, 2007
Crossref 8 words — < 1 %

27 William W. Kirk. "Impact of Different US Genotypes of Phytophthora infestans on Potato Seed Tuber Rot and Plant Emergence in a Range of Cultivars and Advanced Breeding Lines", Potato Research, 05/2009
Crossref 8 words — < 1 %

28 H S Wulanningtyas, S Wulandari, Sudarsono, A Kasim, M S Lestari, Y Baliadi. "Effectiveness of rice 8 words — < 1 %

straw with biodecomposer and biofertilizer application in new land clearing in Merauke, Papua", IOP Conference Series: Earth and Environmental Science, 2021

Crossref

29 Pujono H R, S Kukuh, R Evizal, Afandi, A Rahmat. "The effect of POME application on production and yield components of oil palm in Lampung, Indonesia", IOP Conference Series: Earth and Environmental Science, 2021

8 words — < 1%

Crossref

30 epdf.pub

Internet

8 words — < 1%

31 cloudfront.escholarship.org

Internet

6 words — < 1%

32 M Alwi, W A Yusuf, A Fahmi. "Improving productivity of rice yield on tidal swampland using soil amendment", IOP Conference Series: Earth and Environmental Science, 2021

6 words — < 1%

Crossref

33 Pradeep Kumar Srivatsava, G. R. Maruthi Sankar, P. Vijaya Kumar, S. P. Singh, N. Rani, Archana Singh, V. K. Agarwal. "Effects of Organic and Inorganic Fertilizers on Soil and Plant Nutrients and Yields of Pearl Millet and Wheat under Semi-arid Inceptisols in India", Communications in Soil Science and Plant Analysis, 2015

6 words — < 1%

Crossref

EXCLUDE QUOTES ON

EXCLUDE MATCHES

OFF

EXCLUDE BIBLIOGRAPHY ON