

THE EFFECT OF SILICA AND MANGANESE APPLICATION ON RICE GROWTH AND YIELD

By Paul Benyamin Timotiwu and Maya Maeistia Dewi

THE EFFECT OF SILICA AND MANGANESE APPLICATION ON RICE GROWTH AND YIELD

Paul Benyamin Timotiwu¹⁾ and Maya Maeistia Dewi

Faculty of Agriculture University of Lampung

Jl. Sumantri Brojonegoro No.1 Bandar Lampung 35145 Bandar Lampung Indonesia

¹⁾ Corresponding author Phone: +62-8154039127 E-mail: paul.timotiwu@gmail.com

Received: May 15, 2014 / Accepted: August 7, 2014

ABSTRACT

The application of silica and manganese through the leaves can be directly absorbed by the rice plants for growth and yield. This study aimed to determine the effect of the increasing silica and manganese concentration and their interaction on plant growth and rice yield. The greenhouse research was conducted in the Seed and Plant Breeding Laboratory, Agriculture Faculty, Lampung University from June–September 2013. The treatments were arranged in a 5x2 factorial consisting of five concentrations of Si [(0, 50, 100, 150, and 200 ppm), and two concentrations of Mn (0 and 5ppm)] in a randomized complete block design with three replications. The results showed that increasing concentration of Si increased the green of leaves, the number of productive tillers, dry weight, filled grains number, and decreased angle of the tiller and the number of empty grains. The application of Mn increased the growth and rice yield, except angle of tillers and the number of empty grains. In addition, the effect of Mn depended on the level of Si concentration. Application of Mn with increasing Si concentration increased plant growth and rice yield.

Keywords: manganese, rice, silica

INTRODUCTION

Rice crops (*Oryza sativa* L.) is a producer of staple food as a source of carbohydrate for most of the world's population. In Indonesia, almost 95% of the population consumes rice as the staple food. Every year the demand for rice is increasing with the increase of population. On the other hand, the increase of rice yield is

stagnant which mandates an effort to increase the yield to meet the demand for rice.

Indonesia has sought to increase yield of rice through many ways, one of which is intensification cultivation through fertilization. Fertilization is the provision of materials, both to the soil and the plants intended to supplement for the nutrients that can improve soil fertility and increase the quantity and quality of crops (Thompson and Troeh, 2005).

During the growth period, rice plants require a balanced nutrition to support their survival. In addition to macro nutrients, there are several other micro nutrients, such as silica (Si), which is also beneficial to the rice crop. Some studies suggested that Si has some important roles to the rice crop. The application of Si improves yields through increased efficiency of photosynthesis and induces resistance to pests and diseases (Makarim *et al.*, 2007).

Another essential micro nutrient, manganese (Mn) is also beneficial for the growth of rice. Manganese plays a role in the physiological processes in plants as an activator of several enzymes and a catalisator in splitting water molecules during photolysis process of photosynthesis (Gardner *et al.*, 1985; Humphrise, 2006; and Aref, 2012). According to Sutedjo (2008) and Agustina (2011), Mn could maintain the condition of the green leaves, especially on older leaves. With an addition of Mn, the old leaves were expected to be able to photosynthesize which resulted in the increase of yield potential.

Both Si and Mn are expected to directly increase the production of rice in order that the rice demand could be met. This study aimed to determine the effect of increasing concentration of silica and manganese, as well as their interaction on the rice growth and yield.

Accredited SK No.: 81/DIKTI/Kep/2011

<http://dx.doi.org/10.17503/Agrivita-2014-36-2-p182-188>




Table 2. Recapitulation of the effect of an increase in the concentration of Si and the addition of Mn on rice yield

Comparison	Variable		
	PG	EG	RY
-----% difference-----			
Manganese			
P1:Mn ₀ vs Mn ₁	13.04**	-57.58**	1.78**
Silica			
P2:Si-Linier	**	**	**
P3: Si-Quadratic	**	nd	*
Interaction			
P4:P1×P2	**	**	**
P5:P1×P3	nd	*	nd
The response of Mn			
P6:Si ₀ :Mn ₀ vs Mn ₁	3.69nd	-107.34**	1.20nd
P7:Si ₁ :Mn ₀ vs Mn ₁	11.33**	-55.14**	3.23nd
P8:Si ₂ :Mn ₀ vs Mn ₁	14.53**	-49.52**	10.85**
P9:Si ₃ :Mn ₀ vs Mn ₁	16.82**	-35.57**	12.76**
P10:Si ₄ :Mn ₀ vs Mn ₁	17.43**	-32.58**	14.42**
The response curve of Si			
P11:Mn ₀ :Si-Linier	**	**	**
P12:Mn ₀ :Si-Quadratic	-	**	-
P13:Mn ₁ :Si-Linier	**	tn	**
P14:Mn ₁ :Si-Quadratic	-	tn	-

Remarks: Mn₀= 0 ppm Manganese; Mn₁= 5 ppm Manganese; Si₀= 0 ppm Silica; Si₁= 50 ppm Silica; Si₂= 100 ppm Silica; Si₃= 150 ppm Silica; Si₄= 200 ppm Silica; PG= filled grains number; EG=the empty grains number; RY= rice yield. * and ** significant at $\alpha = 5\%$ and 1%, nd= not significant at $\alpha = 5\%$

This finding was similar to the result of Junior *et al.* (2010), the addition of 2 mmol l⁻¹ Si in a hydroponic solution was to increase the efficiency of photosynthesis of rice plants by strengthening the leaf angle at 19°. The strengthened leaf angle as a result of Si application permitted the leaves to grow more upright and elongate well. This resulted in more efficient absorption of sunlight for photo-synthesis. The increased efficiency in the absorption of sunlight permitted photosynthetic activity to proceed smoothly.

When sunlight was absorbed by the leaves, a photolysis process of splitting water molecules happened. Millaleo *et al.* (2010) mentioned that the role of Mn as a catalyst in the photolysis process would indirectly increase the activity of photo-synthesis. Increased photosynthetic activity affected the concentration of chlorophyll in the leaf tissues. Chlorophyll concentration determined leaf green index measured as SPAD value. The greater the chlorophyll concentration was, the greater the value of SPAD would be, which measured the greater leaf green level. The result showed that the addition of 5 ppm Mn increased the green level of leaves of rice, as compared to those without Mn addition (Figure 2). Working with beans, Santoz *et al.* (2013) reported that a decline in the rate of photosynthesis and the SPAD values as much as 18% and 70%, respectively in the leaves resulted from Mn deficiency. It proved that Mn participated in enhancing photosynthetic activity through increasing the concentration of chlorophyll. These conditions made the leaves remain green even though they were at senescence age.

The positive effects of increased concentrations of Si and Mn applications on the rice plants were also evident from the increase in the number of productive tillers (Figure 3) and dry weight (Figure 4). The increasing concentration to 200 ppm accompanied by the addition of 5 ppm Mn increased organic compounds synthesized by plants through the increased photosynthesis. This synthesis produced starches, lipids, and proteins which were used by the plant for the division process and the increase of cell size, which resulted in the increase on the number of tillers (Salisbury and Ross, 1991 and Taiz and Zeiger, 2010). Assimilated products were then used to increase the supply of organic materials in the plant tissues in order to increase their dry weight. Junior *et al.* (2010) found that the addition of 2 mmol l⁻¹ accompanied by an increase in Mn concentration up to 10 $\mu\text{mol l}^{-1}$ can increase dry matter production of rice. Similar result was found in cucumber plants grown with Si supply, Maksimovi *et al.* (2012) described that root and shoot dry weight increased significantly as the concentration of Mn in the nutrient solution was 0.5 μM .




Figure 1. Relationship between elevated concentrations of Si with and without Mn on angle of tillers




Figure 2. Relationship between elevated concentrations of Si with and without Mn on the level of greenness of leaves




Figure 3. Relationship between elevated concentrations of Si with and without Mn the number of productive tillers




Figure 4. Relationship between elevated concentrations of Si with and without Mn on dry weight




Figure 5. Relationship between elevated concentrations of Si with and without Mn on the number of filled grains




Figure 6. Relationship between elevated concentrations of Si with and without Mn on the number of empty grains




Figure 7. Relationship between elevated concentrations of Si with and without Mn on rice yield

Increasing assimilated products as an indicator of photosynthetic efficiency improvement due to the increased concentration of Si and Mn addition increased the number of filled grains (Figure 5). Assimilated products of photosynthesis translocated from the leaves into the grain to increase the number of filled grains. The increasing number of filled grains and the reducing number of empty grains were shown at Figure 6. Increasing the number of filled grains led to an increase in rice production (Figure 7).

Overall, the results demonstrated that the interaction between the Si and Mn increased the growth and production of rice. Concentration of Si with Mn application increased the efficiency of photosynthesis beneficial to rice yield and growth.

CONCLUSIONS AND SUGGESTION

CONCLUSIONS

The increase of Si concentration to 200 ppm increased the green level of leaves, the number of productive tillers, dry weight, the number of filled grains, and rice yield, and reduced the angle of tillers and the number of empty grains.

The addition of 5 ppm Mn increased the growth and rice yield when compared to without Mn as indicated by all variables, except the angle of tillers and the number of empty grains.

Rice growth and yield increased with the increasing concentration of Si with the addition of 5 ppm Mn.

SUGGESTION

To support the efforts to increase rice production, the research toward limiting factors such as the environment is necessary. Since this study was done in a greenhouse, environmental factors could be controlled in a good fashion, so that the results obtained were not able to describe the yield significantly compared to the study conducted in the field. Therefore, similar field study on rice needs to be done to obtain more realistic results.

ACKNOWLEDGEMENTS

Many thanks are addressed to Dr. Ir. Saiful Hikam, M.Sc., Lampung University for manuscript correction of and final improvement of the language.

REFERENCES

- Agustina, L. 2011. Micro nutrient I (Fe, Mn, Zn, Cu, B, Mo dan Cl) importance of nutrient deficiency and toxicity (in Indonesian). Post-Graduate Program University of Brawijaya. Malang. p. 25-32.
- Aref, F. 2012. Manganese, iron, and copper contents in leaves of maize plant (*Zea mays* L.) grown with different boron and zinc micronutrients. African Journal of Biotechnology 11(4): 896-903.
- Gardner, F.P., R.B. Pearce and R.L. Mitchell. 1985. Physiology of crop plants. The Iowa State University Press. Iowa. p. 98-129.

Paul Benyamin Timotiwi: *The Effect of Silica and Manganese Application*.....

- Humphrise, J.M. 2006. Handbook of plant nutrition. Edited by Allen V. Barker and David J. Pilbeam. CRC Press. New York. p. 351–366.
- Júnior, L.A.Z., R.L.F. Fontes, J.C.L. Neves, G.H. Korndörfer and V.T. de Ávila. 2010. Rice grown in nutrient solution with doses of manganese and silicon. *R. Bras. Ci Solo.* 34: 1629–1639.
- Makarim, A.K., E. Suhartatik and A. Kartohardjono. 2007. Silicon: Essential nutrient in the system of rice production (in Indonesian). Science and Technology (IPTEK) in Crops. 2 (2): 195–204.
- Maksimovic', J.D., M. Mojovic', V. Maksimovic', V. Römheld and M. Nikolic. 2012. Silicon ameliorates manganese toxicity in cucumber by decreasing hydroxyl radical accumulation in the leaf apoplast. *Journal of Experimental Botany*: 1–10.
- Millaleo, R., M. Reyes-Díaz, A.G. Ivanov, M.L. Mora and M. Alberdi. 2010. Manganese as essential and toxic element for plants: Transport, Accumulation and Resistance Mechanisms. *Journal of Soil Science Plant Nutrition* 10(4): 476–494.
- Salisbury, F.B. and C.W. Ross. 1991. Plant physiology. Fourth Edition. Wadsworth Publishing Company. Belmont, California. p. 207–224.
- Santoz, E.F., B.J. Zanchim, A.G. de Campos, R.F. Garrone and J.L. Junior. 2013. Photosynthesis rate, chlorophyll content, and initial development of physic nut without micronutrient fertilization. *R. Bras. Ci.Solo.* (37): 1334–1342.
- Sutedjo, M.M. 2008. Fertilizer and how to fertilize (in Indonesian). Rineka Cipta. Jakarta. pp. 177
- Taiz, L. and E. Zeiger. 2010. Plant physiology. Fifth Edition. Sinaeur Associates Inc. Massachusetts, USA. p. 161–397.
- Thompson, L.M. and F.R. Troeh. 2005. Soil and soil fertility. Sixth Edition. Blackwell Publishing, Iowa. pp. 489.

THE EFFECT OF SILICA AND MANGANESE APPLICATION ON RICE GROWTH AND YIELD

ORIGINALITY REPORT

17%
SIMILARITY INDEX

PRIMARY SOURCES

1	Paul B Timotiwu, Agustiansyah ., Ermawati ., Suci Amalia. "The Effects of Foliar Boron and Silica Through the Leaves on Soybean Growth and Yield", Journal of Agricultural Studies, 2018 <small>Crossref</small>	263 words — 10%
2	www.itraveldeals.in <small>Internet</small>	44 words — 2%
3	fr.scribd.com <small>Internet</small>	33 words — 1%
4	repository.unib.ac.id <small>Internet</small>	23 words — 1%
5	academic.oup.com <small>Internet</small>	20 words — 1%
6	Luiz Antônio Zañão Júnior, Renildes Lúcio Ferreira Fontes, Júlio César Lima Neves, Gaspar Henrique Korndörfer et al. "Rice grown in nutrient solution with doses of manganese and silicon", Revista Brasileira de Ciência do Solo, 2010 <small>Crossref</small>	17 words — 1%
7	journal.ipb.ac.id <small>Internet</small>	11 words — < 1%

8 scialert.net
Internet

9 words — < 1%

9 "Correlation, path analysis and stress indices studies of Saltol introgressed lines of rice for salinity tolerance", ELECTRONIC JOURNAL OF PLANT BREEDING, 2020
Crossref

7 words — < 1%

10 Paul A. Counce, Bobby R. Wells, Richard J. Norman. "Simulated Hail Damage to Rice: I. Susceptible Growth Stages", Agronomy Journal, 1994
Crossref

7 words — < 1%

11 uad.portalgaruda.org
Internet

6 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF