


# Sesbagrandidiflorain A and B: isolation of two new 2-arylbenzofurans from the stem bark of *Sesbania grandiflora*

*By* Noviany Noviany



## Sesbagrandidflorain A and B: isolation of two new 2-arylbenzofurans from the stem bark of *Sesbania grandiflora*

Noviany Noviany, Arif Nurhidayat, Sutopo Hadi, Tati Suhartati, Muhammad Aziz, Neny Purwitasari & Iman Subasman

To cite this article: Noviany Noviany, Arif Nurhidayat, Sutopo Hadi, Tati Suhartati, Muhammad Aziz, Neny Purwitasari & Iman Subasman (2018) Sesbagrandidflorain A and B: isolation of two new 2-arylbenzofurans from the stem bark of *Sesbania grandiflora*, *Natural Product Research*, 32:21, 2558-2564, DOI: [10.1080/14786419.2018.1425858](https://doi.org/10.1080/14786419.2018.1425858)

56

To link to this article: <https://doi.org/10.1080/14786419.2018.1425858>



View supplementary material [↗](#)



Published online: 16 Jan 2018.



Submit your article to this journal [↗](#)



Article views: 86



View Crossmark data [↗](#)



## Sesbigrandiflorain A and B: isolation of two new 2-arylbenzofurans from the stem bark of *Sesbania grandiflora*

Noviany Noviany<sup>a</sup>, Arif Nurhidayat<sup>a</sup>, Sutopo Hadi<sup>a</sup>, Tati Suhartati<sup>a</sup>, Muhammad Aziz<sup>b</sup> ,  
Neny Purwitasari<sup>c</sup> and Iman Subasman<sup>d</sup> 

<sup>a</sup>Faculty of Mathematics and Natural Sciences, Department of Chemistry, University of Lampung, Bandar Lampung, Indonesia; <sup>b</sup>Institute of Innovative Research, Tokyo Institute of Technology, Tokyo, Japan; <sup>c</sup>Faculty of Pharmacy, Department of Pharmacognosy and Phytochemistry, University of Airlangga, Surabaya, Indonesia; <sup>d</sup>STAI Al-Ihya, Kuningan, Indonesia

### ABSTRACT


Native to tropical Asia, *Sesbania grandiflora* (L.), Pers is a member of the Fabaceae family of flowering plants. All parts of *S. grandiflora* are used in traditional medicine and phytochemical investigations have been conducted on extracts of the leaves, seeds and roots of *S. grandiflora* to provide scientific validation of its properties. However, to date, no study has determined the phytochemical constituents of *S. grandiflora* stem bark. The stem bark powdered of *S. grandiflora* was extracted exhaustively with *n*-hexane, EtOAc and 90% aqueous MeOH sequentially. In this study, we successfully isolated two new 2-arylbenzofurans, sesbigrandiflorain A and B, from the EtOAc stem bark of *S. grandiflora*. The structure elucidation of these compounds was determined by using one- and two-dimensional nuclear magnetic resonance, ultraviolet and infrared spectroscopy and electrospray ionisation time-of-flight mass spectrometry. The finding expands the understanding of the natural constituents of the Fabaceae and, in particular, the *Papilionoideae* genera.

### 54 ARTICLE HISTORY


Received 1 November 2017  
Accepted 3 January 2018

### KEYWORDS

2-arylbenzofuran;  
sesbigrandiflorain; *Sesbania grandiflora*



  Noviany Noviany  [noviany@fmipa.unila.ac.id](mailto:noviany@fmipa.unila.ac.id)

 Supplemental data for this article can be accessed at <https://doi.org/10.1080/14786419.2018.1425858>.

© 2018 Informa UK Limited, trading as Taylor & Francis Group

## 1. Introduction

Plants of the family Fabaceae, particularly species of the Papilionoideae subfamily, have been extensively investigated for their phytochemical and pharmacological properties (Gođevac et al. 2008). Several secondary metabolites have been isolated from members of this family, including alkaloids, non-protein amino acids, flavonoids, isoflavonoids, coumarins, phenylpropanoids, anthraquinones, terpenoids and cyanogenic glycosides (Wink and Mohamed 2003). Among them, isoflavonoids are predominantly found in plants of the Papilionoideae subfamily (Kirmızıbekmez et al. 2015). A large number of new and known isoflavonoids has been characterised from plants in which the majority of these compounds were isolated from the Fabaceae (Dewick 2005; Veitch 2007). Isoflavonoids, with their structural diversity, were reported to possess wide range of biological activities against different strains of bacteria, fungi, viruses, plasmodium and various cancer cell lines (Kraft et al. 2001; Lo et al. 2002; Koyasomboon et al. 2006).

*Sesbania grandiflora* is a member of the Fabaceae family native to tropical Asia, including India, Malaysia, Indonesia, Myanmar and the Philippines. All parts of *S. grandiflora* are used in traditional medicine particularly in south-eastern Asia and India to treat various diseases including bacterial infections. Generally, the root is applied as a poultice to relieve from inflammation and fever. Ground root of *S. grandiflora* var. *coccinea* mixed with water is applied externally as a poultice to treat rheumatic swellings. The bark is used as astringent to cure smallpox. In Philippines, the decoction from the crushed bark is used for the treatment of ulcers in the mouth and alimentary canal. In Java, the local healers use the crushed bark for the treatment of thrush and infantile disorders of the stomach, while in Cambodia, the pounded bark is applied to treat the scabies. The leaves juice is used to treat worms, biliousness, fever, gout, itchiness and leprosy. In Malaysia, the crushed leaves are applied to cure sprains and bruises. While in Ayurveda, the leaves are employed for the treatment of epileptic fits and the fruits are used for the treatment of anaemia, bronchitis, fever and tumours. The leaves and flowers juice is a popular remedy for nasal catarrh and headache, head congestion or stuffy nose. The flowers juice is dropped into the eyes to correct dim visions (Wagh et al. 2009).

Several studies have been conducted on extracts obtained from *S. grandiflora* trees (local name: turi) to provide scientific validation of its properties. The methylene chloride and methanol extracts of aerial parts of *S. grandiflora* have shown to have some antifungal activities (Goun et al. 2003). A subsequent research by Doddola et al. (2008) revealed that the leaf juice of *S. grandiflora* exhibits significant antiurolithiatic activity against calcium oxalate-type stones as well as antioxidant properties. Besides that, Laladhas et al. (2010) had evaluated the flowers of *S. grandiflora* *in vivo* and *in vitro* using different cancer cell lines, disclosed that the flowers may serve as a potential anticancer drug candidate. Moreover, the *n*-hexane extract of *S. grandiflora* seeds, which was investigated by Shareef et al. (2012), showed that it possessed antioxidant, anti-inflammatory, analgesic and antipyretic activities.

The earliest phytochemical investigations of *S. grandiflora* were conducted in the 1960s, and resulted in the isolation of  $\alpha$ -5-methyl-5-pentacosanol for the first time from *S. grandiflora* leaves (Tiwari and Bajpai 1964). Recently, Pollard et al. (2011) reported the isolation of a galactomannan from the seeds of *S. grandiflora*. In addition, Hasan et al. (2012) and Noviany et al. (2012) obtained and evaluated the antituberculosis activity of several phenolic compounds from the roots of *S. grandiflora*.

Nowadays, the discovery of lead compounds for development of new scaffolds of drugs from plants, has received considerable attention. Based on the explanation above, *S. grandiflora* is one of the potential medicinal plants that can be employed as a source of new biologically active compounds. Hence, the phytochemical study on this plant is considered essential. Even though *S. grandiflora* was extensively studied by other researchers for its phytopharmacological potential, especially the leaves, flowers and aerial parts of the plant, no phytochemical studies extensively have been performed on the stem bark of *S. grandiflora*. In terms of finding the lead compounds therefore, in this study, we investigated the phytochemical constituents of the stem bark of *S. grandiflora*. The ethyl acetate (EtOAc) extract of *S. grandiflora* stem bark was subjected to further analysis, which led to the isolation of two new 2-arylbenzofurans.

## 2. Results and discussion

The EtOAc extract of *S. grandiflora* stem bark contained two compounds, **1** and **2** (structures are shown in Figure 1), after repeated Column Chromatography (CC) separation.

Sesbagrandiflorain A (Compound **1**) was obtained as a needle-like yellow crystal. The UV spectrum of Compound **1** in MeOH, with  $\lambda_{\text{max}}$  at 213, 264, 348 and 363 nm, indicated the presence of a conjugated system in the isolated structure (Komatsu et al. 1981). The IR absorption of Compound **1** demonstrated the presence of a hydroxyl group (OH) at  $3407 \text{ cm}^{-1}$ , a saturated aliphatic carbon group (CH) at  $2924 \text{ cm}^{-1}$ , an aldehyde group (CHO) at  $1652 \text{ cm}^{-1}$ , and an olefinic group at  $1438 \text{ cm}^{-1}$ . ESI-TOF-MS of Compound **1** revealed a pseudomolecular ion peak  $[M - H]^+$  at  $m/z$  313.0709, which was consistent with a molecular formula of  $\text{C}_{17}\text{H}_{14}\text{O}_6$ . The structure elucidation of Compound **1** was deduced from detailed analysis of  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectral data aided by 2D NMR experiments such as  $^1\text{H}$ - $^1\text{H}$  COSY, HSQC, HMBC and NOESY.

The  $^1\text{H}$  NMR spectrum of Compound **1** (Table S1) revealed a set of aromatic proton resonances with an AB splitting pattern of the A-ring at  $\delta_{\text{H}}$  6.67 (dd,  $J = 8.4; 2.2 \text{ Hz}$ , H-5), 6.71 (d,  $J = 2.2 \text{ Hz}$ , H-7) and 7.55 (d,  $J = 8.4 \text{ Hz}$ , H-4), as well as a methoxyl group ( $\text{OCH}_3$ ) at  $\delta_{\text{H}}$  3.88 (s) and one proton of the CHO group at  $\delta_{\text{H}}$  9.82 (s) on a benzofuran moiety. Another pair of




Figure 1. The structures of the isolated compounds.

*meta*-coupled aromatic protons were evident at  $\delta_{\text{H}}$  6.34 (d,  $J = 2.2$  Hz, H-4') and 6.68 (d,  $J = 2.2$  Hz, H-6') along with a  $\text{OCH}_3$  group at  $\delta_{\text{H}}$  3.83 (s), which indicated a 1,2,3,5-tetrasubstituted benzene in the B-ring. The  $^{13}\text{C}$  NMR data showed peaks characteristic of arylbenzofuran moiety following the signal observed at  $\delta_{\text{C}}$  133.8 (C-4), 109 (C-5), 100.6 (C-7), 98.9 (C-4') and 157.4 (C-6'). Thus, based on the  $^1\text{H}$  and  $^{13}\text{C}$  NMR data (Table S1), Compound **1** was strongly suggested to be a 2-arylbenzofuran.

Further characterisation of Compound **1** was performed using heteronuclear multiple bond correlation (HMBC) analysis (Figure S1). The placement of the  $\text{OCH}_3$  and OH groups in the A- and B-rings was determined by HMBC. The long-range correlations between H- $\text{OCH}_3$  (A-ring,  $\delta_{\text{H}}$  3.88) and C-6 ( $\delta_{\text{C}}$  160), H- $\text{OCH}_3$  (B-ring,  $\delta_{\text{H}}$  3.83) and C-5' ( $\delta_{\text{C}}$  161.9) and H-OH (B-ring,  $\delta_{\text{H}}$  10.21) and C-1' ( $\delta_{\text{C}}$  107.8)/C-4' ( $\delta_{\text{C}}$  98.9) supported the assigned positions for the  $\text{OCH}_3$  and OH groups in the A and B-rings, respectively. The assignment of the CHO group at C(3) also was determined by HMBC, which displayed a cross-peak between the CHO group at C-3 ( $\delta_{\text{C}}$  119.2) and C-1' ( $\delta_{\text{C}}$  107.8). This conclusion was supported by nuclear Overhauser enhancement (NOE) analysis (Figure S2). In this spectrum, a strong NOE association of H-C(4)/H-CHO(3); H-C(7)/MeO-C(6); H-C(4')/MeO-C(5') and H-C(6')/MeO-C(5'), established the location of the CHO and the  $\text{OCH}_3$  groups, respectively.

From the above spectroscopic analyses, sesbagrandiflorain A was identified as 6-methoxy-2-(2',3'-dihydroxy-5'-methoxyphenyl)-1-benzofuran-3-carbaldehyde (Compound **1**).

Sesbagrandiflorain B (Compound **2**) was also isolated as a needle-like yellow crystal, and its molecular formula was determined as  $\text{C}_{16}\text{H}_{12}\text{O}_6$  from ESI-TOF-MS, which exhibited  $[\text{M} - \text{H}]^+$  at  $m/z$  299.0554. The  $^1\text{H}$  NMR spectral data of Compounds **1** and **2** displayed the same substitution-pattern signals of the 2-arylbenzofuran skeleton except for the hydroxyl group at C-6 on the A-ring. Similar features shown by Compounds **1** and **2** was also supported by comparison of their  $^{13}\text{C}$  NMR spectral data (Table S1). The assignment of the OH group was afforded from the results of HMBC analysis (correlations: HO-C(6)/C(6); H-C(5)/C(6); and H-C(4)/C(6)) and nuclear Overhauser enhancement (NOE) spectroscopy data (NOE associations: H-C(7)/HO-C(6)). The HMBC correlation and the NOE associations of Compound **2** are shown in Figure S3 and S4, respectively.

Compound **2** displayed very similar one- and two-dimensional NMR profiles to those of Compound **1**. The only difference was that Compound **1** had a  $\text{OCH}_3$  group attached at C-6 (A-ring) rather than the OH group found in Compound **2** at the same position. Therefore, Compound **2** was characterised as 6-hydroxy-2-(2',3'-dihydroxy-5'-methoxyphenyl)-1-benzofuran-3-carbaldehyde (sesbagrandiflorain B). This type of 2-arylbenzofuran compound was isolated previously by Tanaka et al. (2004) from the roots of *Erythrina variegata*. This is the first report of the occurrence of 2-arylbenzofurans in the stem bark of *S. grandiflora*.

### 3. Experimental

#### 3.1. General methods

Thin-layer chromatography (TLC) was conducted on pre-coated silica gel 60 GF<sub>254</sub> plates (Merck, Darmstadt, Germany) with an absorbent thickness of 0.25 mm sprayed with  $\text{Ce}(\text{SO}_4)_2$  solution for spot visualisation. In addition, preparative TLC was performed on square glass plates with a side length of 0.2 m coated with 0.5 mm Kieselgel F<sub>254</sub> (Merck), which were

air-dried and used without prior activation. Column chromatography (CC) was performed on silica gel (Kieselgel 60, 70–230 mesh ASTM; Merck).

Nuclear magnetic resonance (NMR) spectra were recorded for specimens dissolved in acetone- $d_6$ , with tetramethylsilane as an internal standard, using 500 MHz ( $^1\text{H}$ ) and 125 MHz ( $^{13}\text{C}$ ) spectrometers (Agilent Technologies, Santa Clara, CA, USA). Electrospray ionisation time-of-flight mass spectrometry (ESI-TOF-MS) was performed using a micrOTOF-Q mass spectrometer (Bruker Daltonics Inc., Billerica, MA, USA). Potassium bromide-type infrared (IR) spectra were recorded using a Nicolet Avatar 360 spectrometer (Thermo Fisher Scientific, Waltham, MA, USA). Finally, ultraviolet (UV) spectra were recorded using a UV-Vis Cary 100 spectrophotometer (Agilent Technologies).

### 3.2. Plant material

Samples of the stem bark of *S. grandiflora* were collected on June 2015 in Luhu Ratu, Bandar Lampung, Indonesia. The identity of the plant specimens (No. N-IV) was confirmed by the Herbarium Bogoriense, Research Centre for Biology, Indonesian Institute of Sciences, Bogor, Indonesia. A voucher specimen was deposited at the herbarium.

### 3.3. Extraction and isolation

Air-dried and powdered samples of the stem bark of *S. grandiflora* (1.5 kg) were extracted exhaustively with *n*-hexane, EtOAc and 90% aqueous methanol (MeOH) sequentially at room temperature. Extraction was repeated three times, and the extracts were filtered and evaporated using a rotary vacuum evaporator at 40°C. The masses of the *n*-hexane, EtOAc and MeOH extracts obtained were 15, 60 and 53 g, respectively. The EtOAc extract was selected for further analysis. The EtOAc extract was fractionated by silica-gel vacuum liquid chromatography and eluted with EtOAc:*n*-hexane with a volume ratio ranging from 0–100%, yielding nine major fractions (Fr.), Fr. E1–E9. Fr. E7 (237 mg) was further fractionated by CC on silica gel, and eluted with an acetone:*n*-hexane (2:98–65:35 v/v) gradient to yield 87 sub-fractions (Fr. E7.1–E7.87). A yellow crystal (Compound 1) was obtained from sub-fractions Fr. E7.38–E7.52 after recrystallisation with acetone:*n*-hexane (3:7 v/v). The total amount of Compound 1 obtained was 5.2 mg. The fractionation of Fr. E3 (325 mg) was performed in the same way, using the eluent of an acetone:*n*-hexane (5:95–4:60 v/v) gradient to yield 42 sub-fractions. The fractions were examined by HPLC, and those with identical profiles were combined to give three major sub-fractions (Fr. E3.1–E3.3). Fr. E3.2 (38 mg) was further purified by preparative TLC (plates: 20 × 7 × 0.5 mm) using an acetone:*n*-hexane (3:7 v/v) gradient to provide Compound 2 (4.9 mg).

#### 3.3.1. Sesbigrandiflorain A (1): 6-methoxy-2-(2',3'-dihydroxy-5'-methoxyphenyl)-1-benzofuran-3-carbaldehyde

$\text{C}_{17}\text{H}_{14}\text{O}_6$ , needle-like yellow crystal, IR (KBr)  $\nu_{\text{max}}$  3407, 2924, 1652, 1438, m.p. 215–216°C, UV (methanol)  $\lambda_{\text{max}}$  213, 264, 348, 363 nm. ESI-TOF-MS  $m/z$  313.0709  $[\text{M} - \text{H}]^+$ , calculated 313.0718, NMR (Acetone- $d_6$ ) 3.83 (1H, s, H-OCH<sub>3</sub>), 3.21 (1H, s, H-OCH<sub>3</sub>), 6.34 (1H, d,  $J = 2.2$  Hz, H-4'), 7.7 (1H, dd,  $J = 2.2, 8.4$  Hz, H-5), 6.68 (1H, d,  $J = 2.2$  Hz, H-6'), 6.71 (1H, d,  $J = 2.2$  Hz, H-7), 7.55 (1H, d,  $J = 8.4$  Hz, H-4), 9.82 (1H, s, H-CHO), 10.21 (1H, s, H-OH),  $^{13}\text{C}$  NMR (Acetone- $d_6$ ): 56.1 (C-OCH<sub>3</sub>), 56.2 (C-OCH<sub>3</sub>), 88.5 (C-6'), 98.9 (C-4'), 100.6 (C-7), 107.8 (C-1'), 109.0 (C-5), 109.3

(C-9), 119.2 (C-3), 133.8 (C-4), 152.8 (C-3'), 157.4 (C-2'), 160.0 (C-6), 162.0 (C-5'), 162.9 (C-8), 164.4 (C-2), 191.1 (CHO) (Table S1).

### 3.3.2. *Sesbagrandiflorain B* (2): 6-hydroxy-2-(2',3'-dihydroxy-5'-methoxyphenyl)-1-benzofuran-3-carbaldehyde

$C_{16}H_{12}O_6$ , needle-like yellow crystal, IR (KBr)  $\nu_{max}$   $cm^{-1}$  3461, 1640, m.p. 172–173°C, UV (methanol)  $\lambda_{max}$  215, 265, 348, 368; ESI-TOF-MS  $m/z$  299.0554  $[M - H]^+$ , calculated 299.0561,  $^{16}O$ MR (Acetone- $d_6$ ) 3.83 (1H, s, -OCH<sub>3</sub>), 6.34 (1H, d,  $J = 2.2$  Hz, H-4'), 6.61 (1H, dd,  $J = 2.2, 8.4$  Hz, H-5), 6.64 (1H, d,  $J = 2.2$  Hz, H-7), 6.67 (1H, d,  $J = 2.2$  Hz, H-24), 7.53 (1H, d,  $J = 8.4$  Hz, H-4), 9.23 (1H, s, H-OH), 9.42 (s, H-OH), 9.97 (1H, s, H-CHO), 10.26 (s, H-OH),  $^{13}C$  NMR (Acetone- $d_6$ ): 56.1 (C-OCH<sub>3</sub>), 88.5 (C-6'), 98.8 (C-12), 104.2 (C-7), 107.9 (C-1'), 108.3 (C-9), 109.2 (C-5), 118.7 (C-3), 133.5 (C-4), 152.8 (C-3'), 157.3 (C-2'), 157.9 (C-6), 161.9 (C-5'), 162.5 (C-8), 164.7 (C-2), 191.3 (CHO) (Table S1).

## 4. Conclusions

In this study, two new 2-arylbenzofuran compounds, namely sesbagrandiflorain A and B, were obtained from the EtOAc extract of *S. grandiflora* stem bark. These compounds are reportedly present in the genus *Sesbania* and other members of the family Fabaceae. The bioactivity of both isolated compounds remains under investigation. Our findings expand our understanding of the natural constituents of the Fabaceae and, in particular, the *Papilionoideae* genera.

## Acknowledgments

The authors would like to thank the Director of Research and Community Services, Directorate General of Higher Education, The Ministry of Research, Technology and Higher Education, Republic of Indonesia that provide fund for this project to be undertaken through Penelitian Produk Terapan Scheme 2017 (No.071/SP2H/LT/DRPM/IV/2017). We also gratefully acknowledge Prof. Yana Maolana Syah from Institute Technology of Bandung, Indonesia, for the NMR spectra. Noviany wish to thank Andi Setiawan, Ph.D from University of Lampung and Mr. Tarso Rudiana from UIN Syarif Hidayatulloh Jakarta for their valuable suggestions.

## 5 Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by the Directorate of Research and Community Services, Directorate General of Higher Education, The Ministry of Research, Technology and Higher Education, Republic of Indonesia [grant number 071/SP2H/LT/DRPM/IV/2017].

## ORCID

Muhammad Aziz  <http://orcid.org/0000-0003-2433-8500>



## References

- Dewick, PM. 2005. Medicinal natural product. (2nd ed.), p.8–166, Chichester: John Wiley & Son, Ltd.
- Doddola S, Pasupulati H, Koganti B, Prasad KVSRG. 2008. Evaluation of *Sesbania grandiflora* for antiurolithiatic and antioxidant properties. *Journal of Natural Medicines*. 62:300–307.
- Gođevac D, Zdunić G, Šavikin K, Vajs V, Menković N. 2008. Antioxidant activity of nine Fabaceae species growing in Serbia and Montenegro. *Fitoterapia*. 79:185–187.
- Goun E, Cunningham G, Chu D, Nguyen C, Miles D. 2003. Antibacterial and antifungal activity of Indonesian ethnomedical plants. *Fitoterapia*. 74:592–596.
- Hasan N, Osman H, Mohamad S, Chong W, Awang K, Zahariluddin ASM. 2012. The chemical components of *Sesbania grandiflora* roots and their antituberculosis activity. *Pharmaceuticals*. 5:882–889.
- Kırmızıbekmez H, Uysal GB, Masullo M, Demirci F, Bağcı Y, Kan Y, Piacente S. 2015. Prenylated polyphenolic compounds from *Glycyrrhiza iconica* and their antimicrobial and antioxidant activities. *Fitoterapia*. 103:289–293.
- Komatsu M, Yokoe I, Shirataki Y. 1981. Studies on the constituents of *Sophora* species. XV. Constituents of the root of *Sophora franchetiana* Dunn. *Chemical & Pharmaceutical Bulletin*. 29:2069–2072.
- Koysomboon S, van Altena IV, Kato S, Chantrapromma K. 2006. Antimycobacterial flavonoids from *Derris indica*. *Phytochemistry*. 67:1034–1040.
- Kraft C, Jenett-Siems K, Siems K, Solis PN, Gupta MP, Bienzle U, Eich E. 2001. Andinermals A-C, antiparasmodial constituents from *Andira inermis*. *Phytochemistry*. 58:769–774.
- Laladhas KP, Cheriyan VT, Puliappadamba VT, Bava SV, Unnithan RG, Vijayammal PL, Anto RJ. 2010. A novel protein fraction from *Sesbania grandiflora* shows potential anticancer and chemopreventive efficacy, *in vitro* and *in vivo*. *Journal of Cell and Molecular Medicine*. 14:636–646.
- Lo WL, Chang F-R, Liaw CC, Wu YC. 2002. Cytotoxic coumaronochromones from the Roots of *Euchresta formosana*. *Planta Medica*. 68:146–151.
- Noviany, Osman H, Keng Chong, W, Awang, K, Manshoor, N. 2012. Isolation and characterisation of l, l'-binaphthalene-2,2'-diol, a new biaryl natural product from *Sesbania grandiflora* root. *Journal of Basic Applied Sciences*. 8:253–256.
- Pollard MA, Fischer P, Windhab EJ. 2011. Characterization of galactomannans derived from legume endosperms of genus *Sesbania* (Faboideae). *Carbohydrate Polymers*. 84:550–559.
- Shareef H, Rizwani GH, Zia-ul-Haq M, Ahmad S, Zahid H. 2012. Tocopherol and phytosterol profile of *Sesbania grandiflora* (Linn.) seed oil. *Journal of Medicinal Plants Research*. 6:3478–3481.
- Tanaka H, Hirata M, Etoh H, Sako M, Sato M, Murata J, Murata H, Darnaedi D, Fukai T. 2004. Six new constituents from the roots of *Erythrina variegata*. *Chemistry & Biodiversity*. 1:1101–1108.
- Tiwari RD, Bajpai RK. 1964. Chemical examination of *Sesbania grandiflora* leaves. II. Isolation and study of a saponin. *Proceedings of the National Academy of Sciences, India Section A: Physical Sciences*. 34:239–244.
- Veitch NC. 2007. Isoflavonoids of the leguminosae. *Natural Product Reports*. 24:417–464.
- Wagh VD, WaghkV, Tandale YN, Salve SA. 2009. Phytochemical, pharmacological and phytopharmaceutics aspects of *Sesbania grandiflora* (Hadga): a review. *Journal of Pharmacy Research*. 2:889–892.
- Wink M, Mohamed GIA. 2003. Evolution of chemical defense traits in the Leguminosae: mapping of distribution patterns of secondary metabolites on a molecular phylogeny inferred from nucleotide sequences of the *rbcl* gene. *Biochemical Systematics and Ecology*. 31:897–917.

# Sesbagrandidflorain A and B: isolation of two new 2-arylbenzofurans from the stem bark of Sesbania grandiflora

ORIGINALITY REPORT

# 23%

SIMILARITY INDEX

## PRIMARY SOURCES

- |   |   |                 |
|---|---|-----------------|
| 1 | <a href="https://journals.plos.org">journals.plos.org</a><br>Internet   | 34 words — 1%   |
| 2 | <a href="#">Edible Medicinal And Non-Medicinal Plants, 2014.</a><br>Crossref  | 28 words — 1%   |
| 3 | <a href="#">Nakano, H.. "Growth inhibitory alkaloids from mesquite (Prosopis juliflora (Sw.) DC.) leaves", Phytochemistry, 200403</a><br>Crossref   | 23 words — 1%   |
| 4 | <a href="#">Luiz Cláudio de Almeida Barbosa. "Synthesis of New Phytogrowth-Inhibitory Substituted Aryl-p-Benzoquinones", Chemistry &amp; Biodiversity, 05/2006</a><br>Crossref  | 22 words — 1%   |
| 5 | <a href="#">B. Naseri, M. Gheitury, M. Veisi. "Infections of bean plant and field soil are linked to region, root rot pathogen, and agro-ecosystem", Hellenic Plant Protection Journal, 2020</a><br>Crossref  | 20 words — 1%   |
| 6 | <a href="#">Natural Products, 2013.</a><br>Crossref   | 19 words — 1%   |
| 7 | <a href="#">Nobuya Katagiri, Makoto Okada, Yoshihiro Morishita, Chikara Kaneko. "Synthesis of chiral spiro 3-oxazolin-5-one 3-oxides (chiral nitrones) via a nitrosoketene intermediate and their asymmetric 1,3-dipolar cycloaddition reactions leading to the EPC synthesis of modified amino acids", Tetrahedron, 1997</a><br>Crossref | 16 words — < 1% |
| 8 | <a href="#">Syah, Y.M.. "Two oligostilbenes, cis- and trans-diptoindonesin B, from Dryobalanops</a>   | 16 words — < 1% |

- 
- 9 [dmd.aspetjournals.org](http://dmd.aspetjournals.org) 16 words — < 1%  
Internet
- 
- 10 [herbs.ph](http://herbs.ph) 16 words — < 1%  
Internet
- 
- 11 Marnix L. J. Brinkman, Floor van der Hilst, André P. C. Faaij, Birka Wicke. "Low-ILUC-risk rapeseed biodiesel: potential and indirect GHG emission effects in Eastern Romania", *Biofuels*, 2018 16 words — < 1%  
Crossref
- 
- 12 Henrik Franzyk, Søren Rosendal Jensen, Carl Erik Olsen, Jon Holbech Rasmussen. "SYNTHESIS OF NOVEL HYDROXYMETHYL SUBSTITUTED ANALOGUES RELATED TO CARBOVIR AND NEPLANOCIN A", *Nucleosides, Nucleotides and Nucleic Acids*, 2002 15 words — < 1%  
Crossref
- 
- 13 A. K. M. Shahidur Rahman. "Antibacterial activity of two limonoids from *Swietenia mahagoni* against multiple-drug-resistant (MDR) bacterial strains", *Journal of Natural Medicines*, 01/2009 15 words — < 1%  
Crossref
- 
- 14 Deni Hardiansyah, Christian Maass, Ali Asgar Attarwala, Berthold Müller, Peter Kletting, Felix M. Mottaghy, Gerhard Glatting. "The role of patient-based treatment planning in peptide receptor radionuclide therapy", *European Journal of Nuclear Medicine and Molecular Imaging*, 2015 14 words — < 1%  
Crossref
- 
- 15 Maiti, B.C.. "The regioselective synthesis of monomethoxynaphthylene diacetates", *Tetrahedron*, 20050214 14 words — < 1%  
Crossref
- 
- 16 Rino Leardini, Daniele Nanni, Antonio Tundo, Giuseppe Zanardi, Gian Franco Pedulli. "Liquid-crystalline quinoline derivatives", *Liquid Crystals*, 1987 14 words — < 1%  
Crossref
- 
- 17 [orca.cf.ac.uk](http://orca.cf.ac.uk)

14 words — &lt; 1%

**18** [worldagroforestry.org](http://worldagroforestry.org)

Internet

14 words — &lt; 1%

**19** [eprints.ums.edu.my](http://eprints.ums.edu.my)

Internet

14 words — &lt; 1%

**20** Puspa Asri, Nyoman, and Diah Agustina Puspita Sari. "Pre-Treatment of Waste Frying Oils for Biodiesel Production", *Modern Applied Science*, 2015.

Crossref

13 words — &lt; 1%

**21** Klara R. Birikh, Pablo L. Bernad, Vadim V. Shmanai, Andrei D. Malakhov, Mikhail S. Shchepinov, Vladimir A. Korshun. "Chapter 21 SNP Detection Using Trityl Mass Tags", Springer Science and Business Media LLC, 2009

Crossref

13 words — &lt; 1%

**22** Alvin, Alfonsus, John A. Kalaitzis, Blandine Sasia, and Brett A. Neilan. "Combined genetic and bioactivity-based prioritization leads to the isolation of an endophyte-derived antimycobacterial", *Journal of Applied Microbiology*, 2016.

Crossref

13 words — &lt; 1%

**23** Wink, M.. "Evolution of chemical defense traits in the Leguminosae: mapping of distribution patterns of secondary metabolites on a molecular phylogeny inferred from nucleotide sequences of the rbcL gene", *Biochemical Systematics and Ecology*, 200308

Crossref

13 words — &lt; 1%

**24** Jakse, R.. "Application of alkyl 3-dimethylamino-2-(1H-indol-3-yl)propenoates in the synthesis of 3-heteroarylindoles", *Tetrahedron*, 20040517

Crossref

12 words — &lt; 1%

**25** Alessio Cimmino, Lucia Maddau, Marco Masi, Benedetto T. Linaldeddu, Gennaro Pescitelli, Antonio Evidente. "Fraxitoxin, a New Isochromanone Isolated from ", *Chemistry & Biodiversity*, 2017

12 words — &lt; 1%

- 
- 26 Bang-Wei Yu, Jian-Guang Luo, Jun-Song Wang, Dong-Ming Zhang, Shi-Shan Yu, Ling-Yi Kong. "Pentasaccharide Resin Glycosides from ", *Journal of Natural Products*, 2011  
Crossref 12 words — < 1%
- 
- 27 [opus.bibliothek.uni-wuerzburg.de](http://opus.bibliothek.uni-wuerzburg.de)  
Internet 12 words — < 1%
- 
- 28 Fátima Churruca. "A New, Expeditious Entry to the Benzophenanthrofurane Framework by a Pd-Catalyzed C- and O-Arylation/PIFA-Mediated Oxidative Coupling Sequence", *European Journal of Organic Chemistry*, 06/2005  
Crossref 10 words — < 1%
- 
- 29 Li-Mei LI. "Chemical Constituents from the Seeds of *Ziziphus jujuba* var. *spinosa* (Bunge) Hu", *Journal of Integrative Plant Biology*, 4/2005  
Crossref 10 words — < 1%
- 
- 30 Zhi Na, Qi-Shi Song, Hua-Bin Hu. "Flavonoids from Twigs of ", *Natural Product Communications*, 2014  
Crossref 10 words — < 1%
- 
- 31 [mdpi.com](http://mdpi.com)  
Internet 10 words — < 1%
- 
- 32 [edumag.mrsu.ru](http://edumag.mrsu.ru)  
Internet 9 words — < 1%
- 
- 33 Qiong Gu, Rui-Rui Wang, Xue-Mei Zhang, Yun-Hua Wang, Yong-Tang Zheng, Jun Zhou, Ji-Jun Chen. "A New Benzofuranone and Anti-HIV Constituents from the Stems of ", *Planta Medica*, 2007  
Crossref 9 words — < 1%
- 
- 34 [f1000research.com](http://f1000research.com)  
Internet 9 words — < 1%
- 
- 35 [hdl.handle.net](http://hdl.handle.net)  
Internet 9 words — < 1%

- 
- 36 [www.thieme-connect.com](http://www.thieme-connect.com) Internet 9 words — < 1%
- 
- 37 H. J. Park, H. K. Jeong, M. I. Chang, M. H. Im et al. "Structure determination of new analogues of vardenafil and sildenafil in dietary supplements", *Food Additives and Contaminants*, 2007 9 words — < 1%  
Crossref
- 
- 38 Ohsaki, Ayumi, Ritsuko Yokoyama, Haruka Miyatake, and Yoshiyasu Fukuyama. "Two Diterpene Rhamnosides, Mimosasides B and C, from *Mimosa hostilis*", *CHEMICAL & PHARMACEUTICAL BULLETIN*, 2006. 9 words — < 1%  
Crossref
- 
- 39 [moam.info](http://moam.info) Internet 9 words — < 1%
- 
- 40 [www.scielo.br](http://www.scielo.br) Internet 9 words — < 1%
- 
- 41 [worldwidescience.org](http://worldwidescience.org) Internet 9 words — < 1%
- 
- 42 Abdelsamed I. Elshamy, Ayman F. Abdel-Razik, Mahmoud I. Nassar, Tahia K. Mohamed, Mohamed A. Ibrahim, Salah M. El-Kousy. " A new gorgostane derivative from the Egyptian Red Sea soft coral ", *Natural Product Research*, 2013 9 words — < 1%  
Crossref
- 
- 43 [www.freepatentsonline.com](http://www.freepatentsonline.com) Internet 8 words — < 1%
- 
- 44 Tsunashi Kamo, Michiko Kashiwabara, Kaori Tanaka, Shinji Ando, Hisao Shibata, Mitsuru Hirota. " Plant growth inhibitory activity of azo- and azoxyformamides from and ", *Natural Product Research*, 2006 8 words — < 1%  
Crossref
- 
- 45 Xue Zhang. "Three new bibenzyl derivatives from *Dendrobium nobile*", *Journal of Asian Natural Products Research*, 1/1/2006 8 words — < 1%  
Crossref
-

46	<a href="http://jos.hueuni.edu.vn">jos.hueuni.edu.vn</a> Internet	8 words — < 1%
47	<a href="http://www.thieme-connect.de">www.thieme-connect.de</a> Internet	8 words — < 1%
48	<a href="http://www.academicjournals.org">www.academicjournals.org</a> Internet	8 words — < 1%
49	<a href="http://link.springer.com">link.springer.com</a> Internet	8 words — < 1%
50	Bathini Thissera, Rizliya Visvanathan, Mohammad A. Khanfar, M. Mallique Qader et al. "Sesbania grandiflora L. Poir leaves: A dietary supplement to alleviate type 2 diabetes through metabolic enzymes inhibition", South African Journal of Botany, 2020 Crossref	8 words — < 1%
51	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet	8 words — < 1%
52	<a href="http://eprints.uthm.edu.my">eprints.uthm.edu.my</a> Internet	8 words — < 1%
53	<a href="http://academica-e.unavarra.es">academica-e.unavarra.es</a> Internet	8 words — < 1%
54	<a href="http://www.em-consulte.com">www.em-consulte.com</a> Internet	8 words — < 1%
55	<a href="http://harmonouniversa.selfip.org">harmonouniversa.selfip.org</a> Internet	8 words — < 1%
56	<a href="http://centaur.reading.ac.uk">centaur.reading.ac.uk</a> Internet	8 words — < 1%
57	"Retraction", Natural Product Research, 05/2010 Crossref	8 words — < 1%
58	Macias, F.A.. "11,16 Oxetane lactones. Spectroscopic evidences and conformational analysis", Tetrahedron, 20060814 Crossref	7 words — < 1%

59 van der Westhuyzen, C.W.. "Effect of substituent structure on pyrimidine electrophilic substitution", Tetrahedron, 20070618 7 words — < 1%  
Crossref

60 Abdel Kader Mansour, Mohga M. Eid, Nasser S. A. M. Khalil. " Synthesis of Some New 2- $\alpha$ - - Arabinopyranosyl-1,2,4-triazines as Potential Antitumor Chemotherapeutics ", Nucleosides, Nucleotides and Nucleic Acids, 2003 7 words — < 1%  
Crossref

61 Jonathan H. Marriott, G. Wynne Aherne, Anthea Hardcastle, Michael Jarman. "SYNTHESIS OF CERTAIN 2'-DEOXYURIDINE DERIVATIVES CONTAINING SUBSTITUTED PHENOXY GROUPS ATTACHED TO C-5'; EVALUATION AS POTENTIAL dUTP ANALOGUES", Nucleosides, Nucleotides and Nucleic Acids, 2001 7 words — < 1%  
Crossref

62 D. P. Chakraborty, K. Krohn, P. Messner, S. Roy, C. Schäffer. "Fortschritte der Chemie organischer Naturstoffe / Progress in the Chemistry of Organic Natural Products", Springer Science and Business Media LLC, 2003 7 words — < 1%  
Crossref

63 Gong, Ting, Dong-Xiao Wang, Yan Yang, Ping Liu, Ruo-Yun Chen, and De-Quan Yu. "A Novel 3-Arylcoumarin and Three New 2-Arylbenzofurans from Mucuna birdwoodiana", CHEMICAL & PHARMACEUTICAL BULLETIN, 2010. 7 words — < 1%  
Crossref

64 Seagufta Afrin, Md. Hossain Sohrab, Md. Shamim Ahmed, Choudhury Mahmood Hasan, Monira Ahsan. "Terpenoids and steroids from the stem bark of Sesbania grandiflora and biological studies of the plant extracts", Pharmacy & Pharmacology International Journal, 2019 6 words — < 1%  
Crossref

65 Gomotsang Bojase, Runner R.T. Majinda, Berhanu A. Gashe, Cornelius C.W. Wanjala. "Antimicrobial Flavonoids from Bolusanthus speciosus", Planta Medica, 2002 6 words — < 1%  
Crossref



---

EXCLUDE QUOTES ON

EXCLUDE MATCHES OFF

EXCLUDE  
BIBLIOGRAPHY ON