

Short Communication: In vitro antimicrobial and antimalarial screening of a crude extract of *Streptomyces* sp. AB8 isolated from Lapindo Mud Volcano Area, Sidoarjo, Indonesia

ACHMAD ARIFIYANTO^{1,✉}, ENDAH SETYANINGRUM¹, NISMAH NUKMAL¹, TITIK NUR AENY²

¹Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Lampung, Jl. Prof. Soemantri Brojonegoro No 1, Gedong Meneng, Rajabasa, Bandar Lampung 35144, Lampung, Indonesia, Tel.: +62-721 704625 Fax. +62-721 704625, ✉email: achmad.arifiyanto@fmipa.unila.ac.id

²Department of Plant Protection, Faculty of Agriculture, Universitas Lampung, Jl. Prof. Soemantri Brojonegoro No 1, Gedong Meneng, Rajabasa, Bandar Lampung 35144, Lampung, Indonesia

Manuscript received: 5 April 2021. Revision accepted: xxx June 2021.

Abstract. Arifiyanto A, Setyaningrum E, Nukmal N, Aeny TN. 2021. Short Communication: In vitro antimicrobial and antimalarial screening of a crude extract of *Streptomyces* sp. AB8 isolated from Lapindo Mud Volcano Area, Sidoarjo, Indonesia. *Biodiversitas* 22: 2817-2823. *Streptomyces* is a potential bacterial genus that has been investigated extensively as a source of natural microbial compounds. Its potential metabolites have been widely developed for pharmaceutical, pathogen control, and other applications in agriculture. This study aimed to determine the ability of the *Streptomyces* sp AB8 crude extract in inhibiting *Plasmodium* and pathogenic microbes. *Streptomyces* was cultured on Gause synthetic media for 10 days. The fermented broth culture media has dissolved in a 1:1 mixture of ethyl acetate and methanol. Biochemical characterization of this isolate has carried out using the standard methods. In-vitro antimalarial activity assay was performed using a chloroquine-sensitive *Plasmodium falciparum* strain 3D7. Fresh type O-positive human erythrocytes were suspended at 4 percent hematocrit in a complete medium to maintain culture. The inhibitory concentration (IC₅₀) was determined using probit analysis. The results showed the extract of *Streptomyces* sp. AB8 contains phenolic and alkaloids. *Streptomyces* sp. AB8 extract can inhibit *Dickeya zae* N-Unila 5, *Dickeya zae* N-Unila 10, *Aspergillus* sp IK3, and *Escherichia coli* growth. The results also showed that the IC₅₀ value of extract against *P. falciparum* 3D7 was 17.56 µg/mL. Further research was needed to determine the types of purified bioactive compounds and their bioactivity.

Keywords: Alkaloids, antimalarial, antimicrobial, phenolic, *Streptomyces*

INTRODUCTION

Malaria is a disease caused by the *Plasmodium* parasite, is one of the deadliest diseases in tropical countries (Hay et al. 2004). It was transmitted by the bite of a female *Anopheles* sp. infected with *Plasmodium*. Malaria in humans was caused by five different *Plasmodium* species, i.e., *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium malariae*, *Plasmodium ovale*, and *Plasmodium knowlesi* (Singh et al. 2013).

Unfortunately, the *Plasmodium* sp. parasite was resistant to antimalarial drugs such as chloroquine, amodiaquine, mefloquine, and artemisinin, according to many studies (Cui et al. 2015). Malaria research was actively being conducted to combat parasite resistance. The analysis of metabolites from different plants has been performed (Alkandahri et al. 2019; Fatmawaty et al. 2017; Okokon et al. 2017; Orabuezea et al. 2020; Zeleke et al. 2017), as well as metabolites from microorganisms to combat multi-drug resistant *Plasmodium*. Metabolites derived from microbes were considered to be more desirable due to their shorter life cycles.

Streptomyces, a Gram-positive bacteria, is a member of the actinobacteria phylum which is known to have antimicrobial, anticancer (El-Naggar and El-Ewasy 2017),

anti-inflammatory (Gomathi and Gothandam 2019), and antioxidant (Li et al. 2018). Several *Streptomyces*, such as *Streptomyces hygroscopicus* subsp. *hygroscopicus* (Fitri et al. 2019; Nugraha et al. 2020), *Streptomyces* SUK10 (Zin et al. 2017), and *Streptomyces spectabilis* BCC 4785 (Isaka et al. 2002) have been studied for their antimalarial activity. *Streptomyces* sp. AB8 was obtained from the Lapindo mud volcano (Arifiyanto et al. 2020), has not been investigated for its antimalarial activity. This study aimed to conduct an initial screening of the antimicrobial and antimalarial activity of *Streptomyces* sp. AB8.

MATERIALS AND METHODS

Culture and fermentation

Streptomyces sp. strain AB8 was rejuvenated on yeast starch agar (YSA) medium, Yeast Starch Agar medium consisted of 2.0 g yeast extract (Oxoid™), 10.0 g soluble starch (Sigma-Aldrich), 15.0 g of agar powder (Sigma-Aldrich), and 1000 mL distilled water adjusted to a pH 7.3. Bacteria were cultured at room temperature for 1–5 days. The bacterial stock was cultured on a slanting agar medium. According to Arifiyanto et al. (2021) and Ezeonu

RESULTS AND DISCUSSION

Antimicrobial activity of *Streptomyces* extract

Streptomyces sp. AB8 bacteria were isolated from rhizosphere soil in the Lapindo mud volcano field, Sidoarjo. Previous research showed that *Streptomyces* sp. AB8 could inhibit the growth of pathogenic bacteria (*Staphylococcus aureus* ATCC 6538P and *Escherichia coli* ATCC 25922) (Arifiyanto et al. 2020). *Dickeya zae* as a causative agent of stem and root rot in agricultural and plantation commodities (Aeny et al. 2020) was also inhibited by *Streptomyces* sp. AB8 (Table 1). The results suggested that crop damage caused by *D. zae* could be controlled by the extract of *Streptomyces* sp. AB8 to overcome economic losses. These bacteria could also inhibit the growth of *Aspergillus* IK3, a cockroach pathogen, and *Escherichia coli*, which was isolated from wastewater around the campus of the University of Lampung.

Streptomyces is a Gram-positive bacteria that belong to the order *Actinobacteriales* with branched mycelium and chain-shaped. These bacteria could grow in a variety of environments, such as rhizosphere, marine debris, weathered leaf litter, plant endophytes, and insect symbionts (Mathew et al. 2020). The initial screening revealed that *Streptomyces* sp. AB8 could utilize glucose, fructose, cellulose, and lactose, galactose, sucrose, chitin, mannan, and amyllum (Table 2) as its carbon source. As a decomposer group of microbes, *Streptomyces* has the ability to degrade complex molecules such as cellulose and mannan (Sasongko et al. 2015). Chitin is the main component of fungal cell walls. *Streptomyces* ability to decompose chitin potentially used to control mold. (Okay et al. 2013).

Biochemical characterization of *Streptomyces* sp. AB8

The development of new drug candidates has been driven by the resistance of pathogenic microbes to antibiotics and the risk to human health they pose. Microbial metabolites are increasingly being used to treat harmful bacteria. *Streptomyces*, a Gram-positive bacteria, have been shown to restrict the growth of various harmful bacteria. This bacterial community was thought to be in the middle of the bacteria-mold spectrum producing superior secondary metabolites such as triterpenoids, flavonoids, and alkaloids, and antibiotics. Gram-negative bacteria and pathogenic fungi were a challenge for humans health (Al-Ansari et al. 2019). Some of them were responsible for infection in humans. Pathogenic bacteria such as *Enterobacter aerogenes* and *Proteus mirabilis* were confirmed to be inhibited by *Streptomyces radiopugnans*, *Streptomyces atacamensis*, *Streptomyces fenghuangensis*, *Streptomyces verrucosiporus*, and *Streptomyces mangrove* isolates (Al-Ansari et al. 2019). In humans, *Aspergillus* genera cause diseases such as localized infections, deadly illnesses, allergic responses, and inhaled conidia (Mousavi et al. 2016). Production of lyase, chitinase, protease, and

cellulase enzymes by *Streptomyces* is suspected of supporting antifungal activity. These enzymes can damage the cell wall of the mold, which was rich in components that were easily degraded by these enzymes. Therefore, *Streptomyces* spp could be used as antifungals (de Lima Procópio et al. 2012).

There are several pathways of antimicrobial mechanisms of *Streptomyces* compounds. In general, antimicrobial mechanisms are the interactions of pathogenic microorganisms' biochemical, genetic, and cellular structures. (Chevrette et al. 2019). Metabolites of *Streptomyces* spp may cause the inhibition of DNA division. Disorders of RNA synthesis affect the ability of pathogens to produce proteins that play an important role in the cellular structure of membranes, functional enzymes, and the cell walls of pathogenic microbes. They were also inhibited pathogenic metabolism and structural breakdown (de Lima Procópio et al. 2012).

Alkaloids and phenols were detected in synthetic Gause broth after the fermentation process (Table 2). Alkaloids are present in a variety of plants and animals, including microbes. Alkaloids extracted from biological compounds had long been used to treat a range of symptoms (Zhou et al. 2013); such as cancer, painkillers, high blood pressure, nervous system damage, Parkinson's disease, and antimalarials were among them. Meanwhile, indole alkaloids isolated from *Streptomyces* sp. CT37 has been demonstrated to inhibit *Candida albicans* ATCC 10231 at low concentrations (Fang et al. 2020).

Table 1. The diameter of the growth inhibition by *Streptomyces* sp. AB8

Microbes	Colony diameter (\bar{x} mm)	Clear zone (\bar{x} mm)
<i>Escherichia coli</i>	0.56	0.40
<i>Dickeya zae</i> N-Unila 5	1.60	0.70
<i>Dickeya zae</i> N-Unila 10	1.50	0.50
<i>Aspergillus</i> sp IK3	0.20	0.10

Table 2. Biochemical characters of *Streptomyces* sp. AB8

Test	Result	Test	Result
Glucose	+	Phenol	+
Fructose	+	Alkaloids	+
Cellulose	+	Flavonoids	-
Lactose	+	Saponin	-
Galactose	+	Triterpenoids	-
Sucrose	+	Anthraquinone glycosides	-
Chitinase	+	Tannin	-
Mannanase	+	Catalase	+
Lipase	-	Indole	+
Amylase	+	Protease	+

Note: + : positive reaction; - : negative reaction




Figure 2. The effect of *Streptomyces* extracts on growth (A), growth inhibition (B), the ratio of growth to inhibition (C), the IC_{50} value of extract against *Plasmodium* parasites (D)

Alkaloids, which have a wide structural variation, are distinguished by the presence of a specific nitrogen atom. The majority of alkaloids have only one nitrogen atom, but others had up to five. This nitrogen was discovered in the form of a primary amine (RNH_2), secondary amine (R_2NH), or tertiary amine (R_3NH) (R_3NH) (Cushnie et al. 2014). Many alkaloids contain oxygen in addition to carbon, hydrogen, and nitrogen (Cushnie et al. 2014). Figure 3 showed that there was H bonded NH at an absorbance range of $3070-3350\text{ cm}^{-1}$. This was presumed as a sign of the presence of alkaloids. The acetate and benzoic groups occurred at 1073.23 cm^{-1} and above, followed by double bond $-CH=CH-$ (cis) at $650-750\text{ cm}^{-1}$.

Phenols also present in the liquid media after the fermentation process (Table 2). It was confirmed by the emergence of spectroscopy bands around 990 until 1060 cm^{-1} . The secondary cyclic alcohols are found at these spectra. Phenols donate hydrogen to react with reactive nitrogen and oxygen species as an antioxidant (Pereira et al. 2009). Phenols also have a variety of biochemical and

pharmacological properties such as antiviral, anticancer, antimalarial, and anti-inflammatory activities.

Antimalarial activity

Streptomyces extract at a concentration of $100\text{ }\mu\text{g/mL}$ was able to inhibit parasite growth up to 70.92%, with an average parasite growth rate of 2.15%. Parasitic growth was suppressed at a rate of 4.31% with an inhibitory percentage of 41.84% at a concentration of $10\text{ }\mu\text{g/mL}$. Extract concentration of $1\text{ }\mu\text{g/mL}$ resulted in parasite growth of 5.96% with an inhibition percentage of 19.23%. While a concentration of $0.1\text{ }\mu\text{g/mL}$ resulted in inhibition of 6.55% with parasite growth of 6.93 and a concentration of $0.01\text{ }\mu\text{g/mL}$ resulted in inhibition of 1% with growth of 7.33%. Overall, the treatment resulted in lower growth compared to the control. The increasing extract concentration was linear with inhibition, and vice versa, parasite growth increased with decreasing concentration (Figure 2). The inhibitory concentration to suppress a half parasites population (IC_{50}) was achieved at $17.56\text{ }\mu\text{g/mL}$.




Figure 3. FT-IR spectra of crude extract produced by *Streptomyces sp AB8*

Several past studies have reported on the mechanism of alkaloids in preventing the growth of plasmodial and microbiological infections. Alkaloids inhibit nucleic acid synthesis by suppressing the enzyme dihydrofolate reductase in cell-free (Raimondi et al. 2019). It can prevent cell division by inhibiting the Z-ring formation in bacteria (Keffer et al. 2013). Another study by Godlewska-Żyłkiewicz et al. (2020) showed that alkaloids are sufficient to initiate bacterial homeostasis and disrupt the stability of the outer membrane and cytoplasmic (Cushnie et al. 2014). Alkaloids were also able to restrain fimbria-dependent biofilm formation in *E. coli* (Notarte et al. 2019).

Streptomyces cellulosa strain TES17 produced phenolic compounds with strong antioxidant properties and effective in damaging lung cancer cell lines (Rani et al. 2018). The antibacterial and antimalarial mechanism of phenolic compounds possibly due to the ability to inhibit respiratory electron transport systems, reduced outer membrane containing lipopolysaccharides, disrupts the structure of the cell wall, and the synthesis of cell wall becomes inefficient, affect the permeability, and disrupt the cell division protein FtsZ-ring (Aldulaimi et al. 2019).

The findings on *Streptomyces* extract's antimalarial activity were used as a starting point. Further study to determine its efficacy using an *ex vivo* or *in vitro* approach is needed (Sinha et al. 2017). Peter's Test or 4-day suppression test, histopathological observation, and cytotoxicity for non-cell-targeted are required (Kifle et al. 2020). A metabolite is categorized as antiplasmodial if this metabolite can suppress parasite growth by more than 30% compared to negative controls after completing a set of *in vivo* tests (Castro et al. 1996). In conclusion, the qualitative test approach was supported by infrared spectroscopic data

indicating the presence of phenolic compounds and alkaloids from the bacterial metabolite *Streptomyces AB8*. The crude extract of *Streptomyces sp AB8* can inhibit *Plasmodium*, with an IC₅₀ value of 17.56 µg/mL. The extract of *Streptomyces sp. AB8* could inhibit the growth of *Dickeya zae* N-Unila 5, *Dickeya zae* N-Unila 10, *Aspergillus sp IK3*, and *Escherichia coli*. These results on antimalarial and antimicrobial are preliminary results.

ACKNOWLEDGEMENTS

We would like to thank everyone who contributes to this research. This work was partly supported by grants from Universitas Lampung, Indonesia (1661/UN26.21/PN/2021). The authors have declared that there was no conflict of interests.

REFERENCES

- Aeny TN, Suharjo R, Ginting C, Hapsoro D, Niswati A. 2020. Characterization and host range assessment of *Dickeya zae* associated with pineapple soft rot disease in east Lampung, Indonesia. *Biodiversitas* 21 (2): 587-595. DOI: 10.13057/biodiv/d210221.
- Agustina D, Surtiningsih T, Manuhara YS, Arifiyanto A, Malewa M. 2019. Study of Cellulase Activity Produced by *Penicillium sp.*, *Aspergillus niger* and *Trichoderma viride* on *Imperata cylindrica* (L.) Beauv. *Enrichment Media*. IOP Conf Ser Earth Environ Sci 253 (1): 012016. DOI: 10.1088/1755-1315/253/1/012016.
- Al-Ansari M, Alkubaisi N, Vijayaragavan P, Murugan K. 2019. Antimicrobial potential of *Streptomyces sp.* to the Gram-positive and Gram-negative pathogens. *J Infect Public Health* 12 (6): 861-866. DOI: 10.1016/j.jiph.2019.05.016
- Aldulaimi O, Drijfhout F, Uche FI, Horrocks P, Li W-W. 2019. Discovery, synthesis and antibacterial evaluation of phenolic

- compounds from *Cylicodiscus gabunensis*. BMC Compl Altern Med 19 (1): 183. DOI: 10.1186/s12906-019-2589-2.
- Alkandahri M, Berbudi AN, Vicahyani Utami AS. 2019. Antimalarial activity of extract and fractions of *Castanopsis costata* (Blume) A.DC. Avicenna J Phytomed 9 (5): 474-481.
- Arifiyanto A, Afriani H, Putri MH, Damayanti B, Riyanto CLR. 2021. The biological prospective of red-pigmented bacteria cultured from contaminated agar media. Biodiversitas J Biol Divers 22 (3): 1152-1159. DOI: 10.13057/biodiv/d220310
- Arifiyanto A, Surtiningsih T, Ni'matuzahroh, Fatimah, Agustina D, Alami N. 2020. Antimicrobial activity of biosurfactants produced by actinomycetes isolated from rhizosphere of Sidoarjo mud region. Biocatalysis Agric Biotechnol 24: 101513. DOI: 10.1016/j.bcab.2020.101513.
- Castro O, Barrios M, Chinchilla M, Guerrero O. 1996. Chemical and biological evaluation of the effect of plant extracts against *Plasmodium berghei*. Revista Biol Trop 44 (2A): 361-367.
- Chevrette MG, Carlson CM, Ortega HE, Thomas C, Ananiev GE, Barns KJ, Book AJ, Cagnazzo J, Carlos C, Flanigan W, Grubbs KJ. 2019. The antimicrobial potential of *Streptomyces* from insect microbiomes. Nat Commun 10 (1): 516. DOI: 10.1038/s41467-019-08438-0.
- Cui L, Mharakurwa S, Ndiaye D, Rathod PK, Rosenthal PJ. 2015. Antimalarial drug resistance: Literature review and activities and findings of the ICEMR network. Am J Trop Med Hyg 93 (Suppl 3): 57-68. DOI: 10.4269/ajtmh.15-0007
- Cushnie TPT, Cushnie B, Lamb AJ. 2014. Alkaloids: an overview of their antibacterial, antibiotic-enhancing and antivirulence activities. Int J Antimicrob Agents 44 (5): 377-386. DOI: 10.1016/j.ijantimicag.2014.06.001
- Darmawan TS, Zahroh TT, Merindasya M, Masfaridah R, Hartanti DA, Arum S, Nurhatika S, Muhibuddin A, Surtiningsih T, Arifiyanto A. 2017. Manganese (Mn) stress toward hyperaccumulators plants combination (HPC) using *Jatropha curcas* and lamtoro gung (*L. leucocephala*) in mycorrhizal addition on soybean (*Glycine max*) seedling stage. AIP Conf Proc 1854 (1): 020027. DOI: 10.1063/1.4985418
- de Lima Procópio RE, da Silva IR, Martins MK, de Azevedo JL, de Araújo JM. 2012. Antibiotics produced by *Streptomyces*. Braz J Infect Dis 16 (5): 466-471. DOI: 10.1016/j.bjid.2012.08.014
- El-Naggar NEA, El-Ewasy SM. 2017. Bioproduction, characterization, anticancer and antioxidant activities of extracellular melanin pigment produced by newly isolated microbial cell factories *Streptomyces glaucescens* NEAE-H. Sci Rep 7: 42129. DOI: 10.1038/srep42129.
- Ezeonu CS, Ejikeme CM. 2016. Qualitative and quantitative determination of phytochemical contents of indigenous Nigerian Softwoods. New J Sci 2016: 1-9. DOI: 10.1155/2016/5601327.
- Fang Q, Maglangit F, Mugat M, Urwald C, Kyeremeh K, Deng H. 2020. Targeted isolation of indole alkaloids from *Streptomyces* sp. CT37. Molecules 25 (5): 1108. DOI: 10.3390/molecules25051108.
- Fatmawaty, Rosmalena, Amalia A, Syafitri I, Prasasty VD. 2017. Antimalarial effect of flamboyant (*Delonix regia*) bark and papaya (*Carica papaya* L) leaf ethanolic extracts against *Plasmodium berghei* in mice. Biomed Pharmacol J 10 (3): 1081-1089. DOI: 10.13005/bpj/1206.
- Fitri LE, Alkarimah A, Cahyono AW, Lady WN, Endharti AT, Nugraha RY. 2019. Effect of metabolite extract of *Streptomyces hygroscopicus* subsp. *hygroscopicus* on *Plasmodium falciparum* 3D7 in Vitro. Iran J Parasitol 14 (3): 444-452.
- Godlewska-Żyłkiewicz B, Świsłocka R, Kalinowska M, Golonko A, Świdorski G, Arciszewska Ż, Nalewajko-Sieliwoniuk E, Naumowicz M, Lewandowski W. 2020. Biologically active compounds of plants: Structure-related antioxidant, microbiological and cytotoxic activity of selected carboxylic acids. Materials 13 (9): 4454. DOI: 10.3390/ma13194454.
- Gomathi A, Gothandam KM. 2019. Investigation of anti-inflammatory and toxicity effects of mangrove-derived *Streptomyces rochei* strain VITGAP173. J Cell Biochem 120 (10): 17080-17097. DOI: 10.1002/jcb.28969
- Hay SI, Guerra CA, Tatem AJ, Noor AM, SR. 2004. The global distribution and population at risk of malaria: past, present, and future. Lancet Infect Dis 4 (6): 327-336. DOI: 10.1016/S1473-3099(04)01043-6
- Isaka M, Jaturapat A, Kramyu J, Tanticharoen M, Thebtaranonth Y. 2002. Potent in vitro antimalarial activity of meta cyclo prodigiosin isolated from *Streptomyces spectabilis* BCC 4785. Antimicrob Agents Chemother 46 (4): 1112-1113. DOI: 10.1128/AAC.46.4.1112-1113.2002.
- Keffer JL, Huecas S, Hammill JT, Wipf P, Andreu JM, Bewley CA. 2013. Chrysophaentins are competitive inhibitors of FtsZ and inhibit Z-ring formation in live bacteria. Bioorg Med Chem 21 (18): 5673-5678. DOI: 10.1016/j.bmc.2013.07.033.
- Kifle ZD, Adinew GM, Mengistie MG, Gurmu AE, Enyew EF, Goshu BT, Amare GG. 2020. Evaluation of antimalarial activity of methanolic root extract of *Myrica salicifolia* a rich (Myricaceae) against *Plasmodium berghei*-infected mice. J Evid-Based Integr Med 25: 1-12. DOI: 10.1177/2515690X20920539.
- Li C, Ji C, Tang B. 2018. Purification, characterization and biological activity of melanin from *Streptomyces* sp. FEMS Microbiol Lett 365 (19): 1-8. DOI: 10.1093/femsle/fny077.
- Lin YB, Wang XY, Fang H, Ma YN, Tang J, Tang M, Wei GH. 2012. *Streptomyces shaanxiensis* sp. nov., a blue pigment-producing streptomycete from sewage irrigation soil. Intl J Syst Evol Microbiol 62 (8): 1725-1730. DOI: 10.1099/ijs.0.029959-0.
- Mathew BT, Torky Y, Amin A, Mourad AH, Ayyash MM, El-Keblawy A, Hilal-Alnaqbi A, AbuQamar SF, El-Tarabily KA. 2020. Halotolerant marine rhizosphere-compentent actinobacteria promote *Salicornia bigelovii* growth and seed production using seawater irrigation. Front Microbiol 11. DOI: 10.3389/fmicb.2020.00552.
- Mousavi B, Hedayati MT, Hedayati N, Ilkit M, Syedmousavi S. 2016. *Aspergillus* species in indoor environments and their possible occupational and public health hazards. Curr Med Mycol 2 (1): 36-42. DOI: 10.18869/acadpub.cmm.2.1.36.
- Notarte KI, Devanadera MK, Mayor AB, Cada MC, Pecundo MH, Macabeo AP. 2019. Toxicity, antibacterial, and antioxidant activities of fungal endophytes *Colletotrichum* and *Nigrospora* spp. Isolated from *Uvaria grandiflora*. Philippine J Sci 148: 505-512.
- Nugraha RY, Faratisha IF, Mardhiyyah K, Ariel DG, Putri FF, Winarsih S, Sardjono TW, Fitri LE. 2020. Antimalarial properties of isoquinoline derivative from *Streptomyces hygroscopicus* subsp. *hygroscopicus*: an in silico approach. BioMed Res Intl. DOI: 10.1155/2020/6135696.
- Okay S, Özdal M, Kurbanoglu EB. 2013. Characterization, antifungal activity, and cell immobilization of a chitinase from *Serratia marcescens* MO-1. Turk J Biol 37 (6): 639-644. DOI: 10.3906/biy-1208-45.
- Okokon J, Antia B, Mohanakrishnan D, Sahal D. 2017. Antimalarial and antiparasitoid activity of husk extract and fractions of *Zea mays*. Pharm Biol 55 (1): 1394-1400. DOI: 10.1080/13880209.2017.1302966.
- Orabueza CI, Ota DA, Coker HA. 2020. Antimalarial potentials of *Stemonocoleus micranthus* Harms (Leguminosae) stem bark in *Plasmodium berghei* infected mice. J Tradit Compl Med 10 (1): 70-78. DOI: 10.1016/j.jtcm.2019.03.001.
- Pereira DM, Valentão P, Pereira JA, Andrade PB. 2009. Phenolics: From chemistry to biology. Molecules 14 (6): 2202-2211. DOI: 10.3390/molecules14062202.
- Putri MH, Handayani K, Setiawan WA, Damayanti B, Ratih CL, Arifiyanto A. 2021. Screening of extracellular enzymes on *Serratia marcescens* strain MBC1. Jurnal Riset Biologi dan Aplikasinya 3 (1): 23-29. [Indonesian]
- Quintino L, Namislo A, Davidsson M, Breger LS, Kavanagh P, Avallone M, Elgstrand-Wettergren E, Isaksson C, Lundberg C. 2019. DHFR inhibitors: Reading the past for discovering novel anticancer agents. Molecules 24 (6): 1-19. DOI: 10.3390/molecules24061140.
- Rani R, Arora S, Kaur J, Manhas RK. 2018. Phenolic compounds as antioxidants and chemopreventive drugs from *Streptomyces cellulosa* strain TES17 isolated from rhizosphere of *Camellia sinensis*. BMC Compl Altern Med 18 (1): 82. DOI: 10.1186/s12906-018-2154-4.
- Rosa E, Ekowati CN, Handayani T, Ikhsanudin A, Apriliani F, Arifiyanto, A. 2020. Characterization of entomopathogenic fungi as a natural biological control of American cockroaches (*Periplaneta americana*). Biodiversitas 21 (11): 5276-5282. DOI: 10.13057/biodiv/d211131.
- Sasongko A, Yopi Y, Rahmani N, Lisdiyanti P, Saepudin E. 2015. Enzymatic hydrolysis of mannan from Konjac (*Amorphophallus* sp.) using mannanase from *Streptomyces lipmanii* to produce Manno-oligosaccharides. Makara J Sci 19 (3). DOI: 10.7454/mss.v19i3.4850.
- Setyaningrum E, Arifiyanto A, Nukmal N, Aeny TN, Putri MH, Setiawati UN. 2021. In vitro test for inhibition of *Plasmodium falciparum* 3D7 parasites using *Streptomyces hygroscopicus* subsp. *hygroscopicus*

- Strain i18, Isolated from a Pineapple Farm in Lampung. *J Pure Appl Microbiol* 15: 891-896. DOI: 10.22207/jpam.15.2.45.
- Singh BDC. 2013. Human infections and detection of *Plasmodium knowlesi*. *Clin Microbiol Rev* 26 (2): 165-84. DOI: 10.1128/CMR.00079-12.
- Sinha S, Sarma P, Sehgal R, Medhi B. 2017. Development in assay methods for in vitro antimalarial drug efficacy testing: A Systematic Review. *Front Pharmacol* 8: 754. DOI: 10.3389/fphar.2017.00754.
- Sumardi, Farisi S, Ekowati CN, Arifiyanto A, Rahmawati DE. 2020. Halotolerant *Bacillus* sp. For mannan degradation isolated from mangrove ecosystem at Hanura beach Lampung. *J Pure Appl Microbiol* 14 (2): 1237-1244. DOI: 10.22207/JPAM.14.2.18.
- Zelege G, Kebebe D, Mulisa E, Gashe F. 2017. In vivo antimalarial activity of the solvent fractions of fruit rind and root of *Carica papaya* Linn (Caricaceae) against *Plasmodium berghei* in Mice. *J Parasitol Res*. DOI: 10.1155/2017/3121050.
- Zhou H, Yang Y, Zhang J, Peng T, Zhao L, Xu L, Ding Z. 2013. Alkaloids from an endophytic *Streptomyces* sp. YIM66017. *Nat Prod Commun* 8 (10): 1393-1396. DOI: 10.1177/1934578x1300801012.
- Zin NM, Baba MS, Zainal-Abidin AH, Latip J, Mazlan NW, Edrada-Ebel RA. 2017. Gancidin W, a potential low-toxicity antimalarial agent isolated from an endophytic *Streptomyces* SUK10. *Drug Design Dev Ther* 11: 351-363. DOI: 10.2147/DDDT.S121283.