

Plagiarism Checker X Originality Report



Plagiarism Quantity: 13% Duplicate

Date	Monday, June 28, 2021
Words	317 Plagiarized Words / Total 2395 Words
Sources	More than 20 Sources Identified.
Remarks	Low Plagiarism Detected - Your Document needs Optional Improvement.

209 INTERPRETATION OF CURIE POINT DEPTH AND THERMAL GRADIENT BASED ON MAGNETIC ANOMALY DATA AT SOUTHERN SUMATRA GEOTHERMAL AREA Syamsurijal RASIMENG*(1 , 2) , Wawan GUNAWAN A. KADIR (2), Hendra GRANDIS(2), Chalid IDHAM ABDULLAH (3) 1Jurusan Teknik Geofisika, FT UNILA, Jl . S. Brojonegoro No.1, Bandar Lampung 2Program Studi Teknik Geofisika, FTTM ITB, Jl . Ganesha No.10 , Bandung 3Program Studi Geologi, FITB ITB, Jl . Ganesha No.10 Bandung *e-mail : syamsurijal.rasimeng@eng.unila.ac.id Abstract Curie point depth (CPD) is Curie temperature depth where magnetic mineral lost its magneticity. Curie temperature generally higher on mafic frozen rock than on any other rock. because of its higher titano-magnetic content. The magnetic field induction value depends on rock minerals, so it can be used to determine the depth of the rock.

Magnetic field anomaly are series of signals magnitude as the respon of rock composing material under the surface. Power spectrum application for that signals magnitude can be used to determine lower surface rock mass depth including CPD and thermal gradient. Tectonic activity along Sumatra Fault System (SFS) allow the emergence of CPD depth variation and thermal gradient. Southern Sumatera as part of this system become the focus of this research, because it has geothermal potential indicated by numbers of fluid manifest distribution on the surface. Result of thermal gradient analysis obtained from CPD variation informed that value of thermal gradient was 4 0 -7 5 0C/km on shallow CPD area (8 -15 km). That area divided Sumatera island and extended to North West ♦ South East direction .

Geologically, southern Sumatera also part of magmatic arc area as the SFS prod uct, so it is possibly become geothermal prospect area. Key Words : Curie point depth, geothermal, spectrum power, thermal gradient, magnetic method 1. Introduction Curie Point Depth (CPD) defined as Curie temperature depth where magnetic mineral lost its magneticity. Curie temperature generally higher on mafic frozen rock than on any other rock, because of its higher titano-magnetic content. The presence of magnetic mineral such as magnetite (Fe 3O4), ulvospine (Fe2TiO4), hematite (Fe 2O3) and many others as the earth crust composing minerals will increase Curie temperature up to 580 oC (Blakely 1995). CPD depth directly affected by lower surface rocks temperature as consequence of tectonic activity. In consequence, the variation of CPD depth

Sources found:

Click on the highlighted sentence to see sources.

Internet Pages

- 1% <http://repository.lppm.unila.ac.id/view/>
- <1% <https://www.sciencedirect.com/science/ar>
- 5% <https://www.researchgate.net/publication>
- 1% <https://www.researchgate.net/profile/Sya>
- <1% <https://www.sciencedirect.com/science/ar>
- <1% <http://earthsci.org/education/fieldsk/ge>
- <1% https://en.wikipedia.org/wiki/Quantum_tu
- <1% <https://www.researchgate.net/publication>
- <1% <https://www.science.gov/topicpages/c/cur>
- <1% <https://www.sciencedirect.com/science/ar>
- <1% <https://link.springer.com/article/10.100>
- <1% <https://www.researchgate.net/publication>
- 1% <https://dergipark.org.tr/tr/pub/tbtkeart>
- 1% <https://link.springer.com/article/10.100>
- 1% <https://www.researchgate.net/publication>
- 1% <https://link.springer.com/article/10.100>
- 1% <https://cnecsdti.research.gov.ro/wp-cont>
- 1% <https://www.sciencedirect.com/science/ar>
- 1% <https://www.researchgate.net/profile/Alf>

highly depends on the impact caused by the tectonic activity.

CPD analysis on Southern Sumatra area with high tectonic activity as the impact of the clash between Indo-Australia and Eurasia plate is very important to be performed to get more ideas about the relation between tectonic activity and CPD. Shie and Natawidjaja (2000) divided three Sumatra areas according to their tectonic activities on Sumatra fault, they are northern Sumatra, central Sumatra and southern Sumatra. Magnetic methods based on magnetic field induction observation as the response of magnetized rocks below the surface. The value of magnetic field induction depends on the composition of the rock magnetic mineral content. This makes the presence of magnetic minerals as earth crust rocks composer can be used as an indicator of magnetic measurement.

Magnetic method ability to detect lower surface rock mass, including CPT by applying spectrum analysis has been performed by many researchers. Nevertheless, this technique still considered as interesting matter, proved from researches performed by Bansal et al. (2010), Boyksara and Bektas (2007), Dolmaz et al. (2005), El Nabi (2011), Karastathis et al. (2010), Maden (2010), Minea and Vlad (2010) and many others. This research covered magnetic anomaly data spectrum analysis to analyze CPD depth variation and thermal gradient caused by tectonic activity on southern Sumatra area. Thermal circulation that came from earth mantle (asthenosphere) and pass through fault planes on earth crust causing a thermal convection current. The thermal convection current then will cause variation of CPD distribution pattern under the surface.

Seminar Nasional Kebumihan -VII Yogyakarta, 5 September 2013 Fakultas Teknologi Mineral Universitas Pembangunan Nasional "Veteran" Yogyakarta 210 2. Methods Blakely (1995) and Maden (2010) described total magnetic field power spectrum anomaly as, $(\frac{1}{2\pi}) \int_{-\infty}^{\infty} \frac{1}{k} |M(k)|^2 dk$ (1) with, $(\frac{1}{2\pi}) \int_{-\infty}^{\infty} \frac{1}{k} |M(k)|^2 dk$ (2) where is power spectrum as caused of magnetization, C_m is a constant, and are magnetization direction factor and magnetic field direction factor, k_x and k_y are wavenumber on x and y direction. Z_t and Z_b are the upper and lower limit depth of magnetic anomaly. $|M_x|$ and $|M_y|$ are radial symmetrical form where the average radial value from factor and are constant. If magnetization on anomaly $M(x,y)$ is random and not depend on each other then power spectrum as caused of magnetization (M) become constant, and so the total magnetic field anomaly power spectrum became, $(\frac{1}{2\pi}) \int_{-\infty}^{\infty} \frac{1}{k} dk$ (3) with A is a constant.

If the wave length much smaller than twice of magnetization source anomaly thickness, then equation (3) can be described as, $(\frac{1}{2\pi}) \int_{-\infty}^{\infty} \frac{1}{k} dk$ (4) B is a constant and . Power spectrum curve slope of magnetic field anomaly total on equation (4) is upper limit depth Z_t of magnetic anomaly source (Figure 1). Figure 1. Power spectrum curve of magnetic anomaly data, (a). Determination of upper limit anomaly Z_t , (b). Determination of anomaly central depth Z_o (Maden 2010). To predict the depth of Z_o central magnetic anomaly source, equation (3) can be rewritten as, $(\frac{1}{2\pi}) \int_{-\infty}^{\infty} \frac{1}{k} dk$ (5) where C is a constant. For large wave length, equation (5) can be differentiated to, $2\pi \int_{-\infty}^{\infty} \frac{1}{k} dk$ (6) with $2d$ is magnetic anomaly source thickness. Next, by taken a logarithmic form on equation (6), obtained $\{ \frac{1}{k} \}$ (7) D is a constant.

Based on magnetic anomaly power spectrum curve slope on equation (7), central depth (centroid) of magnetic anomaly Z_o can be predicted. Meanwhile, lower limit depth of Z_b magnetic anomaly can be calculated from the relation of, . (8) Next, determination of thermal flow can be calculated based on Maden (2010) description as, (9) q is thermal flow and λ is thermal conduction coefficient If assuming that temperature gradient is

constant. Curie temperature (T_c) can be determined based on value (Z_b) and geothermal gradient such as, [] (10) Equation (10) usage were based on assumption that surface temperature was 0 oC and there was no heat releasing from the source (CPD) to surface. In consequence, equation (9) and (10) resulting a relation between CPD and thermal flow (q) in form of, [] (11) Equation (11) showed the inverse relation of CPD with thermal flow (Stampolidis et.al., 2005) 3.

Result and Discussion Sumatera fault occurred because of the clash between Indo-Australia plate and Eurasia plate along more than 1600 km, started from southern Sunda strait until northern Andaman Sea (Genrich 2000). Bukit Barisan mountains as product of Sumatera fault created geothermal prospect areas (Hermawan and Rezky 2010). The uplift along Bukit Barisan mountains on Middle Miocene resulting volcanic mountains product, which also followed with granite and diorite intrusion. On Pliocen-Plistosen period, magma activities on this area resulting basaltic, andesitic, and dalsit rocks, and also geothermal system (Mulyadi 1995).

Generally, Sumatra tectonic areas formed by Indo-Australia and Eurasia plate subduction that emerged large magmatic arc on Bukit Barisan mountains that started since Early Permian (Cameron, 1980 dalam Amin dkk., 1994) or Mid -Late Permian (Katili 1969, 1972, 1981; Gafoer 1990 dalam Amin dkk., 1994). Research area that lies on magmatic arc composed of pre-Mesozoikum metamorf, Mesozoikum and Kenozoikum frozen rocks, the crumbling of Tersier \diamond Quartener volcano, and also sediment above it. The movement along Sumatera fault varies from 45-60 mm/year on northern Sumatera to only 1 mm/year on Sunda Strait (Sieh and Natawijaya 2000).

2D power spectrum analysis applied on real data which is the compilation data of satellite magnetic, airborne, and marine magnetic (Maus et al. 2009). The next step was data extraction from that magnetic anomaly contour on southern Sumatera 212 described on Figure (4), which covered 6 0 00 \diamond - 00 00 \diamond su atide 102 0 00 \diamond - 1080 00 \diamond eas longitude. Figure 2. Southern Sumatera Dipole Magnetic Anomaly contour, contour grid 0.05 0, contour interval 25 nT, contour plane 5 km above sea level The earth magnetic field came from the earth core through magnetohydrodynamic process. That main magnetic field inducted the rocks under the surface. Some researches concluded that magnetic rock depth lied on the meeting border of crust -mantle which is CPD (Minea and Vlad 2010) .

20x 2 0 coverage area windowing then performed from southern Sumatera dipole magnetic anomaly map (Figure 4). The windowing real data consists of 81 areas. Figure 3. Window area power spectrum 2D as sample for area O14, O21, O31, X2D and X5A. 213 Based on windowing result, 2D radially average power spectrum calculated to get Z_t depth, Z_o depth and Z_b depth or CPD value. CPD depth obtained from radially average power spectrum logarithmic curve calculation using least-square fitting method that described by Dolmaz (2005). This method performed in three steps; (1). Square windowing with area overlapping for 2D power spectrum, (2). Radially average power spectrum logarithmic calculation on each area, (3). Z_b CPD depth estimation of each area according to Z_o centroid depth and Z_1 upper limit depth on each area.

By modifying equation (4), magnet anomaly Z_t upper limit depth can be calculated from 2D radially average power spectrum logarithmic curve. (12) while Z_o magnet anomaly centroid depth obtained from , () | | | | (13) f is wavenumber (km⁻¹). For Z_b CPD depth calculated using equation (8) Radially average power spectrum

was radially shifting logarithmic power spectrum 2D averaged curve . The curve slope of radially average power spectrum 2D related with magnetic anomaly source depth under the surface. These are depth calculation result of radially average power spektrum logarithmic curve from couples of window area. Figure 4. Radially average power spectrum window area O21. $Z_t=8.62\text{km}$., $Z_o=10.63\text{km}$ and $Z_b=12.64\text{km}$ Z_b depth calculation result obtained from logarithmic curve slope then plotted in form of contour map.

These Figures showed contour depth or CPR and thermal gradient value on southern Sumatera area. 214 Figure 5. Southern Sumatera CPD depth contour. Contour Grid 0.050, contour interval 0.5 km, contour plane elevation zero km on mean sea level. Figure 6. CPD depth surface on Southern Sumatera that showed CPD depth increasing on to East direction Based on the calculation of radially averaged power spectrum logarithmic curve on research areas, magnetic anomaly CPD depth divided into three depth categories, shallow (8 \diamond 15km), moderate (15 \diamond 25 km) and deep (25 \diamond 32 km). 215 Figure 7. Southern Sumatera thermal gradient anomaly surface that showed thermal gradient decreasing on to East direction. Thermal gradient anomaly surface showed similar pattern to CPD depth. The thermal gradient on research area has high value on shallow CPD area (40-75 0C/km). Tendency of thermal gradient value relatively decreasing on to east direction of research area.

Figure 8. Thermal gradient anomaly overlay and southern Sumatera geology map. Red contour line represent 40-75 0C/km thermal gradient while blue line represent less than 4 0 0C/km thermal gradient. 216 Thermal gradient anomaly overlay result of Southern Sumatera geology map showed tendency of high thermal gradient value along the magmatic arc area. 4. Conclusion Thermal gradient analysis result obtained from CPD variation informed that high thermal gradient value (4 0 -7 5 0C/km) has shallow CPD depth (8 -15 km). Geologically, the area along Sumatra island magmatic ar c , which is a product of Sumatera fault system, has a high thermal gradient anomaly. Based on th ese informations, this area considered as an potential geothermal prospect area . 5.

References Amin, T.C., Sidarto, S. Santosa, W. Gunawan, 1994, Geology of the Kotaagung quadrangle, Sumater, Geological Research and Development Centre, Bandung. Bansal, A.R., Gabriel, G., dan Dimri, V.P., 2010, Depth To The Bottom Of Magnetic Sources In Germany, EGM 2010 International Workshop, Capri-Italy. Blakely, R.J., 1995, Potential Theory in Gravity and Magnetic Applications, Cambridge University Press, USA. B \diamond y \diamond ksara \diamond A. dan Bektas \diamond \diamond ., 2007, Curie Point Depth Of Inner East Anatolia-Turkey, Geophysical Research Abstracts, Vol. 9. Dolmaz, M.N., Omer, T.U., Hisarli, Z.M., dan Orbay, N., 2005, Curie Point Depth Variations To Infer Thermal Structure Of The Crust At The African-Eurasian Convergence Zone, Sw Turkey, Earth Planets Space, 57, pp. 373 \diamond 383. El Nabi, S.M.A.,

2011, Curie point depth beneath the Barramiya \diamond Red Sea coast area estimated from aeromagnetic spectral analysis, Arab J Geosci- Saudi Society for Geosciences. Genrich, J.F., Bock, Y., McCaffrey, R., Prawirodirdjo, L., Stevens, C.W., Puntodewo, S.S.O., Subarya, C. dan Wdowski, S., 2000, Distribution of slip at the northern Sumatera fault system, journal of geophysical research, vol. 105, No. B12, pp. 28.327-28.341. Hermawan dan Rezky, 2010, The role of Sumatera fault Structure in Appearance of Geothermal Faeture at Cubadak Area, West Sumatera Indonesia, Proceedings World Geothermal Congress, Bali Indonesia. Karastathis, V.K., Papoulia, J., Di Fiore, B., Makris, J., Tsambas, Stampolidis, A., dan , A., 2010, Exploration of the Deep Structure of the Central Greece Geothermal Field by Passive Seismic and Curie Depth Analysis,

72nd EAGE Conference & Exhibition incorporating SPE EUROPEC 2010 Barcelona, Spain.

Katili, J. and F. Hehuwat, 1967, On the occurrence of Large Transcurrent Faults in Sumatra, Indonesia, J. Geosci. Osaka City Univ., 10, 5 [◆](#) 17. Maden, N., 2010, Curie-point Depth from Spectral Analysis of Magnetic Data in Erciyes Stratovolcano Central TURKEY, Pure Appl. Geophys. 167, pp. 349 [◆](#) 358. Maus, S., dkk., 2009, EMAG2: A 2 [◆](#) arc min resolution Earth Magnetic Anomaly Grid compiled from satellite, airborne, and marine magnetic measurements, Geochem. Geophys. Geosyst., 10, Q08005, doi:10.1029/2009GC002471. Minea, M. dan Minea, V.C., 2010, Curie Point depth estimates and correlation with flat-slab subduction in Mexico, Geophysical Research Abstracts, Vol. 13.

217 Mulyadi, 1995, Interpretation of Goelectric structure at Hululais Prospect Area, South Sumatera, Proceedings 17th NZ Geothermal Workshop. Sieh, K. dan Natawijaya, 2000, Neotectonics of the Sumatera fault Indonesia, Journal of Geophysical Research, Vol. 105, No. B12, pp. 28.295-28.326. Stampolidis, A., Kane, I., Tsokas, G.N., dan Tsourlo, P. (2005), Curie Point Depths Of Albania Inferred From Ground Total Field Magnetic Data, Surv Geophys. 26, 461 [◆](#) 480. [View publication stats](#) [View publication stats](#)