GLOBAL JOURNAL of **PURE and APPLIED** MATHEMATICS

Editor-in-chief: Taekyun Kim

Managing Editor: E.-B. Lin



RIP Research India Publications http://www.ripublication.com

Global Journal of Pure and Applied Mathematics

SJR

Scimago Journal & Country Rank Enter Journal Title, ISSN or Publisher Name

Home

Journal Rankings

Country Rankings

s Viz Tools

About Us

Help

Global Journal of Pure and Applied Mathematics

Country	India	5
Subject Area and Category	Mathematics Applied Mathematics	J
	Mathematics (miscellaneous)	H Index
Publisher	Research India Publications	TT INGOX
Publication type	Journals	
ISSN	09739750, 09731768	
Coverage	2010-ongoing	

+Quartiles **Applied Mathematics** Mathematics (miscellaneous) 2011 2012 2013 2014 2015 SJR Citations per document ++0.32 0.24 0.16 0.08 2011 2015 2012 2013 2014 **Total Cites** Self-Cites +

Global Journal of Pure and Applied Mathematics



0

2011

2012

2013

2014

2015



0

2011

2012

2013

← Show this widget in your own website

2014

2015

Just copy the code below and paste within your html code:

<a href="http://www.scimage



Follow us on Twitter

Scimago Lab, Copyright 2007-2016. Data Source: Scopus®

Global Journal of Pure and Applied Mathematics



Global Journal of Pure and Applied Mathematics (GJPAM) Print ISSN : 0973-1768 Online ISSN: 0973-9750

Editor-in-Chief:

Prof. Taekyun Kim, Division of General Education-Mathematics, Kwangwoon University, Seoul 139-704, Republic of South Korea.

Aims and Scope: The *Global Journal of Pure and Applied Mathematics (GJPAM)* is an international journal of high quality devoted to the publication of original research papers from pure and applied mathematics with some emphasis on all areas and subareas of mathematical analysis and their broad range of applications. Areas and subareas of interest include (but are not limited to) approximation theory; statistics; probability; fluid mechanics; Fuzzy mathematics; transport



Research India Publication

theory; mathematical biology, including population dynamics; wave propagation; special functions; algebra and applications; combinatorics; coding theory; fractional analysis; solid mechanics; variational methods; financial mathematics; asymptotic methods; graph theory; fractals; moment theory; scattering theory; number theory and applications; geometry of Banach spaces; topology and applications; complex analysis; stochastic process; bifurcation theory; differential equations; difference equations; dynamical systems; functional differential equations; numerical analysis; partial differential equations; integral equations; operator theory; Fourier analysis; applied harmonic analysis; optimization; wavelets; signal theory; ultrametric analysis; optimal control; fixed-point theory and applications; reaction-diffusion problems, discrete mathematics; automata theory and more...

Submission: Authors are requested to submit their papers electronically to <u>submit@ripublication.com</u> with mention journal title (**GJPAM**) in subject line.

Indexing and Abstracting: The **GJPAM** is abstracted and indexed in the Mathematical Reviews, MathSciNet, Zentralblatt MATH and EBSCO Databases, **ICI, Index Copernicus, This Journal is in UGC Approved Journal List.**

Frequency: Twelve issues per year.

Annual Subscription Price for year 2015: Library/ Institutional: Print : US\$780.00 Online Only: US\$760.00 Print + Online : US\$820.00 Individual/ Personnel: Print US\$390.00 Inside India: Rs.3000.00

Editorial Board

H <u>Publication Ethics and Publication Malpractice Statement</u>

Instruction for Authors

- Ħ. **Special Issue #** <u>Vol.2 No.1-2</u> <u>No.3</u> (2006) **H** <u>Vol.3 No.1 No.2</u> <u>No.3</u> (2007) **H** <u>Vol.4 No.1 No.2 No.3</u> (2008) **H** <u>Vol.5 No.1 No.2</u> <u>No.3</u> (2009) **Vol.6** No.1 No.2 No.3 (2010) **H** <u>Vol.7 No.1 No.2 No.3</u> (2011) **Vol.8** No.1 No.2 No.3 No.4 No.5 (2012) **Vol.9** No.1 No.2 No.3 No.4 No.5 No.6 (2013) **Vol.10 No.1 No.2** <u>No.3 No.4</u> <u>No.5</u> <u>No.6</u> (2014) **Vol.11 No.1 No.2 No.3 No.4 No.5 No.6** (2015) **Vol.12 No.1 No.2 No.3 No.4 No.5 No.6** (2016) **Vol.13 No.1 No.2 No.3 No.4 No.5 No.6 No.7 (2017) ±** <u>Subscription</u> **<u>Call for Papers</u> d** <u>Other Journals</u>
- © All rights are reserved with **Research India Publications** <u>Journals</u> <u>Contact</u>

<u>Home</u>

Global Journal of Pure and Applied Mathematics (GJPAM)

Volume 13 Number 3 (2017)

Contents

Weyl's Theorem for quasi N-class Ak operators pp. 875–884 D. Senthilkumar and R. Murugan

<u>Common fixed points for F-dominated mappings</u> pp. 885–892 Rakesh Batra

The λ-analogue degenerate Changhee polynomials and numbers pp. 893–900 Jin-Woo Park, Gwan-Woo Jang and Jongkyum Kwon

Approximation of fixed points in 2-Hilbert spaces pp. 901–910 N. Faried, H. A. El-Sharkawy and A. Nasr

Some identities of symmetry for the modified degenerate Carlitz q-Bernoulli polynomials under symmetry group of degree n pp. 911–920 Joohee Jeong, Dong-Jin Kang, Hong Kyung Pak and Seog-Hoon Rim

Snowflake Logarithmic Metrics for Face Recognition in the Zernike Domain pp. 921–939 Ahmed J. Ali, Zahir M. Hussain and Mohammed S. Mechee

<u>The Moment Generating Function of the Four-Parameter Generalized F</u> <u>Distribution and Related Generalized Distributions</u> pp. 941-956 Warsono, Dian Kurniasari, Widiarti, Mustofa Usman and Faiz A. M. Elfaki

<u>Fractional Calculus Associated with General Polynomials and Aleph Functions</u> pp. 957-966 Parul Gupta

<u>Local Stability of Prey-Predator with Holling type IV Functional Response</u> pp. 967-980 Rehab Noori Shalan

<u>On Inequality for the Non-Local Fractional Differential Equation</u> pp. 981–993 Tarachand L. Holambe, Mohammed Mazhar-Ul-Haque

Banach contraction principle on Cone Heptagonal Metric Space pp. 995–1008 Abba Auwalu and Ali Denker

Fuzzy Logic Relation Based Stock Market Forecasting Model pp. 1009-1018 G.P.Pandey and Sanjay Sharma

Even Vertex Odd Mean Labeling of Some Graphs pp. 1019-1033 M.Kannan, R.Vikrama Prasad and R. Gopi <u>Coincidences and Common Fixed-Point Theorems in Intuitionistic Fuzzy Metric</u> <u>Space</u> pp. 1035-1047 Piyush Kumar Tripathi and Sajjan Lal Maurya

<u>Water Evaporation algorithm to solve combined Economic and Emission</u> <u>dispatch problems</u> pp. 1049-1067 Venkadesh Rajarathinam and Anandhakumar Radhakrishnan

Bayesian Zero-Inflated Generalized Poisson (t) Spatio-Temporal Modeling for Analyzing the DHF Endemic Area

pp. 1069-1081 Mukhsar, Bahriddin Apabihi, Sitti Wirdhana Ahmad Bakkareng, Asrul Sani and Edi Cahyono

Square Difference Prime Labeling for Some Snake Graphs pp. 1083-1089 Sunoj B.S and Mathew Varkey T.K

The Classification of Quadratic Rook Polynomials of a Generalized Three Dimensional Board pp. 1091-1101 R. Sangeetha and G. Jayalalitha

Syntactic Semihypergroup pp. 1103–1115 Goutam Chowdhury

An Algorithm for Minimal and Minimum Distance - 2 Dominating Sets of Graph pp. 1117-1126 K. Ameenal Bibi, A. Lakshmi and R. Jothilakshmi

Exact solutions of Schwarz KdV and the (2+1)-dimensional Schwarz-KdV equations via the improved tanh-coth method pp. 1127–1133 Cesar A. Gomez S, Juan C. Hernández R. and Hernan Garzón

[UP] © All rights are reserved with Research India Publications Journals Contact

<u>Home</u>

The Moment Generating Function of the Four-Parameter Generalized F Distribution and Related Generalized Distributions

Warsono¹, Dian Kurniasari¹, Widiarti¹, Mustofa Usman¹ and Faiz A. M. Elfaki²

¹Department of Mathematics, University of Lampung, Indonesia. ²Department of Mathematics, Statistics and Physics, College of Arts and Sciences, Qatar University, P. O. Box 2713 Doha, Qatar.

Abstract

The main objective of this article is to show the derivation of the moment generating function of the four-parameter of generalized (G4F) F distribution. Through parameterization of its moment generating function, the behavior in relation to several well-known generalized distributions is presented. By utilizing MacLaurin series expansion and Stirling formula, it is shown that with parameterization of its moment generating function, the generalized F distribution might have special relationship to several well-known generalized distributions, such as generalized beta of the second kind (GB2), generalized log-logistic (G4LL), and generalized gamma (G3G) distributions.

Keywords: Moment generating function; Generalized beta of the second kind distribution; Generalized log-logistic distribution; Generalized gamma; MacLaurin series; Stirling formula

1. INTRODUCTION

As stated by several authors, the generalized F is one of the most well-known generalized distributions in probability modeling. In fitting survival data of carcinoma patients, Ciampi, et al. (1986) discussed the properties and maximum likelihood

inference of the family of the generalized F distribution. Peng, et al. (1998) investigated the application of the generalized F distribution to a mixture model from lymphoma patients. Kalbfleisch and Prentice (2002) discussed the use of the generalized F distribution.

The generalized F distribution is parameterization of other well-known generalized distributions, such as the generalized beta and generalized log-logistic distributions (Pham-Ghia and Duong 1989, Singh 1989). According to Ciampi, et al. (1986) when a random variable X has four-parameter generalized F distribution with degrees of freedom $2m_1$ and $2m_2$, as denoted by X ~ G4F(μ , σ , m_1 , m_2) or G4F(m_1 , m_2), the corresponding probability density function (PDF) of the G4F(μ , σ , m_1 , m_2) can be written in the following form;

$$f_{G4F}(x) = \frac{e^{-\frac{\mu m_1}{\sigma}} x^{\left(\frac{m_1}{\sigma}\right)^{-1}} \left(\frac{m_1}{m_2}\right)^{m_1} \left[1 + \left(\frac{m_1}{m_2}\right) (e^{-\mu} x)^{\frac{1}{\sigma}}\right]^{-(m_1 + m_2)}}{\sigma \frac{\Gamma(m_1) \cdot \Gamma(m_2)}{\Gamma(m_1 + m_2)}}; x \ge 0$$
(1)

where $\mu > 0$, $\sigma > 0$, $m_1 > 0$, $m_2 > 0$, and Γ is gamma function.

It can be shown that if we let $\mu = \ln b + (1/a) \ln(m_1/m_2)$, $\sigma = 1/a$, and B(m_1,m_2) that is beta function, then the PDF of the G4F distribution becomes

$$f_{GB2}(x) = \frac{a}{xB(m_1, m_2)} \frac{\left[\left(\frac{x}{b}\right)^a\right]^{m_1}}{\left[1 + \left(\frac{x}{b}\right)^a\right]^{m_1 + m_2}}$$

which is the PDF of the GB2 distribution as deliberated by McDonald (1984), and McDonald and Richards (1987).

If we let $\mu = (-\beta - \ln (m_1/m_2))(1/\alpha)$ and $\sigma = 1/\alpha$, where and, then the PDF of the G4F distribution can be written in the form

$$f_{_{G4LL}}(x) = \frac{\alpha}{xB(m_1,m_2)} \left[\frac{1}{1 + e^{-(\beta + \alpha \ln(x))}}\right]^{m_1} \left[1 - \frac{1}{1 + e^{-(\beta + \alpha \ln(x))}}\right]^{m_2}$$

which is the PDF of the G4LL distribution considered by Singh, et al.(1997).

Moreover, as noted by several authors (Ciampi, et al. 1986; Peng et al. 1998; Kalbfleisch and Prentice 2002; and Zhou et al. 2011), the generalized F contains several commonly used distributions as special cases or limiting distributions, such as exponential, gamma, lognormal, and Weibull distributions. Recently, Cox (2008) has linked the G4F distribution to the three-parameter generalized (G3G) and the three-parameter generalized log-logistic (G3LL) distributions. To link the G4F distribution to other generalized distributions, he has parameterized the probability density function (PDF) of the G4F distribution.

However, in term of a moment generating function (MGF), flexibilities of the G4F distribution and relation to other generalized distributions are apparently not too readily accesible in all of above literatures. Based on the MGF, an alternative method could be generated for computing moments and characterizing distributions from their MGFs. Based on MGFs, Warsono (2009) demonstrated the relationship between GB2 and G3G distributions, and Warsono (2010) linked the G4LL distribution to the GB2 and G3G distribution. In developing other aspects of the G4F distribution, it seems worthwhile to extend all of above endeavors and to record them in this paper. Therefore, the main objective of this paper is to derive the MGF of the generalized F distribution and relate other generalized distributions, such as generalized beta, generalized log-logistic, and generalized gamma.

The rest of the article is outlined as follows. In Section 2, the explicit moment generating function of the G4F distribution is derived. By using parameterization of the moment generating function of the the G4F, Section 3 provides discussions of the relation between G4F distribution and GB2 and G4LL distributions. Section 4 contains a description of limiting behavior of the moment generating function of the G4F distribution as a general case of the moment generating function of G3G and G3F distributions. Finally, some concluding remarks are noted in section 5.

2. MOMENT GENERATING FUNCTION OF THE GENERALIZED F DISTRIBUTION

In term of a beta function the pdf of GF distribution in equation (1) can be rewritten as:

$$g(\mathbf{x}) = \frac{1}{\sigma \mathbf{x} B(m_1, m_2)} \frac{\left(e^{-\frac{\mu}{\sigma}} \left(\frac{m_1}{m_2}\right) \mathbf{x}^{\frac{1}{\sigma}}\right)^{m_1}}{\left[1 + \left(e^{-\frac{\mu}{\sigma}}\right) \mathbf{x}^{\frac{1}{\sigma}} \left(\frac{m_1}{m_2}\right)\right]^{m_1 + m_2}}; \ \mathbf{x} \ge 0$$
(2)

where $\mu > 0$, $\sigma > 0$, $m_1 > 0$, and , Γ is gamma function.

Theorem 2.1 Let *X* be a random variable of the G4F(μ , σ ,m₁,m₂) distribution, then moment generating function (MGF) of X is given by

$$M_{G4F}(t) = e^{te^{\mu}}$$

Proof:

$$M_{G4F}(t) = E(e^{tX}) = \int_{-\infty}^{\infty} e^{tx} \cdot f(x) dx$$

= $\int_{0}^{\infty} e^{tx} \left(\frac{1}{\sigma x B(m_1, m_2)}\right) \frac{e^{-\frac{\mu m_1}{\sigma}} \left(\frac{m_1}{m_2}\right)^{m_1} x^{m_1/\sigma}}{\left[1 + (e^{-\mu}x)^{1/\sigma} \frac{m_1}{m_2}\right]^{m_1 + m_2}} dx$

By algebra manipulation we may find the following equation

$$=\frac{1}{\sigma B(m_1,m_2)} \int_0^{\infty} \left(\frac{e^{tx}}{x}\right) \frac{\left[e^{-\frac{\mu}{\sigma}}\left(\frac{m_1}{m_2}\right)x^{\frac{1}{\sigma}}\right]^{m_1}}{\left[1+e^{-\frac{\mu}{\sigma}}\left(\frac{m_1}{m_2}\right)x^{\frac{1}{\sigma}}\right]^{m_1+m_2}} dx$$
(3)

By letting $y = e^{-\frac{\mu}{\sigma}} \left(\frac{m_1}{m_2}\right) x^{\frac{1}{\sigma}}$ we may rewrite the equation (3) in the following form

$$M_{G4F}(t) = \frac{1}{\sigma B(m_1, m_2)} \int_0^\infty \left(\frac{\sigma e^{tx}}{y} \right) \frac{y^{m_1}}{(1+y)^{m_1+m_2}} dy$$
$$= \frac{1}{B(m_1, m_2)} \int_0^\infty e^{tx} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy$$
(4)

Making use of a well-known property of MacLaurin series of the e^{tx} function, then equation (4) is given by

$$\begin{split} \mathbf{M}_{G4F}(t) &= \frac{1}{\mathbf{B}(\mathbf{m}_{1},\mathbf{m}_{2})} \int_{0}^{\infty} \left(\sum_{n=0}^{\infty} \frac{(tx)^{n}}{n!} \right) \frac{y^{\mathbf{m}_{1}-1}}{(1+y)^{\mathbf{m}_{1}+\mathbf{m}_{2}}} \, dy \\ &= \frac{1}{\mathbf{B}(\mathbf{m}_{1},\mathbf{m}_{2})} \int_{0}^{\infty} \left(1+tx + \frac{(tx)^{2}}{2!} + \frac{(tx)^{3}}{3!} + \dots \right) \frac{y^{\mathbf{m}_{1}-1}}{(1+y)^{\mathbf{m}_{1}+\mathbf{m}_{2}}} \, dy \end{split}$$

$$\begin{split} &= \frac{1}{B(m_1,m_2)} \int_0^{\sigma} \left(\frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + tx \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + \frac{(tx)^2}{2!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + \frac{(tx)^3}{3!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} + \dots \right) dy \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \int_0^{\sigma} tx \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \int_0^{\sigma} \frac{(tx)^2}{3!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \int_0^{\sigma} \frac{(tx)^3}{3!} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} y^{\sigma} \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} y^{\sigma} \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} y^{\sigma} \cdot \frac{y^{m_1-1}}{(1+y)^{m_1+m_2-\sigma}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} \frac{y^{m_1+\sigma-1}}{(1+y)^{m_1+\sigma+m_2-\sigma}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} \frac{y^{m_1+\sigma-1}}{(1+y)^{m_1+\sigma+m_2-\sigma}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1-1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} \frac{y^{m_1+\sigma-1}}{(1+y)^{m_1+\sigma+m_2-\sigma}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1+1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} \frac{y^{m_1+\sigma-1}}{(1+y)^{m_1+\sigma+m_2-\sigma}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1+1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} \frac{y^{m_1+\sigma-1}}{(1+y)^{m_1+\sigma+m_2-\sigma}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1+1}}{(1+y)^{m_1+m_2}} dy + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \int_0^{\sigma} \frac{y^{m_1+\sigma-1}}{(1+y)^{m_1+\sigma+m_2-\sigma}} dy + \dots \right] \\ &= \frac{1}{B(m_1,m_2)} \left[\int_0^{\sigma} \frac{y^{m_1+1}}{(1+y)^{m_1+m_2}} dy + t \left(\frac{y^{m_1+1}}{(1+y)^{m_1+\sigma+m_2-\sigma}} dy + \dots \right] \\ &= \frac{1}{$$

$$= \frac{1}{B(m_1, m_2)} \left[B(m_1, m_2) + t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \cdot B(m_1 + \sigma, m_2 - \sigma) + \frac{\left(t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \right)^2}{2!} \cdot B(m_1 + 2\sigma, m_2 - 2\sigma) + \frac{\left(t \left(e^{\frac{\mu}{\sigma}} \left(\frac{m_2}{m_1} \right) \right)^{\sigma} \right)^3}{3!} \cdot B(m_1 + 3\sigma, m_2 - 3\sigma) + \dots \right]$$

Therefore, the MGF of the G4F distribution is

$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left(te^{\mu} \left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \cdot \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$
(5)

It is well-known that the Stirling's approximation formula of the gamma function (Spiegel 1968) is given by

$$\Gamma(az+b) \sim \sqrt{2\pi} \cdot e^{-az} (az)^{az+b-\frac{1}{2}}$$

By Stirling's approximation, equation (5) can be expressed as

$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left(te^{\mu} \left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \cdot \frac{\sqrt{2\pi} e^{-m_1} (m_1)^{m_1 + n\sigma - \frac{1}{2}} \cdot \sqrt{2\pi} e^{-m_2} (m_2)^{m_2 + n\sigma - \frac{1}{2}}}{\sqrt{2\pi} e^{-m_1} (m_1)^{m_1 - \frac{1}{2}} \cdot \sqrt{2\pi} e^{-m_2} (m_2)^{m_2 - \frac{1}{2}}}$$
$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left(te^{\mu} \left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \cdot \left(m_1^{\sigma}\right)^n \left(\left(\frac{1}{m_2}\right)^{\sigma}\right)^n}$$
$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left(te^{\mu} \left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \cdot \left(\left(\frac{m_1}{m_2}\right)^{\sigma}\right)^n}$$
(6)

By MacLaurin series, equation (6) can be written as

$$\begin{split} M_{G4F}(t) = & e^{te^{\mu} \left(\frac{m_2}{m_1}\right)^{\sigma} \left(\frac{m_1}{m_2}\right)^{\sigma}} \\ M_{G4F}(t) = & e^{te^{\mu}} \end{split}$$

2.2. Moment Generating Function (MGF) of Three-Parameter G3F Distribution

Based on the gamma distribution (Malik 1967, and Dyer 1982) and generalized beta with three- parameter distribution (Pham-Ghia and Duong 1989) obtained three-parameters of generalized F distribution with the following pdf

$$f_{G3F}(x) = \frac{\alpha^{m_1}}{B(m_1, m_2)} \frac{x^{m_1 - 1}}{\left[1 + \alpha x\right]^{m_1 + m_2}} dx; x > 0, m_1, m_2, \alpha > 0$$

Theorem 2.2 Let *X* be a random variable of the G3F (α ,m₁,m₂) distribution, then moment generating function (MGF) of X is given by

$$M_{G3F}(t) = \sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^n}{n!} \cdot \frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$

Proof:

$$\begin{split} \mathbf{M}_{\rm G3F}(t) &= \mathbf{E} \Big(\mathbf{e}^{t \mathbf{X}} \Big) = \int_{-\infty}^{\infty} \mathbf{e}^{t \mathbf{x}} \cdot \mathbf{f} \Big(\mathbf{x} \Big) \, d\mathbf{x} \\ &= \int_{0}^{\infty} \mathbf{e}^{t \mathbf{x}} \left(\frac{\alpha^{m_1}}{\mathbf{B}(m_1, m_2)} \right) \frac{\mathbf{x}^{m_1 - 1}}{\left[\mathbf{I} + \alpha \mathbf{x} \right]^{m_1 + m_2}} \, d\mathbf{x} \end{split}$$

By algebra manipulation we may find the following equation

$$= \frac{\alpha^{m_1}}{B(m_1, m_2)} \int_0^\infty (e^{tx}) \frac{x^{m_1 - 1}}{[1 + \alpha x]^{m_1 + m_2}} dx$$
(7)

By letting $y = \alpha x$ we may rewrite the equation (7) in the following form

$$M_{X}(t) = \frac{\alpha^{m_{1}}}{B(m_{1}, m_{2})} \int_{0}^{\infty} \left(e^{tx}\right) \frac{\left(\frac{y}{\alpha}\right)^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} \frac{1}{\alpha} dy$$
$$= \frac{1}{B(m_{1}, m_{2})} \int_{0}^{\infty} \left(e^{tx}\right) \frac{(y)^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} dy$$
(8)

Making use of a well-known property of MacLaurin series of the e^{tx} function, then equation (8) can be written as

$$\begin{split} M_{x}(t) &= \frac{1}{B(m_{1},m_{2})} \int_{0}^{\infty} \left(\sum_{n=0}^{\infty} \frac{(tx)^{n}}{n!} \right) \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} dy \\ &= \frac{1}{B(m_{1},m_{2})} \int_{0}^{\infty} \left(1+tx + \frac{(tx)^{2}}{2!} + \frac{(tx)^{3}}{3!} + ... \right) \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} dy \\ &= \frac{1}{B(m_{1},m_{2})} \int_{0}^{\infty} \left(\frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} + tx \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} + \frac{(tx)^{2}}{2!} \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} + \frac{(tx)^{3}}{3!} \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} + ... \right) dy \\ &= \frac{1}{B(m_{1},m_{2})} \left| \int_{0}^{\infty} \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} dy + \int_{0}^{\infty} tx \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} dy + \\ & \int_{0}^{\infty} \frac{(tx)^{2}}{2!} \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} dy + \int_{0}^{\infty} \frac{(tx)^{3}}{3!} \frac{y^{m_{1}-1}}{(1+y)^{m_{1}+m_{2}}} dy + \right| \end{split}$$

$$= \frac{1}{B(m_1, m_2)} \left[B(m_1, m_2) + \left(\frac{t}{\alpha}\right) \cdot B(m_1 + 1, m_2 - 1) + \frac{\left(\frac{t}{\alpha}\right)^2}{2!} \cdot B(m_1 + 2, m_2 - 2) + \frac{\left(\frac{t}{\alpha}\right)^3}{3!} \cdot B(m_1 + 3, m_2 - 3) + \dots \right]$$

$$\begin{split} &= 1 + \frac{t}{\alpha} \cdot \frac{B(m_1 + 1, m_2 - 1)}{B(m_1, m_2)} + \frac{\left(\frac{t}{\alpha}\right)^2}{2!} \cdot \frac{B(m_1 + 2, m_2 - 2)}{B(m_1, m_2)} + \frac{\left(\frac{t}{\alpha}\right)^3}{3!} \cdot \frac{B(m_1 + 3, m_2 - 3)}{B(m_1, m_2)} + \dots \\ &= \sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^n}{n!} \cdot \frac{B(m_1 + n, m_2 - n)}{B(m_1, m_2)} \\ &= \sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^n}{n!} \cdot \frac{\frac{\Gamma(m_1 + n) \cdot \Gamma(m_2 - n)}{\Gamma(m_1 + n + m_2 - n)}}{\frac{\Gamma(m_1) \cdot \Gamma(m_2)}{\Gamma(m_1 + m_2)}} \end{split}$$

$$=\sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^{n}}{n!} \cdot \frac{\Gamma(m_{1}+n) \cdot \Gamma(m_{2}-n)}{\Gamma(m_{1}+m_{2})} \cdot \frac{\Gamma(m_{1}+m_{2})}{\Gamma(m_{1}) \cdot \Gamma(m_{2})}$$
$$=\sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^{n}}{n!} \cdot \frac{\Gamma(m_{1}+n) \cdot \Gamma(m_{2}-n)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})}$$

Therefore, the MGF of the G3F distribution is

$$M_{x}(t) = \sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^{n}}{n!} \cdot \frac{\Gamma(m_{1}+n) \cdot \Gamma(m_{2}-n)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})}$$
(9)

3. THE RELATION OF THE GF DISTRIBUTION WITH GB2 DISTRIBUTION

Warsono (2010) has mathematically derived the MGF of the GB2 distribution using MGF definition. Based on reparameterization of the MGF of the GLF distribution, use passive sentence- The MGF of the GB2 distribution is provided in this section. The reparameterization proposition is stated and proved.

Proposition 3.1 Let X be a random variable having the G4F (μ , σ , m_1 , m_2) moment and

$$\mu = \ln b + \frac{1}{a} \ln \left(\frac{m_1}{m_2} \right)$$
 and $\sigma = \frac{1}{a}$, then X has the GB2(a, b, m_1, m_2) moment.

Proof:

$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left(te^{\mu}\left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$
$$= \sum_{n=0}^{\infty} \frac{\left(te^{\ln b + \frac{1}{a}\ln\left(\frac{m_1}{m_2}\right)}\left(\frac{m_2}{m_1}\right)^{\frac{1}{a}}\right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{a}\right) \cdot \Gamma\left(m_2 - \frac{n}{a}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$

951

$$\begin{split} &=\sum_{n=0}^{\infty} \frac{\left(te^{lnb} e^{\frac{1}{a}ln\left(\frac{m_1}{m_2}\right)} \left(\frac{m_2}{m_1}\right)^{\frac{1}{a}}\right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{a}\right) \cdot \Gamma\left(m_2 - \frac{n}{a}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\ &= \sum_{n=0}^{\infty} \frac{\left(tb \left(\frac{m_1}{m_2}\right)^{\frac{1}{a}} \left(\frac{m_2}{m_1}\right)^{\frac{1}{a}}\right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{a}\right) \cdot \Gamma\left(m_2 - \frac{n}{a}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\ &= \sum_{n=0}^{\infty} \frac{\left(tb\right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{a}\right) \cdot \Gamma\left(m_2 - \frac{n}{a}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \end{split}$$

This is the moment generating function of the GB2 (a,b, m_1 , m_2) stated by Warsono (2010).

4. THE RELATION OF THE GF DISTRIBUTION WITH GLL DISTRIBUTION

Warsono (2010) derived mathematically the MGF of the GLL distribution using definition of MGF's. Based on reparameterization of the MGF of the GF distribution, the MGF of the GLL distribution is provided in this section.

Proposition 4.1 Let X be a random variable having the GF (μ , σ , m_1 , m_2) moment and

$$\mu = \left(-\beta - \ln\left(\frac{m_2}{m_1}\right)\right) \frac{1}{\alpha} \text{ and } \sigma = \frac{1}{\alpha}, \text{ then X has the GLL}(\alpha, \beta, m_1, m_2) \text{ moment.}$$

Proof:

$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left(te^{\mu} \left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$
$$= \sum_{n=0}^{\infty} \frac{\left(te^{\left(-\beta - \ln\left(\frac{m_2}{m_1}\right)\right)\frac{1}{\alpha}} \left(\frac{m_2}{m_1}\right)^{\frac{1}{\alpha}}\right)^n}{n!} \frac{\Gamma\left(m_1 + \frac{n}{\alpha}\right) \cdot \Gamma\left(m_2 - \frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$

$$\begin{split} &=\sum_{n=0}^{\infty} \frac{\left(te^{-\frac{\beta}{\alpha}-\frac{1}{\alpha}ln\left(\frac{m_2}{m_1}\right)}\left(\frac{m_2}{m_1}\right)^{\frac{1}{\alpha}}\right)^n}{n!} \frac{\Gamma\left(m_1+\frac{n}{\alpha}\right) \cdot \Gamma\left(m_2-\frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\ &=\sum_{n=0}^{\infty} \frac{\left(te^{-\frac{\beta}{\alpha}}e^{-\frac{1}{\alpha}ln\left(\frac{m_1}{m_2}\right)}\left(\frac{m_2}{m_1}\right)^{\frac{1}{\alpha}}\right)^n}{n!} \frac{\Gamma\left(m_1+\frac{n}{\alpha}\right) \cdot \Gamma\left(m_2-\frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\ &=\sum_{n=0}^{\infty} \frac{\left(te^{-\frac{\beta}{\alpha}}\left(\frac{m_2}{m_1}\right)^{-\frac{1}{\alpha}}\left(\frac{m_2}{m_1}\right)^{\frac{1}{\alpha}}\right)^n}{n!} \frac{\Gamma\left(m_1+\frac{n}{\alpha}\right) \cdot \Gamma\left(m_2-\frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \\ &=\sum_{n=0}^{\infty} \frac{\left(te^{-\frac{\beta}{\alpha}}\right)^n}{n!} \frac{\Gamma\left(m_1+\frac{n}{\alpha}\right) \cdot \Gamma\left(m_2-\frac{n}{\alpha}\right)}{\Gamma(m_1) \cdot \Gamma(m_2)} \end{split}$$

5. THE RELATION OF THE G4F DISTRIBUTION WITH G3F DISTRIBUTION

Equation (8) has mathematically derived the MGF of the G3F distribution. Based on reparameterization of the MGF of the GF distribution, the MGF of the G3F distribution is provided in this section.

Proposition 5.1 Let X be a random variable having the G4F (μ , σ , m_1 , m_2) moment and

$$\mu = \left(-\ln\left(\frac{m_2}{m_1}\right) - \ln\alpha\right) \text{ and } \sigma = 1, \text{ then } X \text{ has the } G3F(\alpha, m_1, m_2) \text{ moment.}$$

Proof:

$$M_{G4F}(t) = \sum_{n=0}^{\infty} \frac{\left(te^{\mu} \left(\frac{m_2}{m_1}\right)^{\sigma}\right)^n}{n!} \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$

$$=\sum_{n=0}^{\infty} \frac{\left(te^{\left(-\ln\left(\frac{m_2}{m_1}\right)-\ln\alpha\right)}\left(\frac{m_2}{m_1}\right)\right)^n}{n!} \frac{\Gamma(m_1+n)\cdot\Gamma(m_2-n)}{\Gamma(m_1)\cdot\Gamma(m_2)}$$
$$=\sum_{n=0}^{\infty} \frac{\left(te^{-\ln\left(\frac{m_1}{m_2}\right)}e^{-\ln\alpha}\left(\frac{m_2}{m_1}\right)\right)^n}{n!} \frac{\Gamma(m_1+n)\cdot\Gamma(m_2-n)}{\Gamma(m_1)\cdot\Gamma(m_2)}$$
$$=\sum_{n=0}^{\infty} \frac{\left(te^{\ln\frac{1}{\alpha}}\left(\frac{m_2}{m_1}\right)^{-1}\left(\frac{m_2}{m_1}\right)\right)^n}{n!} \frac{\Gamma(m_1+n)\cdot\Gamma(m_2-n)}{\Gamma(m_1)\cdot\Gamma(m_2)}$$
$$=\sum_{n=0}^{\infty} \frac{\left(\frac{t}{\alpha}\right)^n}{n!} \frac{\Gamma(m_1+n)\cdot\Gamma(m_2-n)}{\Gamma(m_1)\cdot\Gamma(m_2)}$$

This is the moment generating function of the $G3F(\alpha, m_1, m_2)$ stated in equation (9).

6. THE MOMENT OF THE GG AS A LIMITING MOMENT OF THE GF DISTRIBUTIONS

In this section, propositions of limiting moment properties of the GF distribution are stated and proved.

Proposition 6.1 The GF(μ,σ,m_1,m_2) distribution converges to the GG distribution as

m₂ tends to
$$\infty$$
 and $\mu = \ln \gamma + \frac{1}{a} \ln(m_1)$, and $\sigma = \frac{1}{a}$.

Proof:

$$\lim_{m_2 \to \infty} M_{G4F}(t) = \lim_{m_2 \to \infty} \sum_{n=0}^{\infty} \frac{\left(t e^{\mu} \left(\frac{m_2}{m_1} \right)^{\sigma} \right)^n}{n!} \cdot \frac{\Gamma(m_1 + n\sigma) \cdot \Gamma(m_2 - n\sigma)}{\Gamma(m_1) \cdot \Gamma(m_2)}$$

$$\begin{split} &= \lim_{m_{2} \to \infty} \sum_{n=0}^{\infty} \underbrace{\left(te^{in\gamma + \frac{1}{a} \ln(m_{1})} \left(\frac{m_{2}}{m_{1}} \right)^{\frac{1}{a}} \right)^{n}}_{n!} \cdot \frac{\Gamma\left(m_{1} + \frac{n}{a}\right) \cdot \Gamma\left(m_{2} - \frac{n}{a}\right)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})} \\ &= \lim_{m_{2} \to \infty} \sum_{n=0}^{\infty} \underbrace{\left(te^{in\gamma} e^{\frac{1}{a} \ln(m_{1})} \left(\frac{m_{2}}{m_{1}} \right)^{\frac{1}{a}} \right)^{n}}_{n!} \cdot \frac{\Gamma\left(m_{1} + \frac{n}{a}\right) \cdot \Gamma\left(m_{2} - \frac{n}{a}\right)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})} \\ &= \lim_{m_{2} \to \infty} \sum_{n=0}^{\infty} \underbrace{\left(t\gamma(m_{1})^{\frac{1}{a}} \left(\frac{m_{2}}{m_{1}} \right)^{\frac{1}{a}} \right)^{n}}_{n!} \cdot \frac{\Gamma\left(m_{1} + \frac{n}{a}\right) \cdot \Gamma\left(m_{2} - \frac{n}{a}\right)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})} \\ &= \lim_{m_{2} \to \infty} \sum_{n=0}^{\infty} \underbrace{\left(t\gamma(m_{2})^{\frac{1}{a}} \right)^{n}}_{n!} \cdot \frac{\Gamma\left(m_{1} + \frac{n}{a}\right) \cdot \Gamma\left(m_{2} - \frac{n}{a}\right)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})} \\ &= \lim_{m_{2} \to \infty} \sum_{n=0}^{\infty} \underbrace{\left(t\gamma(m_{2})^{\frac{1}{a}} \right)^{n}}_{2!} \cdot \frac{\Gamma\left(m_{1} + \frac{1}{a}\right) \cdot \Gamma\left(m_{2} - \frac{1}{a}\right)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})} + \\ &= \lim_{m_{2} \to \infty} \underbrace{\left(t\gamma(m_{2})^{\frac{1}{a}} \right)^{\frac{1}{a}}}_{3!} \cdot \frac{\Gamma\left(m_{1} + \frac{2}{a}\right) \cdot \Gamma\left(m_{2} - \frac{2}{a}\right)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})} + \\ &= \lim_{m_{2} \to \infty} \underbrace{\left(t\gamma(m_{2})^{\frac{1}{a}} \right)^{\frac{1}{a}}}_{3!} \cdot \frac{\Gamma\left(m_{1} + \frac{3}{a}\right) \cdot \Gamma\left(m_{2} - \frac{3}{a}\right)}{\Gamma(m_{1}) \cdot \Gamma(m_{2})} + \dots \\ &= \lim_{m_{2} \to \infty} 1 + (t\gamma) \cdot \frac{\Gamma\left(m_{1} + \frac{1}{a}\right)}{\Gamma(m_{1})} \lim_{m_{2} \to \infty} (m_{2})^{\frac{1}{a}} \frac{\Gamma\left(m_{2} - \frac{1}{a}\right)}{\Gamma(m_{2})} + \dots \end{split}$$

$$\frac{(t\gamma)^{2}}{2!} \cdot \frac{\Gamma\left(m_{1} + \frac{2}{a}\right)}{\Gamma(m_{1})} \lim_{m_{2} \to \infty} \cdot (m_{2})^{\frac{2}{a}} \frac{\Gamma\left(m_{2} - \frac{2}{a}\right)}{\Gamma(m_{2})} + \frac{(t\gamma)^{3}}{3!} \cdot \frac{\Gamma\left(m_{1} + \frac{3}{a}\right)}{\Gamma(m_{1})} \lim_{m_{2} \to \infty} \cdot (m_{2})^{\frac{3}{a}} \frac{\Gamma\left(m_{2} - \frac{3}{a}\right)}{\Gamma(m_{2})} + \dots$$

Using Stirling's approximation formula of the gamma function, the limiting moment property of the G4F(μ , σ , m_1 , m_2) distribution can be written as:

$$\begin{split} &\lim_{m_{2}\to\infty} M_{\chi}(t) \, GF\left(\mu = \ln \gamma + \frac{1}{a} \ln(m_{1}), \sigma = \frac{1}{a}, m_{1}, m_{2}\right) \\ &= \lim_{m_{2}\to\infty} 1 + (t\gamma) \cdot \frac{\Gamma\left(m_{1} + \frac{1}{a}\right)}{\Gamma(m_{1})} \lim_{m_{2}\to\infty} (m_{2})^{\frac{1}{a}} \cdot \frac{1}{(m_{2})^{\frac{1}{a}}} + \frac{(t\gamma)^{2}}{2!} \cdot \frac{\Gamma\left(m_{1} + \frac{2}{a}\right)}{\Gamma(m_{1})} \lim_{m_{2}\to\infty} (m_{2})^{\frac{2}{a}} \cdot \frac{1}{(m_{2})^{\frac{2}{a}}} + \dots \\ &= 1 + (t\gamma) \cdot \frac{\Gamma\left(m_{1} + \frac{1}{a}\right)}{\Gamma(m_{1})} + \frac{(t\gamma)^{2}}{2!} \cdot \frac{\Gamma\left(m_{1} + \frac{2}{a}\right)}{\Gamma(m_{1})} + \\ &= \sum_{n=0}^{\infty} \frac{(t\gamma)^{n}}{n!} \cdot \frac{\Gamma\left(m_{1} + \frac{n}{a}\right)}{\Gamma(m_{1})} \end{split}$$

This result is the MGF of the GG stated by Warsono(2010). Thus, the G4F distribution converges to the GG distribution as m₂ tends to ∞ and $\alpha = a$, and $\beta = -a \ln \left(\gamma \left(m_2 \right)^{\frac{1}{a}} \right)$

7. CONCLUSION

The moment of the generalized F distribution is parameterization of the generalized beta of the second kind (GB2) and the generalized log-logistic (GLL). The moment of the generalized gamma (GG) distribution is the limiting moment of (GF) distribution. Moreover, since the moments of the gamma and exponential distributions are special cases of the moment of the generalized gamma distribution (Warsono 2009), the moments of both special distributions are also special cases of the moments of the GF distribution.

REFERENCES

- [1]. Ciampi, A., S.A. Hogg, and L. Kates. (1986). Regression analysis of censored survival data with the generalized F family an alternative to the proportional hazards model. *Statistics in Medicine* Vol 5: 85-96.
- [2]. Cox, C. (2008). The generalized *F* distribution: An umbrella for parametric survival analysis. *Statistics in Medicine* 27: 4301-4312.
- [3]. Dyer, D. (1982). The convolution of generalized F distribution. J. Am. Statist. Ass. 77: 184-189.
- [4]. Kalbfleisch, J. D. and Prentice, R. L. (2002). *The Statistical Analysis of Failure Time Data*, Wiley, New York.
- [5]. Malik, H. J. (1967). Exact distributions of the quotient of independent generalized gamma variables. *Can. math. BUN.* **10**,463465.
- [6]. McDonald, J.B. (1984). Some generalized functions for the size distribution of income. *Econometrica* 52(3): 647-663.
- [7]. McDonald, J.B. and D.O. Richards. (1987). Hazard rates and generalized beta distributions. *IEEE Transactions reliability*, Vol. R-36: 463-466.
- [8]. Pham-Gia, T. And Q.P. Duong. (1989). The generalized beta- and Fdistributions in statistical Modelling. *Math. Comput. Modelling*. Vol 12(12): 1613-1625.
- [9]. Peng, Y.P., K.B.G. Dear, and J.W. Denham. (1998). A generalized F mixture model for cure rate estimation. *Statistics in Medicine* 17: 813-830.
- [10]. Singh, K.P. (1989). A generalized log-logistic regression model for survival analysis: hazard rate characteristics. *Biometrie-Praximetrie* 29: 63-74
- [11]. Singh, K.P., Warsono, and A.A. Bartolucci. (1997). Generalized log-logistic model for analysis of environmental pollutant data. Proc. Int. Congress on Modeling and Simulation. Hobart, Tasmania. December 8-11 1997.
- [12]. Spiegel, M.R. (1968). Mathematical Handbook of Formulas and Tables. McGraw-Hill Inc.
- [13]. Warsono. (2009). Moment Properties of the Generalized Gamma Distribution. *Proceedings* on Seminar Nasional Sains MIPA dan Aplikasinya, Bandar Lampung: 157-162.
- [14]. Warsono. (2010). Remarks on Moment Properties of Generalized Distributions. Proceedings of the Third International Conference on Mathematics and Natural Sciences (ICMNS 2010). Faculty of Mathematics and Natural Sciences Institut Teknologi Bandung. p. 1123 -1131.
- [15]. Xiaojing Zhou, X., L.Yan, D.R. Prows, and Runqing Yang. (2011). Generalized F accelerated failure time model for mapping survival trait loci. *Genomics* 97: 379-385.