

# The Locating-Chromatic Number of Origami Graphs

*By* Kurnia Muludi

Article

# The Locating-Chromatic Number of Origami Graphs

Agus Irawan<sup>1,2</sup>, Asmiati Asmiati<sup>1</sup>, La Zakaria<sup>1,\*</sup> and Kurnia Muludi<sup>3</sup>

<sup>1</sup> Department of Mathematics, Universitas Lampung, Bandar Lampung 35145, Indonesia; agusirawan814@gmail.com (A.I.); asmiati.1976@fmipa.unila.ac.id (A.A.)

<sup>2</sup> Information System, STMIK Pringsewu, Lampung 35373, Indonesia

<sup>3</sup> Department of Computer Science, Universitas Lampung, Bandar Lampung 35145, Indonesia; kmuludi@fmipa.unila.ac.id

\* Correspondence: lazakaria.1969@fmipa.unila.ac.id; Tel.: +62-812-790-9255

**Abstract** The locating-chromatic number of a graph combines two graph concepts, namely coloring vertices and partition dimension of a graph. The locating-chromatic number is the smallest  $k$  such that  $G$  has a locating  $k$ -coloring, denoted by  $\chi_L(G)$ . This article proposes a procedure for obtaining a locating-chromatic number for an origami graph and its subdivision (one vertex on an outer edge) through two theorems with proofs.

**Keywords:** locating-chromatic number; origami graphs; subdivision

**MSC:** 05C12; 05C15



**Citation:** Irawan, A.; Asmiati, Zakaria, L.; Muludi, K. The Locating-Chromatic Number of Origami Graphs. *Algorithms* **2021**, *14*, 167. <https://doi.org/10.3390/a14060167>

Academic Editor: Frank Werner

Received: 26 April 2021

Accepted: 26 May 2021

Published: 27 May 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The study of the partition dimension of connected graphs was introduced by Chartrand et al. [1,2] with the aim of finding a new method for attacking the problem of determining the metric dimension in graphs. The application of these metric dimensions can be seen in the navigation of a robot modeled by a graph [3], solving the problem of chemical data classification, and determining how to represent a set of chemical compounds in such a way that different compounds have different representations [5,6]. The concept of the locating-chromatic number was first introduced by Chartrand et al. in 2002, with two obtained graph concepts, namely coloring vertices and partition dimensions of a graph [7]. Finding the locating-chromatic number of a graph is one of the interesting (and un-completely solved) problems of graph theory. Let  $G = (V, E)$  be a connected graph; the distance  $d(x, y)$  between two of its vertices  $x$  and  $y$  is the length of the shortest path between them. Let  $c$  be a proper  $k$ -coloring of  $G$  with color  $\{1, 2, \dots, k\}$ , and  $\Pi = \{C_1, C_2, \dots, C_k\}$  be a partition of  $V(G)$  that is induced by the coloring  $c$ . The color code  $c_{\Pi}(v)$  of  $v$  is the ordered  $k$ -tuple  $(d(v, C_1), d(v, C_2), \dots, d(v, C_k))$ , where  $d(v, C_i) = \min \{d(v, x) : x \in C_i\}$  for any  $i \in \{1, 2, 3, \dots, k\}$ . If all distinct vertices of  $G$  have distinct color codes, then  $c$  is called a  $k$ -locating coloring of  $G$ . The locating-chromatic number denoted by  $\chi_L(G)$  is the smallest  $k$  such that  $G$  has a locating  $k$ -coloring. Let  $c$  be a locating  $k$ -coloring on graph  $G(V, E)$ . Furthermore, the locating-chromatic number has been determined for a few graph classes; for example, if  $P_n$  is a path of order  $n \geq 3$  then the locating-chromatic number is 3; for a cycle  $C_n$  if  $n \geq 3$  is odd,  $\chi_L(C_n) = 3$  was obtained, and if  $n$  is even,  $\chi_L(C_n) = 4$  was obtained; for a double star graph  $(S_{a,b}), 1 \leq a \leq b$  and  $b \geq 2$ ,  $\chi_L(S_{a,b}) = b + 1$  was obtained. Let  $\Pi = \{S_1, S_2, \dots, S_k\}$  be the partition of  $V(G)$  induced by  $c$ . A vertex  $v \in G$  is called a dominant vertex if  $d(v, S_i) = 1$ , where  $v \notin S_i$ . Chartrand et al. characterized all graphs of order  $n$  with the locating-chromatic number  $n - 1$  [8].

The problem of determining the locating-chromatic number of any general graph is an NP-hard problem [9]. This means that to determine the locating-chromatic number of any given graph, we need a specific algorithm. In 2012, Baskoro and Purwasih proposed a procedure to obtain the locating-chromatic number of corona products of two graphs [9]. In

2014, Asmiati obtained the locating-chromatic number of a non-homogeneous amalgamation of stars [10]. Moreover, to determine the locating-chromatic number of disconnected graphs, graphs with dominant vertices and graphs of two components have been discussed in [11–13]. In 2019, the characterization of the locating chromatic number of powers of paths and a condition (sharp upper and lower bounds) for the locating chromatic number of powers of cycles were discussed [14] (see [15] for a discussion of the necessary and sufficient conditions for a pair of two specific start graphs to be an odd mean graph). Asmiati et al. determined the locating-chromatic number of some Petersen graphs;  $P(n, 1)$  four for odd  $n \geq 3$  or five for even  $n \geq 4$  were obtained [16], and in [18] results were obtained for certain barbell graphs. Syofyan et al. have succeeded in determining the locating-chromatic number of homogeneous lobsters [18]. In [19], Asmiati obtained the locating-chromatic number for non-homogeneous caterpillar graphs and non-homogeneous firecracker graphs. Next, Irawan and Asmiati in 2018 determined the locating-chromatic number of subdivision firecrackers graphs [11] and in [21] obtained the certain operation of generalized Petersen graphs  $sP(n, 1)$ . In 2014, Behtoui and Anbarloei determined the locating-chromatic number of the joining of two arbitrary graphs [22]. In addition to that, in this article we propose a procedure for obtaining the locating chromatic number for an origami graph and its subdivision (one vertex on an outer edge). The following definition of an origami graph is taken from [23]. Let  $n \in \mathbb{N}$  with  $n \geq 3$ . An origami graph  $O_n$  is a graph with  $V(O_n) = \{u_i, v_i, w_i : i \in \{1, \dots, n\}\}$  and  $E(O_n) = \{u_i w_i, u_i v_i, v_i w_i : i \in \{1, \dots, n\}\} \cup \{u_i u_{i+1}, w_i w_{i+1} : i \in \{1, \dots, n-1\}\} \cup \{u_n u_1, w_n w_1\}$  (see Figure 1 for an example). Meanwhile, a subdivision of an origami graph  $O_n^*$  is a graph with  $V(O_n^*) = \{u_i, v_i, x_i, w_i : i \in \{1, \dots, n\}\}$  and  $E(O_n^*) = \{u_i w_i, u_i v_i, v_i x_i, x_i w_i : i \in \{1, \dots, n\}\} \cup \{u_i u_{i+1}, w_i w_{i+1} : i \in \{1, \dots, n-1\}\} \cup \{u_n u_1, w_n w_1\}$  (see Figure 2 for an example).




Figure 1. An origami graph  $O_5$ .




Figure 2. A subdivision of an origami graph  $O_5^*$ .

## 2. Results and Discussions

Let  $c$  be a locating coloring in a connected graph  $G$  and  $N(q)$  denote the set of neighbor of a vertex  $q$  in  $G$ . If  $p$  and  $q$  are distinct vertices of  $G$  such that  $d(p, w) = d(q, w)$  for all  $w \in V(G) - \{p, q\}$ , then  $c(p) \neq c(q)$ . In particular, if  $p$  and  $q$  are non-adjacent vertices such that  $N(p) = N(q)$ , then  $c(p) \neq c(q)$  [7].

In the following subsection, the locating-chromatic number of origami graphs  $O_n$  and their subdivisions called  $O_n^*$  is described.

### 2.1. Locating-Chromatic Number of Origami Graphs

**Theorem 1.** Let  $O_n$  be an origami graph for  $n \geq 3$ . Then, the locating-chromatic number of  $O_n$ ,

$$\chi_L(O_n) = \begin{cases} 4, & \text{for } n = 3 \\ 5, & \text{otherwise.} \end{cases}$$

**Proof.** Let  $n \in \mathbb{N}$  with  $n \geq 3$ . An origami graph  $O_n$  is a graph with  $V(O_n) = \{u_i, v_i, w_i : i \in \{1, \dots, n\}\}$  and  $E(O_n) = \{u_i w_i, v_i w_i : i \in \{1, \dots, n\}\} \cup \{u_i u_{i+1}, w_i w_{i+1} : i \in \{1, \dots, n-1\}\} \cup \{u_n u_1, w_n w_1\}$ . Next, to prove the theorem, we consider the following two cases:

**Case 1.**  $\chi_L(O_3) = 4$

First, we determine the lower bound of  $\chi_L(O_3)$ . In the origami graphs  $O_n$  for  $n \geq 3$ , there are three adjacent vertices (complete graph with three vertices, denoted by  $K_3$ ); we then need at least 3-locating coloring. Without loss of generality, we assign three colors for any  $K_3$  in  $O_n$  for  $n \geq 3$ , and then the three vertices are dominant vertices. As a result, if we use three colors it is not enough because there are more than one  $K_3$  in  $O_n$  for  $n \geq 3$ . Therefore,  $\chi_L(O_3) \geq 4$ .

Next, we determine the upper bound of  $\chi_L(O_3) \leq 4$ . To show that 4 is an upper bound for the locating-chromatic number for the origami graph  $O_3$  we describe a locating coloring  $c$  using four colors as follows:

$$\begin{aligned} c(u_i) &= i, i = 1, 2, 3. \\ c(v_i) &= \begin{cases} 2, & \text{for } i = 1, 3 \\ 3, & \text{for } i = 2. \end{cases} \\ c(w_i) &= 4, i = 1, 2, 3. \end{aligned}$$

The coloring  $c$  will create the partition  $\Pi$  on  $V(O_3)$ . We shall show that the color codes of all vertices in  $O_3$  are different. We have:  $c_\Pi(u_1) = (0, 1, 1, 1)$ ;  $c_\Pi(u_2) = (1, 0, 1, 1)$ ;  $c_\Pi(u_3) = (1, 1, 0, 1)$ ;  $c_\Pi(v_1) = (1, 0, 2, 1)$ ;  $c_\Pi(v_2) = (2, 1, 0, 1)$ ;  $c_\Pi(v_3) = (2, 0, 1, 1)$ ;  $c_\Pi(w_1) = (1, 1, 2, 0)$ ;  $c_\Pi(w_2) = (2, 1, 1, 0)$ ;  $c_\Pi(w_3) = (1, 1, 1, 0)$ . Since the color codes of all vertices  $O_3$  are different,  $c$  is a locating-chromatic coloring. Thus,  $\chi_L(O_3) \leq 4$ .

**Case 2.**  $\chi_L(O_n) = 5$ , for  $n \geq 4$

To determine the lower bound, we will show that four colors are not enough. For a contradiction, assume that there exists a 4-locating coloring  $c$  on  $O_n$  for  $n \geq 4$ . We assign  $\{c(u_i), c(v_i), c(w_i), c(u_{i+1})\} = \{1, 2, 3, 4\}$ , where  $c(v_i) \neq c(u_{i+1})$  because  $d(v_i, x) = d(u_{i+1}, x)$ ,  $x \in \{u_i, v_i\}$ . Observe that, on  $O_n$  for  $n \geq 4$ , there are  $n$  vertices  $u_i$  whose degree is 5. As a result, at least two vertices  $w_k, w_l, k \neq l$  have the same color codes, which is a contradiction. Therefore,  $\chi_L(O_n) \geq 5$ , for  $n \geq 4$ .

To show the upper bound for the locating-chromatic number of origami graphs  $O_n$  for  $n \geq 4$ , let us differentiate some subcases.

**Subcase 1.** (Odd  $n$ ), for  $\lceil \frac{n}{2} \rceil$  odd,  $n \geq 5$

Let  $c$  be a coloring of origami graph  $O_n$ ,  $\lceil \frac{n}{2} \rceil$  odd, and  $n \geq 5$ ; we make the partition  $\Pi$  of  $V(O_n)$ :

$$\begin{aligned} C_1 &= \{w_i | 1 \leq i \leq n\}; \\ C_2 &= \{u_i | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n-1\}; \end{aligned}$$

$C_3 = \{u_i \mid \text{for even } i, 2 \leq i \leq \lceil \frac{n}{2} \rceil - 1\} \cup \{u_i \mid \text{for even } i, \lceil \frac{n}{2} \rceil + 3 \leq i \leq n - 1\} \cup \{v_i \mid \text{for odd } i, 1 \leq i \leq n\};$

$C_4 = \{u_1\};$

$C_5 = \{u_{\lceil \frac{n}{2} \rceil + 1}\}.$

For  $\lceil \frac{n}{2} \rceil$  odd, the color codes of all the vertices of  $V(O_n)$  are:

$C_1 = \{w_i \mid 1 \leq i \leq n\}.$

For  $i = 1$ , we have:

$$c_{\Pi}(w_i) = (0, 2, 1, i, \lceil \frac{n}{2} \rceil - i + 1).$$

For  $2 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5$  we have:

$$c_{\Pi}(w_i) = (0, 1, 1, i, \lceil \frac{n}{2} \rceil - i + 1).$$

For  $i = \lceil \frac{n}{2} \rceil + 1$  we have:

$$c_{\Pi}(w_i) = (0, 1, 2, n - i + 1, i - \lceil \frac{n}{2} \rceil).$$

For  $\lceil \frac{n}{2} \rceil + 2 \leq i \leq n, n \geq 5$  we have:

$$c_{\Pi}(w_i) = (0, 1, 1, n - i + 1, i - \lceil \frac{n}{2} \rceil).$$

$C_2 = \{u_i \mid \text{for odd } i, 3 \leq i \leq n\} \cup \{v_i \mid \text{for even } i, 2 \leq i \leq n - 1\}.$

For  $i$  odd,  $3 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5$  we have:

$$c_{\Pi}(u_i) = (1, 0, 1, i - 1, \lceil \frac{n}{2} \rceil - i + 1).$$

For  $i$  odd,  $\lceil \frac{n}{2} \rceil + 2 \leq i \leq n, n \geq 5$  we have:

$$c_{\Pi}(u_i) = (1, 0, 1, n - i + 1, i - \lceil \frac{n}{2} \rceil - 1).$$

For  $i$  even,  $2 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 5$  we have:

$$c_{\Pi}(v_i) = (1, 0, 1, i, \lceil \frac{n}{2} \rceil - i + 2).$$

For  $i = \lceil \frac{n}{2} \rceil + 1$ , we have:

$$c_{\Pi}(v_i) = (1, 0, 3, n - i + 2, 1).$$

For  $i$  even,  $\lceil \frac{n}{2} \rceil + 3 \leq i \leq n - 1, n \geq 9$  we have:

$$c_{\Pi}(v_i) = (1, 0, 1, n - i + 2, i - \lceil \frac{n}{2} \rceil).$$

$C_3 = \{u_i \mid \text{for even } i, 2 \leq i \leq \lceil \frac{n}{2} \rceil - 1\} \cup \{u_i \mid \text{for even } i, \lceil \frac{n}{2} \rceil + 3 \leq i \leq n - 1\} \cup \{v_i \mid \text{for odd } i, 1 \leq i \leq n\}.$

For  $i = 1$ , we have:

$$c_{\Pi}(v_i) = (1, 2, 0, i, \lceil \frac{n}{2} \rceil).$$

For  $i$  odd,  $3 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5$  we have:

$$c_{\Pi}(v_i) = (1, 1, 0, i, \lceil \frac{n}{2} \rceil - i + 2).$$

For  $i$  odd,  $\lceil \frac{n}{2} \rceil + 2 \leq i \leq n, n \geq 9$  we have:

$$c_{\Pi}(v_i) = (1, 1, 0, n - i + 2, i - \lceil \frac{n}{2} \rceil).$$

For  $i$  even,  $2 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 5$  we have:

$$c_{\Pi}(u_i) = (1, 1, 0, i - 1, \lceil \frac{n}{2} \rceil - i + 1).$$

For  $i$  even,  $\lceil \frac{n}{2} \rceil + 3 \leq i \leq n - 1, n \geq 9$  we have:

$$c_{\Pi}(u_i) = (1, 1, 0, n - i + 1, i - \lceil \frac{n}{2} \rceil - 1).$$

For  $C_4 = \{u_1\}$ , we have:

$$c_{\Pi}(u_1) = (1, 1, 1, 0, \lceil \frac{n}{2} \rceil - 1).$$

For  $C_5 = \{u_{\lceil \frac{n}{2} \rceil + 1}\}$ , we have:

$$c_{\Pi}(u_{\lceil \frac{n}{2} \rceil + 1}) = (1, 1, 2, \lceil \frac{n}{2} \rceil - 1, 0).$$

Since for  $n$  odd all vertices have different color codes,  $c$  is a locating coloring of origami graphs  $O_n$ , so that  $\chi_L(O_n) \leq 5$ , for  $\lceil \frac{n}{2} \rceil$  odd,  $n \geq 5$ .

**Subcase 2.** (Odd  $n$ ), for  $\lceil \frac{n}{2} \rceil$  even,  $n \geq 7$ .

Let  $c$  be a coloring of origami graph  $O_n$ ,  $\lceil \frac{n}{2} \rceil$  even, and  $n \geq 7$ ; we make the partition  $\Pi$  of  $V(O_n)$  as follows:

$$C_1 = \{w_i | 1 \leq i \leq n\};$$

$$C_2 = \{u_i | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n - 1\};$$

$$C_3 = \{u_i | \text{for even } i, 2 \leq i \leq \lceil \frac{n}{2} \rceil - 2\} \cup \{u_i | \text{for even } i, \lceil \frac{n}{2} \rceil + 2 \leq i \leq n - 1\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n\};$$

$$C_4 = \{u_1\};$$

$$C_5 = \{u_{\lceil \frac{n}{2} \rceil}\}.$$

For  $\lceil \frac{n}{2} \rceil$  even, the color codes of all the vertices of  $V(O_n)$  are:

$$C_1 = \{w_i | 1 \leq i \leq n\}.$$

For  $i = 1$ , we have:

$$c_{\Pi}(w_1) = (0, 2, 1, i, \lceil \frac{n}{2} \rceil - i).$$

For  $2 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 7$  we have:

$$c_{\Pi}(w_i) = (0, 1, 1, i, \lceil \frac{n}{2} \rceil - i).$$

For  $i = \lceil \frac{n}{2} \rceil$ , we have:

$$c_{\Pi}(w_i) = (0, 1, 2, n - i + 1, i - \lceil \frac{n}{2} \rceil + 1).$$

For  $\lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7$  we have:

$$c_{\Pi}(w_i) = (0, 1, 1, n - i + 1, i - \lceil \frac{n}{2} \rceil + 1).$$

$$C_2 = \{u_i | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n - 1\}.$$

For  $i$  odd,  $3 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 7$  we have:

$$c_{\Pi}(u_i) = (1, 0, 1, i - 1, \lceil \frac{n}{2} \rceil - i).$$

For  $i$  odd,  $\lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7$  we have:

$$c_{\Pi}(u_i) = (1, 0, 1, n - i + 1, \lceil \frac{n}{2} \rceil).$$

For  $i$  even,  $2 \leq i \leq \lceil \frac{n}{2} \rceil - 2, n \geq 7$  we have:

$$c_{\Pi}(v_i) = (1, 0, 1, i, \lceil \frac{n}{2} \rceil - i + 1).$$

For  $i = \lceil \frac{n}{2} \rceil$ , we have:

$$c_{\Pi}(v_i) = (1, 0, 3, i, i - \lceil \frac{n}{2} \rceil + 1).$$

For  $i$  even,  $\lceil \frac{n}{2} \rceil + 3 \leq i \leq n - 1, n \geq 7$  we have:

$$c_{\Pi}(v_i) = (1, 0, 1, n - i + 2, i - \lceil \frac{n}{2} \rceil + 1).$$

$C_3 = \{u_i | \text{for even } i, 2 \leq i \leq \lceil \frac{n}{2} \rceil - 2\} \cup \{u_i | \text{for even } i, \lceil \frac{n}{2} \rceil + 2 \leq i \leq n - 1\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n\}$ .

For  $i = 1$  we have:

$$c_{\Pi}(v_i) = (1, 2, 0, i, \lceil \frac{n}{2} \rceil - i + 1).$$

For  $i$  odd,  $3 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 7$  we have:

$$c_{\Pi}(v_i) = (1, 1, 0, i, \lceil \frac{n}{2} \rceil - i + 1).$$

For  $i$  odd,  $\lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7$  we have:

$$c_{\Pi}(v_i) = (1, 1, 0, n - i + 2, i - \lceil \frac{n}{2} \rceil + 1).$$

For  $i$  even,  $2 \leq i \leq \lceil \frac{n}{2} \rceil - 2, n \geq 7$  we have:

$$c_{\Pi}(u_i) = (1, 1, 0, i - 1, \lceil \frac{n}{2} \rceil - i).$$

For  $i$  even,  $\lceil \frac{n}{2} \rceil + 2 \leq i \leq n, n \geq 7$  we have:

$$c_{\Pi}(u_i) = (1, 1, 0, n - i + 1, i - \lceil \frac{n}{2} \rceil).$$

$C_4 = \{u_1\}$ , we have:

$$c_{\Pi}(u_1) = (1, 1, 1, 0, \lceil \frac{n}{2} \rceil - 1).$$

$C_5 = \{u_{\lceil \frac{n}{2} \rceil}\}$ , we have:

$$c_{\Pi}(u_{\lceil \frac{n}{2} \rceil}) = (1, 1, 2, \lceil \frac{n}{2} \rceil - 1, 0).$$

Since for  $n$  odd all vertices have different color codes,  $c$  is a locating coloring of origami graphs  $O_n$ , so that  $\chi_L(O_n) \leq 5$ , for  $\lceil \frac{n}{2} \rceil$  even,  $n \geq 7$ .

**6** **Subcase 3.** (even  $n$ ), for  $\frac{n}{2}$  odd,  $n \geq 6$ .

Let  $c$  be a coloring of origami graph  $O_n$ ,  $\frac{n}{2}$  odd, and  $n \geq 6$ ; we make the partition  $\Pi$  of  $V(O_n)$ :

$$C_1 = \{w_i | 1 \leq i \leq \frac{n}{2} - 1\} \cup \{w_i | \frac{n}{2} + 1 \leq i \leq n\};$$

$$C_2 = \{u_i | \text{for odd } i, 3 \leq i \leq n - 1\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n\};$$

$$C_3 = \{u_i | \text{for even } i, 2 \leq i \leq n\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n - 1\};$$

$$C_4 = \{u_1\};$$

$$C_5 = \{w_{\frac{n}{2}}\}.$$

**5**

For  $\frac{n}{2}$  odd, the color codes of all the vertices of  $V(O_n)$  are:

$$C_1 = \{w_i | 1 \leq i \leq \frac{n}{2} - 1\} \cup \{w_i | \frac{n}{2} + 1 \leq i \leq n\}.$$

For  $i = 1$ , we have:

$$c_{\Pi}(w_i) = (0, 2, 1, i, \frac{n}{2} - i + 1).$$

For  $2 \leq i \leq \frac{n}{2} - 1, n \geq 6$  we have:

$$c_{\Pi}(w_i) = (0, 1, 1, i, \frac{n}{2} - i + 1).$$

For  $\frac{n}{2} + 1 \leq i \leq n, n \geq 6$  we have:

$$c_{\Pi}(w_i) = (0, 1, 1, n - i + 1, i - \frac{n}{2} + 1).$$

$$C_2 = \{u_i | \text{for odd } i, 3 \leq i \leq n - 1\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n\}.$$

For  $i$  odd,  $3 \leq i \leq \frac{n}{2}, n \geq 6$  we have:

**16**

$$c_{\Pi}(u_i) = (1, 0, 1, i - 1, \frac{n}{2} - i + 1).$$

For  $i$  odd,  $\frac{n}{2} + 2 \leq i \leq n - 1, n \geq 6$  we have:

$$c_{\Pi}(u_i) = (1, 0, 1, n - i + 1, i - \frac{n}{2}).$$

For  $i$  even,  $2 \leq i \leq \frac{n}{2} - 1, n \geq 6$  we have:

$$c_{\Pi}(v_i) = (1, 0, 1, i, \frac{n}{2} - i + 2).$$

For  $i$  even,  $\frac{n}{2} + 1 \leq i \leq n - 1, n \geq 6$  we have:

$$c_{\Pi}(v_i) = (1, 0, 1, n - i + 2, i - \frac{n}{2} + 1).$$

$$C_3 = \{u_i | \text{for even } i, 2 \leq i \leq \frac{n}{2}\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n - 1\}.$$

For  $i = 1$ , we have:

$$c_{\Pi}(v_i) = (1, 3, 0, i, \frac{n}{2} - i + 2).$$

For  $i$  odd,  $3 \leq i \leq \frac{n}{2} - 2, n \geq 10$  we have:

$$c_{\Pi}(v_i) = (1, 1, 0, i, \frac{n}{2} - i + 2)$$

For  $i = \frac{n}{2}$ , we have:

$$c_{\Pi}(v_i) = (2, 1, 0, i, 1).$$

For  $i$  odd,  $\frac{n}{2} + 2 \leq i \leq n - 1, n \geq 6$  we have:

$$c_{\Pi}(v_i) = (1, 1, 0, n - i + 2, i - \frac{n}{2} + 1).$$

For  $i$  even,  $2 \leq i \leq \frac{n}{2} - 1, n \geq 6$  we have:

$$c_{\Pi}(u_i) = (1, 1, 0, i - 1, \frac{n}{2} - i + 1).$$

For  $i$  even,  $\frac{n}{2} + 1 \leq i \leq n, n \geq 6$  we have:

$$c_{\Pi}(u_i) = (1, 1, 0, n - i + 1, i - \frac{n}{2}).$$

For  $C_4 = \{u_1\}$ , we have:

$$c_{\Pi}(u_1) = (1, 2, 1, 0, \frac{n}{2} - i + 1).$$

For  $C_5 = \{w_{\frac{n}{2}}\}$ , we have:

$$c_{\Pi}(w_{\frac{n}{2}}) = (2, 1, 1, \frac{n}{2}, 0).$$

Since for  $n$  even all vertices have different color codes,  $c$  is a locating coloring of origami graphs  $O_n$ , so that  $\chi_L(O_n) \leq 5$ , for  $\frac{n}{2}$  odd,  $n \geq 6$ .

**Subcase 4.** (even  $n$ ), for  $\frac{n}{2}$  even,  $n \geq 4$ .

Let  $c$  be a coloring of origami graph  $O_n$ ,  $\frac{n}{2}$  even, and  $n \geq 4$ ; we make the partition  $\Pi$  of  $V(O_n)$  as follows:

$$C_1 = \{w_i | 1 \leq i \leq \frac{n}{2}\} \cup \{w_i | \frac{n}{2} + 2 \leq i \leq n\};$$

$$C_2 = \{u_i | \text{for odd } i, 1 \leq i \leq n - 1\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n\};$$

$$C_3 = \{u_i | \text{for even } i, 2 \leq i \leq \frac{n}{2}\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n - 1\};$$

$$C_4 = \{u_1\};$$

$$C_5 = \{w_{\frac{n}{2}+1}\}.$$

For  $\frac{n}{2}$  even, the color codes of all the vertices of  $V(O_n)$  are:

$$C_1 = \{w_i | 1 \leq i \leq \frac{n}{2}\} \cup \{w_i | \frac{n}{2} + 2 \leq i \leq n\}.$$

For  $i = 1$  we have:

$$c_{\Pi}(w_i) = (0, 2, 1, i, \frac{n}{2} - i + 2).$$

For  $2 \leq i \leq \frac{n}{2}, n \geq 4$  we have:

$$c_{\Pi}(w_i) = (0, 1, 1, i, \frac{n}{2} - i + 2).$$



For  $\frac{n}{2} + 2 \leq i \leq n, n \geq 4$  we have:

$$c_{\Pi}(w_i) = (0, 1, 1, n - i + 1, i - \frac{n}{2}).$$

$C_2 = \{u_i \mid \text{for odd } i, 3 \leq i \leq n - 1\} \cup \{v_i \mid \text{for even } i, 2 \leq i \leq n\}$ .

For  $i$  odd,  $3 \leq i \leq \frac{n}{2} + 1, n \geq 8$  we have:

$$c_{\Pi}(u_i) = (1, 0, 1, i - 1, \frac{n}{2} - i + 2).$$

For  $i$  odd,  $\frac{n}{2} + 3 \leq i \leq n - 1, n \geq 8$  we have:

$$c_{\Pi}(u_i) = (1, 0, 1, n - i + 1, i - \frac{n}{2} - 1).$$

For  $i$  even,  $2 \leq i \leq \frac{n}{2}, n \geq 4$  we have:

$$c_{\Pi}(v_i) = (1, 0, 1, i, \frac{n}{2} - i + 3).$$

For  $i$  even,  $\frac{n}{2} + 2 \leq i \leq n, n \geq 8$  we have:

$$c_{\Pi}(v_i) = (1, 0, 1, n - i + 2, i - \frac{n}{2}).$$

$C_3 = \{u_i \mid \text{for even } i, 2 \leq i \leq n\} \cup \{v_i \mid \text{for odd } i, 1 \leq i \leq n - 1\}$ .

For  $i = 1$ , we have:

$$c_{\Pi}(v_i) = (1, 3, 0, 1, \frac{n}{2} + 1).$$

For  $i$  odd,  $3 \leq i \leq \frac{n}{2} - 1, n \geq 8$  we have:

$$c_{\Pi}(v_i) = (1, 1, 0, i, \frac{n}{2} - i + 3).$$

For  $i = \frac{n}{2} + 1$ , we have:

$$c_{\Pi}(v_i) = (2, 1, 0, i, 1).$$

For  $i$  odd,  $\frac{n}{2} + 3 \leq i \leq n - 1, n \geq 8$  we have:

$$c_{\Pi}(v_i) = (1, 1, 0, n - i + 2, i - \frac{n}{2}).$$

For  $i$  even,  $2 \leq i \leq \frac{n}{2}, n \geq 4$  we have:

$$c_{\Pi}(u_i) = (1, 1, 0, i - 1, \frac{n}{2}).$$

For  $i$  even,  $\frac{n}{2} + 2 \leq i \leq n, n \geq 8$  we have:

$$c_{\Pi}(u_i) = (1, 1, 0, n - i + 1, i - \frac{n}{2} - 1).$$

For  $C_4 = \{u_1\}$ , we have:

$$c_{\Pi}(u_1) = (1, 2, 1, 0, \frac{n}{2}).$$

For  $C_5 = \{w_{\frac{n}{2}}\}$ , we have:

$$c_{\Pi}(w_{\frac{n}{2}}) = (2, 1, 1, \frac{n}{2}, 0).$$

Since for  $n$  even all vertices have different color codes,  $c_{\Pi}$  is a locating coloring of origami graphs  $O_n$ , so that  $\chi_L(O_n) \leq 5$ , for  $\frac{n}{2}$  even,  $n \geq 4$ .  $c_{\Pi}$  completes the proof of Theorem 1.  $\square$

Note that Figure 1 is an example locating coloring for origami graph  $O_5$ .

2.2. Locating-Chromatic Number for Subdivision Outer Edge of Origami Graphs

**Theorem 2.** Let  $O_n^*$  be a subdivision outer edge of origami graphs for  $n \geq 3$ . Then the locating-

$$\text{chromatic number of } O_n^*, \chi_L(O_n^*) = \begin{cases} 4, & \text{for } n = 3 \\ 5, & \text{otherwise.} \end{cases}$$

**Proof.** Let  $O_n^*, n \geq 3$  be a subdivision of an origami graph;  $O_n^*$  is a graph with  $V(O_n^*) = \{u_i, v_i, x_i, w_i : i \in \{1, \dots, n\}\}$  and  $E(O_n^*) = \{u_i w_i, u_i v_i, v_i x_i, x_i w_i : i \in \{1, \dots, n\}\} \cup \{u_i u_{i+1}, w_i w_{i+1} : i \in \{1, \dots, n-1\}\} \cup \{u_n u_1, w_n w_1\}$ . Next, to prove the theorem, we consider the following two cases:

**Case A.**  $\chi_L(O_3^*) = 4$

First, we determine the lower bound of  $\chi_L(O_3^*)$ .

Without loss of generality, we assign  $A = \{c(u_i), c(v_i), c(x_i), c(w_i), c(u_{i+1})\} = \{1, 2, 3\}$ . Then, there are three dominant vertices in  $A$ , while we still have vertices on other  $A$  that must be colored. As a result, there will be two vertices with the same color codes. Thus,  $\chi_L(O_3^*) \geq 4$ .

Next, determine the upper bound of  $\chi_L(O_3^*) \leq 4$ . To show that 4 is an upper bound for the locating-chromatic number for a subdivision outer edge of origami graph  $O_3^*$ , we describe a locating coloring  $c$  using four colors as follows:

$$\begin{aligned} c(u_i) &= i, i = 1, 2, 3. \\ c(v_i) &= \begin{cases} 2, & \text{for } i = 1, 3 \\ 3, & \text{for } i = 2. \end{cases} \\ c(w_i) &= 4, i = 1, 2, 3. \\ c(x_i) &= i, i = 1, 2, 3. \end{aligned}$$

The coloring  $c$  will create the partition  $\Pi$  on  $V(O_3^*)$ . We shall show that the color codes of all vertices in  $O_3^*$  are different. We have:  $c_\Pi(u_1) = (0, 1, 1, 1)$ ;  $c_\Pi(u_2) = (1, 0, 1, 1)$ ;  $c_\Pi(u_3) = (1, 1, 0, 1)$ ;  $c_\Pi(v_1) = (1, 0, 2, 2)$ ;  $c_\Pi(v_2) = (2, 1, 0, 2)$ ;  $c_\Pi(v_3) = (2, 0, 1, 2)$ ;  $c_\Pi(w_1) = (1, 1, 2, 0)$ ;  $c_\Pi(w_2) = (2, 1, 1, 0)$ ;  $c_\Pi(w_3) = (1, 2, 1, 0)$ .  $c_\Pi(x_1) = (0, 1, 3, 1)$ ;  $c_\Pi(x_2) = (3, 0, 1, 1)$ ;  $c_\Pi(x_3) = (2, 1, 0, 1)$ . Since the color codes of all vertices  $O_3^*$  are different,  $c$  is a locating-chromatic coloring. Thus,  $\chi_L(O_3^*) \leq 4$ .

**Case B.**  $\chi_L(O_n^*) = 5$  for  $n \geq 4$

Since a subdivision of origami graphs  $O_n^*$  for  $n \geq 4$  is obtained by origami graph  $O_n$  with one added vertex in edge  $v_i w_i$ , we have  $\chi_L(O_n^*) \geq 5$  for  $n \geq 4$ . The addition of one vertex on the outside does not reduce the number of colors needed because the number of the sets  $B = \{c(u_i), c(v_i), c(w_i), c(u_{i+1})\}$  is the same.

To show the upper bound for the locating-chromatic number for a subdivision outer edge of origami graph  $O_n^*$  for  $n \geq 4$ , let us consider different subcases.

**Subcase a.** (odd  $n$ ), for  $\lceil \frac{n}{2} \rceil$  odd,  $n \geq 5$ .

Let  $c$  be a coloring for a subdivision outer edge of origami graph  $O_n^*$ , for  $\lceil \frac{n}{2} \rceil$  odd, and  $n \geq 5$ ; we make the partition  $\Pi$  of  $V(O_n^*)$ :

$$\begin{aligned} C_1 &= \{w_i | 1 \leq i \leq n\}; \\ C_2 &= \{u_i | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n-1\} \cup \{x_i | \text{for odd } i, 1 \leq i \leq n\}; \\ C_3 &= \{u_i | \text{for even } i, 2 \leq i \leq \lceil \frac{n}{2} \rceil - 1\} \cup \{u_i | \text{for even } i, \lceil \frac{n}{2} \rceil + 3 \leq i \leq n-1\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n\} \cup \{x_i | \text{for even } i, 2 \leq i \leq n-1\}; \\ C_4 &= \{u_1\}; \\ C_5 &= \{u_{\lceil \frac{n}{2} \rceil + 1}\}. \end{aligned}$$

For for  $\lceil \frac{n}{2} \rceil$  odd the color codes of all the vertices of  $V(O_n^*)$  are:

$$c_{\Pi}(u_i) = \begin{cases} 0, & \text{for the second component, odd } i, 3 \leq i \leq n, n \geq 5 \\ & \text{for the third component, even } i, 2 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 5 \\ & \text{for the third component, even } i, \lceil \frac{n}{2} \rceil + 3 \leq i \leq n - 1, n \geq 9 \\ & \text{for the fourth component, } i = 1 \\ 6, & \text{for the fifth component, } i = \lceil \frac{n}{2} \rceil + 1 \\ 2, & \text{for the third component, } i = \lceil \frac{n}{2} \rceil + 1 \\ i - 1, & \text{for the fourth component, } 2 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5 \\ n - i + 1, & \text{for the fourth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 5 \\ \lceil \frac{n}{2} \rceil - i, & \text{for the fifth component, } i = 1 \\ i - \lceil \frac{n}{2} \rceil - 1, & \text{for the fifth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 5 \\ \lceil \frac{n}{2} \rceil - i + 1, & \text{for the fifth component, } 2 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_i) = \begin{cases} 2, & \text{for the first component, } 1 \leq i \leq n, n \geq 5 \\ 0, & \text{for the second component, odd } i, 1 \leq i \leq n, n \geq 5 \\ & \text{for the third component, even } i, 2 \leq i \leq n - 1, n \geq 5 \\ i, & \text{for the fourth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5 \\ n - i + 2, & \text{for the fourth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 5 \\ \lceil \frac{n}{2} \rceil, & \text{for the fifth component, } i = 1 \\ \lceil \frac{n}{2} \rceil - i + 2, & \text{for the fifth component, } 2 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5 \\ i - \lceil \frac{n}{2} \rceil, & \text{for the fifth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_i) = \begin{cases} 0, & \text{for the first component, } 1 \leq i \leq n, n \geq 5 \\ 2, & \text{for the third component, } i = \lceil \frac{n}{2} \rceil \text{ and } i = n \\ \lceil \frac{n}{2} \rceil - i + 1, & \text{for the fifth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5 \\ i - \lceil \frac{n}{2} \rceil, & \text{for the fifth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 5 \\ i, & \text{for the fourth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5 \\ n - i + 1, & \text{for the fourth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(x_i) = \begin{cases} 0, & \text{for the second component, odd } i, 1 \leq i \leq n, n \geq 5 \\ & \text{for the third component, even } i, 2 \leq i \leq n - 1, n \geq 5 \\ i + 1, & \text{for the fourth component, } \lceil \frac{n}{2} \rceil \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5 \\ n - i + 2, & \text{for the fourth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 5 \\ \lceil \frac{n}{2} \rceil - i + 2, & \text{for the fifth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 5 \\ i - \lceil \frac{n}{2} \rceil - 1, & \text{for the fifth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

Since for  $n$  odd all vertices have different color codes,  $c$  is a locating coloring for subdivision of origami graph  $O_n^*$ , so that  $\chi_L(O_n^*) \leq 5$ , for  $\lceil \frac{n}{2} \rceil$  odd,  $n \geq 5$ .

**Subcase b.** (odd  $n$ ), for  $\lceil \frac{n}{2} \rceil$  even,  $n \geq 7$ .

Let  $c$  be a coloring for a subdivision outer edge of origami graph  $O_n^*$ , for  $\lceil \frac{n}{2} \rceil$  even, and  $n \geq 7$ ; we make the partition  $\Pi$  of  $V(O_n^*)$ :

$$C_1 = \{w_i | 1 \leq i \leq n\};$$

$$C_2 = \{u_i | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n - 1\} \cup \{x_i | \text{for odd } i, 1 \leq i \leq n\};$$

$$C_3 = \{u_i | \text{for even } i, 2 \leq i \leq \lceil \frac{n}{2} \rceil - 2\} \cup \{u_i | \text{for even } i, \lceil \frac{n}{2} \rceil + 2 \leq i \leq n - 1\} \cup \{v_i | \text{for}$$

odd  $i, 1 \leq i \leq n\} \cup \{x_i \mid \text{for even } i, 2 \leq i \leq n - 1\}$ ;  
 $C_4 = \{u_1\}$ ;  
 $C_5 = \{u_{\lceil \frac{n}{2} \rceil}\}$ .

For  $\lceil \frac{n}{2} \rceil$  even, the color codes of all the vertices of  $V(O_n^*)$  are:

$$c_{\Pi}(u_i) = \begin{cases} 0, & \text{for the second component, odd } i, 3 \leq i \leq n, n \geq 7 \\ & \text{for the third component, even } i, 2 \leq i \leq \lceil \frac{n}{2} \rceil - 2, n \geq 7 \\ & \text{for the third component, even } i, \lceil \frac{n}{2} \rceil + 2 \leq i \leq n - 1, n \geq 7 \\ & \text{for the fourth component, } i = 1 \\ & \text{for the fifth component, } i = \lceil \frac{n}{2} \rceil \\ 2, & \text{for the third component, } i = \lceil \frac{n}{2} \rceil \\ i - 1, & \text{for the fourth component, } 2 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 7 \\ n - i + 1, & \text{for the fourth component, odd } i, \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7 \\ \lceil \frac{n}{2} \rceil - 1, & \text{for the fourth component, } i = \lceil \frac{n}{2} \rceil \\ \lceil \frac{n}{2} \rceil - i, & \text{for the fifth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 7 \\ i - \lceil \frac{n}{2} \rceil, & \text{for the fifth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise .} \end{cases}$$

$$c_{\Pi}(v_i) = \begin{cases} 0, & \text{for the second component, even } i, 2 \leq i \leq n - 1, n \geq 7 \\ & \text{for the third component, odd } i, 1 \leq i \leq n, n \geq 7 \\ 2, & \text{for the first component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil \\ i, & \text{for the fourth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 7 \\ n - i + 2, & \text{for the fourth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7 \\ \lceil \frac{n}{2} \rceil - i + 1, & \text{for the fifth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 7 \\ i - \lceil \frac{n}{2} \rceil + 1, & \text{for the fifth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise .} \end{cases}$$

$$c_{\Pi}(w_i) = \begin{cases} 0, & \text{for the first component, } 1 \leq i \leq n, n \geq 7 \\ 2, & \text{for the third component, } i = \lceil \frac{n}{2} \rceil - 1 \text{ and } i = n \\ i, & \text{for the fourth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 7 \\ n - i + 1, & \text{for the fourth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7 \\ \lceil \frac{n}{2} \rceil - i, & \text{for the fifth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 7 \\ i - \lceil \frac{n}{2} \rceil + 1, & \text{for the fifth component, } \lceil \frac{n}{2} \rceil \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise .} \end{cases}$$

$$c_{\Pi}(x_i) = \begin{cases} 0, & \text{for the second component, odd } i, \lceil \frac{n}{2} \rceil \leq i \leq n, n \geq 7 \\ & \text{for the third component, even } i, 2 \leq i \leq n - 1, n \geq 7 \\ 1, & \text{for the fourth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil - 1, n \geq 7 \\ \lceil \frac{n}{2} \rceil + i - 2, & \text{for the fourth component, } \lceil \frac{n}{2} \rceil \leq i \leq n, n \geq 7 \\ \lceil \frac{n}{2} \rceil - i + 2, & \text{for the fifth component, } 1 \leq i \leq \lceil \frac{n}{2} \rceil, n \geq 7 \\ i - \lceil \frac{n}{2} \rceil + 2, & \text{for the fifth component, } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise .} \end{cases}$$

Since for  $n$  odd all vertices have different color codes,  $c$  is a locating coloring for a subdivision of the outer edge of origami graph  $O_n^*$ , so that  $\chi_L(O_n^*) \leq 5$ , for  $\lceil \frac{n}{2} \rceil$  even,  $n \geq 7$ .

**Subcase c.** (even  $n$ ), for  $\frac{n}{2}$  odd,  $n \geq 6$ .

Let  $c$  be a coloring for a subdivision outer edge of origami graph  $O_n^*$ , for  $\frac{n}{2}$  odd, and  $n \geq 6$ ; we make the partition  $\Pi$  of  $V(O_n^*)$ :

$$C_1 = \{w_i | 1 \leq i \leq \frac{n}{2} - 1\} \cup \{w_i | \frac{n}{2} + 1 \leq i \leq n\};$$

$$C_2 = \{u_i | \text{for odd } i, 3 \leq i \leq n - 1\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n\} \cup \{x_i | \text{for odd } i, 1 \leq i \leq n - 1\};$$

$$C_3 = \{u_i | \text{for even } i, 2 \leq i \leq n\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n - 1\} \cup \{x_i | \text{for even } i, 2 \leq i \leq n\};$$

$$C_4 = \{u_1\};$$

$$C_5 = \{w_{\frac{n}{2}}\}.$$

For  $\frac{n}{2}$  odd, the color codes of all the vertices of  $V(O_n^*)$  are:

$$c_{\Pi}(u_i) = \begin{cases} 0, & \text{for the second component, odd } i, 3 \leq i \leq n - 1, n \geq 6 \\ & \text{for the third component, even } i, 2 \leq i \leq n, n \geq 6 \\ & \text{for the fourth component, } i = 1 \\ 2, & \text{for the second component, } i = 1 \\ i - 1, & \text{for the fourth component, } 2 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 1, & \text{for the fourth component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ \frac{n}{2} - i + 1, & \text{for the fifth component, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ i - \frac{n}{2}, & \text{for the fifth component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_i) = \begin{cases} 2, & \text{for the first component, } 1 \leq i \leq n, n \geq 6 \\ 0, & \text{for the second component, even } i, 2 \leq i \leq n, n \geq 6 \\ & \text{for the third component, odd } i, 1 \leq i \leq n - 1, n \geq 6 \\ i, & \text{for the fourth component, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 2, & \text{for the fourth component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ \frac{n}{2} - i + 2, & \text{for the fifth component, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ i - \frac{n}{2} + 1, & \text{for component, fifth component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_i) = \begin{cases} 0, & \text{for the first component, } 1 \leq i \leq \frac{n}{2} - 1, n \geq 6 \\ & \text{for the first component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ & \text{for the fifth component, } i = \frac{n}{2} \\ 2, & \text{for the first component, } i = \frac{n}{2} \\ & \text{for the second component, } i = n \\ i, & \text{for the fourth component, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 1, & \text{for the fourth component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ \frac{n}{2} - i + 1, & \text{for the fifth component, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ i - \frac{n}{2} + 1, & \text{for the fifth component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(x_i) = \begin{cases} 0, & \text{for the second component, odd } i, 1 \leq i \leq n-1, n \geq 6 \\ & \text{for the third component, even } i, 2 \leq i \leq n, n \geq 6 \\ i+1, & \text{for the fourth component, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ n-i+2, & \text{for the fourth component, } \frac{n}{2}+1 \leq i \leq n, n \geq 6 \\ \frac{n}{2}-i+2, & \text{for the fifth component, } 1 \leq i \leq \frac{n}{2}-1, n \geq 6 \\ i-\frac{n}{2}+2, & \text{for the fifth component, } \frac{n}{2}+1 \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

Since for  $n$  even all vertices have different color codes,  $c$  is a locating coloring for a subdivision of the outer edge of origami graph  $O_n^*$ , so that  $\chi_L(O_n^*) \leq 5$ , for  $\frac{n}{2}$  odd,  $n \geq 6$ .

**Subcase d.** (even  $n$ ), for  $\frac{n}{2}$  even,  $n \geq 4$ .

Let  $c$  be a coloring of subdivision origami graph  $O_n^*$ , for  $\frac{n}{2}$  even, and  $n \geq 4$ ; we make the partition  $\Pi$  of  $V(O_n^*)$ :

$$\begin{aligned} C_1 &= \{w_i | 1 \leq i \leq \frac{n}{2}\} \cup \{w_i | \frac{n}{2} + 2 \leq i \leq n\}; \\ C_2 &= \{u_i | \text{for odd } i, 3 \leq i \leq n-1\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n\} \cup \{x_i | \text{for odd } i, 1 \leq i \leq n-1\}; \\ C_3 &= \{u_i | \text{for even } i, 2 \leq i \leq n\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n-1\} \cup \{x_i | \text{for even } i, 2 \leq i \leq n\}; \\ C_4 &= \{u_1\}; \\ C_5 &= \{w_{\frac{n}{2}+1}\} \end{aligned}$$

For  $\frac{n}{2}$  even the color codes of all the vertices of  $V(O_n^*)$  are:

$$c_{\Pi}(u_i) = \begin{cases} 0, & \text{for the second component, odd } i, 3 \leq i \leq n-1, n \geq 4 \\ & \text{for the third component, even } i, 2 \leq i \leq n, n \geq 4 \\ & \text{for the fourth component, } i = 1 \\ i-1, & \text{for the fourth component, } 2 \leq i \leq \frac{n}{2} + 1, n \geq 4 \\ n-i+1, & \text{for the fourth component, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ \frac{n}{2}, & \text{for the fifth component, } i = 1 \\ \frac{n}{2} - i + 2, & \text{for the fifth component, } 2 \leq i \leq \frac{n}{2} + 1, n \geq 4 \\ i - \frac{n}{2} - 1, & \text{for the fifth component, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_i) = \begin{cases} 2, & \text{for the first component, } 1 \leq i \leq n, n \geq 4 \\ 0, & \text{for the second component, even } i, 2 \leq i \leq n, n \geq 4 \\ & \text{for the third component, odd } i, 1 \leq i \leq n-1, n \geq 4 \\ n-i+2, & \text{for the fourth component, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ \frac{n}{2} + i, & \text{for the fifth component, } i = 1 \\ \frac{n}{2} - i + 3, & \text{for the fifth component, } 2 \leq i \leq \frac{n}{2} + 1, n \geq 4 \\ i - \frac{n}{2}, & \text{for the fifth component, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_i) = \begin{cases} 0, & \text{for the first component, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ & \text{for the first component, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ & \text{for the fifth component, } i = \frac{n}{2} + 1 \\ 2, & \text{for the first component, } i = \frac{n}{2} + 1 \\ & \text{for the second component, } i = n \\ i, & \text{for the fourth component, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ n - i + 1, & \text{for the fourth component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ \frac{n}{2} - i + 2, & \text{for the fifth component, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ i - \frac{n}{2}, & \text{for the fifth component, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(x_i) = \begin{cases} 0, & \text{for the second component, odd } i, 1 \leq i \leq n - 1, n \geq 4 \\ & \text{for the third component, even } i, 2 \leq i \leq n, n \geq 4 \\ i + 1, & \text{for the fourth component, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 2, & \text{for the fourth component, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ \frac{n}{2} - i + 3, & \text{for the fifth component, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ i - \frac{n}{2} + 1, & \text{for the fifth component, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

Since for  $n$  even all vertices have different color codes,  $c$  is a locating coloring [6] a subdivision outer edge of origami graph  $O_n^*$ , so that  $\chi_L(O_n^*) \leq 5$ , for  $\frac{n}{2}$  even,  $n \geq 4$ . This completes the proof of Theorem 2.  $\square$

Note that Figure 2 is an example locating coloring for a subdivision of the outer edge of origami graph  $O_5^*$ .

### 3. Conclusions

The proving steps of the two theorems we gave earlier show that the locating-chromatic number of origami graphs  $O_n$ ,  $\chi_L(O_n)$  is 4 for  $n = 3$  and 5 for  $n \geq 4$ ; the same result holds for a subdivision of the outer edge of origami graph  $O_n^*$ . This research can be continued so as to determine the locating-chromatic number for some certain operations of origami graphs.

**Author Contributions:** Conceptualization, A.I. and A.A.; methodology, L.Z. and A.A.; software, A.I. and K.M.; validation, L.Z., A.A. and A.I.; formal analysis, L.Z.; investigation, A.I. and A.A.; resources, A.I. and A.A.; data curation, A.A. and L.Z.; writing—original draft preparation, A.I. and A.A.; writing—review and editing, A.I., A.A., L.Z. and K.M.; visualization, A.I. and K.M.; supervision, A.A. and L.Z.; project administration, A.A.; funding acquisition, A.A. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** The authors would like to thank the [12] ctorate of Research and Community Services at Kemendikbud RI for funding this research and the Head of the Institute of Research and Community Services of Lampung University, who supported this research.

**Conflicts of Interest:** The authors confirm that they have no conflict of interest to declare for this publication.

### References

1. Chartrand, G.; Zhang, P.; Salehi, E. On the partition dimension of a graph. *Congr. Numer.* **1998**, *131*, 55–66. [CrossRef]
2. Chartrand, G.; Zhang, P.; Salehi, E. The Partition Dimension of a Graph. *Aequ. Math.* **2000**, *55*, 45–54. [CrossRef]
3. Khuller, S.; Raghavachari, B.; Rosenfeld, A. Landmarks in graphs, Discrete Applied Mathematics. *Discrete Appl. Math.* **1996**, *70*, 217–229. [CrossRef]
4. Saenpholphat, V.; Zhang, P. Conditional resolvability: A survey. *Int. J. Math. Math. Sci.* **2004**, *38*, 1997–2017. [CrossRef]



5. Johnson, M.M. Structure activity maps for visualizing the graph variables arising in drug design. *J. Biopharm. Stat.* **1993**, *3*, 203–236. [[CrossRef](#)]
6. Chartrand, G.; Zhang, P. The theory and applications of resolvability in graphs: A survey. *Congr. Numer.* **2003**, *160*, 47–68.
7. Chartrand, G.; Erwin, D.; Henning, M.A.; Slater, P.J.; Zhang, P. The locating-chromatic number of a graph. *Bull. Inst. Comb. Appl.* **2002**, *36*, 89–101.
8. Chartrand, G.; Erwin, D.; Henning, M.A.; Slater, P.J.; Zhang, P. Graf of order  $n$  with locating-chromatic number  $n - 1$ . *Discret. Math.* **2003**, *269*, 65–79. [[CrossRef](#)]
9. Baskoro, E.T.; Purwasih, I.A. The locating-chromatic number for corona product of graphs. *Shoutheast Asian J. Sci.* **2012**, *1*, 124–134.
10. Asmiati, A. The Locating-Chromatic Number of Non-Homogeneous Amalgamation of Stars. *Far East J. Math. Sci.* **2014**, *93*, 89–96.
11. Welyyanti, D.; Simanjuntak, R.; Uttungadewa, R.S.; Baskoro, E.T. The locating-chromatic number of disconnected graphs. *Far East J. Math. Sci.* **2014**, *94*, 169–182.
12. Welyyanti, D.; Simanjuntak, R.; Uttungadewa, R.S.; Baskoro, E.T. On locating-chromatic number for graphs with dominant vertices. *Procedia Comput. Sci.* **2015**, *74*, 89–92. [[CrossRef](#)]
13. Welyyanti, D.; Simanjuntak, R.; Uttungadewa, R.S.; Baskoro, E.T. Locating-chromatic number for a graph of two components. *AIP Conf. Proc.* **2017**, *1707*, 20–24.
14. Ghanem, M.; Al-Ezeh, H.; Dabbour, A. Locating Chromatic Number of Powers of Paths and Cycles. *Symmetry* **2019**, *11*, 389. [[CrossRef](#)]
15. Sudhakar, S.; Francis, S.; Balaji, V. Odd mean labeling for two star graph. *Appl. Math. Nonlinear Sci.* **2017**, *2*, 195–200. [[CrossRef](#)]
16. Asmiati; Wamilliana, W.; Defriadi; Yulianti, L. On some Petersen graphs having locating chromatic number four or five. *Far East J. Math. Sci.* **2017**, *102*, 769–778. [[CrossRef](#)]
17. Asmiati, A.; Yana, I.K.S.G.; Yulianti, L. On the locating-chromatic number of certain barbell graphs. *Int. J. Math. Math. Sci.* **2018**, *5*. [[CrossRef](#)]
18. Syofyan, D.K.; Baskoro, E.T.; Assiyatun, H. On the locating-chromatic number of homogeneous lobsters. *AKCE Int. J. Graphs Comb.* **2013**, *10*, 245–252.
19. Asmiati, A. On the locating-chromatic numbers of non-homogeneous caterpillars and firecracker graphs. *Far East J. Math. Sci.* **2016**, *100*, 1305–1316. [[CrossRef](#)]
20. Irawan, A.; Asmiati, A. The locating-chromatic number of subdivision firecrackers graphs. *Int. Math. Forum* **2018**, *13*, 485–492. [[CrossRef](#)]
21. Irawan, A.; Asmiati, A.; Suharsono, S.; Muludi, K.; Zakaria, L. Certain operation of generalized Petersen graphs having locating-chromatic number five. *Adv. Appl. Discret. Math.* **2020**, *22*, 83–97. [[CrossRef](#)]
22. Behtoei, A.; Anbarloei, M. The locating chromatic number of the join of graphs. *Bull. Iran. Math. Soc.* **2014**, *40*, 1491–1504.
23. Nabila, S.; Salman, A.N.M. The rainbow connection number of origami graphs and pizza graphs. *Procedia Comput. Sci.* **2014**, *74*, 162–167. [[CrossRef](#)]



# The Locating-Chromatic Number of Origami Graphs

---

ORIGINALITY REPORT

---

# 16%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

- |   |   |               |
|---|---|---------------|
| 1 | <a href="#">docplayer.net</a><br>Internet   | 88 words — 1% |
| 2 | Mohammad Sadegh Es-haghi, Mohsen Abbaspour, Hamidreza Abbasianjahromi, Stefano Mariani. "Machine Learning-Based Prediction of the Seismic Bearing Capacity of a Shallow Strip Footing over a Void in Heterogeneous Soils", Algorithms, 2021<br>Crossref | 75 words — 1% |
| 3 | <a href="#">repo.itera.ac.id</a><br>Internet  | 60 words — 1% |
| 4 | <a href="#">jims-a.org</a><br>Internet  | 55 words — 1% |
| 5 | Muhammed Bello, Nor Muhainiah Mohd Ali. "Group coloring via its geometric structures", AIP Publishing, 2021<br>Crossref   | 49 words — 1% |
| 6 | <a href="#">link.springer.com</a><br>Internet   | 49 words — 1% |
| 7 | B. Rajan, K. T. Sonia, M. Chris Monica. "Conditional Resolvability of Honeycomb and Hexagonal Networks", Mathematics in Computer Science, 2011<br>Crossref  | 47 words — 1% |

8	Edy Tri Baskoro, Devi Imulia Dian Primaskun. "Improved algorithm for the locating-chromatic number of trees", Theoretical Computer Science, 2020 Crossref	42 words — 1%
9	JIMMIE D. LAWSON. "Domains, integration and 'positive analysis'", Mathematical Structures in Computer Science, 2004 Crossref	40 words — 1%
10	connection.ebscohost.com Internet	29 words — < 1%
11	upcommons.upc.edu Internet	29 words — < 1%
12	Notiragayu, L Zakaria. "The Implementation of Digital Text Coding Algorithm Through A Three Dimensional Mapping Derived From Generalized $\Delta\Delta$ -mKdV Equation Using Mathematica", Journal of Physics: Conference Series, 2019 Crossref	26 words — < 1%
13	Shi-Yao Zhu. "Non-classical Features in the Three-level Jaynes-Cummings Model", Journal of Modern Optics, 1989 Crossref	24 words — < 1%
14	www.mysciencework.com Internet	24 words — < 1%
15	g2a2.imm.uran.ru Internet	23 words — < 1%
16	Leon A. Feldman. "Commentary on the Song of Songs", Brill, 1970 Crossref	20 words — < 1%

- 17 [mdpi-res.com](https://mdpi-res.com)  
Internet 20 words — < 1%
- 
- 18 [repository.its.ac.id](https://repository.its.ac.id)  
Internet 19 words — < 1%
- 
- 19 Hyu-Dae Kim, Dong-Ho Cho. "Enhanced power-saving mechanism for broadcast and multicast service in WLAN", IEEE Communications Letters, 2005  
Crossref 17 words — < 1%
- 
- 20 [www.SciRP.org](https://www.SciRP.org)  
Internet 17 words — < 1%
- 
- 21 IMRAN, Muhammad, Syed Ahtsham ul Haq BOKHARY, Ali AHMAD, and Andrea SEMANIČOVÁ-FEŇOVČÍKOVÁ. "On classes of regular graphs with constant metric dimension", Acta Mathematica Scientia, 2013.  
Crossref 16 words — < 1%
- 
- 22 Chartrand, . "Bounds for the Chromatic Number", Discrete Mathematics and Its Applications, 2008.  
Crossref 15 words — < 1%
- 
- 23 PAVEL KURASOV, SERGUEI NABOKO. "Wigner-von Neumann perturbations of a periodic potential: spectral singularities in bands", Mathematical Proceedings of the Cambridge Philosophical Society, 2007  
Crossref 15 words — < 1%
- 
- 24 Omar Salah, Abdulrahim Shamayleh, Shayok Mukhopadhyay. "Energy Management of a Multi-Source Power System", Algorithms, 2021  
Crossref 14 words — < 1%
- 
- 25 Dantas, S.. "Extremal graphs for the list-coloring version of a theorem of Nordhaus and Gaddum", 12 words — < 1%

---

26 H.B. Macêdo Filho, S. Dantas, R.C.S. Machado, C.M.H. Figueiredo. "Biclique-colouring verification complexity and biclique-colouring power graphs", *Discrete Applied Mathematics*, 2015 11 words — < 1%  
Crossref

---

27 [emis.maths.adelaide.edu.au](http://emis.maths.adelaide.edu.au) 11 words — < 1%  
Internet

---

28 [inwpent5.ugent.be](http://inwpent5.ugent.be) 11 words — < 1%  
Internet

---

29 D. Fitriani, A.N.M. Salman. "Rainbow connection number of amalgamation of some graphs", *AKCE International Journal of Graphs and Combinatorics*, 2020 10 words — < 1%  
Crossref

---

30 Edmond W. H. Lee. "Hereditarily finitely based monoids of extensive transformations", *Algebra Universalis*, 08/28/2009 10 words — < 1%  
Crossref

---

31 Haryeni, Debi Oktia, and Edy Tri Baskoro. "Partition Dimension of Some Classes of Homogenous Disconnected Graphs", *Procedia Computer Science*, 2015. 10 words — < 1%  
Crossref

---

32 A. Estrada-Moreno, C. García-Gómez, Y. Ramírez-Cruz, J. A. Rodríguez-Velázquez. "The Simultaneous Strong Metric Dimension of Graph Families", *Bulletin of the Malaysian Mathematical Sciences Society*, 2015 9 words — < 1%  
Crossref

---

33 Ismael G. Yero, Alejandro Estrada-Moreno, Juan A. Rodríguez-Velázquez. "Computing the k -metric dimension of graphs", Applied Mathematics and Computation, 2017 9 words — < 1%  
Crossref

---

34 [digilib.unila.ac.id](http://digilib.unila.ac.id) 9 words — < 1%  
Internet

---

35 A. Mahalanobis, R.R. Muise, S.R. Stanfill, A. Van Nevel. "Design and application of quadratic correlation filters for target detection", IEEE Transactions on Aerospace and Electronic Systems, 2004 8 words — < 1%  
Crossref

---

36 Sridhar Rajagopalan, Vijay V. Vazirani. "Primal-Dual RNC Approximation Algorithms for Set Cover and Covering Integer Programs", SIAM Journal on Computing, 1998 8 words — < 1%  
Crossref

---

37 [media.neliti.com](http://media.neliti.com) 8 words — < 1%  
Internet

---

38 [www.univ-orleans.fr](http://www.univ-orleans.fr) 8 words — < 1%  
Internet

---

39 [yujor.fon.rs](http://yujor.fon.rs) 8 words — < 1%  
Internet

---

40 Alazemi, H.M.K.. "Stochastic modelling and analysis of 802.11 DCF with heterogeneous non-saturated nodes", Computer Communications, 20071210 7 words — < 1%  
Crossref

---

41 Carmen Hernando, Mercè Mora, Ignacio M. Pelayo, Liliana Alcón, Marisa Gutierrez. "Neighbor-locating 7 words — < 1%

coloring: graph operations and extremal cardinalities",  
Electronic Notes in Discrete Mathematics, 2018

Crossref

---

42 Hadjidimos, A.. "More on modifications and improvements of classical iterative schemes for M-matrices", Linear Algebra and Its Applications, 20030501 7 words — < 1%

Crossref

---

43 Varaporn Saenpholphat, Ping Zhang. "Conditional resolvability in graphs: a survey", International Journal of Mathematics and Mathematical Sciences, 2004 6 words — < 1%

Crossref

---

EXCLUDE QUOTES ON  
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF