

# Subdivision of Certain Barbell Operation of Origami Graphs has Locating- Chromatic Number Five

*By* Kurnia Muludi

# Subdivision of Certain Barbell Operation of Origami Graphs has Locating-Chromatic Number Five

Agus Irawan<sup>1\*</sup>, Asmiati<sup>2</sup>, La Zakaria<sup>2</sup>, Kurnia Muludi<sup>3</sup> and Bernadhita Herindri Samodra Utami<sup>1</sup>,

<sup>1</sup>Information system, STMIK Pringsewu, Jl. Wismarini No. 09 Pringsewu, Lampung, Indonesia

<sup>2</sup>Department of Mathematics, Faculty of Mathematics and Natural Sciences, University of Lampung, Jl. Brodjonegoro No.1 Bandar Lampung, Indonesia.

<sup>3</sup>Computer Sciences, Faculty of Mathematics and Natural Sciences, University of Lampung, Jl. Brodjonegoro No.1 Bandar Lampung, Indonesia.

\*Corresponding author

## Abstract

The locating-chromatic number denote by  $\chi_L(G)$ , is the smallest  $t$  such that  $G$  has a locating  $t$ -coloring. In this research, we determined locating-chromatic number for subdivision of certain barbell operation of origami graphs.

## Key words:

locating-chromatic number, subdivision, certain barbell operation, origami graphs.

## 1. Introduction

The concept of partition dimension was introduced by Chartrand et al. [1] as the development of the concept of metric dimension. The application of metric dimension can be found in robotic navigation [2], chemical data classification [26] and the optimization of threat detecting sensors [4]. The locating-chromatic number was first discovered by Chartrand et al. [5] in 2002, with obtained two graph concepts, coloring vertices and partition dimension of a graph. The locating-chromatic number denote by  $\chi_L(G)$ , is the smallest  $t$  such that  $G$  has a locating  $t$ -coloring. Ne<sup>20</sup> investigated the locating-chromatic number for a path graph  $P_n$ , a cycle graph  $C_n$ , and  $k$ -ble star graph  $S_{a,b}$ . Furthermore, Chartrand et al. [6] characterized all graphs order  $n$  with locating-chromatic number  $n - 1$ . Baskoro and Asmiati [7] characterized all trees with locating-chromatic number 3.

The locating-chromatic number of the join of graphs was introduced by B<sup>3</sup>oei and Anbarloei [8]. Purwasih et al. [9], obtained locating-chromatic number for a subdivision of a graph on one edge. For graph with dominant vertices have been studied in [10]. In [11], Asmiati found the locating-chromatic number of non-homogeneous caterpillar and firecrackers graph, [12] certain barbell graphs  $B_{m,n}$  and  $B_{P(n,1)}$ . In 2019, Irawan et al. [13] obtained locating-chromatic number for certain operation of generalized Petersen graphs  $sP(4,2)$ . Furthermore, in [14] determined the locating-chromatic number for  $sP(n,1)$ , origami graphs [15] and certain barbell origami graphs [16]. The locating-chromatic

number of a graph is a newly interesting topic to study cause there is no general theorem for determining the locating-chromatic number of any graph. In this research, we specifying about locating-chromatic number for subdivision of certain barbell operation of origami graphs, called  $B_{O_n}^s$ . This study is a continuation of previous research.

The following definition of the locating-chromatic number of a graph, dominant vertices, origami graph, and certain barbell origami graphs is taken from [5, 17, 18, 16]. We use some theorems that is basics to work out a lower bound of the locating-chromatic number of a graph is taken from [5, 15]. The set of neighbours of a vertex  $l$  in  $G$ , denoted by  $N(l)$ .

**Theorem 1.1.** [5] Let  $c$  be a locating coloring in a connected graph. If  $k$  and  $l$  are distinct vertices of  $G$  such that  $d(k, w) = d(l, w)$  for all  $w \in V(G) - \{k, l\}$ , then  $c(k) \neq c(l)$ . In particular, if  $k$  and  $l$  are non-adjacent vertices of  $G$  such that  $N(k) \neq N(l)$ , then  $c(k) \neq c(l)$ .

**Theorem 1.2.** [15] Let  $O_n$  be an origami graph for  $n \geq 3$ . The locating chromatic number of an origami graphs  $O_n$  is 4 for  $n=3$  and 5 otherwise.

## 2. Results and Discussion

In this section, we will discuss the locating-chromatic number for subdivision of certain barbell operation of origami graphs, denoted by  $B_{O_n}^s$ .

**Theorem 2.1.** Let  $B_{O_n}^s$  be a subdivision of certain barbell operation of origami graphs for  $n \geq 3, s \geq 1$ . Then the locating-chromatic number of  $B_{O_n}^s$  is five,  $\chi_L(B_{O_n}^s) = 5$ .

**Proof.** Let  $B_{O_n}^s$  be a subdivision of certain barbell operation of origami graphs for  $n \geq 3, s \geq 1$ , with  $V(B_{O_n}^s) = \{u_i, u_{n+i}, v_i, v_{n+i}, w_i, w_{n+i} : i \in \{1, \dots, n\}\} \cup \{x_i :$

Manuscript received September 5, 2021

Manuscript revised September 20, 2021

<https://doi.org/10.22937/IJCNSNS.2021.21.9.9>

$i \in \{1, \dots, s\}$  and  $E(B_{O_n}^s) = \{u_i w_i, u_i v_i, v_i w_i, u_i u_{i+1}, w_i u_{i+1}; i \in \{1, \dots, n\}\} \cup \{u_{n+i} w_{n+i}, u_{n+i} v_{n+i}, v_{n+i} w_{n+i}, u_{n+i} u_{n+i+1}, w_{n+i} u_{n+i+1}; i \in \{1, \dots, n-1\}\} \cup \{u_n x_1, x_s u_{n+1}\} \cup \{x_i x_{i+1}; i \in \{1, \dots, s-1\}\}$ .

To prove the theorem, we will be divided into two cases :

**Cases 1. For  $n = 3$**

First, we determine lower bound of  $\chi_L(B_{O_3}^s)$ . Since subdivision of certain barbell operation of origami graphs, containing origami graphs  $O_3$ , then by Theorem 1.2.  $\chi_L(B_{O_3}^s) \geq 4$ . Next, we will show that 4 colors are not enough. Origami graph  $B_{O_3}^s$  there are six complete graph with four vertices, denote by  $K_4$ . Without loss of generality, we assign three colors for any  $K_4$  in  $B_{O_3}^s$ , and then the six vertices are dominant vertices. As a result, if we use four colors it is not enough because there are more than one  $K_4$  in  $B_{O_3}^s$ . So  $\chi_L(B_{O_3}^s) \geq 5$ .

Next, we determined the upper bound of  $\chi_L(B_{O_3}^s) \leq 5$ . To show that  $\chi_L(B_{O_3}^s) \leq 5$ , consider the 5-coloring  $c$  on  $B_{O_3}^s$  as follow,

$$\begin{aligned} C_1 &= \{u_1, w_2, u_6, v_5\}; \\ C_2 &= \{u_4, w_1, w_5\}; \\ C_3 &= \{u_2, v_1, w_3, u_5, v_4, v_6\} \cup \{x_i | \text{for odd } i, i \geq 1\}; \\ C_4 &= \{u_3, v_2, w_4, w_6\} \cup \{x_i | \text{for even } i, i \geq 2\}; \\ C_5 &= \{v_3\}; \end{aligned}$$

The coloring  $c$  will create partition  $\Pi$  on  $V(B_{O_3}^s)$ . We shall show that the color codes of all vertices in  $B_{O_3}^s$  are different. We have  $c_\Pi(u_1) = (0, 2, 1, 1, 1)$ ;  $c_\Pi(u_2) = (1, 1, 0, 1, 2)$ ;  $c_\Pi(u_3) = (1, 2, 1, 0, 1)$ ;  $c_\Pi(u_4) = (1, 0, 1, 1, s + 3)$ ;  $c_\Pi(u_5) = (1, 1, 1, 0, s + 4)$ ;  $c_\Pi(u_6) = ((0, 1, 1, s + 4)$ ;  $c_\Pi(v_1) = (1, 3, 2, 0, 1)$ ;  $c_\Pi(v_2) = (1, 3, 0, 1, 2)$ ;  $c_\Pi(v_3) = (2, 0, 1, 1, 3)$ ;  $c_\Pi(v_4) = (2, 1, 1, 0, s + 4)$ ;  $c_\Pi(v_5) = (0, 1, 2, 1, s + 5)$ ;  $c_\Pi(v_6) = (1, 2, 1, 0, s + 5)$ ;  $c_\Pi(w_1) = (1, 3, 2, 1, 0)$ ;  $c_\Pi(w_2) = (0, 2, 1, 1, 2)$ ;  $c_\Pi(w_3) = (1, 1, 1, 0, 2)$ ;  $c_\Pi(w_4) = (2, 1, 0, 1, s + 4)$ ;  $c_\Pi(w_5) = (1, 0, 2, 1, s + 5)$ ;  $c_\Pi(w_6) = (1, 1, 0, 1, s + 4)$ . For  $s = 1$ , we have  $c_\Pi(x_i) = (i + 1, 1, 1, 0, i + 2)$ . For  $i$  odd,  $i \leq \lfloor \frac{s}{2} \rfloor$ ,  $s \geq 2$ , we have  $c_\Pi(x_i) = (i + 1, i + 1, 1, 0, i + 2)$ . For  $i$  even,  $i \leq \lfloor \frac{s}{2} \rfloor$ ,  $s \geq 2$ , we have  $c_\Pi(x_i) = (i + 1, i + 1, 0, 1, i + 2)$ . For  $i$  odd,  $i > \lfloor \frac{s}{2} \rfloor$ ,  $s \geq 2$ , we have  $c_\Pi(x_i) = (s - i + 2, s - i + 1, 1, 0, i + 2)$ . For  $i$  even,  $i > \lfloor \frac{s}{2} \rfloor$ ,  $s \geq 2$ , we have  $c_\Pi(x_i) = (s - i + 2, s - i + 1, 0, 1, i + 2)$ .

Since the color codes of all vertices  $B_{O_3}^s$  are different, thus  $c$  is a locating coloring. So  $\chi_L(B_{O_3}^s) \leq 5$ .

**Case 2. For  $n \geq 4$**

First, we determine lower bound of  $\chi_L(B_{O_n}^s)$  for  $n \geq 4$ . Since subdivision of certain barbell operation of origami graphs, containing origami graphs  $O_n$ , then by Theorem 1.2 it is clear that  $\chi_L(B_{O_n}^s) \geq 5$ .

To show the upper bound for the locating-chromatic number for subdivision of certain barbell operation of origami graphs  $\chi_L(B_{O_n}^s) \geq 5$  for  $n \geq 4$ . Let us different some subcases.

**Subcase 2.1. (odd  $n$ ), for  $\lfloor \frac{n}{2} \rfloor$  odd,  $n \geq 5$**

Let  $c$  be a coloring for subdivision of certain barbell operation of origami graph  $B_{O_n}^s$ , for  $\lfloor \frac{n}{2} \rfloor$  odd,  $n \geq 5$  we make the partition  $\Pi$  of  $V(B_{O_n}^s)$  :

$$\begin{aligned} C_1 &= \{w_1 | 1 \leq i \leq n\} \cup \{u_{n+1}\}; \\ C_2 &= \{u_i | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n-1\} \cup \{u_{n+i} | \text{for odd } i, 3 \leq i \leq \lfloor \frac{n}{2} \rfloor - 2\} \cup \{u_{n+i} | \text{for odd } i, \lfloor \frac{n}{2} \rfloor + 2 \leq i \leq n\} \cup \{v_{n+i} | \text{for even } i, 2 \leq i \leq n-1\} \cup \{x_i | \text{for } i, i \geq 2\}; \\ C_3 &= \{u_i | \text{for even } i, 2 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1\} \cup \{u_i | \text{for even } i, \lfloor \frac{n}{2} \rfloor + 3 \leq i \leq n-1\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq 27\} \cup \{u_{n+i} | \text{for even } i, 2 \leq i \leq n-1\} \cup \{v_{n+i} | \text{for odd } i, 1 \leq i \leq n\}; \\ C_4 &= \{u_1\} \cup \{w_{n+i} | 1 \leq i \leq n\} \cup \{x_i | \text{for odd } i, i \geq 1\}; \\ C_5 &= \{u_{\lfloor \frac{n}{2} \rfloor + 1}\} \cup \{u_{n+\lfloor \frac{n}{2} \rfloor}\}. \end{aligned}$$

For  $\lfloor \frac{n}{2} \rfloor$  odd  $n \geq 5$ , the color codes of all the vertices of  $V(B_{O_n}^s)$  are :

$$c_\Pi(u_i) = \begin{cases} 0, & \text{for 2nd ordinate, even } i, 3 \leq i \leq n, n \geq 5 \\ 2, & \text{for 3rd ordinate, even } i, 2 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1, n \geq 5 \\ i-1, & \text{for 3rd ordinate, even } i, \lfloor \frac{n}{2} \rfloor + 3 \leq i \leq n-1, n \geq 9 \\ i-1, & \text{for 4th ordinate, } i = 1 \\ i-1, & \text{for 5th ordinate, } i = \lfloor \frac{n}{2} \rfloor + 1 \\ i-1, & \text{for 3rd ordinate, } i = \lfloor \frac{n}{2} \rfloor + 1 \\ i-1, & \text{for 4th ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ n-i+1, & \text{for 4th ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ i - \lfloor \frac{n}{2} \rfloor - 1, & \text{for 5th ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ \lfloor \frac{n}{2} \rfloor - i + 1, & \text{for 5th ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ \lfloor \frac{n}{2} \rfloor - i, & \text{for 5th ordinate, } i = 1 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_i) = \begin{cases} 0, & \text{for } 2^{nd} \text{ ordinate, } 1 \leq i \leq n-1, n \geq 5 \\ 9, & \text{for } 3^{rd} \text{ ordinate, odd } i, 1 \leq i \leq n, n \geq 5 \\ 2, & \text{for } 2^{nd} \text{ ordinate, } i = 1 \\ 3, & \text{for } 3^{rd} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor + 1 \\ i, & \text{for } 4^{th} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ n-i+2, & \text{for } 4^{th} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ \lfloor \frac{n}{2} \rfloor, & \text{for } 5^{th} \text{ ordinate, } i = 1 \\ \lfloor \frac{n}{2} \rfloor - i + 2, & \text{for } 5^{th} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ i - \lfloor \frac{n}{2} \rfloor, & \text{for } 5^{th} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_i) = \begin{cases} 0, & \text{for } 1^{st} \text{ ordinate, } 1 \leq i \leq n, n \geq 5 \\ 2, & \text{for } 2^{nd} \text{ ordinate, } i = 1 \\ 19, & \text{for } 3^{rd} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ i, & \text{for } 4^{th} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ n-i+1, & \text{for } 4^{th} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ \lfloor \frac{n}{2} \rfloor - i + 1, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ i - \lfloor \frac{n}{2} \rfloor, & \text{for } 5^{th} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(u_{n+i}) = \begin{cases} i-1, & \text{for } 1^{st} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ n-i+1, & \text{for } 1^{st} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ 0, & \text{for } 1^{st} \text{ ordinate, } i = 1 \\ & \text{for } 2^{nd} \text{ ordinate, odd } i, 3 \leq i \leq \lfloor \frac{n}{2} \rfloor - 2, n \geq 9 \\ & \text{for } 2^{nd} \text{ ordinate, odd } i, \lfloor \frac{n}{2} \rfloor + 2 \leq i \leq n, n \geq 5 \\ & \text{for } 3^{rd} \text{ ordinate, even } i, 3 \leq i \leq n-1, n \geq 5 \\ & \text{for } 5^{th} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor + 1 \\ \lfloor \frac{n}{2} \rfloor - 1, & \text{for } 1^{st} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ 2, & \text{for } 2^{nd} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ \lfloor \frac{n}{2} \rfloor - i, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1, n \geq 5 \\ i - \lfloor \frac{n}{2} \rfloor, & \text{for } 5^{th} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_{n+i}) = \begin{cases} i, & \text{for } 1^{st} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ n-i+2, & \text{for } 1^{st} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ 0, & \text{for } 2^{nd} \text{ ordinate, even } i, 2 \leq i \leq n-1, n \geq 5 \\ & \text{for } 3^{rd} \text{ ordinate, odd } i, 1 \leq i \leq n, n \geq 5 \\ & \text{for } 2^{nd} \text{ ordinate, } i = 1 \\ 3, & \text{for } 2^{nd} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ \lfloor \frac{n}{2} \rfloor - i + 1, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ i - \lfloor \frac{n}{2} \rfloor + 1, & \text{for } 5^{th} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_{n+i}) = \begin{cases} i, & \text{for } 1^{st} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 5 \\ n-i+1, & \text{for } 1^{st} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 5 \\ 2, & \text{for } 2^{nd} \text{ ordinate, } i = 1 \text{ and } i = \lfloor \frac{n}{2} \rfloor \\ 0, & \text{for } 4^{th} \text{ ordinate, } 29 \leq i \leq n, n \geq 5 \\ \lfloor \frac{n}{2} \rfloor - i, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1, n \geq 5 \\ i - \lfloor \frac{n}{2} \rfloor + 1, & \text{for } 5^{th} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor \leq i \leq n, n \geq 5 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(x_i) = \begin{cases} s-i+1, & \text{for } 1^{st} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ i+1, & \text{for } 1^{st} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ & \text{for } 3^{rd} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor \\ 1, & \text{for } 1^{st} \text{ ordinate, } i = s \\ & \text{for } 2^{nd} \text{ ordinate, odd } i, i \geq 1 \\ & \text{for } 4^{th} \text{ ordinate, even } i, i \geq 2 \\ s-i+2, & \text{for } 3^{rd} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor \\ i + \lfloor \frac{n}{2} \rfloor - 2, & \text{for } 5^{th} \text{ ordinate, } i < \lfloor \frac{s}{2} \rfloor \\ s-i + \lfloor \frac{n}{2} \rfloor + 1, & \text{for } 5^{th} \text{ ordinate, } i \geq \lfloor \frac{s}{2} \rfloor \\ 0, & \text{otherwise.} \end{cases}$$

Since for odd  $n$  all vertices have different color codes,  $c$  is a locating coloring for subdivision of certain barbell operation of origami graphs  $B_{O_n}^s$ , so that  $\chi_L(B_{O_n}^s) \leq 5$ , for  $\lfloor \frac{n}{2} \rfloor$  odd,  $n \geq 5$ .

**Subcase 2.2.** (odd  $n$ ), for  $\lfloor \frac{n}{2} \rfloor$  even,  $n \geq 7$

Let  $c$  be a coloring for subdivision of certain barbell operation of origami graph  $B_{O_n}^s$ , for  $\lfloor \frac{n}{2} \rfloor$  even,  $n \geq 7$  we make the partition  $\Pi$  of  $V(B_{O_n}^s)$ :

$$C_1 = \{w_i | 1 \leq i \leq n\} \cup \{u_{n+1}\};$$

$$C_2 = \{u_i | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n-1\} \cup \{u_{n+i} | \text{for even } i, 2 \leq i \leq \lfloor \frac{n}{2} \rfloor - 2\} \cup \{u_{n+i}\}$$

for even  $i, \lfloor \frac{n}{2} \rfloor + 2 \leq i \leq n-1 \} \cup \{v_{n+i} | \text{for odd } i, 1 \leq i \leq n\}$ ;

$C_3 = \{u_i | \text{for even } i, 2 \leq i \leq \lfloor \frac{n}{2} \rfloor - 2\} \cup \{u_i | \text{for even } i, \lfloor \frac{n}{2} \rfloor + 2 \leq i \leq n-1\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n\} \cup \{u_{n+i} | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_{n+i} | \text{for even } i, 2 \leq i \leq n-1\} \cup \{x_i | \text{for even } i, i \geq 2\}$ ;

$C_4 = \{u_1\} \cup \{w_{n+i} | 1 \leq i \leq n\} \cup \{x_i | \text{for odd } i, i \geq 1\}$ ;

$C_5 = \{u_{\lfloor \frac{n}{2} \rfloor + 1}\} \cup \{u_{n+\lfloor \frac{n}{2} \rfloor}\}$ .

For  $\lfloor \frac{n}{2} \rfloor$  even  $n \geq 7$ , the color codes of all the vertices of  $V(B_{0n}^S)$  are :

$$c_{\Pi}(u_i) = \begin{cases} 0, & \text{for } 2^{\text{nd}} \text{ ordinate, odd } i, 3 \leq i \leq n, n \geq 7 \\ & \text{for } 3^{\text{rd}} \text{ ordinate, even } i, 2 \leq i \leq \lfloor \frac{n}{2} \rfloor - 2, n \geq 7 \\ & \text{for } 3^{\text{rd}} \text{ ordinate, even } i, \lfloor \frac{n}{2} \rfloor + 2 \leq i \leq n-1, n \geq 7 \\ & \text{for } 4^{\text{th}} \text{ ordinate, } i = 1 \\ & \text{for } 5^{\text{th}} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ 2, & \text{for } 3^{\text{rd}} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ i-1, & \text{for } 4^{\text{th}} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 7 \\ n-i+1, & \text{for } 4^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ i - \lfloor \frac{n}{2} \rfloor, & \text{for } 5^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ \lfloor \frac{n}{2} \rfloor - i, & \text{for } 5^{\text{th}} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1, n \geq 7 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_i) = \begin{cases} 0, & \text{for } 2^{\text{nd}} \text{ ordinate, } i=4, n, 2 \leq i \leq n-1, n \geq 7 \\ & \text{for } 3^{\text{rd}} \text{ ordinate, odd } i, 1 \leq i \leq n, n \geq 7 \\ 2, & \text{for } 2^{\text{nd}} \text{ ordinate, } i = 1 \\ 3, & \text{for } 3^{\text{rd}} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ i, & \text{for } 4^{\text{th}} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 7 \\ n-i+2, & \text{for } 4^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ \lfloor \frac{n}{2} \rfloor - i + 1, & \text{for } 5^{\text{th}} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 7 \\ i - \lfloor \frac{n}{2} \rfloor + 1, & \text{for } 5^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_i) = \begin{cases} 0, & \text{for } 1^{\text{st}} \text{ ordinate, } 1 \leq i \leq n, n \geq 7 \\ 2, & \text{for } 2^{\text{nd}} \text{ ordinate, } i = 1 \\ & \text{for } 3^{\text{rd}} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ i, & \text{for } 4^{\text{th}} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 7 \\ n-i+1, & \text{for } 4^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ \lfloor \frac{n}{2} \rfloor - i, & \text{for } 5^{\text{th}} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 7 \\ i - \lfloor \frac{n}{2} \rfloor + 1, & \text{for } 5^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(u_{n+i}) = \begin{cases} i-1, & \text{for } 1^{\text{st}} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 7 \\ n-i+1, & \text{for } 1^{\text{st}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ 0, & \text{for } 1^{\text{st}} \text{ ordinate, } i = 1 \\ & \text{for } 2^{\text{nd}} \text{ ordinate, even } i, 2 \leq i \leq \lfloor \frac{n}{2} \rfloor - 2, n \geq 7 \\ & \text{for } 2^{\text{nd}} \text{ ordinate, even } i, \lfloor \frac{n}{2} \rfloor + 2 \leq i \leq n-1, n \geq 7 \\ & \text{for } 3^{\text{rd}} \text{ ordinate, odd } i, 3 \leq i \leq n, n \geq 7 \\ & \text{for } 5^{\text{th}} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ 2, & \text{for } 2^{\text{nd}} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ \lfloor \frac{n}{2} \rfloor - i, & \text{for } 5^{\text{th}} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1, n \geq 7 \\ i - \lfloor \frac{n}{2} \rfloor, & \text{for } 5^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_{n+i}) = \begin{cases} i, & \text{for } 1^{\text{st}} \text{ ordinate, } 2 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1, n \geq 7 \\ n-i+2, & \text{for } 1^{\text{st}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ 0, & \text{for } 2^{\text{nd}} \text{ ordinate, odd } i, 1 \leq i \leq n, n \geq 7 \\ & \text{for } 3^{\text{rd}} \text{ ordinate, even } i, 2 \leq i \leq n-1, n \geq 7 \\ 2, & \text{for } 3^{\text{rd}} \text{ ordinate, } i = 1 \\ 3, & \text{for } 2^{\text{nd}} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ \lfloor \frac{n}{2} \rfloor - i + 1, & \text{for } 5^{\text{th}} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 7 \\ i - \lfloor \frac{n}{2} \rfloor + 1, & \text{for } 5^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_{n+i}) = \begin{cases} i, & \text{for } 1^{\text{st}} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, n \geq 7 \\ n-i+1, & \text{for } 1^{\text{st}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n, n \geq 7 \\ 2, & \text{for } 2^{\text{nd}} \text{ ordinate, } i = \lfloor \frac{n}{2} \rfloor \\ & \text{for } 3^{\text{rd}} \text{ ordinate, } i = 1 \\ 0, & \text{for } 4^{\text{th}} \text{ ordinate, } 1 \leq i \leq n, n \geq 7 \\ \lfloor \frac{n}{2} \rfloor - i, & \text{for } 5^{\text{th}} \text{ ordinate, } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1, n \geq 7 \\ i - \lfloor \frac{n}{2} \rfloor + 1, & \text{for } 5^{\text{th}} \text{ ordinate, } \lfloor \frac{n}{2} \rfloor \leq i \leq n, n \geq 7 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(x_i) = \begin{cases} s - i + 1, & \text{for } 1^{st} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ i + 1, & \text{for } 1^{st} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ 1, & \text{for } 1^{st} \text{ ordinate, } i = s \\ & \text{for } 3^{rd} \text{ ordinate, odd } i, i \geq 1 \\ & \text{for } 4^{th} \text{ ordinate, even } i, i \geq 2 \\ i, & \text{for } 2^{nd} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor \\ s - i + 2, & \text{for } 2^{nd} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor \\ 2, & \text{for } 3^{rd} \text{ ordinate, } s = 1 \\ i + \lfloor \frac{n}{2} \rfloor - 2, & \text{for } 5^{th} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor \\ s - i + \lfloor \frac{n}{2} \rfloor, & \text{for } 5^{th} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor \\ 0, & \text{otherwise.} \end{cases}$$

Since for odd  $n$  all vertices have different color codes,  $c$  is a locating coloring for subdivision of certain barbell operation of origami graphs  $B_{O_n}^s$ , so that  $\chi_L(B_{O_n}^s) \leq 5$ , for  $\lfloor \frac{n}{2} \rfloor$  even,  $n \geq 7$ .

**Subcase 2.3.** (even  $n$ ), for  $\frac{n}{2}$  odd,  $n \geq 6$

Let  $c$  be a coloring for subdivision of certain barbell operation of origami graph  $B_{O_n}^s$ , for  $\frac{n}{2}$  odd,  $n \geq 6$  we make the partition  $\Pi = V(B_{O_n}^s)$ :

$$\begin{aligned} C_1 &= \{w_i | 1 \leq i \leq \frac{n}{2} - 1\} \cup \{w_i | \frac{n}{2} + 1 \leq i \leq n\} \cup \{u_{n+1}\}; \\ C_2 &= \{u_i | \text{for odd } i, 3 \leq i \leq n - 1\} \cup \{v_i | \text{for even } i, 1 \leq i \leq n\} \cup \{u_{n+i} | \text{for even } i, 2 \leq i \leq n\} \cup \{v_{n+i} | \text{for odd } i, 1 \leq i \leq n - 1\}; \\ C_3 &= \{u_i | \text{for even } i, i \leq n\} \cup \{v_i | \text{for odd } i, 3 \leq i \leq n - 1\} \cup \{u_{n+i} | \text{for odd } i, 3 \leq i \leq n - 1\} \cup \{v_{n+i} | \text{for even } i, 2 \leq i \leq n\} \cup \{x_i | \text{for odd } i, i \geq 1\}; \\ C_4 &= \{u_i\} \cup \{w_{n+i} | 1 \leq i \leq \frac{n}{2} - 1\} \cup \{w_{n+i} | \frac{n}{2} + 1 \leq i \leq n\} \cup \{x_i | \text{for odd } i, i \geq 2\}; \\ C_5 &= \{w_n\} \cup \{w_{\frac{n}{2}}\}. \end{aligned}$$

For  $\frac{n}{2}$  odd  $n \geq 6$ , the color codes of all the vertices of  $V(B_{O_n}^s)$  are:

$$c_{\Pi}(u_i) = \begin{cases} 0, & \text{for } 2^{nd} \text{ ordinate, odd } i, 3 \leq i \leq n - 1, n \geq 6 \\ & \text{for } 3^{rd} \text{ ordinate, even } i, 2 \leq i \leq n, n \geq 6 \\ & \text{for } 4^{th} \text{ ordinate, } i = 1 \\ 2, & \text{for } 3^{rd} \text{ ordinate, } i = 1 \\ i - 1, & \text{for } 4^{th} \text{ ordinate, } 2 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 1, & \text{for } 4^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ \frac{n}{2} - i + 1, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ i - \frac{n}{2}, & \text{for } 5^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_i) = \begin{cases} 3, & \text{for } 2^{nd} \text{ ordinate, } i = 1 \\ 0, & \text{for } 2^{nd} \text{ ordinate, even } i, 2 \leq i \leq n, n \geq 6 \\ & \text{for } 3^{rd} \text{ ordinate, odd } i, 1 \leq i \leq n - 1, n \geq 6 \\ i, & \text{for } 4^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 2, & \text{for } 4^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ \frac{n}{2} - i + 2, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2} - 1, n \geq 6 \\ i - \frac{n}{2} + 1, & \text{for } 5^{th} \text{ ordinate, } \frac{n}{2} \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_i) = \begin{cases} 0, & \text{for } 1^{st} \text{ ordinate, } 1 \leq i \leq \frac{n}{2} - 1, n \geq 6 \\ & \text{for } 1^{st} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ & \text{for } 5^{th} \text{ ordinate, } i = \frac{n}{2} \\ 2, & \text{for } 2^{nd} \text{ ordinate, } i = 1 \\ i, & \text{for } 4^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 1, & \text{for } 4^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ \frac{n}{2} - i + 1, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2} - 1, n \geq 6 \\ i - \frac{n}{2} + 1, & \text{for } 5^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(u_{n+i}) = \begin{cases} i - 1, & \text{for } 1^{st} \text{ ordinate, } 2 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 1, & \text{for } 1^{st} \text{ ordinate, } \frac{n}{2} \leq i \leq n, n \geq 6 \\ 0, & \text{for } 2^{nd} \text{ ordinate, even } i, 2 \leq i \leq n, n \geq 6 \\ & \text{for } 3^{rd} \text{ ordinate, odd } i, 3 \leq i \leq n - 1, n \geq 6 \\ 10, & \text{for } 1^{st} \text{ ordinate, } i = 1 \\ & \text{for } 3^{rd} \text{ ordinate, } i = 1 \\ \frac{n}{2} - i + 1, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ i - \frac{n}{2}, & \text{for } 5^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_{n+i}) = \begin{cases} i, & \text{for } 1^{st} \text{ ordinate, } 2 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 2, & \text{for } 1^{st} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 0, & \text{for } 2^{nd} \text{ ordinate, odd } i, 1 \leq i \leq n - 1, n \geq 6 \\ & \text{for } 3^{rd} \text{ ordinate, even } i, 2 \leq i \leq n, n \geq 6 \\ & \text{for } 2^{nd} \text{ ordinate, } i = 1 \\ 3, & \text{for } 2^{nd} \text{ ordinate, } i = 1 \\ \frac{n}{2} - i + 1, & \text{for } 5^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2} - 1, n \geq 6 \\ i - \frac{n}{2} + 1, & \text{for } 5^{th} \text{ ordinate, } \frac{n}{2} \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_{n+i}) = \begin{cases} i, & \text{for 1}^{st} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 6 \\ n - i + 1, & \text{for 1}^{st} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 2, & \text{for 2}^{nd} \text{ ordinate, } i = 1 \\ 0, & \text{for 4}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2} - 1, n \geq 6 \\ & \text{for 4}^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ \frac{n}{2} - i + 1, & \text{for 5}^{th} \text{ ordinate, } i = \frac{n}{2} \\ i - \frac{n}{2} + 1, & \text{for 5}^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 6 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(x_i) = \begin{cases} s - i + 1, & \text{for 1}^{st} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ i + 1, & \text{for 1}^{st} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ 0, & \text{for 3}^{rd} \text{ ordinate, even } i, i \geq 2 \\ & \text{for 4}^{th} \text{ ordinate, odd } i, i \geq 1 \\ i, & \text{for 2}^{nd} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ s - i + 2, & \text{for 2}^{nd} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ i + \frac{n}{2} - 1, & \text{for 5}^{th} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ s - i + \frac{n}{2}, & \text{for 5}^{th} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ 1, & \text{otherwise.} \end{cases}$$

Since for odd  $n$  all vertices have different color codes,  $c$  is a locating coloring for subdivision of certain barbell operation of origami graphs  $B_{O_n}^s$ , so that  $\chi_L(B_{O_n}^s) \leq 5$ , for  $\frac{n}{2}$  odd,  $n \geq 6$ .

**Subcase 2.4.** (even  $n$ ), for  $\frac{n}{2}$  even,  $n \geq 4$

Let  $c$  be a coloring for subdivision of certain barbell operation of origami graph  $B_{O_n}^s$ ,  $\frac{n}{2}$  even,  $n \geq 4$  we make the partition  $\Pi$  of  $V(B_{O_n}^s)$ :

$$\begin{aligned} C_1 &= \{w_i | 1 \leq i \leq \frac{n}{2} - 1\} \cup \{w_i | \frac{n}{2} + 1 \leq i \leq n\} \cup \{u_{n+1}\}; \\ C_2 &= \{u_i | \text{for odd } i, 1 \leq i \leq n - 1\} \cup \{v_i | \text{for even } i, 2 \leq i \leq n\} \cup \{u_{n+i} | \text{for odd } i, 3 \leq i \leq n\} \cup \{v_{n+i} | \text{for even } i, 1 \leq i \leq n - 1\}; \\ C_3 &= \{u_i | \text{for even } i, 2 \leq i \leq n - 2\} \cup \{v_i | \text{for odd } i, 1 \leq i \leq n - 1\} \cup \{u_{n+i} | \text{for even } i, 3 \leq i \leq n - 1\} \cup \{v_{n+i} | \text{for odd } i, 2 \leq i \leq n\} \cup \{x_i | \text{for odd } i, i \geq 1\}; \\ C_4 &= \{u_n\} \cup \{w_{n+i} | 1 \leq i \leq \frac{n}{2}\} \cup \{w_{n+i} | \frac{n}{2} + 2 \leq i \leq n\} \cup \{x_i | \text{for even } i, i \geq 2\}; \\ C_5 &= \{w_n\} \cup \{w_{n+\frac{n}{2}+1}\}. \end{aligned}$$

For  $\frac{n}{2}$  even  $n \geq 4$ , the color codes of all the vertices of  $V(B_{O_n}^s)$  are:

$$c_{\Pi}(u_i) = \begin{cases} 0, & \text{for 2}^{nd} \text{ ordinate, odd } i, 1 \leq i \leq n - 1, n \geq 4 \\ & \text{for 3}^{rd} \text{ ordinate, even } i, 2 \leq i \leq n, n \geq 4 \\ & \text{for 4}^{th} \text{ ordinate, } i = n \\ i, & \text{for 4}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ n - i, & \text{for 4}^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n - 1, n \geq 4 \\ \frac{n}{2} - i + 1, & \text{for 5}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ i - \frac{n}{2}, & \text{for 5}^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(v_i) = \begin{cases} 0, & \text{for 2}^{nd} \text{ ordinate, even } i, 2 \leq i \leq n, n \geq 4 \\ & \text{for 3}^{rd} \text{ ordinate, odd } i, 1 \leq i \leq n - 1, n \geq 4 \\ i + 1, & \text{for 4}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ n - i + 1, & \text{for 4}^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ \frac{n}{2} - i + 2, & \text{for 5}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2} - 1, n \geq 4 \\ i - \frac{n}{2} + 1, & \text{for 5}^{th} \text{ ordinate, } \frac{n}{2} \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_i) = \begin{cases} 0, & \text{for 1}^{st} \text{ ordinate, } 1 \leq i \leq \frac{n}{2} - 1, n \geq 4 \\ & \text{for 1}^{st} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ & \text{for 5}^{th} \text{ ordinate, } i = \frac{n}{2} \\ 2, & \text{for 1}^{st} \text{ ordinate, } i = \frac{n}{2} \\ & \text{for 3}^{rd} \text{ ordinate, } i = n \\ i + 1, & \text{for 4}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ n - i, & \text{for 4}^{th} \text{ ordinate, } \frac{n}{2} \leq i \leq n - 1, n \geq 4 \\ \frac{n}{2} - i + 1, & \text{for 5}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ i - \frac{n}{2} + 1, & \text{for 5}^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_{\Pi}(w_{n+i}) = \begin{cases} i - 1, & \text{for 1}^{st} \text{ ordinate, } 2 \leq i \leq \frac{n}{2}, n \geq 4 \\ n - i + 1, & \text{for 1}^{st} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ 0, & \text{for 2}^{nd} \text{ ordinate, even } i, 2 \leq i \leq n, n \geq 4 \\ & \text{for 3}^{rd} \text{ ordinate, odd } i, 3 \leq i \leq n - 1, n \geq 4 \\ & \text{for 1}^{st} \text{ ordinate, } i = 1 \\ 2, & \text{for 3}^{rd} \text{ ordinate, } i = 1 \\ \frac{n}{2} - i + 1, & \text{for 5}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ i - \frac{n}{2}, & \text{for 5}^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_n(v_{n+i}) = \begin{cases} i, & \text{for 1}^{st} \text{ ordinate, } 2 \leq i \leq \frac{n}{2} + 1, n \geq 4 \\ n - i + 2, & \text{for 1}^{st} \text{ ordinate, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ 0, & \text{for 2}^{nd} \text{ ordinate, odd } i, 1 \leq i \leq n - 1, n \geq 4 \\ & \text{for 3}^{rd} \text{ ordinate, even } i, 2 \leq i \leq n, n \geq 4 \\ 3, & \text{for 2}^{nd} \text{ ordinate, } i = 1 \\ \frac{n}{2} - i + 3, & \text{for 5}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ i - \frac{n}{2}, & \text{for 5}^{th} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_n(w_{n+i}) = \begin{cases} i, & \text{for 1}^{st} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ n - i + 1, & \text{for 1}^{st} \text{ ordinate, } \frac{n}{2} + 1 \leq i \leq n, n \geq 4 \\ 0, & \text{for 4}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ & \text{for 4}^{th} \text{ ordinate, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ & \text{for 5}^{th} \text{ ordinate, } i = \frac{n}{2} + 1 \\ 2, & \text{for 4}^{th} \text{ ordinate, } i = \frac{n}{2} + 1 \\ \frac{n}{2} - i + 2, & \text{for 5}^{th} \text{ ordinate, } 1 \leq i \leq \frac{n}{2}, n \geq 4 \\ i - \frac{n}{2}, & \text{for 5}^{th} \text{ ordinate, } \frac{n}{2} + 2 \leq i \leq n, n \geq 4 \\ 1, & \text{otherwise.} \end{cases}$$

$$c_n(x_i) = \begin{cases} s - i + 1, & \text{for 1}^{st} \text{ ordinate, } i > \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ i + 1, & \text{for 1}^{st} \text{ ordinate, } i \leq \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ & \text{for 2}^{nd} \text{ ordinate, } i < \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ s - i + 2, & \text{for 2}^{nd} \text{ ordinate, } i \geq \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ 0, & \text{for 3}^{rd} \text{ ordinate, odd } i, i \geq 1 \\ & \text{for 4}^{th} \text{ ordinate, even } i, i \geq 2 \\ i + \frac{n}{2}, & \text{for 5}^{th} \text{ ordinate, } i < \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ s - i + \frac{n}{2} + 1, & \text{for 5}^{th} \text{ ordinate, } i \geq \lfloor \frac{s}{2} \rfloor, s \geq 2 \\ 1, & \text{otherwise.} \end{cases}$$

Since for odd n all vertices have different color codes, c is a locating coloring for subdivision of certain barbell operation of origami graphs  $B_{O_n}^s$ , so that  $\chi_L(B_{O_n}^s) \leq 5$ , for  $\frac{n}{2}$  even,  $n \geq 4$ . This completes the proof of the theorem.  $\square$

**Acknowledgments**

This work was partially supported by a research grant from DRPM Dikti 2021.

**References**

[1] G. Chartrand, P. Zhang, E. Salehi, On the partition dimension of a graph, *Congressus Numerantium*, vol.130, pp.157-168, 1998.

[2] V. Saenpholphat and P. Zhang, Conditional resolvability: a survey, *International Journal of Mathematics and Mathematical Sciences*, vol.38, pp.1997-2017, 2004.

[3] G. Chartrand and P. Zhang, The theory and applications of resolvability in graphs. A survey, *Congr.Numer.*160, pp.47–68, 2003.

[4] M. Johnson, Conditional resolvability: a survey, *Structure-activity maps for visualizing the graph variables arising in drug design*, *Journal of Biopharmaceutical Statistics*, vol.3, no.2, pp.203-236, 1993.

[5] G. Chartrand, D. Erwin, M. A. Henning, P. J. Slater, P. Zhang, The locating-chromatic number of a graphs, *Bulletin of the Institute of Combinatorics and its Applications*, vol. 36, pp.89-101, 2002.

[6] G. Chartrand, D. Erwin, M. A. Henning, P. J. Slater, P. Zhang, Graf of order n with locating-chromatic number n – 1, *Discrete Mathematics*, vol.269, no.1-3, pp.65 – 79, 2003

[7] E. T. Baskoro and Asmiati, Characterizing all trees with locating-chromatic number 3, *Electronic Journal of Graph Theory and Applications (EJGTA)*, vol.1, no.2, pp.109-117, 2013.

[8] A. Behtoei, M. Anbarloei, The locating chromatic number of the join of graphs, *Bulletin of the Iranian Mathematical Society*, vol.40, no.6, pp.1491-1504, 2014.

[9] I. A. Purwasih, E. T. Baskoro, H. Assiyatun, D. Suprijanto, The bounds on the locating-chromatic number for a subdivision of a graph on one edge, *Procedia Computer Science*, vol.74, pp.84 – 88, 2015.

[10] D. Welyanti, E. T. Baskoro, R. Simanjuntak, S. Uttunggadewa, On locating-chromatic number for graphs with dominant vertices, *Procedia Computer Science*, vol.74, pp.89-92, 2015.

[11] Asmiati, On the locating-chromatic numbers of non-homogeneous caterpillars and firecrackers graphs, *Far East Journal of Mathematical Sciences (FJMS)*, vol.100, no.8, pp.1305-1316, 2016.

[12] Asmiati, I. K. G. Yana, and L. Yulianti, On The Locating Chromatic Number of Certain Barbell Graphs, *International Journal of Mathematics and Mathematical Sciences*, pp.1-5, 2018.

[13] A. Irawan, Asmiati, S. Suharsono, K. Muludi, The locating-chromatic number for certain operation of generalized Petersen graphs  $sp(4,2)$ , *Journal of Physics: Conference Series*, 1338, pp.1-6, 2019.

[14] A. Irawan, Asmiati, S. Suharsono, K. Muludi, L. Zakaria, Certain operation of generalized Petersen graphs having locating-chromatic number five, *Advances and Applications in Discrete Mathematics*, vol.24, no.2, pp.83-97, 2020.

[15] A. Irawan, Asmiati, L. Zakaria, K. Muludi, The locating-chromatic number of origami graphs, *Algorithms*, vol.14, no.167, pp.1-15, 2021.

[16] A. Irawan, Asmiati, S. Suharsono, K. Muludi, The Locating-Chromatic Number of Certain Barbell Origami Graphs, *Journal of Physics: Conference Series*, 1750, pp.1-13, 2021.

[17] Asmiati, E. T. Baskoro, Characterizing all graphs containing cycles with locating-chromatic number 3, *AIP Conference Proceedings*, 1450, pp.351-357, 2012.

[18] S. Nabila, A. N. M.Salman, The Rainbow Connection Number of Origami Graphs and Pizza Graphs, *Procedia Computer Science*, 74, pp.162-167, 2015.

# Subdivision of Certain Barbell Operation of Origami Graphs has Locating-Chromatic Number Five

ORIGINALITY REPORT

14%

SIMILARITY INDEX

## PRIMARY SOURCES

1	<a href="http://sinta3.ristekdikti.go.id">sinta3.ristekdikti.go.id</a> Internet	48 words — 1%
2	<a href="http://www.aimspress.com">www.aimspress.com</a> Internet	43 words — 1%
3	<a href="http://hdl.handle.net">hdl.handle.net</a> Internet	41 words — 1%
4	<a href="http://bims.iranjournals.ir">bims.iranjournals.ir</a> Internet	36 words — 1%
5	Roger M. Macfarlane. "Optical Stark spectroscopy of solids", <i>Journal of Luminescence</i> , 2007 Crossref	35 words — 1%
6	Buhani, Musrifatun, D.S. Pratama, Suharso, Rinawati. "Modification of Chaetoceros sp. Biomass with Silica-Magnetite Coating and Adsorption Studies towards Cu(II) Ions in Single and Binary System", <i>Asian Journal of Chemistry</i> , 2017 Crossref	31 words — 1%
7	"Graph Theory", Springer Science and Business Media LLC, 2018 Crossref	24 words — 1%

- 
- 8 G.M. Odell, G. Oster, P. Alberch, B. Burnside. "The mechanical basis of morphogenesis", *Developmental Biology*, 1981 24 words — 1%  
Crossref
- 
- 9 Zehui Shao, S. M. Sheikholeslami, Pu Wu, Jia-Biao Liu. "The Metric Dimension of Some Generalized Petersen Graphs", *Discrete Dynamics in Nature and Society*, 2018 21 words — 1%  
Crossref
- 
- 10 D. Fitriani, A.N.M. Salman. "Rainbow connection number of amalgamation of some graphs", *AKCE International Journal of Graphs and Combinatorics*, 2016 19 words — < 1%  
Crossref
- 
- 11 S. Sahni. "Matrix multiplication of data routing using a partitioned optical passive stars network", *IEEE Transactions on Parallel and Distributed Systems*, 2000 19 words — < 1%  
Crossref
- 
- 12 [hal.archives-ouvertes.fr](http://hal.archives-ouvertes.fr) 19 words — < 1%  
Internet
- 
- 13 Thomas J. Schaefer. "On the complexity of some two-person perfect-information games", *Journal of Computer and System Sciences*, 1978 18 words — < 1%  
Crossref
- 
- 14 Andreas Stein, Edlyn Teske. "Explicit bounds and heuristics on class numbers in hyperelliptic function fields", *Mathematics of Computation*, 2001 14 words — < 1%  
Crossref
- 
- 15 [www.dmlett.com](http://www.dmlett.com) 13 words — < 1%  
Internet

- 
- 16 Varaporn Saenpholpat, Ping Zhang. "On resolving edge colorings in graphs", International Journal of Mathematics and Mathematical Sciences, 2003  
Crossref 12 words — < 1%
- 
- 17 [doaj.org](http://doaj.org)  
Internet 11 words — < 1%
- 
- 18 [fmipa.itb.ac.id](http://fmipa.itb.ac.id)  
Internet 10 words — < 1%
- 
- 19 [kar.kent.ac.uk](http://kar.kent.ac.uk)  
Internet 10 words — < 1%
- 
- 20 Ismael G. Yero, Juan A. Rodríguez-Velázquez. "A note on the partition dimension of Cartesian product graphs", Applied Mathematics and Computation, 2010  
Crossref 9 words — < 1%
- 
- 21 Yero, I.G.. "On the metric dimension of corona product graphs", Computers and Mathematics with Applications, 201105  
Crossref 9 words — < 1%
- 
- 22 [ijlemr.com](http://ijlemr.com)  
Internet 9 words — < 1%
- 
- 23 [www.ejgta.org](http://www.ejgta.org)  
Internet 9 words — < 1%
- 
- 24 Anthony Joseph, Polyxeni Lamprou. "A new interpretation of the Catalan numbers arising in the theory of crystals", Journal of Algebra, 2018  
Crossref 8 words — < 1%
- 
- 25 Cong X. Kang, Eunjeong Yi. "Bounds on the sum of domination number and metric dimension of 8 words — < 1%

- 
- 26 Vivekanandan M., Srinivasan R.. "Locating chromatic number of direct product of some graphs", Malaya Journal of Matematik, 2020  
8 words — < 1%  
Crossref
- 
- 27 arxiv.org  
Internet  
8 words — < 1%
- 
- 28 epdf.pub  
Internet  
8 words — < 1%
- 
- 29 Chartrand, G.. "On hamiltonian colorings of graphs", Discrete Mathematics, 20050228  
7 words — < 1%  
Crossref
- 
- 30 Lothar Papula. "Mathematik für Ingenieure und Naturwissenschaftler Klausur- und Übungsaufgaben", Springer Science and Business Media LLC, 2004  
7 words — < 1%  
Crossref
- 
- 31 Jin-Yi Cai, Zhiguo Fu, Shuai Shao. "Beyond #CSP: A dichotomy for counting weighted Eulerian orientations with ARS", Information and Computation, 2020  
6 words — < 1%  
Crossref
- 
- 32 T. J. Arputha, P. Venugopal, M. Giridaran. "RADIO ANTIPODAL MEAN LABELING OF TRIANGULAR SNAKE FAMILIES", Advances in Mathematics: Scientific Journal, 2020  
6 words — < 1%  
Crossref

---

EXCLUDE QUOTES ON

EXCLUDE MATCHES OFF

EXCLUDE BIBLIOGRAPHY ON