

Tugiyono Tugiyono

Fish Condition Factor as Bioindicator of Water Quality on Mang...

Sources Overview

17%

OVERALL SIMILARITY

1

www.scribd.com
INTERNET

16%

2

Universitas Siliwangi on 2021-09-11
SUBMITTED WORKS

1%

Excluded search repositories:


None

Excluded from document:

- Bibliography
- Quotes
- Citations
- Small Matches (less than 10 words)

Excluded sources:

theskitour.com, internet, 17%



Integrated Sci-Tech :

The Interdisciplinary Research Approach

Volume 1

Editorial board:

Ardian Ulvan

Irza Sukmana

Publisher :

UPT. Perpustakaan Universitas Lampung



**Integrated Sci-Tech :
The Interdisciplinary Research Approach
Volume 1**

ISBN
978 – 602 – 73260 – 1 - 9

Editorial board:
Ardian Ulvan
Irza Sukmana

Cover and Layout :
IC-STAR Team

Publisher :
UPT. Perpustakaan Universitas Lampung

Photo :
Backcover : Pegadung Gigi Hiu Beach, Lampung - photographer : Lukman Hakim

All rights reserved



Editors/Reviewers

Trio Adiono
Deni Noviana
Denni Kurniawan
Diding Suhandy
Warsito
Shirley Savetlana
Edwin Azwar
Joni Agustian
Dewi A. Iryani
Lukmanul Hakim
Gigih Forda Nama
Hery Dian Septama



Preface

This book is all about the interdisciplinary research that integrates engineering, life and applied sciences, medical and biomedical engineering, agriculture engineering and food sciences. The aim was to provide the initial roadmap at a cross section basic research, technological and social developments, processes development, applications integrity, and real-world usage. The genuine motivation for the book was to provide a suitable reference text for those who interested in the multi and interdisciplinary studies which might be beneficial for basic and advance researches, enhancing the curriculum and enriching teaching and learning materials, mostly in the level of postgraduate studies.

In addition, the book was also planned to provide advanced orientation and understanding for related industries and governments to looking across industrial partnerships, business strategic, and policy and regulations. In general, the book is expected to be beneficial for a wide range of readers.

This book consists of twenty five chapters divided into four sections i.e., engineering, life and applied sciences, medical and biomedical engineering, agriculture and food science. Each chapter is a completely self-directed contribution in chained discussion which aims to bring academia, researcher, practitioners and students rise to speed with the novel developments within the particular area.

In order to enhance the reader experience, each book chapter contains its own abstract, instruction, main body, as well as conclusion sections. Moreover, bibliography resources are available at the end of each chapter.

To achieve all these aims and goals, the book should deliver a breadth of information. We are pleased and thankful for all distinguish authors and reviewers for their contribution that have made this book possible. We do hope that you will enjoy this book and find it as a useful guide and reference.

Editorial board:

Ardian Ulvan

Irza Sukmana

Table of Contents

Editors/Reviewers	i
Preface	ii
Table of Contents	iii
SECTION 1 : ENGINEERING	1
Special Contribution	2
Interdisciplinary Research Activities on Disaster Prevention and Mitigation at Kobe University by Forming Collaboration COE	3
Mechanical Characterization of Cells Exposed to Mechanical Loading	4
Analyses of Mobile Positioning Data.....	5
Chapter 1	6
The Utilization of Sorghum Rod Powder as Filler to Enhance Mechanical Strength In Bioplastics Synthesis	6
Chapter 2	13
Radiometric Correlation to Sulphur and Iron Content at BM-179 Kalan-West Kalimantan Uranium Ore	13
Chapter 3	21
Masterplan Road Network in the Border Region of Nunukan Regency of the North Kalimantan Province	21
Chapter 4	31
Migration in The Rural Impact : A Case Study of Bulupitu and Sepanjang Village, Malang Regency, Indonesia	31
Chapter 5	37
Ba _{0.5} Sr _{0.5} TiO ₃ based Photodiode Application as Light Sensor for Automatic Lighting Control Switch	37
Chapter 6	44
Regulation of 12-pulse Rectifier Converter using ANFIS-based Controller in a HVDC Transmission System	44
Chapter 7	54
Estimation of Ground Rod Depth for Effective Performance during Installation in Different Soil Types in Ibadan, South – West Nigeria	54
Chapter 8	64
Reaction Kinetics of Acetic Acid and Ethanol Esterification Catalyzed by ZSM-5 Catalyst	64

Chapter 9	71
Design of The Innovative Clothes Dryer by using "Triz" Approach	
Chapter 10	77
Hydrofoil Boat For Indonesian Waters	
Chapter 11	83
Urban Community Behavioral on the Traffic Light and Implementation of Intelligence Traffic Control System	
Chapter 12	91
K-Means Analysis in Mapping Concept Based on Geographic Information System	
Chapter 13	99
Dynamics of A Re-Parametrization of Two Dimensional Map	
Chapter 14	104
Characterization of Methyl Ester Obtained from Nanochloropsis Occulata and Tetraselmis Chuii by using In-situ and Conventional Method	
Chapter 15	111
Making Photodiode Based on $Ba_{0.5}Sr_{0.5}TiO_3$ Thin Film on P-type Si (100) Substrate with Chemical Solution Deposition (CSD) Method	
Chapter 16	115
Designing Direct Current Electric Circuit for Foster Creative Thinking	
Chapter 17	121
Smart Monitoring Data Centre base on Mini Single Board Computer BCM 2835	
Chapter 18	129
Vocational High School E-Learning Readiness: A Survey for Industrial Knowledge Transfer	
Chapter 19	136
Modeling and Simulation of Solar PV Array Emulator Utilizing Buck Converter with Adaptive Control Base on Neural Network	
Chapter 20	146
Line Balancing by combining given Work Cell and single tasks, a Small Scale Industry case	
Chapter 21	153
Simulation of Type PWR (Pressurised Water Reactor) Reactor Water Temperature using Optimal Discrete Control and D-Pole Assignment Method	
Chapter 22	158
A Game of Arranging Scrambled Letters into Meaningful Words for Young Children using FSA Method	
Chapter 23	165
Design Method of Position and Attitude Controller Using for Quad-rotor System	

Chapter 24	170
Design of Boiler Controller with LAN Based Data Logger	
Chapter 25	182
Tensile Strength Analysis of Concrete-Cellulose Composite from Coconut Coir	
SECTION 2 : AGRICULTURE AND FOOD SCIENCE	190
Special Contribution	191
Bacterial Enzymes with Special Characteristics for Biotechnological Applications	192
Chapter 26	193
A Preliminary Assessment for The Presence of a Crushing Plant in Lampung Timur Regency	
Chapter 27	200
Food Technopreneur A Design of New Curriculum in Indonesia's Higher Education	
Chapter 28	205
Establishing Working Relationship of Food Supplier as Part of effectiveness Food Safety Assessment: Case Study in Indonesia Global Chain Restaurants	
Chapter 29	211
Effect of Manure and Urea on Chemical Properties of Sandy Soil and Physiological Properties of Aloe Vera L. Plant Cultivated in Coastal Sandy Area	
SECTION 3 : MEDICAL SCIENCES AND BIOMEDICAL ENGINEERING	219
Special Contribution	220
A Begin of Robot Supported Human Programming	221
Chapter 30	222
Application Brain Wave for Wheel Robotic Movement using Mindflex	
SECTION 4 : LIFE AND APPLIED SCIENCES	228
Special Contribution	229
Some Examples of Designing Integrated Heterogeneous Catalyst System	230
Chapter 31	231
Mitigation of N_2O and CH_4 emissions from Corn Field using Urea Granulated with Nitrification Inhibitors and Zeolite	
Chapter 32	238
Influence of the Concentration of Ga-doped on the Structural and Optical Properties of ZnO Thin Films	

Chapter 33	244
Histopathology of gill of <i>Pangasius sutchi</i> infected with <i>Aeromonas hydrophila</i> and are cured using Curcumin	
Chapter 34	250
Homeschooling in Lampung Province	
Chapter 35	255
Education as an Earthquake Disaster Mitigation Efforts to Improve Safety in Children Through State Primary Media Comics in The Village New District Labuhan Ratu Bandarlampung Lampung	
Chapter 36	269
Hipotetic Model of Continuous Professional Development of Vocational Lecturer in The Higher Vocational Education In Lampung	
Chapter 37	274
Fish Condition Factor as Bioindicator of Water Quality on Mangrove Ecosystems at Labuhan Maringgai, Indonesia.	

Fish Condition Factor as Bioindicator of Water Quality on Mangrove Ecosystems at Labuhan Maringgai, Indonesia.

Tugiyono¹ and Jani Master²

Abstract— This paper discuss on the use of fish Condition Factor (CF) as a bioindicator of water quality level in mangrove ecosystems at Labuhan Maringgai, Indonesia. The results shows that fish CF, in the research site, are mostly higher than one. The following are the value for each fish species: *Mystus nigricops* (1.23), *Eleutheronema tetradactylus* (0.93), *Valamugil seheli* (1.09), *Mallotus villosus* (1.61), *Paraplagusia blochi* (0.56), *Lutjanus griseus* (2.22), *Epinephelus fuscoguttatus* (1.74), *Lutjanus campechanus* (1.74), *Argyrops bleekeri* (3.07), *Macronemus microchus* (1.19), *Arius saron* (1.03), *Lates calcarifis* (1.23), *Plecus canius* (0.52). Regarding to the measured CF, it is known that the water quality at the concerned mangrove ecosystem is suitable for fish survival (fish CF > 1). Measured water quality at the concerned site was considered as mild to moderately polluted, proved by Pollution Index values of 1.84 to 3.34. nitrate and turbidity, in particular, were failed to meet the water quality standard.

Keywords— Condition factor, Labuhan Maringgai, Mangrove ecosystems, Pollution index

1. INTRODUCTION

Lampung Mangrove Center consist of 700 Ha of mangrove forest located in the village of Margasari, Labuhan Maringgai which is part of East Lampung Regency. The Mangrove Center was established based on the regent letter No. 660/305/04/SK/2005/1546/J.26 /KL/2005 on May 10, 2005. Mangrove forest is a type of forest located in areas influenced by the tidal fluctuation (coastal and estuaries). Plant species that grow in the forest are mostly tolerant to high salinity level. In addition, mangrove ecosystem is a system consisting of organisms (plants and animals) that interact each other as well as with their habitat environmental factors [1,2].

It has been well known that there are two principal functions of mangrove ecosystems, namely the ecological function and economical function. Based on ecologically perspective, mangrove ecosystem play their role as the following a) a source of organic material for aquatic ecosystem, b) food source for marine organisms, c) habitat of marine organisms, d) spawning and nursing ground for marine organisms and e) protect land from abrasion. Mean while in

term of economical point of view, mangrove forests provide a) source of building materials, b) firewood, c) raw materials of paper, d) food, medicine and textile, b) tourism spot [2,3].

It was presented by [4] that mangrove is considered as a very unique ecosystem since in the area it can be found two groups of organisms living and adapt in two difference side of the mangrove ecosystem as their habitat, namely a) terrestrial organisms living in the forest canopy or its land such as birds, snakes and monkey, their activity mostly during low tide so that they do not need special adaptation to see water. b) marine organisms live in the bottom of the tree or in saline water. The dominant animal group, in the area, is a kind of mollusks, shrimp, worms and some other kinds of fish. Mollusks which are often found on the stem or roots of the mangroves is snails (Littorinidae) and oysters (Bivalvia) which are mostly lived in the rooting side of mangroves. It is well known that shrimp and particularly crab are very abundant on the mangrove ecosystem, they live with how to make a hole in the soft substrates, such as mangrove crabs (*Scylla*, sp), crab (*Uca* sp), small crab attaching (*Portunus* sp), soldier crabs (*Mucirtis* sp), stone crabs (*Grapsus* sp) and various ghost crabs (*Dotia*, *Cleistotoma*). They life by eating any kinds of detritus from both the sea and land. They are also supports their life in the ecosystem by filtering both plankton and zooplankton. The mangrove area is suitable location for nursery grounds, feeding grounds, and spawning grounds of most shrimp and also for saltwater fish particularly mullet, before their lives and migrate to the deep sea [3,4,5].

A. Condition Factor (CF)

Condition Factor is the Fulton's formula [weight/length³] x100, which is widely used to determine the nutritional state of fish and it is also used to evaluate the fattiness of fish [6]. The CF can also be used as an indicator of health status or fitness level of particular fish [7]. The CF can also be used as an indicator for monitoring the nutrition and health status of concerned fish populations during a long period of exposure time [8].

The value of CF is influenced by age, gender, season, maturity level of the gonads, the fulfillment of the gastric contents, type of consumed food, and development level of the fish muscles. On particular type of fish, the weight of its gonad is more than 15% of its total weight. In case of female fish, CF value will dramatically decrease after their eggs has been released from their ovaries [9].

¹ F. Department of Biology Faculty of Mathematics and Natural Sciences University of Lampung Indonesia (corresponding author to provide phone—+62(0)8127965562; email: tugiyono64@yahoo.com.au).

² S. Department of Biology Faculty of Mathematics and Natural Sciences University of Lampung Indonesia (email: jaster@unila.ac.id).

B. Pollution Index (PI)

Quality of water is commonly determined by using the pollution index (PI). Pollution index used to determine contaminants compare to the standard of water quality parameters. The pollution index is however only can be measured for specified sites, which then be used to interpret for wider area of water. The method will be able to be directly connected to the water quality standard both based on concerned parameter and particular water usage [10].

II. RESEARCH METHODS

A. Time and place

This research was conducted during period of June-July 2015. The research was performed in the Mangrove Centre located at Margasari, East Lampung Regency, Lampung, Indonesia. Sampling locations are showed at Figure 1.



Figure 1. Sampling location in Mangrove Forest of Margasari Village, East Lampung Regency.

B. Research Approach

1) Diversity And Status Of Fish Condition

Number of fish at different locations were collected during data collection, carried out supported by fishermen of Margasari. The fish were obtained by direct capture on research site or caught by fishermen within the area of mangrove ecosystems. Each type of fish was taken at least five samples. Number of cultivated fish (tilapia/*Oreochromis niloticus* and milkfish/*Chanos chanos*), were also taken as a comparison. All samples were measured their total length, and weight. Mean while species identification of fish was conducted in biology laboratory, the University of Lampung. In addition, water quality sampling were also performed at certain location as presented in Figure 1. The water quality parameters which were analyzed include: turbidity, total suspended solid (TSS), total dissolved solid (TDS), temperature, salinity, nitrate, nitrite, sulfide, ammonia, phosphate, sulfate, chemical oxygen demands (COD), biological oxygen demands (BOD), and dissolved oxygen (DO).

2) Determination Of Condition Factor

Analysis was conducted to define the status of the fish condition. The method was determined based on the value of the Condition Factor (CF), derived from the Fulton's formula, $[\text{gram}(\text{centimeters})^3] \times 100$ [6].

3) Determination of the level of quality of waters with pollution index (PI).

$$PI = \frac{\sqrt{\frac{(C/L)^2_M + (C/L)^2_R}{2}}}{L} \quad (1)$$

L = the concentration of water quality standard
 C = the concentrations of measured parameters
 M = maximum
 R = average
 Evaluation of the value of the PI is :
 $0 \leq PI \leq 1.0$: meet the quality standard (good condition)
 $1.0 \leq PI \leq 5.0$: light polluted,
 $5.0 \leq PI \leq 10$: moderately polluted
 $PI > 10$: heavily polluted [10].

III. RESULTS AND DISCUSSION

A. Diversity Index

Based on the sampling results, it was obtained 15 fish species with diversity index of 2.33. The diversity index 2.33 can be considered that the fish diversity in the concerned mangrove forest ecosystems is in a good category [11]. The fish species which was encountered most frequently (dominant) is *Mystus nigriceps*. The reason could be due to the fact that the main food of the species is a small shrimp which can be easily found in the area of study.

B. Fish Condition Status

Analysis of the status of fish condition was performed based on a value of condition factor. The calculation results of the CF values both at the mangrove ecosystems and at fish cultivation pond, can be seen in Table 2.

Table 2. Value of Fish Condition Factor

Fish Species	Means	SD
<i>Mystus nigriceps</i>	1,229	0,251
<i>Etroplus suratensis</i>	0,833	0,160
<i>Tilapia nilotica</i>	1,009	0,092
<i>Molitor villosus</i>	1,608	0,289
<i>Paraplegusia bleekeri</i>	0,561	0,055
<i>Lutjanus griseus</i>	2,219	0,433
<i>Epinephelus fuscoguttatus</i>	1,420	0,117
<i>Lutjanus campechanus</i>	1,737	0,281
<i>Argyrops bleekeri</i>	3,070	0,143
<i>Macromis microchus</i>	1,191	0,042
<i>Arius sapor</i>	1,033	0,055
<i>Lates calcarifer</i>	1,238	0,057
<i>Polydora cornuta</i>	0,526	0,191
<i>Chanos chanos</i>	0,855	0,039
<i>Oreochromis niloticus</i>	1,980	0,544

Results of the study shows that most fish (10 species) has a value of CF more than one. While the rest three fish species has value of CF less than one, this is most probably due to the fish shape, which is more cylindrical compare to the other species. In particular, [6] mentioned that CF based on Fulton's formula was not suitable for the cylindrical-shaped fish. As a comparison, it was also carried out the CF calculations for cultivated, in ponds, fish (*Chanos chanos* and *Oreochromis niloticus*) with the value of CF are 1.98 and 0.85 respectively. The value was then used as an evaluation indicator of health and fitness conditions of fish [7]. Fish could be considered in a healthy condition if there are enough food and fish seem fatter/fleshy, so fish tends to be heavier compare to its visual length. Fish in those conditions will have a value of CF higher than one. On the other hands, fish that consume less food, skinny, and slender fish has a value of condition factor of less than one [12]. The CF value affected by many factors such as the condition of the organism itself, environmental conditions, the number of organisms, as well as the availability of food. The better condition of the environment, the higher factor conditions will be [13]. As stated by [14] waters within mangrove ecosystems can provide a natural system in maintaining fishery productivity.

Based on the collected CF values, it is proved that mangrove forest ecosystem is very essential in supporting the growth of fish populations. Similar research also indicated that the value of CF on two species of Tilapia namely *Tilapia zillii* and *Oreochromis aeneus* in full strength sea water was less than one (0.86; 0.53). On the other hands, it was more than one (2.81; 3.46) in fresh water. Both of these species grow well in fresh water [15]. Similar research conducted by [16] at PT. Gunung Madu Sugar Factory was concluded that value of CF of Tilapia (*Oreochromis latipes*) were decreased at wastewater treatment pond compared to the fresh water ponds.

Pollution Index (PI)

The calculation results of the pollution index can be found in Figure 2.




Figure 2. Value of Pollution Index (PI)

The water quality within mangrove ecosystems at the research site was considered to be, based on the PI values, a light polluted to moderately polluted. The criteria is based on the marine life quality standard according to [17]. The water quality parameters that exceed the standard are nitrates and

turbidity. Nitrate, as an indicator of fertility levels, could be sourced from the decomposition of organic materials such as domestic waste, detritus of mangrove plants and other water plants [18].

IV. CONCLUSION

Mangrove forest ecosystem at the village of Margasari Village is strongly support the life of the fish populations in the area. Fish populations, based on the value of diversity index, is considered in a good condition ($H=2.55$). The condition status of the fish at the concerned site was mostly in a healthy condition and fleshy ($CF > 1$). In addition to that, Water quality conditions at the concerned site was considered in the high fertility rate and plenty of natural food available for the fish population.

V. ACKNOWLEDGEMENTS

The authors would like to thank Directorate General of Higher Education, The Ministry of National Education, Indonesia, for providing us financial support for this research, (contract number: 156/UN26/S/LPPL/2015). Our gratitude also goes to Lampung Mangrove Center for facilitating us during field research, and Dr. Endro P. Wahono for his comments and suggestions during manuscript preparation.

REFERENCES

- [1] Anonymous, *Profile Of Lampung Mangrove Center. The Team Of Lampung Mangrove Team Center. Research Institutions University Of Lampung, Bandar Lampung Indonesia, 2010*, pp. 1-19, (Indonesian).
- [2] A. Kusanti, *Mangrove Forest Rehabilitation Masterplan of Coastal East Lampung*, University of Lampung, Bandar Lampung, 2008, pp. 1-10, (Indonesian).
- [3] D.G. Bergen, *Ecosystems and Natural Resources of coastal and Ocean Management as well as Principle (Synopsis)*, The Center Studies Coastal and Ocean Resources, Bogor Agricultural University 2002, pp. 16-25, (Indonesian).
- [4] A. Nourji, *Sea archipelago*. CV. Jambatan, Jakarta, 2005, pp. 12-25, (Indonesian).
- [5] Tugyono, 2010. "Fertility evaluation of ecosystems of the Coastal Waters in the village of Srimasari sub-district of Labuhan Maringgaj Regency of East Lampung Province of Lampung". *Proceedings of the National conference of Limnology 3 th*, ISBN 987-978-8163-14-2, 2010, pp. 23-28, (Indonesian).
- [6] Z. Lucky, *Methods for the diagnosis of fish diseases*. Amerind Publishing Co. Pvt. Ltd. New Delhi, 1977, pp.10-21.
- [7] C.B. Ermadillo, M.N. Druay, E.T. Marraigan, A.C. Emata, 2000. Salinity tolerance of larvae of the mangrove red snapper (*Lutjanus argentimaculatus*) during ontogeny. *Aquaculture* 190, 155-167.
- [8] M.T. Hoque, F.M. Yusoff, A.T. Law, M.A. Syed, 1998. Effect of hydrogen sulphide on liver somatic index and Fulton's Condition Factor in *Micropogonias undulatus*. *Journal of Fish Biology* 52, 23-30
- [9] PSM, C.Bamham, A. Baxter, *Condition Factor, K, for Salmonid Fish*. Department Primary Industries, State of Victoria, ISSN: 1440-2254, 2003, pp.1-5.
- [10] Anonymous, *The Decision of the Minister of State for the Environment no. 113 th 2003*, about the Determination of the Status of Water Quality Guidelines, Jakarta Indonesia, 2003, pp.1-15, (Indonesian).
- [11] Samangan Tjahjono, *Procedure for prediction and assessment of impacts on vegetation. A collection of national conference papers: Forecasting methodology in the analysis of impacts on environmental Assessment*. PPLH cooperation PPLH-LP, IPB with BK-PSL and BAPEDAL, Bogor, 1992, pp.1-12, (Indonesian).

- [12] H. Richter, C. Luckstadt, U.L. Focken, K.Becker, 2000. An Improved Procedure to assess fish condition on the basis of length-weight relationships. *Arch.Fish.Res.* 48 (3):226-235.
- [13] M.I. Effendia, *Fisheries Biology Methods*. Foundation Of Dewi Sri 199, pp. 15-23, (Indonesian).
- [14] I. Nagelkerken, S.J.M. Elaber, S. Bouillon, P. Green, M. Haywood, I.O. Kurton, J.O. Meynecke, J. Pavlik, H.M. Penrose, A. Sasikumar, and P.J. Somerfield, 2008. The habitat functions of mangroves for terrestrial and marine fauna. *Aquatic Botany* 89: 155-185.
- [15] A. Nebemsiyah, J.D. Maganra, C. Rumbaha, 2012. Length-Weight relationship and Condition Factor of Tilapia species grown in marine and fresh water pond. *Agric. Sci. J.N.Am.*, 20(2): 117-124.
- [16] Tegriyono, N. Cahyani, F. Supriyanto, dan S. Hadi 2011. Biomonitoring of Effects Following Exposure of Fish to Sugar Refinery Effluent. *Modern Applied Science* 2:39-45.
- [17] Anonymous, *The Decision of the Minister of State for the Environment no. 31 in 2004*, about Standard of sea water Quality, Jakarta Indonesia 2004, pp. 1-10, (Indonesian).
- [18] Das W. Connell, G.J. Miller, *Chemistry and Ecotoxicology of Pollution*. Wiley-Interscience Publication. John Wiley and Sons Canada. 1984, pp. 161-191.