

Novriyandi Hanif

Jurnal 2018-3.pdf

Sources Overview

14%

OVERALL SIMILARITY

1	"Polybrominated diphenyl ethers with broad spectrum antibacterial activity from the Indonesian marine sponge Lamellodysidea herbac...	5%
	CROSSREF	
2	1library.net	2%
	INTERNET	
3	www.mdpi.com	2%
	INTERNET	
4	Universitas Brawijaya on 2018-12-27	1%
	SUBMITTED WORKS	
5	www.ncbi.nlm.nih.gov	<1%
	INTERNET	
6	Anna Beker van Woudenberg, André Wolterbeek, Lindsey te Brake, Cor Snel et al. "A category approach to predicting the developmen...	<1%
	CROSSREF	
7	ijpsr.com	<1%
	INTERNET	
8	Alejandro M. S. Mayer, Aimee J. Guerrero, Abimael D. Rodríguez, Orazio Tagliatela-Scafati et al. "Marine Pharmacology in 2014–201...	<1%
	CROSSREF	
9	Hong Bai, Wen-Wen Kong, Chang-Lun Shao, Yun Li, Yun-Zhang Liu, Min Liu, Fei-Fei Guan, Chang-Yun Wang. "Zebrafish Embryo Toxicity ...	<1%
	CROSSREF	
10	publicatio.bibl.u-szeged.hu	<1%
	INTERNET	
11	Yuki Hitora, Rika Maeda, Koyo Honda, Yusaku Sadahiro, Yuji Ise, Esther D. Angkouw, Remy E.P. Mangindaan, Sachiko Tsukamoto. "Neo...	<1%
	CROSSREF	
12	www.degruyter.com	<1%
	INTERNET	

Excluded search repositories:

None

Excluded from document:

Bibliography

Quotes

Citations

Small Matches (less than 10 words)

Excluded sources:

japsonline.com, internet, 57%

"Ichthyotoxic principles against zebrafish embryos from the Indonesian marine sponge *Neopetrosia chaliniformis*", *Journal of Applied Pharmaceutical Science*, 2018, crossref, 57%

m.japsonline.com, internet, 13%

A. Murni, P. Yuliansih, K. Mohamad, M. Kita, N. Hanif. "A glucosylated steroid from the active fraction of Indonesian *Ficus deltoidea* leaves", *AIP Publishing*, 2020, crossref, 10%

repository.lppm.unila.ac.id, internet, 6%

Novriyandi Hanif, Anggia Murni, Marie Yamauchi, Masahiro Higashi, Junichi Tanaka. "A New Trinor-guaiane Sesquiterpene from an Indonesian Soft Coral sp ", *Natural Product Communications*, 2015, crossref, 4%

Universitas Brawijaya on 2018-12-30, submitted works, 1%

Universitas Brawijaya on 2018-12-14, submitted works, 1%

www.semanticscholar.org, internet, 1%

University of Edinburgh on 2021-04-28, submitted works, 0%

"Handbook of Anticancer Drugs from Marine Origin", *Springer Science and Business Media LLC*, 2015, crossref, 0%



Ichthyotoxic principles against zebrafish embryos from the Indonesian marine sponge *Neopetrosia chaliniformis*

Novriyandi Hanif^{*}, Resi Ardianti¹, Peni Ahmadi², Andi Setiawan^{3,4}, Kusdiantoro Mohamad⁵, Nicole J. de Voogd⁶, Anggia Murni⁷, Junichi Tanaka²

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Bogor 16680, Indonesia.

²Department of Chemistry, Biology and Marine Science, Faculty of Sciences, University of the Ryukyus, Okinawa 903-0213, Japan.

³Department of Chemistry, Faculty of Mathematics and Natural Sciences, Lampung University, Bandar Lampung 35145, Indonesia.

⁴UPT Integrated Laboratory of Innovation and Technology Center, Lampung University, Bandar Lampung 35145, Indonesia.

⁵Department of Anatomy, Physiology, and Pharmacology, Faculty of Veterinary Medicine, Bogor Agricultural University, Bogor 16680, Indonesia.

⁶Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands.

⁷Tropical Biopharmaca Research Center, Bogor Agricultural University, Bogor 16128, Indonesia.

ARTICLE INFO

Article history:

Received on: 10/12/2017

Accepted on: 06/02/2018

Available online: 31/08/2018

Key words:

Oxaquinolizidine,
Neopetrosia chaliniformis,
NMR, molecular modeling,
zebrafish embryos.

ABSTRACT

The chemical ecology of *Neopetrosia chaliniformis* has never been studied before. A study on the extract of the Indonesian marine sponge *N. chaliniformis* for the ichthyotoxicity including its teratogenic effect against zebrafish (*Danio rerio*) embryos led to the identification of oxaquinolizidine alkaloids araguspongines C (1) and D (2). Their structure and conformation were determined on the basis of spectroscopic method and molecular modeling analysis. The ichthyotoxicity araguspongines fraction against zebrafish embryos showed LC₅₀ 4.3 µg/ml (death, 48-hours post fertilization (hpf)) and 3.6 µg/ml (death, 72 hpf) as well as observation of its teratogenic effects including death, coagulation egg, tail, notochord, heart malformation, and yolk-sac edema at 24, 48, and 72 hpf. Because 76% cell death malformations in zebrafish embryos was observed, araguspongine rich fraction may serve as antimetabolic-like drug in the treatment of cancer.

INTRODUCTION

Marine sponges of the genus *Neopetrosia* have been the subjects of numerous chemical investigations as they contain unique molecules with diverse biological activities (Skropeta and Wei, 2013; Blunt *et al.*, 2013). In spite of the fact, its chemical ecology particularly ichthyotoxicity using zebrafish embryos (*Danio rerio*) has never been studied before. This aspect can trigger the discovery of bioactive molecules either as new molecules or possessing new activities. The zebrafish embryos (*D. rerio*) are powerful and versatile model for vertebrate development (Grunwald and Eisen, 2002), disease modeling (Lieschke and

Currie, 2007), and drug discovery (Zon and Peterson, 2005) because of its morphological and physiological similarity to mammals including *Homo sapiens* (Aston *et al.*, 2017), the existence of many genomic tools, and the ease with which large, phenotype-based screens can be performed (Zon and Peterson, 2005; Kari *et al.*, 2007). More specifically, the teratogenic effect of zebrafish (*D. rerio*) embryos and mammalian are found to have significant correlation (Kari *et al.*, 2015) suggesting for an initial target identification of bioactive molecules. Zebrafish embryos toxicity including teratogenic effect have been used to screen a novel microtubule inhibitor from the synthetic library (Moon *et al.*, 2002).

The metabolites isolated so far from *Neopetrosia* sponges can be grouped into three chemical classes: 1) alkaloids including tetrahydroisoquinolines (Oku *et al.*, 2003), polycyclic amines (Liang *et al.*, 2015; Li *et al.*, 2011; Sorek *et al.*, 2007), and pyridine

*Corresponding Author

Novriyandi Hanif, Department of Chemistry, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Bogor 16680, Indonesia. E-mail: nhanif@apps.ipb.ac.id

nucleosides (Shubina *et al.*, 2015), 2) tricyclic peptides (William *et al.*, 2005), and 3) quinones including pentacyclic hydroquinones (Leone *et al.*, 2008) and sesquiterpene benzoquinones (Winder *et al.*, 2011). Their crude extracts and isolated compounds exhibited various bioactivities such as cytotoxic (Oku *et al.*, 2003; Liang *et al.*, 2015; Li *et al.*, 2011; Sorek *et al.*, 2007; Shubina *et al.*, 2015; William *et al.*, 2005; Winder *et al.*, 2011), antibiotic (Leone *et al.*, 2008), antiplasmodial (Wei *et al.*, 2010), antibacterial (Liu *et al.*, 2004), and antifungal activities (Qaralleh *et al.*, 2010) to name a few. From the standpoint of chemistry, *Neopetrosia* metabolites e.g., polycyclic amines have challenged researchers to work because of difficulties in their purification, structure elucidation, and instability of the molecules (Oku *et al.*, 2003; Morinaka *et al.*, 2011).

As the marine sponge *N. subtriangularis* was found to contain symbiotic microbes (Freeman *et al.*, 2011), the symbionts could be the true producers of the bioactive molecules isolated from the host *Neopetrosia* sponges. In our continuing interests on bioactive molecules from marine organisms (Hanif *et al.*, 2015), we examined the Indonesian marine sponge *N. chaliniformis* collected off Lampung, Indonesia. As its *n*-BuOH extract showed potent activity against zebrafish embryos (LC₅₀ 61 µg/ml), the extract was separated to give an active fraction showing ichthyotoxicity against zebrafish embryos. Further chromatographic separation resulted in the isolation of a macrocyclic oxoquinolizidine alkaloid known as araguspongines C (1) and D (2) in which chemical and new biological result of active fraction are reported in this article.

MATERIAL AND METHODS

General

Optical rotation was obtained with a Jasco P-1010 digital polarimeter. NMR spectra were recorded on a Bruker Avance III 500 spectrometer (500 MHz for ¹H and 125 MHz for ¹³C). Chemical shifts (δ) were referenced to TMS or CDCl₃ signals (¹H δ 7.26, ¹³C δ 77.16). All chemical shifts (δ) are given in ppm and coupling constants (*J*) are in Hz. Multiplicities of ¹³C NMR data were determined by DEPT experiments. ESIMS data were obtained on a Jeol JMS-T 100LP mass spectrometer. High performance liquid chromatography (HPLC) separations were carried out on a Hitachi L-6000 pump fitted with a Shodex RI-101 refractive index and a SPD-20A Shimadzu UV detectors. A column used for HPLC was Cosmosil 5C₈-MS (4.6 × 250 mm), and analytical TLC was performed on Merck silica gel 60 F254 plates and visualized with Dragendorff reagent. All solvents used were reagent grade.

Animal material

The marine sponge was collected by hand using scuba off Lampung Bay, Indonesia in September 2014. The sponge was identified as *Neopetrosia chaliniformis* by one of us (NJdV). A voucher specimen has been deposited at the Department of Chemistry, Bogor Agricultural University, Indonesia (Code NH-G1-0001-14-a).

Extraction and isolation

The marine sponge specimen (wet, 700 g) stored in EtOH was extracted three times using MeOH (3 × 500 ml). The

combined extracts were concentrated under reduced pressure, and the residue was partitioned using *n*-hexane and 90% aqueous MeOH. The latter layer was partitioned further between CH₂Cl₂ and 50% aqueous MeOH. Finally, the 50% aqueous MeOH layer was concentrated and added with H₂O. It was then extracted with *n*-BuOH. All layers (*n*-hexane, CH₂Cl₂, *n*-BuOH) were checked for their activity against zebrafish embryos, and we found the *n*-BuOH extract as the most toxic (LC₅₀ 61 µg/ml). Purification of the extract (4.03 g) using silica gel 60 (0.040–0.063 mm) with gradient elution CHCl₃-MeOH gave six fractions A–F. The most ichthyotoxic fraction F (15.4 mg, LC₅₀ 4.3 µg/ml) was further purified using HPLC C₈ with eluent H₂O:MeOH:NH₄OH (2:7:0.3) to give araguspongine C (1) (0.5 mg) and araguspongine D (2) (0.4 mg).

Zebrafish assay

Fertilized and normally developed embryos of the zebrafish *Danio rerio* were used for bioassay. The method used for zebrafish embryos assay was modified from Bai *et al.* (2016). For the first screening of the extracts, the zebrafish embryos were used at 72 hpf and exposed to the sample for 24 hours. Observation of the embryos was based on whether it is dead or live. To determine more precisely, the embryos (0 hpf) were used and the LC₅₀ was determined at 48 hpf. Moreover, teratogenic effect was observed at 0, 24, 48, 72, and 96 hpf exposed with a series of sample concentrations at 0.5, 1, 2, 4, 6, and 8 µg/ml. The number of embryos used for the first screening was 10, while for the latter experiments was 20 for each well in 24 well microplates. Inverted microscope IX70 (Olympus) with objective power 1x–5x was used for observation of teratogenic effect. DMSO (1%) was used to improve the solubility of samples. Endpoints were assessed using binary responses (dead/live). Embryos and larvae were considered dead when coagulation had occurred and/or no heartbeat was detected. The LC₅₀ was calculated by using statistical software SPSS IBM 23.0. The experiment was repeated for three times.

Conformational analysis

Conformational searches were performed with Spartan'14 (Wavefunction Inc.) using a commercially available PC. Energy-minimized molecule was obtained from calculation using MMFF with further optimization using semi-empirical method AM1. Initial conformers was set to 10,000, then the obtained conformer was optimized with AM1 semi-empirical method.

RESULT AND DISCUSSION

Two oxoquinolizidine alkaloids araguspongines C (1) and D (2) have been isolated from the active fraction F [LC₅₀ 4.3 µg/ml (death, 48 hpf) and 3.6 µg/ml (death, 72 hpf)] showing positive results with Dragendorff reagent. Compound 1 was obtained as a glass with specific rotation [α]_D²⁶ + 43° (0.042, MeOH). Its molecular formula was determined as C₂₈H₅₀N₂O₄ by observing its HRESIMS ion at *m/z* 479.38588 [M+H]⁺ (calcd. for C₂₈H₅₁N₂O₄, Δ +1.0 mmu) indicating five degrees of unsaturation. Since the ¹³C NMR spectrum showed only 14 signals, it was suggested that 1 is a symmetric molecule. The ¹H NMR (CDCl₃, 500 MHz) of 1 showed key signals at δ 3.56 (H₂, t, *J* = 10.6); 1.76 (H₂-3a, m);

1.07 (H₂-3b, m); 3.11 (H4-a, brt, $J = 12.4$); 2.97 (H4-b, dd, $J = 13.6$, 4.1); 2.35 (H6-a, dd, $J = 8.5$, 2.2); 3.04 (H6-b, td, $J = 11.8$, 2.6); 1.80 (H₂-7a, m); 1.52 (H₂-7b, m); 1.40 (H₂-8a, m); 1.30 (H₂-8b, m); 4.07 (H-10, s); 1.62 (H₂-11a, m); 1.30 (H₂-11b, m); 1.31 (H₂-15a, m); 1.55 (H₂-15b, m); 1.32 (H₂-16a, m); 1.55 (H₂-16b, m), while its ¹³C NMR (CDCl₃, 125 MHz) exhibited major signals at δ 76.7 (CH, C2), 26.2 (CH₂, C3), 52.6 (CH₂, C4), 44.2 (CH₂, C6), 21.0 (CH₂, C7), 22.7 (CH₂, C8), 70.9 (C, C9), 90.5 (CH, C10), 38.7 (CH₂, C11), 29.8 (CH₂, C12), 32.4 (CH₂, C13), 31.6 (CH₂, C14), 25.1 (CH₂, C15), 36.5 (CH₂, C16). All the above data are in agreement with those reported for araguspongine C (Kobayashi *et al.*, 1989; Orabi *et al.*, 2002). The relative stereochemistry of **1** was confirmed by coupling constants, NOEs, and conformational analyses using Spartan '14 as in Figure 1 which was 2*R**, 9*R**, 10*S**, 2'*R**, 9'*R**, 10'*S*'* (Figure 2).




Fig. 1: Key NOESY correlations, analysis of coupling constants, and calculated distance (Å) of **1** between two selected hydrogens for energy-minimized molecule obtained from calculated MMFF and AMI semi-empirical methods.




Fig. 2: Structure of **1** and **2** with its relative stereochemistry.

Compound **2** isolated after **1** was eluted and analyzed for C₂₈H₅₀N₂O₂ by HRESIMS indicating it to be a bis-1-oxaquinolizidine alkaloid with the absence of two hydroxy groups at C9 and C9' (δ_C 40.56). The ¹H NMR (CDCl₃, 500 MHz) of **2** showed key signals at δ 3.35 (H2, t, $J = 10.6$); 1.68 (H₂-3a, m); 1.47 (H₂-3b, m); 2.18 (H4-a, ddd, $J = 3.3$, 11.9, 11.9); 2.94 (H4-b, dd, $J = 11.5$, 1.5); 1.99 (H6-a, ddd, $J = 11.7$, 11.7, 3.0); 2.76 (H6-b, d, $J = 11.5$); 1.65 (H₂-7a, m); 1.58 (H₂-7b, m); 1.68 (H₂-8a, m); 1.25 (H₂-8b, m); 1.61 (H-9, m); 3.06 (H-10, d, $J = 8.4$); 1.59 (H₂-11a, m); 1.34 (H₂-11b, m); 1.40 (H₂-12a, m); 1.18 (H₂-12b, m); 1.30 (H₂-13a, m); 1.18 (H₂-13b, m); 1.26 (H₂-14a, m); 1.16 (H₂-14b, m); 1.58 (H₂-15a, m); 1.35 (H₂-15b, m); 1.60 (H₂-16a, m); 1.40 (H₂-16b, m), while its ¹³C NMR exhibited primary signals at δ 75.3 (CH, C2), 32.3 (CH₂, C3), 54.3 (CH₂, C4), 54.1 (CH₂, C6), 24.9 (CH₂, C7), 29.0 (CH₂, C8), 40.6 (CH, C9), 95.9 (CH, C10), 36.2 (CH₂, C11), 28.8 (CH₂, C12), 31.6 (CH₂, C13), 25.2 (CH₂, C14), 25.3 (CH₂, C15), 35.4 (CH₂, C16). The observed specific optical rotation of **2** was $[\alpha]_D^{26.7} + 12.1^\circ$ (0.033, MeOH). The relative configuration of **2** has been analyzed using NOE and coupling constants (Kobayashi *et al.*, 1998) to conclude the structure as in **2** (Figure 1).

Table 1: Teratogenic effect of F against zebrafish embryos.

Abnormality	Σ^a	[%] ^b
Body axis	18	19
Brain	0	0
Caudal fin	0	0
Circulation	3	3.2
Eye	0	0
Heart	14	15
Jaw	0	0
Notochord	12	13
Otic vesicle	7	0
Pigmentation	0	0
Somite	0	0
Trunk	0	0
Yolk-sac	16	17
Embryo with cell death	72	76

^aThe number of embryos affected teratogenic effect with all of concentration for 96 hpf.

^bPercentage malformation obtained from total embryos affected teratogenic effect with all of concentration for 96 hpf.

Due to the small amount of **1** and **2**, we did not determine each compound for teratogenic activity against zebrafish, but the activity of the fraction F containing **1** and **2** was evaluated against zebrafish embryos from 0 to 96 hpf. The fraction F exhibited ichthyotoxicity included teratogenic effect such as cells death, coagulation of eggs, tail, notochord, and heart malformation as well as yolk-sac edema (Figure 3, Table 1). One major malformation was concluded as zebrafish embryos with cell death (76%) (Table 1). Those parameters were in agreement with the endpoint used for assessing the embryo toxicity (Bai *et al.*, 2016). At 24 hpf, normal embryos began with cells death at 2 $\mu\text{g/ml}$ and had an increasing number of cells death (initial coagulation) at 4 $\mu\text{g/ml}$ (48 hpf). Body axis malformation was further observed at

2 $\mu\text{g/ml}$ (48 hpf) and tail malformation was seen at 4 $\mu\text{g/ml}$ (72 hpf). Notochord malformation was monitored at 1 $\mu\text{g/ml}$ (96 hpf). The remaining observation for 96 hpf was heart malformation and yolk-sac edema at 2 $\mu\text{g/ml}$. As the fraction F showed cells death at the embryo development (2 $\mu\text{g/ml}$, 24 hpf) and led coagulation

of eggs, the araguspongine-rich fraction F could be used as an antimitotic agent (van Vuuren *et al.*, 2015). Furthermore, the ichthyotoxicity fraction of the sponge *N. chaliniformis* is also benefit for the understanding of the chemoeological effects of marine chemical defensive substances.




Fig. 3: Zebrafish embryos morphology at 24, 48, 72, 96 hpf with the corresponding concentration at 1, 2, 4, and 6 $\mu\text{g/ml}$.

CONCLUSION

Araguspongines C (1) and D (2) have been isolated from the Indonesian marine sponge *Neopetrosia chaliniformis*. Compounds 1 and 2 were characterized by NMR, HRMS and their optical rotation data as well as comparison with literature values. Furthermore, compound 1 was analyzed its stereochemistry using NMR and molecular modeling. The active fraction F containing araguspongine molecules was shown to have ichthyotoxicity included teratogenic effect against zebrafish embryos, which could be used as antimitotic and chemoeological agents. This is the first report that araguspongines C (1) and D (2) were isolated from *N. chaliniformis* and have a new biological activity of active fraction against zebrafish (*Danio rerio*) embryos.

REFERENCES

- Aston M, Dankert, EN, Alam SkK, Hoepner LH. Fishing for cures: the allure of using zebrafish to develop precision on oncology therapies. *npj Precision Oncology*, 2017; 1; 39; 1-14.
- Bai H, Kong WW, Shao CL, Li Y, Liu YZ, Liu M, Guan FF, Wang CY. Zebrafish embryo toxicity microscale model for ichthyotoxicity evaluation of marine natural products. *Mar Biotechnol*, 2016; 18; 264-270.
- Blunt JW, Copp BR, Keyzers RA, Munro MHG, Prinsep MR. Marine Natural Products. *Nat Prod Rep*, 2013; 30; 237-323 and other reviews of this series.
- Freeman CJ, Thacker RW. Complex interactions between marine sponges and their symbiotic microbial communities. *Limnol Oceanogr*, 2011; 56; 1577-1586.

Grunwald DJ, Eisen JS. Headwaters of the zebrafish – emergence of a new model vertebrate. *Nat Rev Genet*, 2002; 3; 717-724.

Hanif N, Murni A, Yamauchi M, Higashi M, Tanaka J. A new trinor-guaiane sesquiterpene from an Indonesian soft coral *Anthelia* sp. *Nat Prod Commun*, 2015; 10: 1907-1910.

Kobayashi M, Kawazoe K, Kitagawa I. Araguspongines B, C, D, E, F, G, H, and J, new vasodilative bis-1-oxaquinolizidine alkaloids from an okinawan marine sponge, *Xestospongia* sp. *Chem Pharm Bull*, 1989; 37: 1676-1678.

Kobayashi M, Miyamoto Y, Aoki S, Murakami N, Kitagawa I, In Y, Ishida T. Isomerization of dimeric 2,9-disubstituted 1-oxa-quinolizidine alkaloids and structural revision of araguspongines B and E, isolated from a marine sponge of *Xestospongia* sp. *Heterocycles*, 1998; 47: 195-203.

Leone PD, Carroll AR, Towerzey L, King G, McArdle BM, Kern G, Fisher S, Hooper JNA, Quinn RJ. Exiguaquinol: a novel pentacyclic hydroquinone from *Neopetrosia exigua* that inhibits *Helicobacter pylori* MurI. *Org Lett*, 2008; 10: 2585-2588.

Li Y, Qin S, Guo YW, Gu YC, van Soest RWM. 9'-Epi-3 β ,3' β -dimethylxestospongin C, a new macrocyclic diamine alkaloid from the Hainan Sponge *Neopetrosia exigua*. *Planta Med*. 2011; 77: 179-181.

Liang Z, Sulzmaier FJ, Yoshida WY, Kelly M, Ramos JW, Williams PG. Neopetrocyclamines A and B, polycyclic diamine alkaloids from the sponge *Neopetrosia cf. exigua*. *J Nat Prod*, 2015; 78:543-547.

Lieschke GJ, Currie PD. Animal models of human disease: Zebrafish swim into view. *Nat Rev Genet*, 2007; 8; 353-367.

Liu H, Mishima Y, Fujiwara T, Nagai H, Kitazawa A, Mine Y, Kobayashi H, Yao X, Yamada J, Oda T, Namikoshi M. Isolation of araguspongine M, a new stereoisomer of an araguspongine/xestospongin alkaloid, and dopamine from the marine sponge *Neopetrosia exigua* collected in Palau. *Mar Drugs*, 2004; 2: 154-163.

Morinaka BI, Molinski TF. Xestoproxamines A-C from *Neopetrosia proxima*. Assignment of absolute stereostructure of bis-piperidine alkaloids by integrated degradation-CD analysis *J Nat Prod*, 2011; 74: 430-440.

Moon HS, Jacobson EM, Khersonsky SM, Luzung MR, Walsh DP, Xiong W.; Lee JW, Parikh PB, Lam JC, Kang TW, Rosania GR, Schier AF, Chang YT. A novel microtubule destabilizing entity from orthogonal synthesis of triazine library and zebrafish embryo screening. *J Am Chem Soc*, 2002; 124: 11608-11609.

Oku N, Matsunaga S, van Soest RWM, Fusetani N. Renieramycin J. A highly cytotoxic tetrahydroisoquinoline alkaloid, from a marine sponge *Neopetrosia* sp. *J Nat Prod*, 2003; 66: 1136-1139.

Orabi KY, El Sayed KA, Hamann MT, Dunbar DC, Al Said MS, Higa T, Kelly M. Araguspongines K and L, new bioactive bis-1-oxaquinolizidine N-Oxide alkaloids from Red Sea specimens of *Xestospongia exigua*. *J Nat Prod*, 2002; 65: 1782-1785.

Shubina LK, Makarieva TN, Yashunsky DV, Nifantiev NE, Denisenko VA, Dmitrenok PS, Dyshlovoy SA, Fedorov SN, Krasokhin VB, Jeong SH, Han J, Stonik VA. Pyridine nucleosides neopetrosides A and B from a marine *Neopetrosia* sp. sponge. Synthesis of neopetroside A and its β -riboside analogue. *J Nat Prod*, 2015; 78:1383-1389.

Skropeta D, Wei L. Recent advances in deep-sea natural products. *Nat Prod Rep*, 2014; 31: 999-1025.

Sorek H, Rudi A, Benayahu Y, Kashman Y. Njaoamines G and H, two new cytotoxic polycyclic alkaloids and a tetrahydroquinolone from the marine sponge *Neopetrosia* sp. *Tetrahedron Lett*. 2007; 48: 7691-7694.

Wei X, Nieves K, Rodriguez AD. Neopetrosiamine A, biologically active bis-piperidine alkaloid from the Caribbean Sea sponge *Neopetrosia proxima*. *Bioorg Med Chem Lett*, 2010; 20: 5905-5908.

William DE, Austin P, Diaz-Marero AR, van Soest R, Matainaho T, Roskelley CD, Roberge M, Andersen RJ. Neopetrosiamides, peptides from the marine sponge *Neopetrosia* sp. that inhibit amoeboid invasion by human tumor cells. *Org Lett*, 2005; 7: 4173-4176.

Winder PL, Baker HL, Linley P, Guzman EA, Pomponi SA, Diaz MC, Reed JK, Wright AE. Neopetrosiquinones A and B, sesquiterpene benzoquinones isolated from the deep-water sponge *Neopetrosia* cf. *proxima*. *Bioorg Med Chem*, 2011; 19: 6599-6603.

Qaralleh H, Idid S, Saad S, Susanti D, Taher M, Khleifat K. Antifungal and antibacterial activities of four Malaysian sponge species (Petrosiidae). *J Med Mycol*, 2010; 20: 315-320.

van Vuuren RJ, Visagie MH, Theron AE. Antimitotic drug in the treatment of cancer. *Cancer Chemother Pharmacol*, 2015; 76:1101-1112.

Zon LI, Peterson R. *In vivo* drug discovery in the zebrafish. *Nat Rev Drug Discovery*, 2005; 4: 35-44.

How to cite this article:

Hanif N, Ardianti R, Ahmadi P, Setiawan A, Mohamad K, de Voogd NJ, Murni A, Tanaka J. Ichthyotoxic Principles against Zebrafish Embryos from the Indonesian Marine Sponge *Neopetrosia chaliniformis*. *J App Pharm Sci*, 2018; 8(08): 044-048.