



ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id>

[JoF] Manuscript ID: jof-1612807 - Submission Received

1 message

Editorial Office <jof@mdpi.com>

Thu, Feb 10, 2022 at 5:14 PM

Reply-To: jof@mdpi.com

To: John Hendri <john.hendri@fmipa.unila.ac.id>

Cc: Andi Setiawan <andi.setiawan@fmipa.unila.ac.id>, Fendi Setiawan <fdsetiawan05@gmail.com>, Ni Luh Gede Ratna Juliasih <niluhratna.juliasih@fmipa.unila.ac.id>, Widyastuti Widyastuti <widyaindawione@gmail.com>, Aspita Laila <aspita.laila@fmipa.unila.ac.id>, "Wawan A. Setiawan" <wawan.as@fmipa.unila.ac.id>, "Fernandy M. Djailani" <fernandydjailani@ung.ac.id>, Mulyono Mulyono <mulyono@fmipa.unila.ac.id>, Masayoshi Arai <araim@phs.osaka-u.ac.jp>

Dear Professor Hendri,

Thank you very much for uploading the following manuscript to the MDPI submission system. One of our editors will be in touch with you soon.

Journal name: Journal of Fungi

Manuscript ID: jof-1612807

Type of manuscript: Article

Title: Fungicide activity of Culture Extract from *Kocuria palustris* 19C38A1 against *Fusarium oxysporum*

Authors: Andi Setiawan, Fendi Setiawan, Ni Luh Gede Ratna Juliasih, Widyastuti Widyastuti, Aspita Laila, Wawan A. Setiawan, Fernandy M. Djailani, Mulyono Mulyono, John Hendri *, Masayoshi Arai *

Received: 10 February 2022

E-mails: andi.setiawan@fmipa.unila.ac.id, fdsetiawan05@gmail.com, niluhratna.juliasih@fmipa.unila.ac.id, widyaindawione@gmail.com, aspita.laila@fmipa.unila.ac.id, wawan.as@fmipa.unila.ac.id, fernandydjailani@ung.ac.id, mulyono@fmipa.unila.ac.id, john.hendri@fmipa.unila.ac.id, araim@phs.osaka-u.ac.jp

You can follow progress of your manuscript at the following link (login required):

https://susy.mdpi.com/user/manuscripts/review_info/87d3cfe460ed25d55c2d8390c1a00a63

The following points were confirmed during submission:

1. JoF is an open access journal with publishing fees of 2000 CHF for an accepted paper (see <https://www.mdpi.com/about/apc/> for details). This manuscript, if accepted, will be published under an open access Creative Commons CC BY license (<https://creativecommons.org/licenses/by/4.0/>), and I agree to pay the Article Processing Charges as described on the journal webpage (<https://www.mdpi.com/journal/jof/apc>). See <https://www.mdpi.com/about/openaccess> for more information about open access publishing.

Please note that you may be entitled to a discount if you have previously received a discount code or if your institute is participating in the MDPI Institutional Open Access Program (IOAP), for more information see <https://www.mdpi.com/about/ioap>. If you have been granted any other special discounts for your submission, please contact the JoF editorial office.

2. I understand that:

a. If previously published material is reproduced in my manuscript, I will provide proof that I have obtained the necessary copyright permission. (Please refer to the Rights & Permissions website: <https://www.mdpi.com/authors/rights>).

b. My manuscript is submitted on the understanding that it has not been published in or submitted to another peer-reviewed journal. Exceptions to this rule are papers containing material disclosed at conferences. I confirm that I will inform the journal editorial office if this is the case for my manuscript. I confirm that all authors are familiar with and agree with submission of the contents of the manuscript. The journal editorial office reserves the right to contact all authors to confirm this in case of doubt. I will provide email addresses for all authors and an institutional e-mail address for at least one of the co-authors, and specify the name, address and e-mail for invoicing purposes.

If you have any questions, please do not hesitate to contact the JoF editorial office at jof@mdpi.com

Kind regards,
JoF Editorial Office
St. Alban-Anlage 66, 4052 Basel, Switzerland
E-Mail: jof@mdpi.com
Tel. +41 61 683 77 34
Fax: +41 61 302 89 18

*** This is an automatically generated email ***



ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id>

[JoF] Manuscript ID: jof-1612807 - Manuscript Resubmitted

1 message

JoF Editorial Office <jof@mdpi.com>

Thu, Mar 3, 2022 at 2:05 PM

Reply-To: William Zhang <william.zhang@mdpi.com>, JoF Editorial Office <jof@mdpi.com>

To: John Hendri <john.hendri@fmipa.unila.ac.id>

Cc: Andi Setiawan <andi.setiawan@fmipa.unila.ac.id>, Fendi Setiawan <fdsetiawan05@gmail.com>, Ni Luh Gede Ratna Juliasih <niluhratna.juliasih@fmipa.unila.ac.id>, Widyastuti Widyastuti <widyaindawione@gmail.com>, Aspita Laila <aspita.laila@fmipa.unila.ac.id>, "Wawan A. Setiawan" <wawan.as@fmipa.unila.ac.id>, "Fernandy M. Djailani" <fernandydjailani@ung.ac.id>, Mulyono Mulyono <mulyono@fmipa.unila.ac.id>, Masayoshi Arai <araim@phs.osaka-u.ac.jp>

Dear Professor Hendri,

Thank you very much for resubmitting the modified version of the following manuscript:

Manuscript ID: jof-1612807

Type of manuscript: Article

Title: Fungicide activity of Culture Extract from *Kocuria palustris* 19C38A1 against *Fusarium oxysporum*

Authors: Andi Setiawan, Fendi Setiawan, Ni Luh Gede Ratna Juliasih, Widyastuti Widyastuti, Aspita Laila, Wawan A. Setiawan, Fernandy M. Djailani, Mulyono Mulyono, John Hendri *, Masayoshi Arai *

Received: 10 February 2022

E-mails: andi.setiawan@fmipa.unila.ac.id, fdsetiawan05@gmail.com, niluhratna.juliasih@fmipa.unila.ac.id, widyaindawione@gmail.com, aspita.laila@fmipa.unila.ac.id, wawan.as@fmipa.unila.ac.id, fernandydjailani@ung.ac.id, mulyono@fmipa.unila.ac.id, john.hendri@fmipa.unila.ac.id, araim@phs.osaka-u.ac.jphttps://susy.mdpi.com/user/manuscripts/review_info/87d3cfe460ed25d55c2d8390c1a00a63

A member of the editorial office will be in touch with you soon regarding progress of the manuscript.

Kind regards,

JoF Editorial Office

Postfach, CH-4020 Basel, Switzerland

Office: St. Alban-Anlage 66, CH-4052 Basel

Tel. +41 61 683 77 34 (office)

Fax +41 61 302 89 18 (office)

E-mail: jof@mdpi.com<https://www.mdpi.com/journal/jof/>

*** This is an automatically generated email ***



ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id>

[JoF] Manuscript ID: jof-1612807 - Revised Version Received

1 message

JoF Editorial Office <jof@mdpi.com>

Thu, Mar 3, 2022 at 3:03 PM

Reply-To: william.zhang@mdpi.com

To: John Hendri <john.hendri@fmipa.unila.ac.id>

Cc: Andi Setiawan <andi.setiawan@fmipa.unila.ac.id>, Fendi Setiawan <fdsetiawan05@gmail.com>, Ni Luh Gede Ratna Juliasih <niluhratna.juliasih@fmipa.unila.ac.id>, Widyastuti Widyastuti <widyaindawione@gmail.com>, Aspita Laila <aspita.laila@fmipa.unila.ac.id>, "Wawan A. Setiawan" <wawan.as@fmipa.unila.ac.id>, "Fernandy M. Djailani" <fernandydjailani@ung.ac.id>, Mulyono Mulyono <mulyono@fmipa.unila.ac.id>, Masayoshi Arai <araim@phs.osaka-u.ac.jp>, JoF Editorial Office <jof@mdpi.com>

Dear Professor Hendri,

Thank you very much for providing the revised version of your paper:

Manuscript ID: jof-1612807

Type of manuscript: Article

Title: Fungicide activity of Culture Extract from *Kocuria palustris* 19C38A1 against *Fusarium oxysporum*

Authors: Andi Setiawan, Fendi Setiawan, Ni Luh Gede Ratna Juliasih, Widyastuti Widyastuti, Aspita Laila, Wawan A. Setiawan, Fernandy M. Djailani, Mulyono Mulyono, John Hendri *, Masayoshi Arai *

Received: 10 February 2022

E-mails: andi.setiawan@fmipa.unila.ac.id, fdsetiawan05@gmail.com, niluhratna.juliasih@fmipa.unila.ac.id, widyaindawione@gmail.com, aspita.laila@fmipa.unila.ac.id, wawan.as@fmipa.unila.ac.id, fernandydjailani@ung.ac.id, mulyono@fmipa.unila.ac.id, john.hendri@fmipa.unila.ac.id, araim@phs.osaka-u.ac.jphttps://susy.mdpi.com/user/manuscripts/review_info/87d3cfe460ed25d55c2d8390c1a00a63

We will continue processing your paper and will keep you informed about the status of your submission.

Kind regards,

Mr. William Zhang

Assistant Editor

E-Mail: caroline.song@mdpi.com

Skype live:.cid.b3c7f2da5fa8b6c

As a side note, JoF is organizing three awards for young investigators in 2021. Would be great if you will help spread the news to your lab members if suitable: <https://www.mdpi.com/journal/jof/awards>

--

MDPI Branch Office, Wuhan, Jingan St, 430064 Wuhan, China

Tel.: +86 027 8780 8658; Fax: +86 027 8761 2588

JoF Editorial Office

E-Mail: jof@mdpi.com<http://www.mdpi.com/journal/jof/>

News:

JoF received a new Impact Factor 5.816 (2021 Journal Citation Reports®)

To submit to the journal, click here: <https://www.mdpi.com/journal/jof>

JoF 2021 Young Investigator Awards - open for nomination: 31 October 2021

<https://www.mdpi.com/journal/jof/awards>

Follow us on Twitter: @JoF_MDPI

MDPI - Multidisciplinary Digital Publishing Institute

<http://www.mdpi.com/>



ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id>

Set Up Your Profile at SciProfiles.com

1 message

JoF Editorial Office <jof@mdpi.com>

Tue, Mar 8, 2022 at 8:17 AM

Reply-To: info@sciprofiles.com, william.zhang@mdpi.com

To: Andi Setiawan <andi.setiawan@fmipa.unila.ac.id>

Cc: William Zhang <william.zhang@mdpi.com>

Dear Dr. Setiawan,

Congratulations on your paper being accepted for publication in JoF.

To enhance your visibility, as well as that of co-authors, we would like to invite you to set up profiles on SciProfiles (<https://sciprofiles.com>), which will allow us to add a link for each author to their permanent homepage in the final version of the paper.

You can use the link below to register or login using your MDPI credentials:

https://sciprofiles.com/author_registration/87d3cfe460ed25d55c2d8390c1a00a63/UXhiVVpvVjFpclJYSFpGdmlwbUxYVnIGODBQNDdBclFmaU1HTG9xcEVVVT0=

After you have finished setup, please send us the link to your profile within two days, so that all articles you publish with MDPI will be linked to your SciProfiles account, where colleagues and peers will be able to see all of your publications, as well as your other academic contributions.

When filling in your profile:

1. Please use the email address that is associated with your publications if possible;
2. Please add a photo to help colleagues identify you easily and to enhance the visibility of your profile;
3. Since this is your public profile that others will see, please add your publications and key projects. Your publication list can be easily prepared by verifying your ORCID;
4. Please complete all sections of your profile in as detailed as possible, so that other researchers can find you.

Many thanks in advance for your collaboration. If you have any questions, please feel free to contact info@sciprofiles.com.

Kind regards,

Dr. Shu-Kun Lin, Founder and President

<https://sciprofiles.com/profile/2>



ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id>

[JoF] Manuscript ID: jof-1612807; doi: 10.3390/jof8030280. – Promotion Tips

1 message

jof@mdpi.com <jof@mdpi.com>

Wed, Mar 9, 2022 at 7:03 PM

Reply-To: william.zhang@mdpi.com

To: andi.setiawan@fmipa.unila.ac.id, fdsetiawan05@gmail.com, niluhratna.juliasih@fmipa.unila.ac.id, widyastuti.unila@gmail.com, aspita.laila@fmipa.unila.ac.id, wawan.as@fmipa.unila.ac.id, fernandydjailani@ung.ac.id, mulyono@fmipa.unila.ac.id, john.hendri@fmipa.unila.ac.id, araim@phs.osaka-u.ac.jp

Cc: jof@mdpi.com, william.zhang@mdpi.com

Dear Authors,

Congratulations! Your manuscript has been published in JoF.

Thank you for choosing our journal. Your article is already available, open access, to everyone free of charge via the journal website. Promoting your work is an important part of the post-publication process which will increase the visibility, impact and citation of your work. We encourage you to promote and share your article widely.

We have some tips about what you can do to promote your article here:

<https://www.mdpi.com/authors/promoting>

To promote discoverability, views and downloads, you can:

(1) Social Media

- Share your article on various social networks, such as Facebook, LinkedIn, Mendeley and Twitter by clicking the link on the webpage of your article. Tagging @MDPIOpenAccess as well as key people in your field might lead to retweets: Your publication may become viral on social media!
- Send a short text (up to 200 characters) to the MDPI assistant editor who was your contact for the publication of your manuscript, who will then post it on the journal Twitter account.
- Write a blog post to explain the meaning and possible outcomes of your research. This will lead to higher engagement of your research community. Ask your institution or society to post your paper on their social media accounts and to include a story about your paper in their newsletters.

(2) Link Share

- Share the article link directly with colleagues and peers in your field.
- Add a link to your article in your email signature.
- Update your personal and institutional websites by adding the title of your article and a link to it.

(3) Academic Research-sharing Platforms

- Set up your profile on academic research-sharing platforms, such as SciProfiles, ResearchGate, Academia.edu or Google Scholar, and add a short summary of your article.
- Register an ORCID author identifier and add the article information to your profile.
- Deposit your article in repositories (such as those run by your university) to make your research more discoverable.

(4) Conferences

- Present your publication at conferences in the form of a presentation or a poster. You may participate in conferences relevant for your field organized by MDPI on Sciforum.net.

Please keep up with any journal updates and news via our twitter account

(https://twitter.com/JoF_MDPI) or LinkedIn page
(<https://www.linkedin.com/company/76120972/admin/>).
Find out more about MDPI on our website: <https://www.mdpi.com/about>.

Keep in touch!

Kind regards,
JoF Editorial Office
St. Alban-Anlage 66, 4052 Basel, Postfach, CH-4020 Basel, Switzerland
Tel.: +41 61 683 77 34; Fax: +41 61 302 89 18
E-Mail: jof@mdpi.com
<https://www.mdpi.com/journal/jof/>

Disclaimer: MDPI recognizes the importance of data privacy and protection. We treat personal data in line with the General Data Protection Regulation (GDPR) and with what the community expects of us. The information contained in this message is confidential and intended solely for the use of the individual or entity to whom they are addressed. If you have received this message in error, please notify me and delete this message from your system. You may not copy this message in its entirety or in part, or disclose its contents to anyone.



ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id>

[JoF] Manuscript ID: jof-1612807; doi: 10.3390/jof8030280. Paper has been published.

1 message

jof@mdpi.com <jof@mdpi.com>

Wed, Mar 9, 2022 at 7:03 PM

Reply-To: william.zhang@mdpi.com, jof@mdpi.com

To: andi.setiawan@fmipa.unila.ac.id, fdsetiawan05@gmail.com, niluhratna.juliasih@fmipa.unila.ac.id, widyastuti.unila@gmail.com, aspita.laila@fmipa.unila.ac.id, wawan.as@fmipa.unila.ac.id, fernandydjailani@ung.ac.id, mulyono@fmipa.unila.ac.id, john.hendri@fmipa.unila.ac.id, araim@phs.osaka-u.ac.jp

Cc: billing@mdpi.com, website@mdpi.com, jof@mdpi.com, latic@mdpi.com, william.zhang@mdpi.com

Dear Authors,

We are pleased to inform you that your article "Fungicide Activity of Culture Extract from *Kocuria palustris* 19C38A1 against *Fusarium oxysporum*" has been published in JoF and is available online:

Abstract: <https://www.mdpi.com/2309-608X/8/3/280>PDF Version: <https://www.mdpi.com/2309-608X/8/3/280/pdf>

The meta data of your article, the manuscript files and a publication certificate are available here (only available to corresponding authors after login):

https://susy.mdpi.com/user/manuscripts/review_info/87d3cfe460ed25d55c2d8390c1a00a63

Please take a moment to check that everything is correct. You can reply to this email if there is a problem. If any errors are noticed, please note that all authors must follow MDPI's policy on updating published papers, found here: <https://www.mdpi.com/ethics#16>.

To encourage open scientific discussions and increase the visibility of published articles, MDPI recently implemented interactive commenting and recommendation functionalities on all article webpages (side bar on the right). We encourage you to forward the article link to your colleagues and peers.

We encourage you to set up your profile at www.SciProfiles.com, MDPI's researcher network platform. Articles you publish with MDPI will be linked to your SciProfiles page, where colleagues and peers will be able to see all of your publications, citations, as well as your other academic contributions. Please also feel free to send us feedback on the platform that we can improve it quickly and make it useful for scientific communities.

You can also share the paper on various social networks by clicking the links on the article webpage. Alternatively, our Editorial Office can post an announcement of your article on our Twitter channel, please send us a text of up to 200 characters with spaces. Please note that our service Scitations.net will automatically notify authors cited in your article. For further paper promotion guidelines, please refer to the following link: <https://www.mdpi.com/authors/promoting>.

We would be happy to keep you updated about new issue releases of jof. Please enter your e-mail address in the box at <https://www.mdpi.com/journal/jof/toc-alert/> to receive notifications. After issue release, a version of your paper including the issue cover will be available to download from the article abstract page.

To order high quality reprints of your article in quantities of 25-1000, visit: <https://www.mdpi.com/2309-608X/8/3/280/reprints>

We support the multidisciplinary preprint platform /Preprints/, which permanently archives full text documents and datasets of working papers in all subject areas. Posting on the platform is entirely free of charge, and full details can be viewed at <http://www.preprints.org>.

We are dedicated to providing an outstanding publishing service, and we invite you to complete our author satisfaction survey <https://www.surveymonkey.com/r/authorfeedbackmdpi>. The survey contains 20 short questions and will only take a couple of minutes to complete.

To help us improve our English editing service, provided as part of MDPI's editorial process, please take a few minutes to participate in the following survey: <https://www.surveymonkey.com/r/DCNXBW7> (for English editing service).

Thank you for choosing JoF to publish your work, we look forward to receiving further contributions from your research group in the future.






Kind regards,

--

MDPI
Postfach, CH - 4020 Basel, Switzerland
Office: St. Alban-Anlage 66, 4052 Basel, Switzerland
Tel. +41 61 683 77 34
Fax: +41 61 302 89 18
E-mail: website@mdpi.com
<https://www.mdpi.com/>

Article

Fungicide Activity of Culture Extract from *Kocuria palustris* 19C38A1 against *Fusarium oxysporum*

Andi Setiawan ¹, Fendi Setiawan ¹, Ni Luh Gede Ratna Juliasih ¹, Widyastuti Widyastuti ¹, Aspita Laila ¹, Wawan A. Setiawan ², Fernandy M. Djailani ³, Mulyono Mulyono ¹, John Hendri ^{1,*}
and Masayoshi Arai ^{4,*}

- ¹ Department of Chemistry, Faculty of Mathematics and Natural Science, Lampung University, Bandar Lampung 35145, Indonesia; andi.setiawan@fmipa.unila.ac.id (A.S.); fdsetiawan05@gmail.com (F.S.); niluhratna.juliasih@fmipa.unila.ac.id (N.L.G.R.J.); widyastuti.unila@gmail.com (W.W.); aspita.laila@fmipa.unila.ac.id (A.L.); mulyono@fmipa.unila.ac.id (M.M.)
- ² Department of Biology, Faculty of Mathematics and Natural Science, Lampung University, Bandar Lampung 35145, Indonesia; wawan.as@fmipa.unila.ac.id
- ³ Department of Fish Processing, Faculty of Fisheries and Marine Science, Gorontalo State University, Gorontalo 96128, Indonesia; fernandydjailani@ung.ac.id
- ⁴ Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamada-oka, Suita, Osaka 565-0871, Japan
- * Correspondence: john.hendri@fmipa.unila.ac.id (J.H.); araim@phs.osaka-u.ac.jp (M.A.); Tel.: +62-81-2792-7379 (J.H.); +81-6-6879-8215 (M.A.)

Abstract: Secondary metabolites of actinomycetes are a potential source of bioactive compounds in the agricultural sector. This study aimed to determine the fungicidal properties of extracts of marine organism-derived actinomycetes. Actinomycetes were isolated from marine organisms using agar media with 1% colloidal chitin in artificial seawater. Then, the isolates were cultured on liquid media with 1% colloidal chitin in artificial seawater under static conditions for 14 days. The culture was extracted, the fungicide properties were evaluated using the microtiter 96-well plate method, and the influence of inhibition was visualized using apotome and SEM. Finally, the active extract was analyzed using LCMSMS. In the present study, 19 actinomycetes were isolated from marine organisms, and the isolates were examined with regard to their antifungal activities. Of these nineteen isolates, the isolate 19C38A1 was picked out from the rest. Hence, it showed significant control towards *F. oxysporum*. The prospective strain 19C38A1 was determined to be *Kocuria palustris* 19C38A1. The extract 19C38A1 was shown to cause damage to cell integrity, indicated by the shrinking form, and inhibited germination in the *F. oxysporum*; subsequently, the chemical characteristics of the compound produced by the potential isolate 19C38A1 indicated the presence of benzimidazole compounds in the active fraction of C38BK2FA. These results indicate that actinomycetes derived from marine organisms near the coast of Oluhuta, Tomini Bay, Gorontalo, related to strain 19C38A1, are not widely known as sources of valuable fungicides. This preliminary information is important, as it can be used as a basis for further development in the search for fungicides derived from marine actinomycetes.

Keywords: marine-derived actinomycetes; antifungal; *Fusarium oxysporum*; colloidal chitin



Citation: Setiawan, A.; Setiawan, F.; Juliasih, N.L.G.R.; Widyastuti, W.; Laila, A.; Setiawan, W.A.; Djailani, F.M.; Mulyono, M.; Hendri, J.; Arai, M. Fungicide Activity of Culture Extract from *Kocuria palustris* 19C38A1 against *Fusarium oxysporum*. *J. Fungi* **2022**, *8*, 280. <https://doi.org/10.3390/jof8030280>

Academic Editor: Gary A. Strobel

Received: 10 February 2022

Accepted: 8 March 2022

Published: 9 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Losses of agricultural products are generally caused by crop diseases, most of which are caused by soil-borne fungi, *Fusarium oxysporum*. This fungus is the main cause of the decline in the productivity of agricultural commodities such as bananas and cereal-based crops [1]. This is because plants are damaged by this fungus, and in addition, the rate of the spread of the fungus is relatively high [2]. Currently, synthetic fungicides are still used to overcome this obstacle. The application of pesticides increased by 3.5 million tons by 2020 [3]. However, the intensive use of synthetic fungicides has an impact on environmental pollution and the spread of plant disease outbreaks [4]. Given the risks posed by synthetic

fungicides and the increasing global preference for safer and environmentally friendly alternatives, new and valuable fungicides must be developed [5].

Based on a literature review, azole groups were found to be one of the active compounds commonly found in synthetic fungicides [6]. The active azole moiety targets the cytochrome P450 enzyme, known as sterol 14 α -demethylase (CYP51) or lanosterol 14 α -demethylase, and inhibits the initial step in ergosterol biosynthesis [7]. More specifically, the benzimidazole group has broad-spectrum activity; it interacts with β -tubulin and inhibits hyphal growth [8]. The benzimidazole group is a strong inhibitor of tubulin polymerization and exerts antifungal activity by targeting the β -tubulin subunit of microtubules, which means cell division is not carried out, leading to cell death.

In recent years, some researchers have looked for antifungal compounds derived from microorganisms. Actinomycetes are potential sources of active metabolites with very diverse structures and different activities [9]. The results of a recent study reported that *Streptomyces* sp. SY195 is able to produce antifungal compounds that inhibit the growth of *Candida albicans* in Gauze liquid media [10]. It has also been reported that *Streptomyces sporoclivatus* NBRC 100,767 cultivated on ISP3 media was able to produce fungicidal compounds that could inhibit the growth of *F. oxysporum* [11]. However, to date, information regarding azole fungicides, especially those produced from rare actinomycetes cultivated on liquid chitin colloid media, is still scarce.

In this study, we aimed to isolate actinomycetes from marine organisms and to screen, purify, and characterize culture extracts that display antifungal activities. A selected isolate was chosen for the further analysis of fungicide activity. In addition, shrimp shell waste was used as a source of colloidal chitin, which would be used as main a carbon source in actinomycetes growth media.

2. Materials and Methods

2.1. Biomaterials

2.1.1. Preparation Shrimp Shell Waste and Colloid Chitin

Shrimp shell waste was collected from free-market Gudang Lelang, Teluk Betung Bandar Lampung. The shrimp shells were washed with tap water to remove impurities such as dust and soil. The shrimp shells were then dried overnight in an oven at 60 °C. The dried shrimp shells were demineralized using HCl 2 N, stirred at 30 °C for 2 h, and then neutralized with distilled water. After that, the dried, demineralized residue was deproteinated with NaOH 2.5%, stirred at 90 °C for 2 h, and then neutralized using distilled water until a pH of 7 was reached. The pellet was dried in an oven at 60 °C overnight to obtain the chitin flake. Chitin was transformed into colloidal chitin by concentrated HCl being added and it being stirred for 2 h; then, it was neutralized with distilled water to a pH of 7 and stored in a freezer at -20 °C until it was used in further experiments [12].

2.1.2. Actinomycetes Isolates

Samples of marine organisms were collected from Tomini Bay, Gorontalo, Indonesia (0°25'11.9" N 123°08'31.8" E), in August 2019. Small pieces of marine organisms were rinsed with sterile artificial seawater (ASW). The pieces were placed into plates containing colloidal chitin agar medium supplemented with 25 μ g/mL of cycloheximide and 25 μ g/mL of nalidixic acid, and they were cultured at room temperature for 14 days. The isolation of actinomycetes was carried out via the dilution method using agar media, 1% colloid chitin, and ASW (g L^{-1}) consisting of 27.0 g of NaCl, 5.6 g of $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 1.5 g of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1.0 g of KNO_3 , 0.07 g of K_2HPO_4 , 6.6 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and 0.04 of NaHCO_3 [13–15].

2.1.3. Pathogenic Fungi

The phytopathogen *F. oxysporum* used in this study was collected from the Microbiology Laboratory, Department of Biology, Lampung University. *Malassezia Globosa* were collected from the Department of Parasitology, University of Indonesia. Both fungi were maintained in potato dextrose agar (PDA) media.

2.2. Cultivation

The inoculum of actinomycetes was prepared using 100 mL of liquid colloidal chitin containing 1% in artificial seawater (ASW) in a 500 mL culture flask, which was incubated for 7 days under static conditions. Seven-day-old inoculum was transferred into a liquid fermentation medium containing 500 mL of 1% colloidal chitin in ASW in a 2000 mL Erlenmeyer flask. The culture was incubated for 14 days at room temperature under static conditions.

2.3. Screening of Antifungal Activity

The high-throughput screening method was used to determine the antifungal activity using 96-well plates, and absorbance was measured using a Hospitex plate reader at OD₆₃₀. Seven-day-old fungi were cultured on potato dextrose agar (PDA). Spores were harvested using a moistened sterile cotton swab; before being used, the cotton swab was moistened in a tube containing a drop of Tween 80 in potato dextrose broth (PDB) and transferred into a 10 mL inoculum tube containing PDB. The tube was vortexed for 15 s. The heavy particles were allowed to settle down for 3–5 min. The upper homogenous suspension was separated from heavy particles into a new, sterile tube. Inoculum was adjusted to 0.5 Mc Farland standard turbidity equal to (10⁶ CFU/mL) or (OD₆₃₀ = 0.08–0.1) [16]. All *n*-butanol extracts were made for 10 mg/mL of stock solution. Ketoconazole was used as a positive control, 12.5% MeOH was used as a solvent control, and wells without fungus were used as a control of contamination. Each well contained 80 µL of PDB media, 100 µL of fungal suspension at 1.0 × 10⁵ CFU (colony-forming units)/mL, and 20 µL of extract solution [17]. The plate was incubated for 48 h at room temperature. The % of inhibition was measured as follows:

$$\% \text{ of Inhibition} = \frac{\text{OD Control} - \text{OD Treatment}}{\text{OD Control}} \times 100\%$$

2.4. Characterization of Selected Actinomycetes

2.4.1. Morphological Study

The arrangement of spore ornament structure was examined using cover slip culture medium and observed in 400 M microscopically [18]. A clean sterile coverslip was planted at a tan angle of 45° in an agar plate and was cultured for 7 days. The coverslip was removed from the agar plate using a tweezer and placed upwards on a clean glass slide. The coverslip was observed under the Zeiss Axio Imager Microscope (400×). A scanning electron microscope (SEM) study was performed based on mycelium and spore ornament identification. The inoculum of selected actinomycetes was prepared in a 1% colloidal chitin liquid medium in artificial seawater and incubated under static conditions. After 7 days, the shrimp shell was placed onto a clean Petri dish. One milliliter of actinomycetes suspension was added to moisten the one gram of shrimp shell. Then, the culture was incubated for 7 days under static conditions. After the incubation, a small piece of shrimp shell was cut using the SLEE Disposable Blades microtome. The sample was prepared by it placing on aluminum stubs, which were fixed with carbon adhesive tabs. Then, gold plating was completed in 20 min. The gold-plated metal stub was observed using SEM EVO with 10 kV electron high voltage, Carl Zeiss EVO MA 10, Oberkochen, Germany.

2.4.2. Phylogenetic Analysis of Strain 19C38A1

Genomic DNA was extracted following the genomic Wizard® Genomic DNA KIT protocol (cat. No. A1120, Promega, Madison, WI, USA). PCRs of 16S rDNA sequences were completed using a Sensoquest Sensodirect thermocycler from Germany. PCRs were performed using a forward primer: 5'-AGA GTT TGA TCM TGG CTC AG-3' [19]. Additionally, a reverse primer: 5'-CCG TAC TCC CCA GGC GGG G-3' [20], which amplified 810 bp, was used. The PCR reactions were completed using a 2G Fast ReadyMix Kit (cat. No. KK5102, Merck, Taufkirchen, Germany). The PCR reactions were carried out at a total volume of 25 µL, containing 5 µL of DNA template (50 ng/L), 12.5 µL of 2G Fast ReadyMix, 6.5 µL of RNase-free water, 0.5 µL of forward primer, and 0.5 µL of the reverse primer.

Amplification was carried out in 35 cycles as follows: denaturation for 60 s at 92 °C, primer annealing for 60 s at 54 °C, and polymerization for 90 s at 72 °C. Bands were detected with the Qiaxcel Advanced instrument (Qiagen, Hilden, Germany) according to the protocol, then sequenced using the Sanger method. The results of the sequencing were analyzed phylogenetically using Mega version 11 software.

2.5. Scale-Up Cultivation and Extraction

The selected isolate 19C38A1 was cultured (twice) in 2 L of 1% colloidal chitin liquid medium for 14 days under static conditions, and the biomass was harvested using a centrifuge at 7000 rpm, 4 °C, for 10 min to separate the supernatant and pellet. The filtrate was partitioned using *n*-butanol, and the organic fraction was concentrated via evaporation at 40 °C under reduced pressure. The organic fraction was fractionated using silica gel 60 (0.063–0.200 mm, Merck KgaA, Darmstadt, Germany) open-column chromatography using *n*-hexane and EtOAc. Thin-layer chromatography (TLC) was conducted using silica gel F₂₅₄ (aluminium sheets, Merck KgaA, Darmstadt, Germany) and specific reagent Dragendorff's (Solution 1: 1.7 g of bismuth nitrate with 80 mL of water and 20 mL of glacial acetic acid; Solution 2: KI solution (50% *w/v*, 100 mL) with glacial acetic acid), and ninhydrin (2% *w/v* in 10 mL of ethanol).

2.6. Characterization

2.6.1. Microscopy Analysis of Fungicide Activity

To verify the inhibitory ability of the potent isolate extract against *F. oxysporum*, further microscopy analysis was carried out using an Apotome Microscope and SEM EVO Carl Zeiss EVO MA 10, Oberkochen, Germany. At first, spore germination was observed in the *F. oxysporum* population, which occurred by transferring 20 µL of the 48 h incubation results on a glass slide. Samples were observed at 400 M magnification. Conidia were considered germinated if the length of the germ tube was wider than the conidia. Further analysis used SEM to visualize the surface morphology of the hyphae of *F. oxysporum* and to determine the possible mechanism between the crude extract of potent isolate and *F. oxysporum*. For microscopic analysis, the 96-well plates used were modified with carbon tape. The wells were given a mixture of 100 µL of extract and 100 µL of *F. oxysporum* (10⁶ spores/mL). *F. oxysporum* (10⁶ spores/mL) was used as a control. At the edge of the well, sterile carbon tape was attached. The plate was then incubated for 48 h at 30 °C. In the step prior to SEM analysis, the carbon band with adhering hyphae was carefully removed and washed three times with 0.1 M PBS [21]. Samples were fixed with 10% (*v/v*) formaldehyde overnight at room temperature, then rinsed three times with 0.1 M PBS and dehydrated in graded ethanol (30% for 10 min, 50% for 10 min, 70% for 10 min, 90% for 10 min, and 100% for 1 h). Samples were air dried, mounted on aluminum stubs using double-sided carbon tape, sputtered with gold, and visualized using SEM EVO Carl Zeiss EVO MA 10, Oberkochen, Germany.

2.6.2. Liquid Chromatography–Mass Spectrometry (LC-MS)

The identification and characterization of fungicide compounds produced by isolate 19C38A1 were carried out using LC-MSMS Positive Mode. The active fraction was dissolved in methanol and analyzed via LC-MSMS analysis, which was equipped with the ACQUITY UPLC[®] H-Class System (Waters, Beverly, MA, USA), ACQUITY UPLC[®] HSS C18 column (1.8 µm 2.1 × 100 mm) (Waters, Beverly, MA, USA), and Xevo G2-S Qtof Mass Spectro (Waters, Beverly, MA, USA).

3. Results and Discussion

3.1. Sample Collection and Isolation of Actinomycetes

Actinomycetes were isolated from 17 marine organisms collected from near the coast of Oluhuta, Tomini Bay, Gorontalo, Indonesia (0°25'11.9'' N 123°08'31.8'' E). Samples of marine organisms were collected via Self-Contained Underwater Breathing Apparatus (SCUBA) diving at depths of 5–20 m. Tomini Bay waters are located on the equator covering three provinces,

namely North Sulawesi, Gorontalo, and Central Sulawesi. Tomini has high marine biodiversity, evidenced by the variety of fish species, coral reefs, seagrass beds, and mangroves. Tomini Bay is the second-largest bay in Indonesia after Cendrawasih Bay. Tomini geologically formed around six million years ago and has an area of 70.020 km². Tomini is rich in geological history, and this small region has a very high level of endemism [22].

In this study, 19 actinomycetes associated with marine organisms were isolated from 15 sponges, 1 tunicate, and 1 macroalga. The presence of actinomycetes in marine environments has been reported to be capable of producing bioactive metabolites with diverse activities [23]. Furthermore, the presence of actinomycetes has been found in a range of marine organisms such as sponges [24], tunicates [25], and macroalgae [26].

3.2. Screening of Antifungal Activity

The results of the bioactivity test of the actinomycetes extract against *F. oxysporum* and *M. Globosa* are shown in Table 1. Almost all the extracts had inhibitory activity levels below 50%, except for isolate 19C38A1 from the sponge, which had an inhibitory activity level of 54%.

Table 1. Screening of antifungal assay.

No	Sample Code	Phylum	Isolate Actinomycetes	% Inhibition at 1.0 mg/mL	
				<i>F. oxysporum</i>	<i>M. globosa</i>
1	01A07	Porifera	19A07A1	0	5
2	01A13	Porifera	19A13A3	14	17
3	01A15	Porifera	19A15A1	0	3
4	01A18	Porifera	19A18A1	0	0
5	01B19	Tunicate	19B19A1	0	32
			19B19A2	3	8
6	01B20	Porifera	19B20A1	42	6
7	01B21	Porifera	19B21A1	1	36
8	02C30	Porifera	19C30A1	0	26
9	02C32	Porifera	19C32A1	0	31
10	02C33	Porifera	19C33A2	45	39
11	02C34	Porifera	19C34A1	0	32
12	02C35	Porifera	19C35A1	0	28
13	02C36	Porifera	19C36A1	0	46
14	02C38	Porifera	19C38A1	54	7
15	03D40	Macroalgae	19D40A3	0	29
16	03D41	Porifera	19D41A1	0	37
			19D41A2	0	13
17	03D46	Porifera	19D46A1	0	27

Several studies have reported that sponge-derived actinomycetes can produce secondary metabolite compounds with a variety of therapeutic activities; isolates such as *Gordonia*, *Kocuria*, *Nocardia*, *Micrococcus*, *Micromonospora*, and *Microbacterium* have been successfully isolated from marine sponges [27]. Moreover, tunicate was also reported to be a host for several types of actinomycetes such as genus of *Actinomadura*, *Aeromicrobium*, *Arthobacter*, *Brevibacterium*, *Curtobacterium*, *Gordonia*, *Kocuria*, *Micrococcus*, *Micromonospora*, *Nocardia*, *Nocardiopsis*, *Saccharopolyspora*, *Salinispora*, *Solwaraspora*, and *Verrucosipora* [28]. However, extract isolates 19B19A1 and 19B19A2 from tunicate showed relatively weak activity. The same weak activity was observed in extract of isolate 19D40A3 from macroalgae. In addition, the presence of actinomycetes associated with macroalgae such as *Rhodococcus*, *Nonomuraeae*, *Microbispora*, *Isopterocola*, *Nocardiopsis*, and *Microbacterium* have been reported [29]. Based on the screening data, it appears that actinomycete 19C38A1 has the potential to produce fungicides. Therefore, isolate 19C38A1 was selected for further research related to understanding its morphology and phylogenetic properties, as well as characterization of the metabolites produced.

3.3. Characterization of Selected Actinomycetes

3.3.1. Morphological Study

The morphology of the isolated 19C38A1 grown for 7 days on 1% colloidal chitin media is shown in Figure 1a,b. In addition, the surface of the 19C38A1 isolate on the

shrimp shell was very clearly observed, as shown in Figure 1c. The results of observations (Figure 1a) showed white colonies with aerial mycelium, in accordance with the general characteristics of actinomycetes [30]. Further observation using a light microscope at 400× magnification (M) showed the presence of spore fragmentation, which characterizes the family Micrococcaceae [31]. Advanced analysis using SEM enabled us to observe aerial mycelia that displayed a straight spore chain with a diameter of 2 microns (Figure 1c) [32].




Figure 1. (a) Isolate 19C38A1 in colloidal chitin agar media 1%; (b) visualization of 19C38A1, with light microscopic scale 400×; (c) SEM image of aerial hyphae isolate 19C38A1, bar 2 µm.

3.3.2. Phylogenetic Analysis of Isolates 19C38A1

Sequencing of the 16S rDNA gene indicated the genus *Kocuria*. The new isolate, *Kocuria palustris* 19C38A1, was identified via sequencing with a similarity percentage of 99.77% and was successfully registered in GenBank with access number LC659429. Based on phylogenetic analysis, it is known that this isolate is known as *K. palustris* (Figure 2). According to the literature, *K. palustris* belongs to a rare group of actinomycetes that have been obtained from marine habitats [33]. This group of rare actinomycetes actually reflects a potential source of active metabolites that have not been studied much in terms of the structure and properties of their activities. Moreover, the presence of Micrococcaceae is common in terrestrial and marine sources. Although Micrococcaceae are widely distributed in sponges, there is very little information regarding natural products of bioactive compounds, especially from the genus *Kocuria*, associated with sponges [34].




Figure 2. Phylogenetic tree using maximum likelihood method and Hasegawa–Kishino–Yano model of 23 *Kocuria* representatives and *Micrococcus luteus* (Acc. No. AJ536198) as an outgroup. Bootstrap values (1000 resamples) are given in percentages at the nodes of the tree. The isolate *Kocuria palustris* 19C38A1 was presented.

3.4. Scale-Up Cultivation and Extraction

To produce the biomass of actinomycetes, isolate 19C38A1 was cultured in liquid fermentation in two 2 L Erlenmeyer flasks. After 2 weeks of cultivation, actinomycetes were formed in surface media, but they were clumsy, had an insoluble structure, and were not pigmented. The culture media was centrifuged to separate the solids and the filtrate. Then, the filtrate was partitioned with *n*-butanol. The *n*-butanol extract of 19C38A1 was evaporated under reduced pressure to afford crude C38B extract (1.2 gr). The C38B extract was subjected to several steps of open-column chromatography using SiO₂ as a stationary phase, and it was eluted with *n*-hexane: EtOAc (linear gradient) to obtain a 4 mg active fraction of C38BK2FA. The results of the TLC analysis, as shown in Figure 3a, confirm the presence of orange spots from Dragendorff's specific reagent, indicating the presence of alkaloid. The same was observed for the orange spots on the C38BK2FA fraction, as shown in Figure 3b, with a higher orange color intensity and an elongated shape. This indicates the presence of several alkaloid components in the C38BK2FA fraction. Further tests on the inhibitory activity showed that the C38BK2FA fraction had 88% inhibition against the growth of *F. oxysporum* at a concentration of 1 mg/mL.




Figure 3. TLC result of (a) crude C38B extract with eluent *n*-hexane: EtOAc (1:1); (b) active C38BK2FA fraction with eluent *n*-hexane: DCM (10:1).

3.5. Characterization

3.5.1. Microscopic Analysis of Fungicide Activity

Analysis based on observations using apotome microscopy showed clear differences in fungal cell morphology between the control and *F. oxysporum* treated with C38BK2FA extract. It was seen that in the control, *F. oxysporum* grew regularly, with elongated and branching hyphae (Figure 4a). In the treatment of the C38BK2FA extract sample, it was seen that there was prominent inhibition; the compound extract had an effect on mycelial growth and conidia germination, which was characterized by the absence of apical cell elongation (Figure 4b). It was found that the bioactive compounds present in the C38BK2FA extract created a stressful environment that inhibited conidia germination and hyphal growth. Further observations using SEM enabled us to obtain an overview of the effect of damage on the morphology of hyphae and conidia. In the control, it was seen that the hyphae growth was regular, which displayed a tubular structure, flat width, smooth surface, and elongated shape. The conidia showed a characteristic cube shape and a flat bottom (Figure 4c). On the other hand, after being treated with the addition of C38BK2FA extract, the hyphae structure was wrinkled and distorted (Figure 4d). Changes in the structure of hyphae and conidia were correlated with the assay for the inhibition of conidia germination. Our observations of this fungicidal effect are similar to those of Koch and Loffer [35], who observed the effect of actinomycete bioactive compounds on phytopathogens.




Figure 4. (a) Control of *F. oxysporum* in apotome microscopy observation 400 M; (b) *F. oxysporum* treated with crude C38BK2FA extract observed under apotome microscopy 400 M; (c) control of *F. oxysporum* observed using SEM 10 K x; (d) *F. oxysporum* treated with crude C38BK2FA extract observed using SEM 15 K x.

3.5.2. Liquid Chromatography–Mass Spectrometry

The results of the structural analysis of the active C38BK2FA fraction using ESI-TOF-MS showed the presence of 10 chromatogram peaks, as shown in Figure 5. Meanwhile, the interpretation of the mass spectroscopic data showed that the four chromatogram peaks at retention times 13.65 (1), 14.00 (2), 14.4 (3), and 15.80 (4) minutes had molecular peak ions $[M + H]^+$ at m/z 367.2429, 365,2344, and m/z 367.2344, with the molecular formula $C_{22}H_{28}N_4O$, and the molecular peak ion $[M + H]^+$ at m/z 447.3065 with the molecular formula $C_{23}H_{38}N_6O_3$, respectively. The data show that the active C38BK2FA fraction had nitrogen content, and this was confirmed by the data from the previous TLC test results. Based on the database on Chemspider, the molecular formula $C_{22}H_{28}N_4O$ is an alkaloid group of benzimidazole with the name N-(4-Methoxybenzyl)-1-[2-(1-piperidinyl) ethyl]-1H-benzimidazole-2-amine. Meanwhile, the compound with the molecular formula $C_{23}H_{38}N_6O_3$ is a triazole compound with the name Methyl 1-[(3R,5S)-5-(cycloheptylcarbamoyl)-1-(1-ethyl-4-piperidinyl)-3-pyrrolidinyl]-1H-1,2,3-triazole-4-carboxylate. Based on the data obtained, it can be concluded that the fungicidal activity of the active C38BK2FA fraction is due to the presence of benzimidazole compounds.

Compounds in the benzimidazole group have unique amphoteric properties and contain two nitrogen atoms. This means this group of compounds can be acidic and basic. Moreover, the benzimidazole ring compound has two tautomeric forms, in which a hydrogen atom is located on two nitrogen atoms. Then, electron-rich nitrogen heterocycles can accept or donate protons and easily form various weak interactions. Compounds in the benzimidazole group easily bind to therapeutic targets, which makes compounds in the benzimidazole group broad-spectrum agents [36]. Benzimidazole derivatives play an important role in controlling various fungal pathogens [37].




Figure 5. ESI TOF MSMS spectrum fraction C38BK2FA, $[M + H]^+ = m/z$ 365.2344.

Benzimidazole belongs to a group of broad-spectrum systemic fungicides that have been used to control plant diseases caused by *Fusarium* spp. by interacting with tubulin and inhibiting hyphae growth. The mechanism of action is related to the inhibition of fungal growth through impaired tubulin polymerization [38]. In addition, the presence of azole group compounds triggers fluidity, asymmetry, and integrity in the fungal cell membrane [39]. Triazole group compounds are also known to be active against most *Candida* species. However, this group of compounds did not display activity on most fungi such as *Aspergillus* spp., *Fusarium* spp., and *Mucormycetes* [40].

4. Conclusions

These findings concluded that the C38BK2FA extract fraction of *Kocuria palustris* 19C38A1 displayed 88% inhibition of the growth rate of *F. oxysporum* at a concentration of 1mg/mL. This activity can be caused by the presence of components of the benzimidazole group of alkaloid extracts. This pioneering information is very important for use in further research related to the development of new biofungicides that are environmentally friendly and sustainable.

Author Contributions: J.H., A.S. and M.A. conceived and designed the experiments; A.S., W.W., E.S., W.A.S., A.L., F.M.D. and N.L.G.R.J. performed the experiments; all authors analyzed the data; J.H., A.S., M.M. and M.A. wrote and revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Deputy of Research Strengthening and Development and the Ministry of Research and Technology/National Agency for Research and Innovation of the Republic of Indonesia, Basic Research with grant no. 3972/UN26.21/PN/2021 to A.S. This research was also partially supported by the Platform Project for Supporting Drug Discovery and Life Science Research (Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS)) from the Japan Agency for Medical Research and Development (AMED) (grant no. JP21am0101084), the Kobayashi International Scholarship Foundation, and a Grant-in-Aid for Scientific Research B (grant nos. 18H02096, 17H04645, and 21H02069) from the Japan Society for the Promotion of Science (JSPS) to M.A.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to thank the Deputy of Research Strengthening and Development and the Ministry of Research and Technology/National Agency for Research and Innovation of the Republic of Indonesia for the Basic Research grant. The authors also thank the support of the World Class Professor (WCP) program 2021, the UPT LTSIT Laboratory, the University of Lampung for providing scanning electron microscopy facilities, the Department of Biology Lampung University for providing the *Fusarium oxysporum*, and the Department of Parasitology University of Indonesia, Jakarta to providing *Malassezia globosa*.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Juroszek, P.; von Tiedemann, A. Climate change and potential future risks through wheat diseases: A review. *Eur. J. Plant Pathol.* **2013**, *136*, 21–33. [\[CrossRef\]](#)
2. Nicolopoulou-Stamati, P.; Maipas, S.; Kotampasi, C.; Stamatis, P.; Hens, L. Chemical pesticides and human health: The urgent need for a new concept in agriculture. *Front. Public Health* **2016**, *4*, 148. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Sharma, A.; Kumar, V.; Shahzad, B.; Tanveer, M.; Sidhu, S.G.P.; Handa, N.; Kohli, S.K.; Yadav, P.; Bali, A.S.; Parihar, R.D.; et al. Worldwide pesticide usage and its impacts on ecosystem. *SN Appl. Sci.* **2019**, *1*, 1446. [\[CrossRef\]](#)
4. Zhang, W. Global pesticide use: Profile, trend, cost/benefit and more. *Proc. Int. Acad. Ecol. Environ. Sci.* **2018**, *8*, 1–27.
5. Wallace, R.L.; Hirkala, D.L.; Nelson, L.M. Mechanisms of action of three isolates of *Pseudomonas fluorescens* active against postharvest grey mold decay of apple during commercial storage. *Biol. Control* **2018**, *117*, 13–20. [\[CrossRef\]](#)
6. Jørgensen, L.N.; Heick, T.M. Azole Use in Agriculture, Horticulture, and Wood Preservation—Is It Indispensable? *Front. Cell. Infect. Microbiol.* **2021**, *806*, 730297. [\[CrossRef\]](#)
7. Zhou, Y.; Xu, J.; Zhu, Y.; Duan, Y.; Zhou, M. Mechanism of action of the benzimidazole fungicide on *Fusarium graminearum*: Interfering with polymerization of monomeric tubulin but not polymerized microtubule. *Phytopathology* **2016**, *106*, 807–813. [\[CrossRef\]](#)
8. Hollomon, D.W.; Butters, J.A.; Barker, H.; Hall, L. Fungal β -tubulin, expressed as a fusion protein, binds benzimidazole and phenyl carbamate fungicides. *Antimicrob. Agents Chemother.* **1998**, *42*, 2171–2173. [\[CrossRef\]](#)
9. Jose, P.A.; Maharshi, A.; Jha, B. Actinobacteria in natural products research: Progress and prospects. *Microbiol. Res.* **2021**, *246*, 126708. [\[CrossRef\]](#)
10. Yi, W.; Qin, L.; Lian, X.Y.; Zhang, Z. New antifungal metabolites from the Mariana trench sediment-associated actinomycete *Streptomyces* sp. SY1965. *Mar. Drugs* **2020**, *18*, 385. [\[CrossRef\]](#)
11. Cao, P.; Li, C.; Wang, H.; Yu, Z.; Xu, X.; Wang, X.; Zhao, J.; Xiang, W. Community structures and antifungal activity of root-associated endophytic actinobacteria in healthy and diseased cucumber plants and *Streptomyces* sp. HAAG3-15 as a promising biocontrol agent. *Microorganisms* **2020**, *8*, 236. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Hsu, S.C.; Lockwood, J. Powdered chitin agar as a selective medium for enumeration of actinomycetes in water and soil. *Appl. Microbiol.* **1975**, *29*, 422–426. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Alkausar, R.; Juliasih, N.L.; Buhani, B.; Syaifei, S.; Nurachman, Z.; Setiawan, A. Chromatographic isolation of β -phycoerythrin from *Porphyridium cruentum* using ceramic hydroxyapatite. *AAFL Bioflux* **2021**, *14*, 1351–1358.
14. Shamikh, Y.I.; El Shamy, A.A.; Gaber, Y.; Abdelmohsen, U.R.; Madkour, H.A.; Horn, H.; Hassan, H.M.; Elmaidomy, A.H.; Alkhalifah, D.H.M.; Hozzein, W.N. Actinomycetes from the Red Sea Sponge *Coscinoderma mathevisi*: Isolation, Diversity, and Potential for Bioactive Compounds Discovery. *Microorganisms* **2020**, *8*, 783. [\[CrossRef\]](#)
15. Setiawan, A.; Widyastuti, W.; Irawan, A.; Wijaya, O.S.; Laila, A.; Setiawan, W.A.; Juliasih, N.L.G.R.; Nonaka, K.; Arai, M.; Hendri, J. Solid state fermentation of shrimp shell waste using *Pseudonocardia carboxydivorans* 18A13O1 to produce bioactive metabolites. *Fermentation* **2021**, *7*, 247. [\[CrossRef\]](#)

16. Al-Hatmi, A.; Curfs-Breuker, I.; de Hoog, G.S.; Meis, J.F.; Verweij, P.E. Antifungal susceptibility testing of *Fusarium*: A practical approach. *J. Fungi* **2017**, *3*, 19. [[CrossRef](#)]
17. Qi, D.F.; Zou, L.P.; Zou, D.B.; Chen, Y.F.; Gao, Z.F.; Feng, R.J.; Zhang, M.; Li, K.; Xie, J.; Wang, W. Taxonomy and broad-spectrum antifungal activity of *Streptomyces* sp. SCA3-4 isolated from rhizosphere soil of *Opuntia stricta*. *Front. Microbiol.* **2019**, *10*, 1390.
18. Goodfellow, M.; Kämpfer, P.; Busse, H.-J.; Trujillo, M.E.; Suzuki, K.; Ludwig, W.; Whitman, W.B. (Eds.) *Bergey's Manual of Systematic Bacteriology*, 2nd ed.; Springer: New York, NY, USA, 2012.
19. Heuer, H.; Krsek, M.; Baker, P.; Smalla, K.; Wellington, E.M.H. Analysis of actinomycete communities by specific amplification of genes encoding 16S rRNA and gel-electrophoretic separation in denaturing gradients. *Appl. Environ. Microb.* **1997**, *63*, 3233–3241. [[CrossRef](#)]
20. Baskaran, R.; Mohan, P.M.; Sivakumar, K.; Kumar, A. Antimicrobial Activity and Phylogenetic Analysis of *Streptomyces parvulus* Dosmb-D105 Isolated from the Mangrove Sediments of Andaman Islands. *Acta Microbiol. Immunol. Hung.* **2016**, *63*, 27–46. [[CrossRef](#)]
21. Deepthi, B.V.; Poornachandra Rao, K.; Chennapa, G.; Naik, M.K.; Chandrashekara, K.T.; Sreenivasa, M.Y. Antifungal Attributes of *Lactobacillus plantarum* MYS6 against Fumonisin Producing *Fusarium proliferatum* Associated with Poultry Feeds. *PLoS ONE* **2016**, *11*, e0155122. [[CrossRef](#)]
22. Ministry of Environment (MOE). *Description Map Marine Ecoregion Indonesia*; Ministry of Environment, Deputy of Environmental Management: Jakarta, Indonesia, 2013.
23. Hameş-Kocabaş, E.E.; Ataç, U.Z.E.L. Isolation strategies of marine-derived actinomycetes from sponge and sediment samples. *J. Microbiol. Methods* **2012**, *88*, 342–347. [[CrossRef](#)] [[PubMed](#)]
24. Cheng, C.; MacIntyre, L.; Abdelmohsen, U.R.; Horn, H.; Polymenakou, P.; Edrada-Ebel, R.; Hentschel, U. Biodiversity, antitrypanosomal activity screening, and metabolomics profiling of actinomycetes isolated from Mediterranean sponges. *PLoS ONE* **2015**, *10*, e0138528. [[CrossRef](#)]
25. Dou, X.; Dong, B. Origins and bioactivities of natural compounds derived from marine ascidians and their symbionts. *Mar. Drugs* **2019**, *17*, 670. [[CrossRef](#)] [[PubMed](#)]
26. Uzair, B.; Mena, F.; Khan, B.A.; Mohammad, F.V.; Ahmad, V.U.; Djeribi, R.; Mena, B. Isolation, purification, structural elucidation and antimicrobial activities of kocumarin, a novel antibiotic isolated from actinobacterium *Kocuria marina* CMG S2 associated with the brown seaweed *Pelvetia canaliculata*. *Microbiol. Res.* **2018**, *206*, 186–197. [[CrossRef](#)]
27. Granito, R.N.; Custódio, M.R.; Rennó, A.C.M. Natural marine sponges for bone tissue engineering: The state of the art and future perspectives. *J. Biomed. Mater. Res. Part B Appl. Biomater.* **2016**, *105*, 1717–1727. [[CrossRef](#)]
28. Donia, M.S.; Wang, B.; Dunbar, D.C.; Desai, P.V.; Patny, A.; Avery, M.; Hamann, M.T. Mollamides B and C, cyclic hexapeptides from the Indonesian Tunicate *Didemnum molle*. *J. Nat. Prod.* **2008**, *71*, 941–945. [[CrossRef](#)]
29. Pereira, L. Macroalgae. *Encyclopedia* **2021**, *1*, 177–188. [[CrossRef](#)]
30. Yaradoddi, J.S.; Kontro, M.H.; Ganachari, S.V.; Banapurmath, N.R.; Oli, A.; Katti, A.S.; Sulochana, M.B. Actinobacteria in Marine Environments. In *Actinobacteria. Rhizosphere Biology*; Yaradoddi, J.S., Kontro, M.H., Ganachari, S.V., Eds.; Springer: Singapore, 2021; pp. 21–38. [[CrossRef](#)]
31. Huang, J.; Huang, Y. *Lentzea tibetensis* sp. nov., a novel Actinobacterium with antimicrobial activity isolated from soil of the Qinghai–Tibet Plateau. *Int. J. Syst. Evol. Microbiol.* **2021**, *71*, 004976. [[CrossRef](#)]
32. Acharyabhata, A.; Kandula, S.K.; Terli, R. Taxonomy and Polyphasic Characterization of Alkaline Amylase Producing Marine Actinomycete *Streptomyces rochei* BTSS 1001. *Int. J. Microbiol.* **2013**, *2013*, 276921. [[CrossRef](#)]
33. Ballav, S.; Kerkar, S.; Thomas, S.; Augustine, N. Halophilic and halotolerant actinomycetes from a marine saltern of Goa, India producing anti-bacterial metabolites. *J. Biosci. Bioeng.* **2015**, *119*, 323–330. [[CrossRef](#)]
34. Palomo, S.; González, I.; de la Cruz, M.; Martín, J.; Tormo, J.R.; Anderson, M.; Hill, R.T.; Vicente, F.; Reyes, F.; Genilloud, O. Sponge-derived *Kocuria* and *Micrococcus* spp. as sources of the new thiazolyl peptide antibiotic kocurin. *Mar. Drugs* **2013**, *11*, 1071–1086. [[CrossRef](#)] [[PubMed](#)]
35. Koch, E.; Löffler, I. Partial characterization of the antimicrobial activity of *Streptomyces antimycoticus* FZB53. *J. Phytopathol.* **2009**, *157*, 235–242. [[CrossRef](#)]
36. Gaba, M.; Mohan, C. Development of drugs based on imidazole and benzimidazole bioactive heterocycles: Recent advances and future directions. *Med. Chem. Res.* **2015**, *25*, 173–210. [[CrossRef](#)]
37. Lucas, G.B.; Campbell, C.L.; Lucas, L.T. Diseases Caused by Soilborne Fungi. In *Introduction to Plant Diseases: Identification and Management*; Springer: Boston, MA, USA, 1992; pp. 162–191.
38. Kara, M.; Oztas, E.; Ramazanoğulları, R.; Kouretas, D.; Nepka, C.; Tsatsakis, A.M.; Veskoukis, A.S. Benomyl, a benzimidazole fungicide, induces oxidative stress and apoptosis in neural cells. *Toxicol. Rep.* **2020**, *7*, 501–509. [[CrossRef](#)] [[PubMed](#)]
39. Ruma, Y.N.; Keniya, M.V.; Tyndall, J.D.A.; Monk, B.C. Characterisation of *Candida parapsilosis* CYP51 as a Drug Target Using *Saccharomyces cerevisiae* as Host. *J. Fungi* **2022**, *8*, 69. [[CrossRef](#)] [[PubMed](#)]
40. Seyedmousavi, S.; Verweij, P.E.; Mouton, J.W. Isavuconazole, a broad-spectrum triazole for the treatment of systemic fungal diseases. *Expert Rev. Anti-Infect. Ther.* **2015**, *13*, 9–27. [[CrossRef](#)]