

ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id>

Manuscript Revision Request, Invoice, Presentation Video Guidance for JESSD 159

2 messages

International Symposium JESSD <symposium.jessd@ui.ac.id> To: andi.setiawan@fmipa.unila.ac.id Cc: niluhratna.juliasih@fmipa.unila.ac.id, john.hendri@fmipa.unila.ac.id Tue, Sep 14, 2021 at 11:43 AM

Dear Andi Setiawan

Department of Chemistry, Faculty of Mathematics and Natural Science, Bandar Lampung

Thank you for submitting your manuscript "Screening Extract EtOAc Sponge-derived Fungi" with Manuscript ID **JESSD 159**. Congratulations, We are delighted to inform you that your paper on the 2nd International Symposium of Earth, Energy, Environmental Science, and Sustainable Development has been accepted for presentation at the symposium. The editorial team and a group of expert reviewers have assessed your submission and feel that it has the potential to be published in **IOP Conference Series: Earth and Environmental Science** (https://www.scimagojr.com/journalsearch.php?q=26916&tip=sid&clean=0) and the next process is the revision article and publication payment.

First. We would like to invite you to revise the paper and resubmit it for further review. Please see the attached reviewer comments for further details about the necessary revisions and also the similarity check results. We ask that you submit the revised version of your manuscript. **Please note:**

- 1. Your revised manuscript **should be accompanied** by a summary of your responses to the reviewers' comments.
- 2. Highlight your revision in the manuscript using a red font color
- 3. Send your revised manuscript by replying to this email and CC to symposium.jessd@gmail.com
- 4. Your revised results will be sent to the reviewer, but will not change the decision that your article has been accepted. Revisions are needed to ensure that all substantial suggestions and inputs have been carried out. It is very possible that there are still corrections and inputs, but the editor gives a tolerance limit of up to 2 rounds of review
- Once your manuscript is moved to publishing, our production editor will keep you informed of your article's progress in the production process. You will also receive proof of your manuscript for final review (galley proof).
- 6. You have 3 days to respond to this revise and resubmit request ending on **September 17th, 2021**, after which point we will presume that you have withdrawn your submission.

Second, publication payment. Please check the Invoice and payment guidelines (attached). For the proceedings, the maximum number of pages is 6 (short paper) and an additional fee will be charged if more than the specified page. The invoice will be revised if there is a request from the authors, including if you need a waiver. Please check on https://symposiumjessd.ui.ac.id/registration-fee/

Third, if the revised article and payment process is done, we are able to proceed with the Notice of Acceptance (LoA), Receipt, and publication process with the publisher.

Fourth, In order to be included in the publication, we need you to provide the following requirements for your presentation materials:

- 1. Presentation video, Make sure the video length doesn't exceed 9 minutes (2 minutes for opening-closing and 7 minutes for main material).please check these guidelines https://symposiumjessd.ui.ac.id/guideline-virtual-presentantions/. Also, an example video presentation from one of speaker here: https://drive.google.com/file/d/18_ApcENQYhPY7XIS0noZNFTk3DiCo-M3/view
- 2. Name of the author who will attend the virtual symposium

Please provide all the requirements by sending us the email with the subject "**JESSD 159** Presentation Materials" in 3 days after you received this, which is before **September 17th, 2021**. Send the materials to symposium.jessd@ui.ac.id and CC to symposium.jessd@gmail.com (mandatory).

Please make sure that we can have access to all the materials you provided. Also, we would like to ask for your permission to archive your presentation material on the 2nd JESSD Symposium website and YouTube channel and the public will be able to access it after the symposium.

unila.ac.id Mail - Manuscript Revision Request, Invoice, Presentation Video Guidance for JESSD 159

Fifth, before submitting your final revised article and payment process with us, we have to clarify our conditions to prevent any inconvenience during the whole publication process.

- 1. The JESSD Symposium acts as a link between authors, reviewers, and publishers. We help authors to improve the quality of their papers through input from reviewers who are experts in their fields.
- 2. The publisher has full responsibility to decide whether your paper is accepted or not. So in advance, we have to affirm from the beginning that: The JESSD Symposium's committee can't guarantee that the submitted paper in our conference would be accepted and published by the publisher.
- 3. In addition, we can't guarantee if in the future the proceedings are discontinued from Scopus. What we guarantee is that all papers received in this symposium will be submitted to the publisher and if accepted by the publisher, the proceeding and book chapter will be submitted for inclusion in leading Abstracting and Indexing services, including Web of Science, Scopus, CrossRef, Google Scholar. More information about our Terms and Conditions available on our official website.

Always include these two information in the subject line for all your communications with us regarding your paper:

- 1. Your manuscript ID
- 2. Information subject:
- a. Paper submission
- b. Paper revision
- c. Symposium receipt
- d. Publication receipt
- e. Article status

This will help us to respond to you most quickly and accurately. Please contact us if you have any questions or concerns. If in 1x24 hours you do not receive a reply from this email, try to contact us through symposium.jessd@gmail.com or contact our admin through Whatsapp (+62 819 3259 2196)

We're excited to move forward with your submission. Thank you for your interest, and we look forward to working with you on a successful conference.

Sincerely,

Herdis Herdiansyah 2nd JESSD Symposium Chair School of Environmental Science, Universitas Indonesia scholarhub.ui.ac.id/jessd | https://symposiumjessd.ui.ac.id/

6 attachments

- Format-Author Revision-JESSD-159.docx 32K
- Similarity Manuscript ID_JESSD-159.pdf 1576K
- Symposium Payment Guidelines.pdf
- Article Sample IOP.docx 1562K
- Template Guideline (IOP).docx 471K
- Invoice bank transfer-JESSD 159.pdf

ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id> To: osy.datul@gmail.com

Tolong dipelajari [Quoted text hidden]

6 attachments

Format-Author Revision-JESSD-159.docx

Tue, Sep 14, 2021 at 6:13 PM

32K

- Similarity Manuscript ID_JESSD-159.pdf
- Symposium Payment Guidelines.pdf 133K
- Article Sample IOP.docx 1562K
- Template Guideline (IOP).docx 471K
- Invoice bank transfer-JESSD 159.pdf 64K

12/15/21, 9:27 PM



ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id>

Your article from 2nd JESSD Symposium in IOP: EES is published

2 messages

International Symposium of JESSD <symposium.jessd@ui.ac.id> Wed, Dec 15, 2021 at 8:09 PM To: Widyastuti Unila <widyastuti.unila@gmail.com> Cc: John Hendri <john.hendri@fmipa.unila.ac.id>, Andi Setiawan <andi.setiawan@fmipa.unila.ac.id>

Dear Widyastuti

I hope you stay healthy and safe. Your paper entitled Application SEM-EDX in Biodegradation of Seafood Wastes by Sponge-Derived Actinomycetes 19C38A1 in Solid Fermentation with Manuscript ID JESSD-179 is published on IOP: Earth and Environmental Science, please check this link: https://iopscience.iop.org/issue/1755-1315/940/1. We will inform you once your paper is listed in Google Scholar and Scopus.

However, the publisher can't guarantee that your article will be listed on Scopus; it depends on Scopus's committee, as we have informed on the website (flyer and invitation). Also, we can't not guarantee that IOP: Earth and Environmental Science as a publisher will not be discontinued from Scopus. Beside Scopus, your article will be submitted for inclusion in leading Abstracting and Indexing services, including Web of Science, Scopus, CrossRef, Google Scholar. We will keep you updated.

It's a pleasure for us to work with you, however I would like to apologize if during the process there are any inconveniences. I hope we can meet again in the 3rd JESSD Symposium which is estimated to be in August or September 2022. More info will be updated soon.



Herdis Herdiansyah 2nd JESSD Symposium Chair School of Environmental Science, Universitas Indonesia scholarhub.ui.ac.id/jessd | https://symposiumjessd.ui.ac.id/

ANDI SETIAWAN <andi.setiawan@fmipa.unila.ac.id> To: International Symposium of JESSD <symposium.jessd@ui.ac.id>

Wed, Dec 15, 2021 at 9:26 PM

Dear Herdis Herdiansyah,

Thank you very much for your information.

Sincerely, Andi Setiawan.

[Quoted text hidden]

https://mail.google.com/mail/u/2/?ik=3b6409cac5&view=pt&search=all&permthid=thread-f%3A1719217739687706365&simpl=msg-f%3A1719217... 1/1

https://unila.turnitin.com/viewer/submissions/oid:7867:13072527/print?locale=en

IOP Conference Series: Earth and Environmental Science

PAPER • OPEN ACCESS

Application SEM-EDX in Biodegradation of Seafood Wastes by Sponge-Derived Actinomycetes 19C38A1 in Solid Fermentation

To cite this article: Widyastuti et al 2021 IOP Conf. Ser.: Earth Environ. Sci. 940 012048

You may also like

 Droduction of soybean in the application of <u>Rhizobium spp. and Actinomycetes spp.</u>
 D U Zainuddin, E Syam'un and A Dachlan

<u>Isolation and Identification of Bacteria and Actinomycetes Isolated from Wilting Banana Plants (Musa Sp.)</u>
 Saryono, P. Finna, P. Usman et al.

ellulolytic and Xylanolytic Actinomycetes selection to degrade Lignocellulosic biomass of Robusta coffee pulp (Coffee canephora) E Putri, Y Rukayadi, T C Sunarti et al.

View the article online for updates and enhancements.



the Electrochemical Society dvancing solid state & electrochemical science & technology

241st ECS Meeting

May 29 – June 2, 2022 Vancouver • BC • Canada Extended abstract submission deadline: **Dec 17, 2021**

Connect. Engage. Champion. Empower. Accelerate. Move science forward



This content was downloaded from IP address 110.137.38.189 on 15/12/2021 at 14:17

JESSD 2021

IOP Publishing

IOP Conf. Series: Earth and Environmental Science 940 (2021) 012048 doi:10.1088/1755-1315/940/1/012048

Application SEM-EDX in Biodegradation of Seafood Wastes by Sponge-Derived Actinomycetes 19C38A1 in Solid Fermentation

Widyastuti^{1,2}, J Hendri¹ and A Setiawan^{1*}

¹Department of Chemistry, Faculty of Mathematics and Natural Science, University of Lampung, Bandar Lampung, 35145, Lampung, Indonesia. ORCID ID 0000-0002-8119-0701

²Doctoral Program, Faculty of Mathematics and Natural Science, University of Lampung, Bandar Lampung, 35145, Lampung, Indonesia

¹widyastuti.unila@gmail.com, ¹john.hendri@fmipa.unila.ac.id, ^{1*}andi.setiawan@fmipa.unila.ac.id

Abstract. Biodegradation of chitin by microorganisms can produce derivative products that have economic value. This research aims to apply SEM-EDX analysis in observing the biodegradation process of seafood industrial waste by actinomycetes. Shrimp shells, cuttlefish bones and fish scales were obtained from the free market. In the early stages, the SEM-EDX spectrum analysis of the substrate showed almost the same carbon, oxygen, and mineral compositions. While the surface of each substrate is quite varied. On the second day of the fermentation, SEM image analysis showed that the growth rate of actinomycetes on each substrate was significantly different. The difference in growth was supported by SEM imaging data which showed damage to the surface of each substrate. Further analysis of the degradation products by HPLC on the second and third days showed the formation of glucosamine. It suspected that actinomycetes can break down shrimp shell waste into glucosamine. This information is very important as the basis for further research related to the optimization of the glucosamine and chitooligosaccharide (COS) production process using the solid fermentation method.

1. Introduction

The demand for seafood to date tends to increase [1]. This obviously has the consequence of increasing seafood product industrial waste. The low value of seafood product waste raises problems related to distribution, transportation, waste handling, and environmental degradation. Efforts to utilize the availability of seafood product industrial waste as a source of chitin and chitosan have been carried out. Seafood industry waste is chemically processed to extract chitin. Furthermore, chitin products can be converted into chitosan and its derivative products which have a higher economic value. This conventional method is quite effective but can reduce the quality of the surrounding environment [2]. In line with technological developments, solid waste treatment such as seafood industrial waste can be overcome by solid-state fermentation (SSF) methods. Currently, the application of the SSF method is a reliable alternative because the process is relatively cheaper when compared to other fermentation methods. In addition, the SSF process is more environmentally friendly and easy to perform. The



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOL Published under licence by IOP Publishing Ltd 1

JESSD 2021	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 940 (2021) 012048	doi:10.1088/1755-1315/940/1/012048

success of SSF is strongly influenced by the type of microorganisms utilized, the selection of media, and environmental conditions such as pH, minerals, and humidity [3].

Actinomycetes are decomposing microorganisms that play a very important role in the ecosystem [4]. Actinomycetes have also been widely utilized in solid fermentation processes for agro-industrial waste treatment. However, in the development of conventional optimization techniques in the fermentation process, it takes time because of the many parameters that must be measured. Considering this, this research aims to utilize the Scanning Electron Microscope-Electron Diffraction-X (SEM-EDX) analysis method in the biodegradation research of seafood industrial waste by actinomycetes. The application of SEM-EDX has been widely utilized in the biomedical field, but it is still very limited for researches on the development of the solid fermentation field. EDX microanalytic information contains semi-qualitative and semi-quantitative information. Furthermore, SEM imaging data can also be performed quickly [5].

This research will describe the stages of actinomycetes ability to convert chitin in seafood waste into its derivative products. Supported by HPLC analysis for the resulting biodegradation products, the incorporation of the data obtained will accelerate optimal conditioning in the solid fermentation process. This research will certainly be very useful as a first step in the development of solid fermentation processes.

2. Method

2.1. Materials

2.1.1. Isolate Actinomycetes

In this research, the isolate actinomycetes 19C38A1 was selected from the deposit of Unit Pelayanan Teknis Laboratorium Terpadu dan Sentra Inovasi Teknologi (UPT LTSIT) University of Lampung, which was obtained from the isolation of the 19C38 sponge off the coast of Oluhuta, Gorontalo. The isolates were juvenile in ISP2 media and 1% chitin colloid.

2.1.2. Substrate for Solid State Fermentation

Shrimp shells, cuttlefish bones, and fish scales were obtained from free-market Lempasing, Bandar Lampung, while fish scales were obtained from the Pringsewu market. Fresh waste samples were washed with water to separate from impurities and then dried at 55°C overnight. Dry waste samples were stored in the refrigerator for further experiments.

2.2. Procedure

All work and equipment related to microorganisms was carried out under sterile conditions.

2.2.1. Microscopic analysis.

The characteristics of the isolate 19C38A1 were examined microscopically using the cover glass culture method [6] Briefly, a sterile coverslip was inserted at an angle of 45° in the center for 2% chitin agar. After 4 days of incubation at 32°C, the cover slip was removed and placed on a clean slide. The coverslip was finally observed under a light microscope (100X), using a Zeiss Observer A1 microscope.

2.2.2. Solid state fermentation.

The SSF process was carried out using waste substrates of shrimp shells, cuttlefish bones and fish scales. Each substrate weighing 10 grams was placed in a petri dish. Then the media was inoculated with 1 ml of actinomycetes suspension and incubated statically at 32°C. Observation of actinomycetes growth was carried out every day and observed using SEM. While the results of biodegradation were analyzed using high performance liquid chromatography (HPLC).

JESSD 2021

IOP Publishing doi:10.1088/1755-1315/940/1/012048

IOP Conf. Series: Earth and Environmental Science 940 (2021) 012048

2.2.3. Analysis SEM-EDX.

At the initial stage, the respective surface and composition were analyzed by SEM-EDX. The next stage, actinomycetes growth and substrate surface growth were observed using SEM. For analysis purposes, the gold coating process was carried out for 15 minutes. Once gold coated, the specimen could be examined with an electron microscope (SEM-EDX Zeiss EVO MA 10). Gold coated specimens on stub were observed by SEM at a high electron (EHT) voltage of 10 kV, using a secondary electron detector. The surface is scanned at low magnification until a growth line is clearly detected. The surface area with a clear and intact spore structure was then selected and examined at a higher magnification. Suitable surface in the preparation is ready to be photographed [7].

2.2.4. Analysis Glukosamine

The results of solid fermentation were analyzed using HPLC 20A Shimadzu based on the glucosamine formed. Fermentation of 10 grams of solid waste was extracted with 10 ml of distilled water, then the extraction results were centrifuged at 8000 rpm. After that, 1 mL of the filtrate was added with 4 mL of absolute ethanol (EtOH), then centrifuged at 8000 rpm to separate the precipitate formed. The precipitate is then redissolved with distilled water and ready for analysis. Glucosamine was analyzed chromatographically using a C18 Shimadzu column and acetonitrile (CH₃CN: H₂O) mobile phase flowed in a gradient (30% to 70%) for 10 minutes, the column temperature was adjusted with CTO 20A at 40°C, the results of the separation were detected with the PDA detector LC20AD.

3. Results and discussion

3.1. Isolate actinomycetes 19C38

Isolate 19C38A1 is one of the actinomycetes collections deposited at UPT LTSIT, Universitas Lampung. This isolate was obtained from the sponge coded 19C38 showed in Figure 1a. The sponge was collected from the waters off the coast of Oluhuta, Gorontalo in 2019. Macroscopic observations on isolate 19C38A1 appeared gray, and microscopic observations using a light microscope into Figure 1b showed nocardioform mycelium substrate characteristics (mycelium that decomposed into bacilli and *cocci*) [8]. The results of the chitinolytic test of isolate 19C38A1 in Figure 1c observed a clear zone on the outside around the isolated colony. This indicates the ability of isolate 19C38A1 to utilize chitin as a carbon source in its life cycle. Or it can be said that 19C38A1 isolate was able to degrade chitin biomaterial. The ability of actinomycetes to produce has been previously reported, but researches on the application of chitinase from actinomycetes are still very rare.



Figure 1. (a) Under water image sponge 19C38; (b) mikroscopic 19C38A1; (c) Chitinolitic clear zone.

3.2. Solid State Fermentation Substrate (SSF)

3.2.1. Substrate. In addition to the type of microorganism, another determining factor that supports the success of SSF application is the type of substrate to be selected. In this research, it focused on the utilization of substrates from seafood industry waste, including shrimp shells, cuttlefish bones, and

JESSD 2021	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 940 (2021) 012048	doi:10.1088/1755-1315/940/1/012048

fish scales. The three types of waste are sources of chitin biopolymer. The ability of microorganisms to produce chitinase that can degrade chitin has been previously reported by Sneha Ja et al. [9]. Theoretically, substrates obviously containing chitin can be degraded by chitinase-producing microorganisms. In the degradation process, chitin can be converted into its derivative products such as chitosan and chitoligosaccharides. At the initial stage of characterization, the three types of substrates utilized were analyzed using SEM-EDX.

The results of observations using SEM-EDX are shown in Figures 2a, 2b, and 2c. The semiqualitative and quantitative data of the EDX spectrum shown each substrate has a similar composition of Carbon (C), Oxygen (O) while the mineral content (Ca, Mg, Zn) was quite varied. This difference in composition will certainly affect the growth rate of microorganisms on the surface of the substrate. As it has been reported that the variation of actinomycetes media composition such as carbon, oxygen, nitrogen, pH, temperature and metal ions affect the production of chitinase.



Figure 2a. SEM-EDX spectrum of shrimp shell substrate

ries 53.91	[wt.%] [at.%] 53.91 62.39		
	53.91 62.39		
	Show where the second s	8.51	
ries 42.39	42.39 36.83	7.51	
ries 3.71	3.71 0.79	0.36	
ries 0.00	0.00 0.00	0.00	
ries 0.00	0.00 0.00	0.00	1
tal: 100.00	100.00 100.00		
	Ca		Zn
	ries 0.00 ries 0.00	ries 0.00 0.00 0.00 ries 0.00 0.00 0.00 stal: 100.00 100.00 100.00	ries 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.



Figure 2b. SEM-EDX spectra of cuttlefish bone substrate

Further analysis using SEM imaging on each substrate showed that the surface of each substrate was very varied. This difference is obviously related to differences in mineral composition which consequent differences in the physical and chemical properties of the substrate.

JESSD 2021 IOP Publishing IOP Conf. Series: Earth and Environmental Science 940 (2021) 012048 doi:10.1088/1755-1315/940/1/012048



Figure 2c. SEM-EDX spectrum of fish scale substrate

3.2.2. Solid State Fermentation. The results of the fermentation process in a solid-state against 3 types of waste consisting of shrimp shells, strong bones, and fish scales using isolate 19C38 are shown in Figures 3, 4, and 5.



Figure 3. Isolate 19C38A1 on shrimp shell surface (a) observation on 2nd day; (b) shrimp shell surface after washing



Figure 4. Isolate 19C38A1 on cuttlefish bone surface (a) observation on 2nd day; (b) cuttlefish bone surface after washing

5

https://unila.turnitin.com/viewer/submissions/oid:7867:13072527/print?locale=en

JESSD 2021

IOP Publishing

IOP Conf. Series: Earth and Environmental Science 940 (2021) 012048 doi:10.1088/1755-1315/940/1/012048



Figure 5. Isolate 19C38A1 on the surface of fish scales (a) observations on 2nd day; (b) fish scale after washing

The results of SEM imaging on the second day, shown in Figure 3a, it was observed that actinomycetes produced spores and hyphae on the shrimp shell substrate, in Figure 4a observed the presence of hyphae on the cuttlefish bone substrate, whereas in Figure 5a there was no growth of actinomycetes. This indicated that the substrate affects the growth rate of actinomycetes. Further analysis of SEM imaging after each washing process observed damage to the surface of the shrimp shell substrate as depicted in Figure 3b and partial damage to the cuttlefish bone in Figure (4b) and no damage was observed on the surface of fish scales in Figure 5b. This indicated the growth rate of actinomycetes related to the ability of the mycelium to enter the substrate. To obtain more complete information from the results of the biotransformation process, the next experiment will focus on the production of chitin degradation analysis on shrimp shell media.

3.3. Glucosamine Analysis

The results of the analysis of waste biodegradation products observed on days 0, 2 and 3, are shown in Figure 6. The chromatogram at the initial conditions (day 0) showed that the substrate did not contain glucosamine (Figure 6a). The results of further observations were observed that the chromatogram peaks at retention time (rt) 2,593 minutes, and 2,567 minutes (Figures 6c and 6d) respectively on the second and third days. The peak that appeared was similar to the retention time with standard glucosamine (rt = 2,584) min (Fig. 6b). The interpretation of glucosamine peaks on day 2 and day 3 showed a continuous pattern of enzyme degradation. This can be seen from the decrease in the peak intensity of glucosamine on day 3 and the difference in the chromatogram pattern with increasing degradation products to form more polar compounds.



6

https://unila.turnitin.com/viewer/submissions/oid:7867:13072527/print?locale=en

JESSD 2021 IOP Publishing IOP Conf. Series: Earth and Environmental Science 940 (2021) 012048 doi:10.1088/1755-1315/940/1/012048



Figure 6. Glucosamine test results of SFF (a) control; (b) measurement Standard glucosamine; (c) the 2^{nd} day measurement; (d) 3^{rd} day measurement.

4. Conclusion

Based on the data obtained, it can be concluded that the application of SEM-EDX in the SSF process is very helpful to provide microanalysis data in a semi-micro-qualitative and semi-micro quantitative manner quickly. This information will be very important as the basis for developing the SSF research.

Acknowledgements

The authors would like to thank to The Deputy of Research Strengthening and Development, The Ministry of Research and Technology/National Agency for Research and Innovation of the Republic of Indonesia, for Basic Research with Grand No. 4233/UN26.21/PN/2020. The authors also thank the UPT LTSIT Laboratory, the University of Lampung for providing scanning electron microscopy facilities, and thanks to Fernady Djailani for collecting samples, and Masayoshi Arai for photographing underwater marine organisms.

References

- Mao X, Guo N, Sun J and Xue C 2017 Comprehensive utilization of shrimp waste based on biotechnological methods A review J. Cleaner Prod. 143 814–823
- [2] Madalano M, Vassalini I, and Alessandri I 2020 Green Routes for the Development of Chitin/Chitosan Sustainable Hydrogels Sustain Chem. 1 325-344
- [3] Brzezinska M M, Walczak E, Lalke-Porczyk and W. Donderski 2010 Utilization of shrimp-shell waste as a substrate for the activity of chitinases produced by microorganisms *Plolish Journal of Environmental Studies* 10 177-182
- [4] Barka E A, Vatsa P, Sanchez L, Gaveau-Vaillant, N Jacquard, C Klenk, H. P. Clément, C Ouhdouch Y, and Van Wezel GP 2016 Taxonomy Physiology and Natural Products of Actinobacteria *Microbiol Mol. Biol. Rev.* 80 1–43
- [5] Scimeca M, Bischetti S, Lamsira H K, Bonfiglio R and Bonanno E, 2018 Energy Dispersive X ray (EDX) microanalysis: A powerful tool in biomedical research and diagnosis Eur J Histochem 62 1 2841
- [6] Goodfellow M, Kampfer P and Dusse H J 2012 Bergey's Manual of Systematic Bacteriology".
 - 2nd New York NY USA Springer
- [7] Kumar V, Bharti A, Gusain O, and Bisht G S, 2011 Scanning Electron Microscopy of Streptomycetes Without Use of Any Chemical Fixatives Scanning
- [8] T Bennur, A R Kumar, S Zinjarde and V Javdekar 2015 Nocardiopsis species Incidence, ecological roles and adaptations *Microbiological Research* 174 33-47
- [9] Sneha Jha, Hasmukh A Modi and Chaitanya K J, 2016 Characterization of extracellular chitinase produced from Streptomyces rubiginosus isolated from rhizosphere of Gossypium sp. Cogent Food & Agriculture 2 1 13