

Sumardi Sumardi

# 2-Characterization of protease from bacillus sp. on.pdf

Sources Overview

13%

OVERALL SIMILARITY

1	Universitas Sumatera Utara on 2022-01-20	4%
	SUBMITTED WORKS	
2	download.garuda.kemdikbud.go.id	2%
	INTERNET	
3	eprints.untirta.ac.id	1%
	INTERNET	
4	jurnal.ugm.ac.id	<1%
	INTERNET	
5	adoc.pub	<1%
	INTERNET	
6	Universitas Airlangga on 2020-09-24	<1%
	SUBMITTED WORKS	
7	web-2.oauife.edu.ng	<1%
	INTERNET	
8	Sriwijaya University on 2020-05-20	<1%
	SUBMITTED WORKS	
9	Afriani ., Arnim ., Yetti Marlida, Yuherman .. "Isolation and Characterization of Lactic Acid Bacteria Proteases from Bekasam for use as...	<1%
	CROSSREF	
10	D Agustina, T Surtiningsih, Y S W Manuhara, A Arifiyanto, M Malewa. " Study of Cellulase Activity Produced by sp., and on (L.) Beauv. E...	<1%
	CROSSREF	
11	biodiversitas.mipa.uns.ac.id	<1%
	INTERNET	
12	bmcmedgenomics.biomedcentral.com	<1%
	INTERNET	
13	A R Ramadhan, Z Bachruddin, Widodo, Y Erwanto, C Hanim. "Isolation and selection of proteolytic lactic acid bacteria from colostrum ...	<1%
	CROSSREF	
14	www.lib.unair.ac.id	<1%
	INTERNET	
15	Birla Institute of Technology on 2015-07-06	<1%
	SUBMITTED WORKS	
16	pdfs.semanticscholar.org	<1%
	INTERNET	
17	pertambangan.fst.uinjkt.ac.id	<1%
	INTERNET	

18	www.hindawi.com	<1%
INTERNET		
19	www.omicsonline.org	<1%
INTERNET		
20	www.smujo.id	<1%
INTERNET		

Excluded search repositories:

None

Excluded from document:

- Bibliography
- Quotes
- Citations
- Small Matches (less than 10 words)

Excluded sources:

Sumardi, Rochmah Agustrina, Christina Nugroho Ekowati, Yovita Selvie Pasaribu. " Characterization of protease from sp. on medium containing FeCl exposed to magnetic field 0.2 mt ", IOP Conference Series: Earth and Environmental Science, 2018, crossref, 69%

Universitas Sumatera Utara on 2021-06-16, submitted works, 6%

123dok.com, internet, 6%

ejournal.undip.ac.id, internet, 5%

Y S Soeka, M Ilyas. "Production and characterization of cellulases derived from saprophytic fungi *Penicillium bilaiae* InaCC F16", IOP Conference Series: Earth and Environmental Science, 2020, crossref, 5%

iopscience.iop.org, internet, 4%

id.123dok.com, internet, 3%

D Nirmala, P Yudha, D Cahyanto. "The effect of pH and incubation time on rude protease enzymes activity of *Bacillus mycoides* from anchovy isolates (*Stolephorus* sp.)", IOP Conference Series: Earth and Environmental Science, 2021, crossref, 3%

Chao Zhang, Xiao-Jun Lan, Xue-Tao Bai. "Study on the biological hazards and control measures of the Dunhuang manuscripts", IOP Conference Series: Earth and Environmental Science, 2021, crossref, 3%

Chairudin, Agustiner, J Permadi. "Efficacy of application time of *Penicillium* sp. suspension on White Root Fungus (*Rigidoporus lignosus*) in Nutmeg (*Myristica fragrans*)", IOP Conference Series: Earth and Environmental Science, 2021, crossref, 3%

Zulyusri, Desyanti, Y Hidayat. "Pathogenicity of entomopatogenic fungi isolates from infected pest crop against dry wood termites *Cryptotermes* sp. (Ioptera: Kalotermitidae)", Journal of Physics: Conference Series, 2021, crossref, 2%

Norazwina Zainol, Emine Seydametova, Jailani Salihon, Peter Convey. "Screening of significant factors affecting pravastatin production by *Penicillium* sp. ESF21P", IOP Conference Series: Materials Science and Engineering, 2020, crossref, 2%

D A Svetlov, A A Piksaykina, M V Vildyaeva, N V Piksaykin. "Biodestruction of Buildings and Constructions and Improvement of Their Longevity and Environmental Friendliness at the Basis of Application of Bio-Resistant Materials", IOP Conference Series: Materials Science and Engineering, 2021, crossref, 2%

M Pratika, M Ananda, I N Suwastika. "Protease activity from bacterial isolates of *Nepenthes maxima* reinw. ex nees", Journal of Physics: Conference Series, 2021, crossref, 2%

N A Sidorova, N A Chechkova. "Results of primary screening of proteolytically active microorganisms that are promising for the transformation of organic pollution", IOP Conference Series: Earth and Environmental Science, 2021, crossref, 2%

I Indrawati, N Rossiana, T. R Hidayat. " Antibacterial Activity of Bacterial Endophytes from Kupa Plant Miq. (Merr & Perry) Against Pathogenic Bacteria ", IOP Conference Series: Earth and Environmental Science, 2018, crossref, 2%

Ratna Stia Dewi, Hana. "Screening of microfungi from spent mushroom for decolorizing and removing heavy metals from batik effluent and its toxicity", IOP Conference Series: Earth and Environmental Science, 2021, crossref, 2%

"The 4th International Conference on Biological Sciences and Biotechnology", IOP Conference Series: Earth and Environmental Science, 2019, crossref, 2%

Susiana Purwantisari, Achmadi Priyatmojo, Retno Peni Sancayaningsih, Rina Sri Kasiamdari, Kadarwati Budihardjo. "Lignification on Potatoes by Application of *Trichoderma viride*", IOP Conference Series: Earth and Environmental Science, 2020, crossref, 2%

I Irdawati, S Syamsuardi, A Agustien, Y Rilda. " Screening of Thermophilic Bacteria Produce Xylanase from Hot Spring South Solok ", IOP Conference Series: Materials Science and Engineering, 2018, crossref, 2%

G A K Sutariati, L O S Bande, A Khaeruni, Muhidin, L Mudi, R M Savitri. " The effectiveness of preplant seed bio-invigoration techniques using sp. CKD061 to improving seed viability and vigor of several local upland rice cultivars of Southeast Sulawesi ", IOP Conference Series: Earth and Environmental Science, 2018, crossref, 2%

D Iswantini, N Nurhidayat, H Ferit. " Alcohol Dehydrogenase of strain for Measuring Alcohol Electrochemically ", IOP Conference Series: Earth and Environmental Science, 2017, crossref, 2%

Muhammad Aiman Faiz Mohd Zaidi, Mohammad Najmi Masri, Wee Seng Kew. "Microbiologically Influenced Corrosion of Iron by Nitrate Reducing *Bacillus* Sp", Journal of Physics: Conference Series, 2021, crossref, 2%

Mohammed Fadhil Abood, Haider Mousa Hamzah, Dhafer Fakri Al-Rawii. "Production of Cellulase and bioethanol by ethanol-tolerant coculture of *Bacillus cereus* and *Fusarium solani*", *Journal of Physics: Conference Series*, 2021, [crossref](#), 2%

M I Said. "Potential development of poultry feather waste resources as raw material in industry: A review", *IOP Conference Series: Earth and Environmental Science*, 2020, [crossref](#), 2%

N Zainol, S R Rahim. " Effect of garlic solution to sp. removal ", *IOP Conference Series: Materials Science and Engineering*, 2018, [crossref](#), 2%

Haiping Zou, Xiaohong Yu, HongLiang Wu. "Effect of sintering insulation time on the properties and microstructure of graphite-copper composite", *Journal of Physics: Conference Series*, 2021, [crossref](#), 2%

S W Suciwati, P Manurung, S Sembiring, R Situmeang. "Comparative study of *Cladophora* sp. cellulose by using FTIR and XRD", *Journal of Physics: Conference Series*, 2021, [crossref](#), 2%

A A Lukyanov. "Micromycetes in ampelocenosis soil under different keeping system", *Journal of Physics: Conference Series*, 2020, [crossref](#), 2%  
[china.iopscience.iop.org](#), internet, 1%

[www.science.gov](#), internet, 1%

[repository.lppm.unila.ac.id](#), internet, 0%

[digilib.unila.ac.id](#), internet, 0%

PAPER • OPEN ACCESS

# Characterization of protease from *bacillus* sp. on medium containing FeCl<sub>3</sub> exposed to magnetic field 0.2 mt

To cite this article: Sumardi *et al* 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **130** 012046

View the [article online](#) for updates and enhancements.

## You may also like

- [Keratinase-producing fungi from local environmental samples of Far South Thailand and their efficiency in hydrolyzing keratinous wastes](#)  
Abdullah D. Dalee, Mustakim Chehama, Khosiya Sali et al.

14 [Study of Cellulase Activity Produced by \*Penicillium\* sp., \*Aspergillus niger\* and \*Trichoderma viride\* on \*Imperata cylindrica\* \(L.\) Beauv. Enrichment Media](#)  
10 D Agustina, T Surtiningsih, Y S W Manuhara et al.

17 [The Effect of pH and incubation time on crude protease enzymes activity of \*Bacillus mycoides\* from anchovy isolates \(\*Stolephorus\* sp.\)](#)  
6 D Nirmala, P Yudha and D Cahyanto

## Characterization of protease from *bacillus* sp. on medium containing FeCl<sub>3</sub> exposed to magnetic field 0.2 mt

Sumardi<sup>1\*</sup>, Rochmah Agustrina<sup>1</sup>, Christina Nugroho Ekowati<sup>1</sup> and Yovita Selvie Pasaribu<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty mathematic and natural science, University of Lampung

Jalan Prof. Dr. Soemantri Brodjonegoro No. 1 Bandar Lampung, Lampung 35145

\*Email: [sumardi\\_bio@yahoo.co.id](mailto:sumardi_bio@yahoo.co.id) Telephone: +6285216391087 Fax : +62-721-704625

**Abstract** This purpose of this research is to determine the character of the protease enzymes from *Bacillus* sp. on media content of FeCl<sub>3</sub> exposed to 0.2 mT magnetic field. The data obtained were analyzed descriptively. The result showed that protease enzyme without Fe resulted in the highest activity at pH 8, temperature 30°C with the addition of activator Mn<sup>2+</sup>, and V<sub>max</sub> of 0.28 U/ml, and K<sub>m</sub> of 4.60 U/ml. The protease enzyme on media without magnetic field exposure and containing Fe yielded the highest activity at pH 8, temperature 30°C with the addition of activator Mn<sup>2+</sup>, and V<sub>max</sub> of 0.33 U/ml, and K<sub>m</sub> of 5.64 U/ml. The protease enzyme on medium with magnetic field exposure and use Fe as inductors have the highest activity at pH 9, the temperature of 55°C with the addition of activator Mn<sup>2+</sup>, and V<sub>max</sub> of 0.35 U/ml, and K<sub>m</sub> 10.04 U/ml.

**Keywords:** determine, inductor

### 1. Introduction

The production of the enzyme with high activity can be achieved by optimizing of medium conditions such as pH, temperature, carbon and nitrogen sources, or by providing additional treatment such as exposure to a magnetic field of the medium [1]. Ion Fe is an essential nutrient for almost all microorganisms in growth because, of its role as a cofactor in some enzymes.

Iron or Fe is one of the ferromagnetic metal ions. Fe is an essential nutrient for optimal growth of a cell. Fe acts as a cofactor for several enzymes is required in the biochemical processes of cell, respiratory and photosynthesis reaction [2].

Fe in the form of Fe<sup>2+</sup> is more easily utilized by bacteria as Fe<sup>2+</sup> is soluble in water. If the Fe in the form of Fe<sup>3+</sup> the user is first converted to Fe<sup>2+</sup> with the rest of the energy that has been used for growth [3]. Magnetic field exposure on bacterial medium containing Fe metal will cause the magnetization in the medium to motion moment - the moment of the dipole into the direction of the magnetic field. The change in direction dipole moment Fe suspected that would cause changes in enzyme activity is controlled by the bacteria.

Based on the research Selfiana et al [4] demonstrated that exposure to 0.2 mT magnetic field for 10 minutes on Fe metal ions affect the activity of *Bacillus* sp. to produce a protease enzyme. The addition of the metal ions Fe in salt form FeCl<sub>3</sub> as much as a 0.01% exposed to the magnetic field 0.2 mT for



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

10 minutes yields the highest enzyme activity 0.06 U/ml.

So far, the protease enzyme results of these studies have not been characterized. Characterization needs to be done to determine the optimum pH and temperature, the metal ions that can increase the activity of the enzyme, as well as chemical kinetics optimal protease enzyme so that to obtain high enzyme activity. Therefore, characterization studies protease from *Bacillus* sp. on media containing  $\text{FeCl}_3$  was exposed to a magnetic field of 0.2 mT necessary.

## 2. Method

### 2.1. The Culture of *Bacillus* sp.

Rejuvenation culture is done by scraping the ose culture of *Bacillus* sp., *Bacillus* sp which is a collection of the laboratory of microbiology FMIPA UNILA. Then grown on a modified medium composition comprising Mendel's with skim; 0.5 g; Yeast extract 0.35 g; Water Tryptone 0.35 g; NaCl 0.2 g;  $\text{KH}_2\text{PO}_4$  0.245 g;  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  0.035 g;  $(\text{NH}_4)_2\text{SO}_4$  0.175 g in 100 mL of distilled water and agar 1.5 g as the compactor. Later, it was tilt incubated at 37°C in an incubator for two days.

### Proteolytic test on a modified solid media Mendel's

$\text{FeCl}_3$  which has been exposed by the magnetic field 0.2 mT for 10 minutes is used as an inductor in the growth media. The proteolytic test consisted of three treatments as follows:

a. **Treatment of media without inductors ( $\text{FeCl}_3$ ).**

As a control treatment, a modified solid media Mendel has not given exposure to a magnetic field and are not given the inductor.

b. **Media treatment with the inductor ( $\text{FeCl}_3$ ).**

Treatment Negative is treated using a modified solid media Mendel's and given inductor  $\text{FeCl}_3$  neither the media nor the inductors are being exposed to a magnetic field.

c. **Media treatment with exposure to a magnetic field in the inductor ( $\text{FeCl}_3$ ).**

Positive treatment is treated using a modified solid media Mendel's and given inductor  $\text{FeCl}_3$ .  $\text{FeCl}_3$  was exposed to 0.2 mT magnetic field for 10 minutes before use.

Observations existence of a clear zone formed around colonies of bacteria do when the bacterial cultures were incubated for 18 hours at a temperature of 37°C. Colonies of bacteria and a clear zone formed around colonies of bacteria measured in diameter and further defined Proteolytic Index (PI):

$$\text{PI} = \frac{\text{the diameter of clear zone} - \text{the diameter of colony}}{\text{the diameter of the colony}} \quad (1)$$

### 2.2. Enzyme Production

Enzyme production is done by the respective treatment in the media without the addition of agar. Protease enzyme production is done by inoculating 5 ml starter *Bacillus* sp. in 45 ml liquid media Mendel's modified in Erlenmeyer 250 ml with the same treatment as the proteolytic test.

All treatments shake culture was incubated in an incubator at 120 rpm at a temperature of 40 ° C with a 24-hour long incubation time that is adjusted with previous studies. Extraction is done by centrifuging the protease enzyme of bacterial growth media at a speed of 10,000 rpm for 10 min at 4°C.

### 2.3. Protease Activity Test

0.1 ml protease sample of crude extract was added to the mixture using a 0.5 ml casein substrate as follows: A total of 0.5 ml of phosphate buffer pH 7.0:0.1 M was added and incubated at 39 °C for 10 minutes. After wards plus 0.5 ml of 0.1 M TCA then incubated at 39 °C for 10 minutes, and centrifuged at 10,000 rpm at 4 °C for 10 minutes. Then 0.375 ml of the supernatant was taken and added with 1:25 ml of Na<sub>2</sub>CO<sub>3</sub> 0.4 M and 0:25 ml Folin reagent, incubated at 37 °C for 20 minutes [5].

Observations were carried out by measuring the Optical Density (OD) at a wavelength of 578 nm. Blank value calculation is done in the same way, where the protease sample was replaced with distilled water. While the standard value calculation is done by replacing the protease samples with tyrosine 5 mM.

Protease activity was calculated in units of PU (protease units) per ml of the enzyme extract. One protease unit is defined as the amount of enzyme that can produce one μmol of tyrosine per minute on measurement conditions were measured using the following formula:

$$PU = \frac{Asp - Asbl}{Ast} \times \frac{1}{T} \quad (2)$$

Description:

PU :	Unit Activity Protease (Units/ml)
Asp :	Value Absorbance sample
Ast :	Value Absorbance standard
Abl :	Value Absorbance Blank
T :	Time

### 2.4. Determination of optimum pH

Effect of pH on the enzyme activity is tested by reacting a solution of crude extract enzyme and 0.25% incubated casein substrate at 39 °C in a buffer 0.05M different series, namely: pH 4.5 (citrate buffer), pH 6.7 (phosphate buffer), pH 8.9 (tris-HCl buffer ), pH 10,11,12 (glycine-NaOH buffer).

### 2.5. Determination of Optimum Temperature

The effect of temperature on the enzyme activity is tested by reacting a solution of crude extract enzyme at optimum pH with temperature variation test 25, 30, 35, 40, 45, 50 55, 60, 65, and 70 °C.

### 2.6. Effect of Metal Ion on Protease Activity

Effect metal ions Ca<sup>2+</sup>, Mn<sup>2+</sup>, Cu<sup>2+</sup>, Mg<sup>2+</sup>, Fe<sup>3+</sup> in the form of a salt of each - each CaCl<sub>2</sub>, MnCl<sub>2</sub>, CuCl<sub>2</sub>, MgCl<sub>2</sub>, and FeCl<sub>3</sub> as an activator. As an inhibitor of the protease activity used is metal ion chelating compounds, ethylene diamine tetraacetic acid (EDTA) by reacting a solution of crude extract of enzymes, enzyme substrates used as casein 0.25% with one mM of the metal ion. The enzyme is incubated with the metal ions and inhibitors for 10 minutes at a temperature and pH optimum then tested the activity of the enzyme.

### 2.7. Determination of K<sub>m</sub> and V<sub>max</sub>

Determination of K<sub>m</sub> and V<sub>max</sub> is done by testing the activity enzyme at its optimum temperature and pH variation casein different substrate concentrations of 0%, 0.5%, 1%, 1.5%, 2% and 2.5%. The results of the activity test then tabulated and made the curve relationship between concentration and specific activity of the enzyme casein. Once it is put in the Lineweaver-Burk linear equations. Km value and Vmax is obtained by the formula [6]:

$$\frac{1}{V_o} = \frac{k_m}{V_{maks}} \frac{1}{[S]} + \frac{1}{V_{maks}} \quad (3)$$

Specification:

$V_o$  : initial velocity on the substrate concentration [S]


$V_{max}$  : Maximum Velocity

$K_m$  : Constanta of Michaelis-Menten enzyme for a particular substrate


### 3. Results and Discussions

#### 3.1. *Bacillus* sp. protease production


Proteolytic activity of *Bacillus* sp. on solid media Mendel's modified evident from the formation of a clear zone around bacterial colonies growing. With the protease enzyme produced by *Bacillus* sp., casein in the *Bacillus* sp media. The hydrolyzed into smaller peptides and amino acids, their degradation process is characterized by the formation of a clear zone around the colony.



**Figure 1.** The test results of proteolytic media without  $\text{FeCl}_3$  and no magnetic exposure. Isolate *Bacillus* sp. showed small the clear zone in protease medium



**Figure 2.** The test results on the media proteolytic treatment with Fe without exposure to magnetic fields. Isolate *Bacillus* sp. showed the clear zone in protease medium.



**Figure 3.** The test results on the media proteolytic treatment with Fe and exposure to magnetic fields. Isolate *Bacillus* sp. showed the clear zone in protease medium.

Proteolytic index value (PI) at the highest media *Bacillus* sp. after incubation for 18 hours of media treatment obtained by Fe outcome magnetic field exposure is equal to 5.62. The highest protease enzyme activity of 0.10 U/ml was obtained from the same media (Table 1).

**Table 1.** Comparison of proteolytic index and Protease Activity each treatment

Treatment media	Proteolytic Index (PI)	Protease Activity (U/ml)
Without Fe and no magnetic field exposure	0.54	0.08
without magnetic field exposure and containing Fe	2.65	0.07
with magnetic field exposure and containing Fe	5.62	0.10

The area that formed a clear zone indicates that the bacteria in a culture that has a high ability to change the substrate contained in a growth medium [7]. Proteolytic activity of bacteria is influenced by several factors given during treatment. The factors that influence proteolytic activity for each isolates are pH, substrate and enzyme concentration, temperature and the presence of activators and inhibitors. The high value of the proteolytic index on media treatment is given Fe results of magnetic field exposure is suspected because  $\text{FeCl}_3$  which has been exposed by the magnetic field has induced a higher power and become more positive effect on the proteolytic activity of *Bacillus* sp.


Fe ions in salt  $\text{FeCl}_3$  which is ferromagnetic have magnetic properties that the direction of the dipole moments of the movement direction of the external magnetic field. As a result of rectification movement can survive even if the external magnetic field has no [8].

If  $\text{FeCl}_3$  which has been exposed by the magnetic field is added to the bacterial growth media that is stored on the magnetic properties of media expected to affect the growth of bacteria grown on these

media. As reported by Sudarti et al [9] that the magnetic field can directly influence the metabolic activity of cells one to produce the enzyme. Farisna and Zulaikha [10] explains that *Bacillus* sp. able to grow on solid medium containing Fe as Fe metal ions required for the metabolism of bacterial cells even though in small amounts.

### 3.2. Determination of optimum pH

The optimum pH test results protease enzyme growth of *Bacillus* sp media prove that the treatment medium without Fe, pH optimum pH 8 to achieve the protease activity of 0.11 U/ml. While the optimum pH of protease enzyme in the treatment medium by Fe without exposure to magnetic fields reaches pH 8 with protease activity of 0.15 U/ml. As well as the optimum pH of the protease enzyme in the treatment by the medium Fe magnetic field exposure results reaches a pH of 9 with protease activity of 0.11 U/ml (Figure 4).



**Figure 4.** The activity of protease in difference pH

*Bacillus* sp isolates that it was inoculated in the three media treatments resulted in active proteases in the range of pH 8 and 9. Several studies have shown that the optimum activity of proteolytic enzymes varies from pH 8-10 [11]. Isolates of *Bacillus* sp. isolated from the digestive tract of chicken, protein degradation can grow at pH 7-9 [12]. Protease *licheniformis* B-05 LHSB also has an optimum pH 9. Some protease from genus *Bacillus* other has an optimum pH range 8-10 [13].


The optimum pH difference in the three treatment medium is thought to occur because of the influence of pH on the enzyme active site. The research result Ikehara et al [14], suggests that exposure to magnetic fields ELF has the effect of a reversible bond NH and CN of a peptide bond, and changing the secondary structure of the beta cells and alpha-helices in proteins, so easily have been the attachment of H<sup>+</sup> and OH<sup>-</sup> on the side active enzymes.

At pH 4-7 protease enzyme activity tend to be smaller than the optimum pH for the number of H<sup>+</sup> ions that interfere with the enzyme active site, while at pH 10-12 decreased protease enzyme activity for enzyme functional groups disturbed by the many OH<sup>-</sup> that causes the enzyme denatured.

When FeCl<sub>3</sub> which has been exposed by the magnetic field is added in bacterial growth media will cause a difference in the number of H<sup>+</sup> and OH<sup>-</sup> ions in the enzyme active site. Thus, there is a difference in the pH of the enzyme treated with magnetic field exposure are not given exposure [9].

### 3.3. Determination of optimum Temperature

Each enzyme has the maximum activity at a certain temperature; the enzyme activity will increase with increasing temperatures up to optimum temperature is reached. The optimum temperature of the protease enzyme in the treatment medium without Fe reaches a temperature of 30°C with protease activity of 0.12 U/ml. Meanwhile, the optimum temperature of the protease enzyme treatment media by Fe without exposure to the magnetic field reaches a temperature of 30°C with protease activity of 0.44 U/ml. As well, the optimum temperature of the protease enzyme treatment given media exposure results Fe magnetic field reaches a temperature of 55°C with protease activity of 0.13 U/ml (Figure 5).



**Figure 5.** The activity of protease in difference temperature

Further increase in temperature will cause decreased enzyme activity. At temperatures lower than the optimum temperature, the enzyme activity is also low, due to lower activation energies available. The energy needed to create the conditions of an active complex level, both of the molecules of enzyme or substrate molecules.

Protease enzyme was produced on all three medium treatment has the optimum temperature 30 and 55°C. This is confirmed by the results of an intensive search Fitriani [15], where he reported isolates of protease *Bacillus* KUB B19 CC BPPT actively working at a temperature of 30 to 65°C with the optimum temperature at 60°C. Based on the optimum temperature of the protease is classified as thermophilic protease active means to work at high temperatures. Thermophilic proteins that are different characteristics in its structure compared to mesophilic proteins. Thermophilic proteins have more salt bridge structure coupled with the presence of the main chain hydrogen bonds are plentiful when compared to mesophilic proteins. Also, this protein has an amino acid residue more hydrophobic than mesophilic proteins. This will lead to increased activity of enzyme proteins at high temperatures [16]


The difference between the values of the activity of Fe media with media treatment without treatment by Fe without exposure to a magnetic field is thought to occur due to exposure to a magnetic field causes a change of Ca ion movement<sup>2+</sup> and the increasing rate of movement of the ion. Such conditions lead to changes in transport in the cell membrane that affect cellular metabolism activity. It can have an impact on the process of cell growth. Increased bacterial growth will result in differences in enzyme activity value [17].

The temperature difference optimum media treatment by Fe the result of exposure to magnetic fields with the protease enzyme in media other treatments, allegedly caused by treatment  $\text{FeCl}_3$  which is exposed to a magnetic field causes motion moments of the dipole direction of the magnetic field from the outside so that the influence of electromagnetic waves that pass through it. While the results of research Ikehara et al. [14], suggests that exposure to ELF magnetic fields have reversible effects on the NH and CN bond of peptide bonds, and changing the secondary structure of the beta cells and alpha-helical proteins. As a result is the release of bonds of constituent proteins and changes in the molecular structure.


### 3.4. Effect of Metal Ion and Specific Inhibitor

Addition of  $\text{Mn}^{2+}$  ions to a final concentration of 5 mM can raise the activity of the protease enzyme in the treatment medium without Fe into 0.27 U/ml. The addition of  $\text{Mn}^{2+}$  ions on the addition of  $\text{Mn}^{2+}$  ion concentration of 1 mM protease enzyme activity in the media treatment by Fe without exposure to a magnetic field, increasing to 0.18 U/ml. But with a final concentration of 5 mM was not able to raise the activity of the protease enzyme in media exposure treatment without magnetic field and added inductors Fe. The same thing happened on the media treatment of the protease enzyme by Fe result of exposure to magnetic fields. The addition of  $\text{Mn}^{2+}$  ions on the addition of  $\text{Mn}^{2+}$  ion concentration of 1 mM protease enzyme activity increased to positive treatment 0.30 U/ml. But, with a final concentration of 5 mM was not able to raise the activity of the protease enzyme in the treatment with magnetic field exposure and added inductor Fe (Figures 6 and 7).

The addition of EDTA inhibitor compounds with a final concentration of 5 mM greatly decreases the activity of the protease enzyme in the three media such treatment, until there is no activity at all three enzymes (Figure 6 and 7).



**Figure 6.** Effect of metal ions and specific inhibitor of the protease enzyme activity in a concentration of 1 mM



**Figure 7.** Effect of metal ions and specific inhibitor of the protease enzyme activity in concentrations of 5 mM

Test results show that a strong activator for the protease enzyme of isolates of *Bacillus* sp. the third was inoculated on media treatment is  $Mn^{2+}$  at a final concentration of 5 mM able to enhance the catalytic activity of the protease enzyme of *Bacillus* sp. The influence of the addition of metal ions from a few studies have been done, such as the activity measurement results with the addition of metal ions on the substrate casein, metal ion  $Mn^{2+}$  (5 mM) is a potent activator of protease *Bacillus epidermidis*, which can increase the activity of protease respectively 3 and 2 times folding of the protease control [18]. For *Bacillus* A1 strain  $Mn$  metal ion<sup>2+</sup> as an activator can increase the enzyme activity according to the research it Adinarayana et al [19]. Inhibitors are compounds that can inhibit the enzyme activity. Extra activity inhibitor compound is one method to identify the enzyme. Differences in protease activity value on each concentration allegedly caused by the difference itself. Enzymes in treatment media which has the highest enzyme activity at a concentration of 1 mM decreased in concentration 5mM, allegedly can not grow optimum conditions of high metal concentrations, but the optimum conditions a low concentration.

Effect of protease inhibitors on the activity was shown in Figure 10, EDTA is a chelating metal (metalloprotease inhibitor) 5 mM slow down to up do not occur enzyme activity in it. The addition of 2 mM EDTA in protease of *Bacillus thermophilic* activity strain HS08 lowered relative to respectively 6% and 4% [20]. The results of analysis show that the protease activity of *Bacillus* sp. perfect inhibited by EDTA so that it can be concluded that these proteases are classified into metalloprotease protein [21]. Chantawannakul [22] reported a protease inhibitor that inhibits the fibrinolytic activity of strains *Bacillus* fermented soybeans 38 isolates from Thailand, namely 1.10 fenantrolin and identified as metalloprotease.

### 3.5. Protease Kinetics Chemistry

Determination of the value of  $K_m$  and  $V_{max}$  is useful to determine the enzyme kinetics that can be known how much bonding enzymes and enzyme substrate and how fast it can perform the activity. Determination of the value of  $V_{max}$  and  $K_m$  is done by measuring the initial velocity at various concentrations of protease enzyme substrate.

From the research that has been conducted shows that the increasing concentration of substrates will increase the reaction rate. However, at a certain concentration limits, no increases the reaction rate even if the substrate concentration is enlarged. This is because at a certain substrate concentration limit, all active parts have been met by the substrate or has been saturated with the substrate [23]. In this study, obtained the value of  $K_m$  and  $V_{max}$  of the enzyme protease in the treatment medium without Fe each - each amounted to 4.60 mM and 0.28 U/ml. It shows one ml of the enzyme can produce products as much as 0.28 units. As well, the value of  $K_m$  and  $V_{max}$  of the enzyme protease on media treatment by Fe without exposure to magnetic fields, respectively - each of 5.64 mM and 0.33 U/ml. Show one ml of the enzyme can produce products as much as 0.33 units. Meanwhile, the value of  $K_m$  and  $V_{max}$  of the enzyme protease media Fe treatment by exposure to magnetic fields result, each - each of 10.04 mM and 0.35 U/ml. This means that one ml of the enzyme can produce products as much as 0.35 units (Table 2).

**Table 2.** Comparison of  $K_m$  and  $V_{max}$  values between each treatments

Treatment media	$K_m$ (U/ml)	$V_{maks}$ (U/ml)
Without Fe and no magnetic field exposure	4.60	0.28
without magnetic field exposure and containing Fe	5.64	0.33
with magnetic field exposure and containing Fe	10.04	0.35

According to Kovacs et al [24], exposure to magnetic fields on a protein enzyme trypsin can produce different ranges in a redox reaction due to changes in the shape of the helix peptide bonds and hydrogen bonds, thus increasing the  $K_m$  significantly. A value of  $K_m$  are high indicates a low affinity for the substrate and instead of.

In general, the crude extract of the enzyme will give a high value of  $K_m$ . It can be caused by the presence of contaminants other enzymes which also have an affinity for the substrate. The higher the value of  $K_m$ , the concentration required to achieve half of the maximum reaction rate is also higher. Maximum velocity ( $V_{max}$ ) will be achieved when all of enzyme molecule has interacted with the substrate or enzyme has been saturated with the substrate, so that an increase in substrate concentration will not increase the rate of reaction [25].

#### 4. Conclusion

From the discussion in this study, the character of the protease enzyme of *Bacillus* sp on media content of  $FeCl_3$  in said the magnetic field of 0.2 mT. Where the characters are, among others: the optimum pH and temperature influence of metal ions and inhibitors, as well as the value of enzyme kinetics.

#### Acknowledgments

The authors gratefully acknowledge that the research is supported by University of Lampung of Indonesia ([www.unila.ac.id](http://www.unila.ac.id)). The support is under the research grant of Hibah Pascasarjana-UNILA of Year 2017 Contract Number 809/UN26.21/PP/2017.

## References

- [1] Mubarik NR (2001) Imobilisasi Protease *Bacillus subtilis* ATCC 6633 menggunakan Matriks Gel Poliakrilamida. *Jurnal Hayati* 8(1): 11-14.
- [2] Shiyang H, Y Feng, H Ren, and Y Zhang (2011) The Impact of Iron Oxide Magnetic Nanoparticle on the Soil Bacterial Community. *School of Biological Science and Medical Engineering*. 11: 1408 – 1417.
- [3] Sutariningsih E (2012) *Potensi Pengguna Magnetic Pengguna Fe dan Hg*. Fakultas Biologi Universitas Gajah Mada. Yogyakarta.
- [4] Selfiana I, Sumardi, Agustina R (2016) The Effect of Metal Ions Fe and Zn Exposed to Magnetic Field 0.2 mT on The Production of Protease in *Bacillus* sp..*Prosiding USR International Seminar on Food Security*. 2016(1):73-74.
- [5] Bergmeyer HV and Grassl (1983) *Method of Enzymatic Analysis 2*. Verlag Chemie, Weinheim.
- [6] Wahyuntari B dan Hendrawati H (2012) Properties of an Extracellular Protease of *Bacillus megaterium* DSM 319 as Depilating Aid of Hides. *Microbiology Indonesia*. 6(2). doi : 10.5454/mi.6.2.4.
- [7] Vijayaraghavan, Ponnuswamy and Samuel GPV (2013) A simple method for the detection of protease activity on agar plates using bromocresolgreen dye. *Journal Biochemistry Technology* 4(3): 628-630.
- [8] Sipahutar WS (2015) *Efek Waktu Wet Milling dan Suhu Annealing Terhadap Sifat Fisis, Monostruktur dan Magnet Dari Flakes NdFeB*. Prosiding Fisika Universitas Sumatra Utara.
- [9] Sudarti, Nurhayati, E Ruriani, and VT Hersa (2014) Prevalence of Salmonella Typhimurium on Gado-Gado Seasoning by Treatment of Extremely Low Frequency (ELF) Magnetic Field. *Artikel-ELF-Salmonella*. Jember University.
- [10] Farisna ST and E Zulaikha (2015) *Resistensi Bacillus Endogenik Kalimas Surabaya terhadap Logam Besi (Fe)*. Jurusan Biologi FMIPA Institut Teknologi Sepuluh Nopember.
- [11] Soeka YS and Sulistiani (2014) Karakterisasi Protease *Bacillus subtilis* A1 InaCC B398 Yang Diisolasi Dari Terasi Samarinda. *Berita Biologi Bidang Biologi*. Puslit-LIPI. 13(2).
- [12] Nurhayati, Ninik and Sumaryanto (2012) Protease production by *Bacillus subtilis* ATCC 1633. *Proceeding The Is' Conferece on Industrial Enzyme and Biotechnology*. 201-205.
- [13] Johnvesl B, and Naik GR (2001) Study on production of thermostable alkaline protease from thermophilic and alkaliphilic *Bacillus* sp. JB99 in a chemically defined medium, *Journal. Process Biochemistry*., 37: 139-144.
- [14] Ikehara T, Yamaguchi H, Hosokawa K, Miyamoto H, and Aizawa K (2003) Effects of ELF Magnetic Fields on Membrane Protein Strucutre of Living HeLa Cells Studied by Fourier Transform Infrared Spectroscopy. *Bioelectromagnetics* 24(7): 457-464.
- [15] Fitriani S (2013) Purifikasi Parsial dan Karakterisasi Enzim Protease dari Isolat B19 KUB BPPT CC.[skripsi]. Bogor(ID) :Institut Pertanian Bogor.
- [16] Sadeghi A, Shahidi F, Mortazavi SA, and Mahalati N (2008) Evaluation of Different Parameters Effect on Maltodextrin Production by  $\alpha$ -amilase Termamyl 2-x. *World Applied Sciences Journal*. 3 (1): 34-39.
- [17] Kristinawati, Andika (2015) Pengaruh Lama Paparan Medan Magnet *Extremly Low Frequency* Terhadap pH dan Kadar Air Pada Proses Pembuatan Keju Jenis *Cream Cheese*. [skripsi]. Jember(ID) :Universitas Jember.
- [18] Baehaki A (2012) Kolagenase *Bacillus licheniformis* F11 asal Palembang dan aplikasinya pada pembuatan peptida kolagen bioaktif. [tesis]. Bogor (ID) :Institut Pertanian Bogor.
- [19] Adinarayana K, P Ellaiah and DS Prasad (2003) Purification and partial characterization of thermostable serine alkaline protease from a newly isolated *Bacillus subtilis* PE-1. *American Association of Pharmaceutical Scientists Pharmaceutical Sciences Technology*. 4(4):440-448.

- [20] Guang-rong H, Wei L, Dai DH, Wei-lian H (2011) Purification and characterization of a novel extracellular chitinase from thermophilic *Bacillus* sp. Hu1, *African. Journal. Biotechnology*. 10:2476-2485.
- [21] Teufel P and Gotz F (1993) Characterization of an extracellular metalloprotease with elastase activity from *Staphylococcus epidermidis*. *Journal. Bacteriology*. 175(13):4218-4224.
- [22] Chantawannakul P.(2001) Characterization of proteases of *Bacillus* strain 38 isolated from traditionally fermented soybean in Northern Thailand. *Science Asia* 28:241-248.
- [23] Kumar D and Bhalla TC (2004) Purification and Characterization of a Small Size Protease from *Bacillus* sp. APR-4. *Journal Experiment Biology* 42: 515-517.
- [24] Kovacs, Phillip E, Richard LV, and Pedro JJA (1997) The Effect of Static Magnetic Fields on Biological System : Implications for Enhanced Biodegradation. *Critical Reviews in Environment Science and Technology*. 27(4): 319-382.
- [25] Trismilah and Lutfi A (2009) Membran polyethersulfone dan regenerated cellulose untuk ultrafiltrasi: Pengaruh pH terhadap proses ultrafiltrasi xilanase. *Jurnal Sains Dan Teknologi Indonesia*. 11(2): 76-83.