

PAPER NAME

**Antioxidant and Cytotoxic Activities of M
elinjo (Gnetum gnemon L.) Seed Fraction
s on HeLa Cell Line**

AUTHOR

Asep Sukohar

WORD COUNT

4637 Words

CHARACTER COUNT

25799 Characters

PAGE COUNT

6 Pages

FILE SIZE

522.5KB

SUBMISSION DATE

Jul 28, 2022 12:15 PM GMT+7

REPORT DATE

Jul 28, 2022 12:16 PM GMT+7

● **12% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 9% Internet database
- 7% Publications database
- Crossref database
- Crossref Posted Content database
- 6% Submitted Works database

● **Excluded from Similarity Report**

- Bibliographic material
- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources

Antioxidant and Cytotoxic Activities of Melinjo (*Gnetum gnemon* L.) Seed Fractions on HeLa Cell Line an *In Vitro*

Asep Sukohar¹, Suharyani^{1,2,*}, Sutyarso², Hendri Busman², Nuning Nurcahyani², Evi Kurniawaty¹

Asep Sukohar¹, Suharyani^{1,2,*},
Sutyarso², Hendri Busman², Nuning
Nurcahyani², Evi Kurniawaty¹

¹Faculty of Medicine, University of Lampung,
INDONESIA.

²Magister Biologi, ¹⁰Faculty of Mathematics
and Natural Sciences, University of Lampung,
INDONESIA.

Correspondence

Suharyani

Faculty of Medicine, Magister Biology,
Faculty of Mathematics and Natural
Sciences, University of Lampung,
INDONESIA.

E-mail: suharyanisodarmo@gmail.com

History

- Submission Date: 19-01-2022;
- Review completed: 14-04-2022;
- Accepted Date: 27-05-2022.

³ DOI : 10.5530/pj.2022.14.71

Article Available online

<http://www.phcogj.com/v14/i3>

Copyright

© 2022 Phcogj.Com. This is an open-
access article distributed under the terms
of the Creative Commons Attribution 4.0
International license.

ABSTRACT

Introduction: Medicinal plants have been investigated for possible anti-cancer effects. One of them is *Gnetum gnemon* L. (melinjo). This study aims determined *in vitro* antioxidant activity and the cytotoxic effects of polar, semipolar and non polar melinjo seed fractions against HeLa cell line. **Methods:** The melinjo seed were extracted with ethanol as a solvent. Then, the fractionation was done using liquid-liquid extraction method with three different polarity solvent. Cytotoxic activity was carried out using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) colorimetric assay in HeLa cell lines at concentrations ranging from 25 to 400 µg/mL. Antioxidant activity was determined by the diphenyl picryl hidrazil (DPPH) radical scavenging method. **Results:** Phytochemical screening indicated the presence of phyto-constituents like flavonoids, terpenoid and tanin. The DPPH scavenging activity by the melinjo seed aqueous, ethyl acetate and n-hexane fraction was 733,12 ± 18,95 µg/mL; 68,40 ± 1,9 µg/mL and 2035,70 ± 65,59 µg/mL, respectively. The cytotoxic activity of the melinjo seed fractions showed that the ethyl acetate was the most active fraction against HeLa cell line with IC₅₀ value 45,27 µg/mL. **Conclusion:** In this study, we have observed that the melinjo seed fractions exhibited antioxidant and cytotoxic activity against HeLa cell lines. This is presumably due to the content of phytochemicals and stilbenoids such as resveratrol and gnetin C. Melinjo seeds are more potent as anticancer Compared with other plants that also contain RSV such as grape extract (*Vitis vinifera* L.) against lung cancer cells (A549). From the three fractions, the ethyl acetate fraction had the highest antioxidant and cytotoxic effect compared to the water and n-hexane fractions. *Gnetum gnemon* L. can be considered as a potential source of anticancer agents. However, more research is needed to determine the mechanism of action.

Key words: Antioxidant, Cytotoxic, Melinjo, Anti-cancer, HeLa.

INTRODUCTION

The main causative factors in the induction of various chronic and degenerative diseases including cancer is oxidative stress.¹ Oxidative stress is imbalance condition between the formation and elimination of reactive oxygen species (ROS). There is an increase in ROS production and decreased antioxidant capability of the cell.² Reactive oxygen species (ROS) are chemically reactive molecules in cell and associated with various biological processes, including cell proliferation, differentiation and programmed cell death.³ Antioxidants nutrients have been shown to be involved in ROS detoxification.⁴ Antioxidants are divided into three groups, namely the phenol, the amin and the amino-phenols.⁵

¹⁵ Cervical cancer is one of the most common female cancers worldwide. It is caused by human papilloma virus (HPV) infection.⁶ Based on the viruses ability to promote the proliferation of infected cells and lead to the malignant transformation, HPV can be subdivided into three classifications. There are low, intermediate and high risk oncogenic potentials.⁷ ¹⁴ Persistence of infection is more common with the high-risk oncogenic HPV types, causing cervical cancer cases in about 99,7%.^{6,8}

Cancer treatment involves several approaches which include: surgical intervention, chemotherapy and radiation therapy or often a combination.⁹ The chemotherapeutic drugs is

severely limited because of their side effects. On the other hand, natural compounds have the potential to selectively exert cytotoxic effect on cancer cells without affecting normal cells.¹⁰ Natural compound are complex chemical molecules present in various parts of plant. They have pharmacological or biological activities for the treatment of cancer and other diseases.¹¹ One of the plant that have natural compound is melinjo (*Gnetum gnemon* L.). It is known that melinjo seed extract (MSE) contains trans-resveratrol (tRV); isorhapontigenin; gnetin C, gneunosides A, C, and D and gnetin L.¹² Melinjo seed extract (MSE) has been reported to have a broad spectrum of pharmacological effects such as anticancer,¹⁰ inhibitory angiogenesis,¹³ antibacteria,¹⁴ antioxidant.¹⁵ However, its possible effects of melinjo seed fraction against cervical cancer remain uncertain. The liquid-liquid extraction method aims to classify compounds based on the level of polarity.¹⁶ This study will determine the antioxidant activity, cytotoxic and anti-proliferative of melinjo seed fractions on HeLa cervical cancer cell line. Antioxidant activity is associated with anti-proliferative activity (cytotoxic activity) on cancer cells¹⁷ and calculated as half maximal inhibitory concentration (IC₅₀) value.

MATERIAL AND METHOD

Materials

Red melinjo seeds was collected from Tanjung Agung Village, Teluk Betung District, Pesawaran,

Cite this article: Sukohar A, Suharyani, Sutyarso, Busman H, Nurcahyani N, Adjeng ANT, et al. Antioxidant and Cytotoxic Activities of Melinjo (*Gnetum gnemon* L.) Seed Fractions on HeLa Cell Line an *In Vitro*. Pharmacogn J. 2022;14(3): 559-564.

Lampung, Indonesia in March 2021. HeLa cells are collection from the Cytogenetics and Cell Culture laboratory, Faculty of Medicine, Padjadjaran University, Bandung, Indonesia.

Determination and preparation of melinjo seed fraction

Plant identified has been done by Botanical Laboratory, Faculty of Mathematics and Natural Sciences, University of Lampung. Melinjo seed was sorted, washed, and drained, then the peel was separated from its seeds. The seeds were air-dried then extracted according to¹⁸ with modified. Dried melinjo seeds were macerated with ethanol (Merck, New York, USA.) as an extraction solvent with ratio 1:3 w/v for 2 days. The extract was evaporated in 40°C, 50rpm with rotary evaporator. The MSE was further separated by liquid-liquid extraction. The thick extract was dissolved in warm water at a ratio of 1:10, then partitioned with n-hexane and ethyl acetate. The results that obtained ethyl acetate fraction and water fraction than evaporated in 40°C, 50rpm with rotary evaporator.

Phytochemical analysis

The qualitative study of phytochemicals aims to identify chemical constituents that have pharmacological activity, namely alkaloids, flavonoids, saponins, tannins and terpenoids. The test is based on visual observation between color changes and/or the formation of a precipitate after the addition of specific reagents.¹⁹

Antioxidant activity determination

Antioxidant activity was determined using spectrophotometer method modified from.⁹ Two milliliter samples were mixed with 2 mL 2,2-diphenyl-1-picrylhydrazyl (DPPH) solution in ethanol. The mixture was incubated at room temperature (25 ± 2°C) in dark room for 30 min. Ascorbic acid was used as positive control. Absorbance was read at 516 nm using a spectrophotometer UV/VIS Perkin Elmer Lambda 25 and the percentage inhibition was calculated with following equation:

$$\text{DPPH radical scavenging (\%)} = \frac{[\text{Abscontrol} - \text{Abssample}]}{\text{Abscontrol}} \times 100$$

Where Abs control is the absorbance of Ascorbic acid and Abs sample is the absorbance of the melinjo seed fractions. IC₅₀ value is the concentration of the melinjo seed fractions required to inhibit 50% of DPPH radical scavenging. A scatter graph was plotted to obtain IC₅₀ value.

Cell culture and cytotoxicity assay

HeLa cells were cultured in Roswell Park Memorial Institute (RPMI) 1640 medium supplemented with 10% fetal bovine serum and 100 U/ml of penicillin streptomycin. The cells were cultured at 37°C in an incubator with 5% CO₂. The cytotoxic effect of the melinjo seed fraction on the HeLa cells was assessed using MTT assay. Briefly, cells were seeded in a 96-multiwell plates (2 × 10⁴ cells/well), and treated in triplicate with various concentrations of melinjo seed fractions (25–400 µg/ml) for 24 h and then incubated with MTT reagents for 2 h at 37°C. Absorbance was measured at 550 nm. The minimum concentration of fraction that was toxic to cancer cells was recorded as the effective drug concentration. The IC₅₀ values, concentrations which reduce the absorbance of treated cells by 50%, were graphically obtained from the dose response curves.

Statistical analysis

All assays were carried out in triplicates and data presented as mean ± standard deviation. EXCEL package 2016 was used for the analyses of mean, standard deviation, and percentage inhibition while linear regression analysis was used to determine the IC₅₀. To determine

the effect of concentration on viability cell between treatment analyzed using ANOVA followed by LSD test with a 95% confidence level.

RESULT

Total yield of melinjo seed fractions

The total yield of melinjo seed fraction by using the solvents, namely, aqueous, ethyl acetate and n-hexane were 0,6 g; 0,18 g; 0,18 g (weight/weight), respectively with reference to the air dried plant material.

Phytochemical analysis

Preliminary screening of melinjo seed fractions aqueous, ethyl acetate and N-hexane showed the presence of diversity of phytochemical constituents. The aqueous and ethyl acetate fraction has flavonoids, terpenoid and tannin. And the n-hexane fraction only has terpenoid.

Antioxidant activity

In this study, the different solvent for partition between aqueous, ethyl acetate and N-hexane seed fraction of *Gnetum gnemon* L. were subjected to DPPH free radical scavenging assay. The antioxidant capacity of the fraction was compared with ascorbic acid as the standard antioxidant. The IC₅₀ value of ascorbic acid was 5.42 µg/mL with a correlation coefficient (R²) 0.999. The DPPH scavenging activity by the *Gnetum gnemon* seed aqueous, ethyl acetate and N-hexane fraction is presented in figure 1). The results showed that there is significant difference (p<0,05) between the obtained DPPH scavenging activities of the three fractions based on one way Anova test. The ethyl acetate fraction has highest antioxidant activity compare to the aqueous and N-hexane fraction.

Effect of *Gnetum gnemon* L. seed fractions on viability HeLa cell line

The result of MTT assays revealed that the *Gnetum gnemon* L. seed fractions decreased the percent viability cells but different extent. Ethyl acetat was found more cytotoxic than the aqueous and N-hexana fractions toward HeLa cancer cell line. The IC₅₀ values of aqueous, ethyl acetat and N-hexana of *Gnetum gnemon* L. seed fractions against HeLa cancer cell line are represented in (Table 1). The results show that the ethyl acetat fraction of melinjo seed has highest cytotoxic activity.

This fractions also revealed the morphological changes and shrinkage of cell leading to cell death (Figure 2). Some apoptotic cells indicated with shrinkage and irregular shape, while the HeLa cell shape was polygonal and attaching to the matrix.




Figure 1: The values presented are mean ± standard deviation, n = 3. The results show that there is a significant difference based on the One Way Anova test at a confidence level of 5%.




Figure 2: Morphological changes showing of HeLa cell line. (a) control cell; (b) treatment with aqueous fraction; (c) treatment with ethyl acetate fraction; (d). treatment with n-hexane fraction (magnification was 100x).




Figure 3: Cells viability after treatment with *Gnetum gnemon* L. seed fractions.

Table 1: IC₅₀ value of *Gnetum gnemon* L. seed fractions against HeLa cell line (µg/mL).

Melinjo seed fraction	IC ₅₀ value (µg/mL)	
Aqueous fraction	No data	
Ethyl acetate fraction	45,27	<i>p</i> <0,05*
N-hexane fraction	209,06	
Doxorubicin	2,38	

The values presented are mean ± standard deviation, n = 3. The results show that there is a significant difference based on the One Way Anova test at a confidence level of 5%.

Reduction of cell viability depend on dose reduction was reported in cell treated with all fraction. The most potent in reduction cell viability was ethyl acetate fraction. Treated with *Gnetum gnemon* L. seed ethyl acetate fraction significantly enhanced of the mortality of cancer cells at the low concentration (50 µg/mL) (Figure 3).

DISCUSSION

The use of herbal medicines in the treatment of cancer due to its diversity phyto-metabolic content has received attention.²⁰ Some

active compounds including flavonoids, diterpenoids, triterpenoids and alkaloids are known to have anticancer effects.²¹ Based on the results of phytochemical tests on the melinjo seed fraction found the chemical compounds including flavonoid, terpenoids and tannins. This compound can deactivate free radicals by donating hydrogen atoms to free radicals.²² Hydrogen atom transfer is dominant to scavenge radical.²³ The role of antioxidants is the interactions depend on oxidative free radicals. The discoloration of DPPH indicates the antioxidant scavenging of the sample such as phenolic compounds, especially phenolic acids and flavonoids.²⁴ There is a relationship between ROS with oncogene function and suppression function. ROS Reduction activity will decrease MAPK activity, then decrease cFos and cJun activity. C-fos and cJun activities will have an impact decrease the activity of CDK 4 and 6 which in turn causes inhibition of the G1 phase which plays a major role in cell cycle proliferation. This can cause balance instability and predominately cause apoptosis in cells due to oncogene suppression.²⁵

Based on the results of the cytotoxic test conducted on the melinjo seed fraction of HeLa cervical cancer cells, the three fractions (polar, semipolar and non-polar) were able to kill HeLa cells indicated by decrease in cell viability (%). The IC₅₀ value of the ethyl acetate and n-hexane fractions obtained is <1000 µg/mL. Extracts that have IC₅₀ values <1000 µg/mL have potential as anti-cancer.²⁶ If compared with previous studies that explored the anticancer potential of *Gnetum gnemon* L. in other cancer cells such as HT-29, Colon-26, MCF-7, DU145, PC-3 and PTEN-CaP8, the IC₅₀ value was 35-39 g/mL.¹⁰ This difference could be due to the content of secondary metabolites such as gnetin C (GC) and resveratrol (RSV) in the sample. The cytotoxic effects of plant are usually caused by the secondary metabolites contained in them. Several ingredients such as alkaloids, polyphenols, saponins, tannins, flavonoids and stilbenoids (GC and RSV) were found in melinjo (*Gnetum gnemon* L.).^{27,28} Research with other plants that also contain RSV, such as in the study of²⁹ which explored grape extract (*Vitis vinifera* L.) against lung cancer cell line (A549), the IC₅₀ value was 61,21±1.45µg/mL. It can be concluded that melinjo seeds have more potential as anti-cancer. As seen in the Table. 1, the ethyl acetate fraction had the highest cytotoxicity effect. This possibility is because in the ethyl acetate fraction contain several active compounds such as flavonoids, triterpenoids, and alkaloids that have been shown to have anticancer effects.²¹ The ethyl acetate fraction gave a higher cytotoxic effect due to the less polar flavonoid compounds. Ethyl acetate is able to dilute less polar flavonoid compounds such as isoflavones, flavonones, methylated flavones and flavonols.²² In addition, in the melinjo seeds contain stilbenoid such as resveratrol and gnetin C. According to the research of,³⁰ the resveratrol content in ethyl acetate solvent was 686 mg/kg while in water and n-hexane solvent was not detected.

Melinjo seed fractions also be able to cell destruction based on their morphology (Figure 2). Normal HeLa cells are polygonal or round with extensive cytoplasm.³¹ After being given the treatment, the cells became spherical and disconnected from each other and showed cellular shrinkage and nuclear condensation.³² Cell shrinkage as a result of cell dehydration. Loss of intracellular fluid including K⁺ and Cl⁻ causes apoptotic cell death under various inducers of apoptosis.³³ Figure 3 shows that the number of living cells decreases as the concentration and duration of exposure increase. This is presumably because the ethanol extract of melinjo seeds also contains resveratrol compounds, because resveratrol is a compound that is easily soluble in alcohol.³⁴ The study stated that the viability of HeLa cells treated with 20 mol/L resveratrol was reduced after 24 hours³⁴ and significantly reduced after 48 hours.³⁴

Resveratrol can induce apoptosis through intrinsic and extrinsic apoptotic pathways. Based on research by,³⁵ after treatment with resveratrol, mitochondrial membrane disturbances and apoptosis-

related markers occurred, such as increase Bax/Bcl-2 ratio, and the form of caspase-8 and caspase-3. Resveratrol activates both Fas ligand-mediated and mitochondrial apoptosis in HL-60 cells by increasing Bax expression, release of cytochrome C into the cytosol.³⁶ This is in line with research conducted by³⁷ which stated that there was a decrease in mitochondrial membrane potential in HPV 16 positive cervical cancer cell lines. In addition, there was an increase in p53 expression leads to caspase-9 activation. Caspase-9 is a mitochondrial initiator in the intrinsic apoptotic pathway activated by septameric apoptosomes by mitochondrial release of cytochromes C, Apaf-1 and pro-caspase-9 and subsequent activation of caspase-3. Furthermore, activated caspase-9 can cleave and activate caspase-3. Caspase-3 as a caspase effector initiates the degradation process of apoptotic processes such as cell shrinkage, membrane blebbing, DNA fragmentation and the formation of apoptotic bodies (small cell units). Research by³⁸ proved an increase in caspase-3 and caspase-9 in DU145 prostate cancer cells treated with RSV. In addition to resveratrol, there is a secondary metabolite Gnetin C (GC), which is a component in the extract of melinjo seeds. It is known that the amount of GC in melinjo seeds is 28.0 mg/g.³⁹ Gnetin C has the ability to suppress endothelial cell function related to angiogenesis,¹³ inhibit proliferation, migration.⁴⁰ Based on the *in-vivo* test, it was able to induce apoptosis through the caspase-3/7-independent mechanism (extrinsic pathway).¹⁰

Based on phytochemical tests, melinjo seed fraction contains tannin compounds that play role in inducing phosphorylation of the tumor suppressor p53. It can increase the expression of target genes such as p21 and BAX.⁴¹ The inhibition of the G1/S phase transition is mainly dependent on the p21 and p27 levels. Tannins are able to inhibit the expression of p21 and p27.⁴² Protein 27 is a protein that binds to cyclin and CDK so that there is an obstacle to the S phase. Tannins are able to inhibit the expression of cyclin D1, cyclin E and CDK-4.⁴¹ On the other hand tannins can also change the expression of mitochondrial pore factors Bax, Bcl-2 and Bcl-XL.⁴²

CONCLUSIONS

In the present study, we have observed that the melinjo seed fractions exhibited antioxidant and cytotoxic activity against HeLa cell lines. When compared with other plants that also contain RSV such as grape extract (*Vitis vinifera* L.) against lung cancer cells (A549), it can be concluded that melinjo seeds have more potential as anti-cancer. From the three fractions, the ethyl acetate fraction had the highest antioxidant and cytotoxic effect compared to the water and n-hexane fractions. Further studies are needed to determine the required mechanism of action.

ACKNOWLEDGEMENTS

Authors are thanks to Institute for Research and Community Service (LPPM) University of Lampung which had fully funded this research through DIPA BLU 2021 – Terapan Scheme and to Cell Culture and Cytogenetic Laboratory, Faculty of Medicine, Padjadjaran University, which had assisted the implementation of this research.

CONFLICTS OF INTEREST


None.

REFERENCES

- Oirere EK, Anusooriya P, Malarvizhi D, Raj CA. Research Article Antioxidant, Cytotoxic and Apoptotic Activities of Crude Extract. Int J Pharm Sci Rev Res. 2016;36(6):28-34.
- Di Domenico F, Foppoli C, Coccia R, Perluigi M. Antioxidants in cervical cancer: Chemopreventive and chemotherapeutic effects of polyphenols. Biochim Biophys Acta Mol Basis Dis. 2012;1822:(5)737-747.
- Liu RM, Li YB, Liang XF, Liu HZ, Xiao JH, Zhong JJ. Structurally related ganoderic acids induce apoptosis in human cervical cancer HeLa cells: Involvement of oxidative stress and antioxidant protective system. Elsevier Ltd. 2015;240.
- Gupta RK, Singh N. Morinda citrifolia (Noni) alters oxidative stress marker and antioxidant activity in cervical cancer cell lines. Asian Pacific J Cancer Prev. 2013;14(8):4603-4606.
- Sukohar A. Effects of caffeine against expression on Mir-423-3p in Cell Lines Hep-G2. Biomed Pharmacol J vol. 2018;11(1):429-435.
- Okunade KS. Human papillomavirus and cervical cancer. J Obstet Gynaecol. 2019;1-7.
- Tulay P, Serakinci N. The role of human papillomaviruses in cancer progression. J Cancer Metastasis Treat. 2016;2:201-213.
- Burd EM. Human Papillomavirus and Cervical Cancer. Clin Microbiol Rev. 2003;16(1):1-17.
- Wangsawat N. Antioxidant activity and cytotoxicity against cancer cell lines of the extracts from novel *Xylaria* species associated with termite nests and LC-MS analysis. Antioxidants. 2021;10(10).
- Narayanan NK. Antitumor activity of melinjo (*Gnetum gnemon* L.) seed extract in human and murine tumor models *in vitro* and in a colon-26 tumor-bearing mouse model *in vivo*. Cancer Med. 2015;4(11):1767-1780.
- Ghaffari T. Natural Phytochemicals Derived from Gymnosperms in the Prevention and Treatment of Cancers. Int J Mol Sci. 2021;22(12):6636.
- Tani H. Pharmacokinetics and safety of resveratrol derivatives in humans after oral administration of melinjo (*Gnetum gnemon* L.) seed extract powder. J Agric Food Chem. 2014;62(8):1999-2007.
- Kunimasa K. Resveratrol derivative-rich melinjo (*Gnetum gnemon* L.) seed extract suppresses multiple angiogenesis-related endothelial cell functions and tumor angiogenesis. Mol Nutr Food Res. 2011;55(11):1730-1734.
- Parhusip AJN, Angel V, Anugrahati NA, Honga J. Ethyl Acetate Extract of Red Melinjo (*Gnetum Gnemon* L.) Peel as Antibacterial Compound. Trends Microbiol. 2020;194:189-194.
- Thi An D, Trung LQ, Yamada K, Nakao S, Takami A. Stilbene derivatives from melinjo extract have antioxidant and immune modulatory effects in healthy individuals. Integr Mol Med. 2015;2(6):405-413.
- Sukohar A, Setiawan S, Wirakusumah F, Sastramihardja H. Isolasi dan karakterisasi senyawa sitotoksik dan asam klorogenat dari biji kopi robusta lampung. J Med Planta. 2011;1(4):11-25.
- Sammar M, Farich BABU, Rayan I, Falah M, Rayan A. Correlation between cytotoxicity in cancer cells and free radical - scavenging activity: *In vitro* evaluation of 57 medicinal and edible plant extracts. Oncol Lett. 2019;18:6563-6571.
- Kato E, Tokunaga Y, Sakan F. Stilbenoids isolated from the seeds of melinjo (*Gnetum gnemon* L.) and their biological activity. J Agric Food Chem. 2009;57(6):2544-2549.
- Sasidharan S. Extraction, Isolation and Characterization of Bioactive Compounds From Plants Extract. Lect Notes Math. 2011;8(1):1-10.
- Nerli RB, Jalalpure SS, Hiremath MB, Neelagund S, Aladakatti R. *In vitro* antioxidant and anticancer activity of *Leea indica* leaf extracts on human prostate cancer cell lines. Integr Med Res. 2017;6(1):79-87.
- Elufioye TO, Abdul AA, Moody JO. Cytotoxicity Studies of the Extracts, Fractions and Isolated Compound of *Pseudocedrela kotschy* on Cervical Cancer (HeLa), Breast Cancer (MCF - 7) and Skeletal Muscle Cancer (RD) Cells. Pharmacognosy Res. 2017;9:46-50.
- Annisaa E, Sasikirana W, Ekawati N, Rahmania I, Dini E. Correlation Between Antioxidant and Cytotoxic Activity of Parijoto (*Medinilla speciosa* Blume) Fractions in 4T1 Cell Line. Indones J Cancer Chemoprevention. 2021;12(1):21-27.

23. He S, Yan X. From Resveratrol to Its Derivatives: New Sources of Natural Antioxidant. *Curr Med Chem*. 2013;20:1005-1017.
24. Eddebbagh M, Messaoudi M, Abourriche A, Berrada M, Attaleb M, Bennamara A. Correlation of the Cytotoxic and Antioxidant Activities of Moroccan Pomegranate (*Punica granatum*) with Phenolic and Flavonoid Contents. *J Pharm Pharmacol*. 2016;4:511-519.
25. Sukohar A, Muhartono. Comparative effects of chlorogenic acid and doxorubicin against expression of caspase-3 in cell lines Hep-G2. *J Chem Pharm Res*. 2015;7(1):187-192.
26. Prayong P, Barusru S, Weerapreeyakul N. Cytotoxic activity screening of some indigenous Thai plants. *Fitoterapia*. 2008;79(7-8):598-601.
27. Parhusip JN, Sitanggang BA. Antimicrobial Activity of Melinjo Seed and Peel Extract (*Gnetum gnemon*) Against Selected Pathogenic Bacteria. *Microbiol Indones*. 2011;5(2):103-112.
28. Santoso M, Naka Y, Angkawidjaja C, Yamaguchi T, Matoba T, Takamura H. Antioxidant and DNA damage prevention activities of the edible parts of *Gnetum gnemon* and their changes upon heat treatment. *Food Sci Technol Res*. 2010;16(6):549-556.
29. Rosa E, Dela, Yuswandi, Michelle, Siregar, Tagor M. Antioxidant activity of melinjo seeds and skin extracts (*Gnetum gnemon* L.). *FaST- J. Sains dan Teknol*. 2020;4(1):92-98.
30. Hafidz KA, Puspitasari N, Azminah, Yanuar A, Artha Y, Mun'Im A. HMG-CoA reductase inhibitory activity of *Gnetum gnemon* seed extract and identification of potential inhibitors for lowering cholesterol level. *Journal of Young Pharmacists*. 2017;9(4):559-565.
31. Hutomo S. Perubahan morfologi sel HeLa setelah paparan ekstrak etanolik *Curcuma longa*. *Majalah Kedokteran Gigi Indonesia*. 2016;2(1):1-5.
32. Haron NH. *In Vitro* Cytotoxic Activity of *Clinacanthus nutans* Leaf Extracts Against HeLa Cells. *Asian Pacific J Cancer Prev*. 2019;20(2014):601-609.
33. Ukui MN, Himizu TS, Kada YO. Normotonic Cell Shrinkage Induced by Na⁺ Deprivation Results in Apoptotic Cell Death in Human Epithelial HeLa Cells. *J Physiol Sci*. 2006;56(5):335-339.
34. Fei Q. Molecular Mechanism of Resveratrol's Lipid Membrane Protection. *Sci Rep*. 2018;8(1):1-12.
35. Li L. Resveratrol suppresses human cervical carcinoma cell proliferation and elevates apoptosis via the mitochondrial and p53 signaling pathways. *Oncol Lett*. 2018;15(6):9845-9851.
36. Fan Y. Resveratrol induces autophagy-dependent apoptosis in HL-60 cells. *BMC Cancer*. 2018;18(1):1-10.
37. García-Zepeda SP, García-Villa E, Díaz-Chávez J, Hernández-Pando R, Gariglio P. Resveratrol induces cell death in cervical cancer cells through apoptosis and autophagy. *Eur J Cancer Prev*. 2013;22(6):577-584.
38. Chin YT. Anti-proliferative and gene expression actions of resveratrol in breast cancer cells *in vitro*. *Oncotarget*. 2014;5(24):12891-12907.
39. Tani H. Pharmacokinetics and safety of resveratrol derivatives in humans after oral administration of melinjo (*Gnetum gnemon* L.) seed extract powder. *J Agric Food Chem*. 2014;62(8):1999-2007.
40. Espinoza JL, Inaoka PT. Gnetin-C and other resveratrol oligomers with cancer chemopreventive potential. *Ann N Y Acad Sci*. 2017;1403(1):5-14.
41. Youness RA, Kamel R, Elkasabgy NA, Shao P, Farag MA. Recent advances in tannic acid (gallotannin) anticancer activities and drug delivery systems for efficacy improvement; a comprehensive review. *Molecules*. 2021;25(6):20.
42. Darwin P. Tannic acid inhibits the Jak2/STAT3 pathway and induces G1/S arrest and mitochondrial apoptosis in YD-38 gingival cancer cells. *Int J Oncol*. 2015;47(3):1111-1120.

GRAPHICAL ABSTRACT



ABOUT AUTHORS



Asep Sukohar is a Professor, Lecture, Faculty of Medicine, University of Lampung.



Sutyarso is a Professor, Lecture, Magister Biology, Faculty of Mathematics and Natural Sciences, University of Lampung.



Hendri Busman is a Lecture, Magister Biology, Faculty of Mathematics and Natural Sciences, University of Lampung.



Nuning Nurcahyani is a Lecture, Magister Biology, Faculty of Mathematics and Natural Sciences, University of Lampung.



Evy Kurniawaty, Lecture, Faculty of Medicine, University of Lampung.



Suharyani, Master, Magister Biology, Faculty of Mathematics and Natural Sciences, University of Lampung.

Cite this article: Sukohar A, Suharyani, Sutyarso, Busman H, Nurcahyani N, Adjeng ANT, et al. Antioxidant and Cytotoxic Activities of Melinjo (*Gnetum gnemon* L.) Seed Fractions on HeLa Cell Line an *In Vitro*. *Pharmacogn J.* 2022;14(3): 559-564.

● 12% Overall Similarity

Top sources found in the following databases:

- 9% Internet database
- Crossref database
- 6% Submitted Works database
- 7% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	mdpi.com Internet	1%
2	jocpr.com Internet	1%
3	Venkata Naga Anantha Sandhya Rani Nandyala, Kothapalli Bonnoth Ch... Crossref	<1%
4	hindawi.com Internet	<1%
5	F. O. Jimoh, A. A. Adedapo, A. A. Aliero, A. J. Afolayan. " Polyphenolic ... Crossref	<1%
6	Fouzia HMIMID. "Phytochemical screening chemical composition and t... Crossref	<1%
7	mts.intechopen.com Internet	<1%
8	Curtin University of Technology on 2018-03-13 Submitted works	<1%

- 9 **Universiti Sains Malaysia on 2018-10-19** <1%
Submitted works

- 10 **Tati Suhartati, Yandri, Jhons F. Suwandi, Sutopo Hadi. "Two Flavan De...** <1%
Crossref

- 11 **W A Tanod, U Yanuhar, Maftuch, D Wahyudi, Y Risjani. "DPPH scavengi...** <1%
Crossref

- 12 **kb.psu.ac.th:8080** <1%
Internet

- 13 **Sridevi I. Puranik, Murigendra B. Hiremat, Rajendra B. Nerli, Shridhar C...** <1%
Crossref

- 14 **cmr.asm.org** <1%
Internet

- 15 **cyberleninka.org** <1%
Internet

- 16 **daneshyari.com** <1%
Internet

- 17 **Prashanth K.B. Nagesh, Pallabita Chowdhury, Elham Hatami, Sonam K...** <1%
Crossref

- 18 **University of Bradford on 2010-11-11** <1%
Submitted works

- 19 **sites.kowsarpub.com** <1%
Internet

- 20 **fasebj.org** <1%
Internet

- 21 **onlinesciencepublishing.com** <1%
Internet
-
- 22 **japsonline.com** <1%
Internet
-
- 23 **Hiroko Tani, Susumu Hikami, Sanae Iizuna, Maiko Yoshimatsu et al. " P...** <1%
Crossref
-
- 24 **Mohammed Abdul Mujeeb, Ankalabasappa Vedamurthy, Arun Kashivis...** <1%
Crossref

● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)

EXCLUDED SOURCES

phcogj.com	93%
Internet	
mail.phcogj.com	14%
Internet	
spandidos-publications.com	5%
Internet	
repository.lppm.unila.ac.id	4%
Internet	
pasca.unila.ac.id	4%
Internet	
S Andriani, E L Widiastuti, N Nurcahyani, E rosa, H Busman. "Cytotoxic activity ...	3%
Crossref	
S Andriani, E L Widiastuti, N Nurcahyani, E rosa, H Busman. " Cytotoxic activity...	3%
Crossref	
Sultan Agung Islamic University on 2020-11-26	3%
Submitted works	
researchonline.ljmu.ac.uk	2%
Internet	

Shridhar C. Ghagane, Sridevi I. Puranik, Vijay M. Kumbar, Rajendra B. Nerli et ...	2%
Crossref	
pdffox.com	2%
Internet	
bmccancer.biomedcentral.com	2%
Internet	
Yingying Fan, Jen-Fu Chiu, Jing Liu, Yan Deng, Cheng Xu, Jun Zhang, Guanwu ...	2%
Crossref	
South Dakota Board of Regents on 2019-04-16	2%
Submitted works	
biomedpharmajournal.org	2%
Internet	
Niwana Wangsawat, Lutfun Nahar, Satyajit D. Sarker, Cherdchai Phosri et al. "...	2%
Crossref	
Areeya Suchantabud, Teeraporn Katisart, Chusri Talubmook. "Chronic Toxicit...	2%
Crossref	
tessera.spandidos-publications.com	2%
Internet	
link.springer.com	2%
Internet	
Angeles University Foundation on 2021-11-17	2%
Submitted works	
Universitas Jenderal Achmad Yani on 2022-03-11	2%
Submitted works	

Academic Library Consortium on 2020-08-10	2%
Submitted works	
Sriwijaya University on 2021-02-05	2%
Submitted works	
Padjadjaran University on 2022-07-10	2%
Submitted works	
Universitas Indonesia on 2021-01-05	2%
Submitted works	
Academic Library Consortium on 2020-08-10	2%
Submitted works	
Universitas Airlangga on 2020-06-07	2%
Submitted works	
School of Business and Management ITB on 2022-06-28	2%
Submitted works	
Padjadjaran University on 2022-07-10	2%
Submitted works	
Academic Library Consortium on 2021-08-11	2%
Submitted works	
Universitas Airlangga on 2020-04-11	2%
Submitted works	
Universitas Indonesia on 2021-12-07	2%
Submitted works	
Emmanuel College on 2020-10-15	2%
Submitted works	

Academic Library Consortium on 2020-01-30	2%
Submitted works	
repository.ubaya.ac.id	2%
Internet	
Universitas Indonesia on 2021-04-21	2%
Submitted works	
Universitas Indonesia on 2021-04-21	2%
Submitted works	
Universitas Indonesia on 2021-08-18	2%
Submitted works	
School of Business and Management ITB on 2022-06-24	2%
Submitted works	
School of Business and Management ITB on 2021-12-20	2%
Submitted works	
docksci.com	2%
Internet	
Universitas Indonesia on 2020-03-20	2%
Submitted works	
res.mdpi.com	2%
Internet	
nature.com	2%
Internet	
Academic Library Consortium on 2020-12-01	2%
Submitted works	

ijcc.chemoprev.org	2%
Internet	
Universitas Sebelas Maret on 2021-03-29	2%
Submitted works	
Surabaya University on 2020-11-28	2%
Submitted works	
School of Business and Management ITB on 2020-11-10	2%
Submitted works	
Academic Library Consortium on 2020-02-12	2%
Submitted works	
Surabaya University on 2020-01-31	2%
Submitted works	
Universitas Indonesia on 2021-12-07	2%
Submitted works	
Universitas Indonesia on 2021-12-07	2%
Submitted works	
Universitas Indonesia on 2019-01-23	2%
Submitted works	
Universitas Indonesia on 2018-10-29	2%
Submitted works	
Oluwakemi Charles-Okhe, Michael A. Odeniyi, Titilayo O. Fakeye, Omonike O. ...	2%
Crossref	
Oluwakemi Charles-Okhe, Michael A. Odeniyi, Titilayo O. Fakeye, Omonike O. ...	2%
Crossref	

School of Business and Management ITB on 2022-05-24	2%
Submitted works	
repository.unair.ac.id	2%
Internet	
Narayanan, Narayanan K., Kazuhiro Kunimasa, Yukio Yamori, Mari Mori, Hidek...	1%
Crossref	
Narayanan K. Narayanan, Kazuhiro Kunimasa, Yukio Yamori, Mari Mori et al. " ...	1%
Crossref	
Hendri Busman, Sutyarso Sutyarso, Salman Farisi, Fukrapti Fukrapti, Aulia Rik...	1%
Crossref	
Venkata Siva Satyanarayana Kantamreddi, V. Thirumala Veni, G. Y. S. K. Swa...	1%
Crossref	
Vidya Shivram Patil, Nutan Padmnabh Malpathak. "Micro-morphoanatomical ...	1%
Crossref	
Tom Mathew Kalappurayil, Benny Pulinilkumthadathil Joseph. "A Review of ...	1%
Crossref	
Mitchell Henry Wright, Joseph Sirdarta, Alan White, Anthony Carlson Greene,...	1%
Crossref	
Fatma Sri Wahyuni, Daud Ahmad Israf Ali, Nordin Hj. Lajis, Dachriyanus D. "An...	1%
Crossref	
frontiersin.org	1%
Internet	
Ru-Ming Liu, Ying-Bo Li, Xiang-Feng Liang, Hui-Zhou Liu, Jian-Hui Xiao, Jian-J...	1%
Crossref	

Liu, Ru-Ming, Ying-Bo Li, Xiang-Feng Liang, Hui-Zhou Liu, Jian-Hui Xiao, and Ji...	1%
Crossref	
Nusaibah Zahratunnisa, Berna Elya, Arikadia Noviani. "Inhibition of Alpha-Gluc...	1%
Crossref	
researchsquare.com	1%
Internet	
WWW.MDPI.COM	1%
Internet	
Adeolu A Adedapo. "Assessment of the medicinal potentials of the methanol ...	1%
Crossref	
Adeolu A Adedapo. "Antioxidant activities and phenolic contents of the metha...	1%
Crossref	
Adeolu A Adedapo. "Antibacterial and antioxidant properties of the methanol ...	1%
Crossref	
atlantis-press.com	1%
Internet	
phcogres.com	1%
Internet	
Amanda Listiyani, Berna Elya, Nuraini Puspitasari. "Antioxidant Activity and Li...	<1%
Crossref	
readkong.com	<1%
Internet	
spandidos-publications.com	<1%
Internet	

Universitas Negeri Jakarta on 2019-04-22	<1%
Submitted works	
Universitas Negeri Jakarta on 2019-04-09	<1%
Submitted works	
liawbf.pixnet.net	<1%
Internet	
Sayoni Nag, Krishna Das Saha. "Chitosan-Decorated PLGA-NPs Loaded with T...	<1%
Crossref	
Mohammed Abdul Mujeeb, Ankalabasappa Vedamurthy, Arun Kashivishwanat...	<1%
Crossref	
phytopharmajournal.com	<1%
Internet	
jcreview.com	<1%
Internet	
A.J. Afolayan, F.O. Jimoh, M.O. Sofidiya, S. Koduru, F.B. Lewu. " Medicinal Pot...	<1%
Crossref	
thieme-connect.com	<1%
Internet	
ijppr.com	<1%
Internet	
International Islamic University Malaysia on 2012-05-29	<1%
Submitted works	
mafiadoc.com	<1%
Internet	

ijpsr.com	<1%
Internet	
<hr/>	
Sonja Veljović, Mile Veljović, Ninoslav Nikićević, Saša Despotović, Siniša Rad...	<1%
Crossref	
<hr/>	
Ubon Ratchathani University on 2022-05-12	<1%
Submitted works	
<hr/>	
academicjournals.org	<1%
Internet	
<hr/>	
University of Bradford on 2010-10-07	<1%
Submitted works	
<hr/>	
Tani, Hiroko, Susumu Hikami, Sanae Iiduna, Maiko Yoshimatsu, Takashi Asa...	<1%
Crossref	