

PAPER NAME

**35.pdf**

AUTHOR

**Mahfut Mahfut**

WORD COUNT

**3587 Words**

CHARACTER COUNT

**19368 Characters**

PAGE COUNT

**17 Pages**

FILE SIZE

**5.7MB**

SUBMISSION DATE

**Sep 10, 2022 3:03 PM GMT+7**

REPORT DATE

**Sep 10, 2022 3:05 PM GMT+7**

### ● 16% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 16% Internet database
- 5% Publications database
- Crossref database
- Crossref Posted Content database
- 3% Submitted Works database

### ● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources
- Manually excluded text blocks

# Current Research Trends in Biological Science

Vol. 1



Book Publisher  
International

# **1** Current Research Trends in Biological Science

**Vol. 1**



# **Current Research Trends in Biological Science**

**Vol. 1**

3 India ■ United Kingdom

  
**Book Publisher  
International**

## **Editor(s)**

**Dr. Sławomir Borek**

Assistant Professor,  
Department of Plant Physiology, Faculty of Biology, Adam Mickiewicz University, Poznań,  
Poland.

Email: borek@amu.edu.pl;

**FIRST EDITION 2020**

**ISBN 978-93-89816-64-8 (Print)**

**ISBN 978-93-89816-65-5 (eBook)**

**DOI: 10.9734/bpi/crtbs/v1**



# Contents

---

<b>Preface</b>	i
<b>Chapter 1</b> <b>Plant Passive Immunity: Micromorphological and Biochemical Features of the Maloideae (Rosaceae) External Tissues</b> Alexander S. Voronkov, Tamara Kh. Kumachova and Tatiana V. Ivanova	1-16
<b>Chapter 2</b> <b>Unraveling Siderophore Producing Bacteria from Plant Rhizosphere</b> B. Soundaryaa and S. Aruna Sharmili	17-28
<b>Chapter 3</b> <b>Population Dynamics of Five Important Commercial Fish Species in the Sundarbans Ecosystem of Bangladesh: Recent Advancement</b> Md. Golam Mustafa, Imran Ahmed and Mohammad Ilyas	29-44
<b>Chapter 4</b> <b>Insect Vectors Associated to Dumpsites in Gombe Metropolis, Nigeria, Western Africa</b> E. Abba, I. Lamogo, A. Rejoice, A. Jemimah and K. P. Yoriyo	45-54
<b>Chapter 5</b> <b>Micropropagation of Narcissus tazetta 'Chinensis' and Its Relation to Secondary Metabolites Content: Brief Overview</b> Hamdy M. Abdel-Rahman, A. M. F. Al-Ansary, K. N. Rashed and A. A. Rizkalla	55-66
<b>Chapter 6</b> <b>The Role of Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) in Recognition of Pseudomonas aeruginosa in Lung Epithelial Cells: Detailed Study</b> Ekong, Mercy Okon, Tarh, Jacqueline Ebob and Iroegbu, Christian Ukwuoma	67-75
<b>Chapter 7</b> <b>Effect of Potassium Alum on the Mycoflora of Postharvest Spoilage of Solanum lycopersicum L. (Tomato)</b> Lawrence O. Amadi, Eunice J. Udoh, Rebecca U. Thompson, Ruth G. Benjamin and Felicia W. Nmomo	76-86
<b>Chapter 8</b> <b>Morphological Character Variations in Lasiodiplodia Species: Pathogen of Inflorescence Dieback in Cashew Growing Ecologies of Nigeria</b> Dele Omoyele Adeniyi	87-94
<b>Chapter 9</b> <b>Ameliorating Effect of Moringa against Liver and Kidney Injury Induced by Monosodium Glutamate: Brief Overview</b> Rehab M. El-Gharabawy, Amira S. Ahmed and Thara I. Al-Adhahd	95-105
<b>Chapter 10</b> <b>A Formula of Bacterial Pathogenicity (Theoretical Review): Recent Development</b> Yurii V. Ezepechuk	106-112
<b>Chapter 11</b> <b>Cultivation of Mint (Mentha x gracilis) in Agroforestry System</b> Dalva Paulus	113-123

---

---

<b>Chapter 12</b> <b>The Details of Revamp Studies on Morpho-Histology of the Male Reproductive System of <i>Halys dentatus</i> Fabricious (Hemiptera: Pentatomidae)</b> Jyoti H. Gangurde	124-130
<b>Chapter 13</b> <b>Protease Composition in Tissue Extracts of Hydrobionts from Antarctic Region: Recent Study</b> Nataliia Raksha, Tetiana Halenova, Tetyana Vovk, Oleksii Savchuk and Lydmila Ostapchenko	131-138
<b>Chapter 14</b> <b>Effectiveness Test of Orchid Mycorrhizal Isolate (<i>Ceratorhiza</i> and <i>Trichoderma</i>) Indonesia and Its Role as a Biofertilizer: Critical Overview</b> Mahfut	139-145

---



## Preface

1 This book covers all areas of biological research. The contributions by the authors include Maloideae; pathogenic fungi; polyphenols; fruits; rhizosphere; carboxylate; PGPR; *Escherichia fergusonii*; *Musca domestica*; *Anopheles* spp. *Aedes* spp; *Periplaneta americana*; alkaloids; callus induction; daffodils; *Narcissus tazetta*; cystic fibrosis; cytokines; *Pseudomonas aeruginosa*; postharvest; mycotoxin; tomato; inhibition; cashew; pigmentation; sporulation; pycnidia; monosodium glutamate; liver; kidney; *Moringa oleifera*; bacterial pathogenicity; adhesions; invasions; toxins; bio-molecules; infectious diseases; prokaryotic and eukaryotic organisms; medicinal plants; essential oil; linalool; intercropping; male reproductive system; marine hydrobionts; protease inhibitors; collagenolytic and trypsin-like activities; Orchid mycorrhizal; ceratorhiza; trichoderma; *Phalaenopsis amabilis* etc. This book contains various materials suitable for students, researchers and academicians in the field of biological research.



---

# Effectiveness Test of Orchid Mycorrhizal Isolate (*Ceratorhiza* and *Trichoderma*) Indonesia

Mahfut<sup>1\*</sup>

DOI: 10.9734/bpi/crtbs/v1

---

## ABSTRACT

**Aim:** The existence of Orchid Mycorrhizal Fungi (OMF) has a role to stimulate growth and support the supply of orchid nutrition as a biofertilizer agent. This study aimed to determine the association of mycorrhizal with *Phalaenopsis amabilis* (L.) Blume which was carried out through the effectiveness test of two Indonesian orchid mycorrhizal isolates i.e. *Ceratorhiza* and *Trichoderma*.

**Study Design:** This study consisted of 4 treatments. Each treatment was repeated 3 times, each repetition of 5 plantlets, so that the total plantlet used was 60.

**Place and Duration of Study:** Laboratory of Plant Biotechnology, Department of Biology, Universitas Gadjah Mada, Indonesia, between June 2017 and April 2018.

**Methodology:** The method of inoculating orchid mycorrhizal by placing a plantlet in a petri dish containing orchid mycorrhizal for 1, 2, 3 and 4 days. Then plantlets are grown on sterile moss growing media and acclimatized in a greenhouse. Observation of each treatment is carried out every day for the next month. Observation variables include the number of initial and final roots, the number of live and dead roots, and the number of living and dead plants.

**Results:** The results of the orchid mycorrhizal induction test showed that the *Ceratorhiza* inoculation treatment showed a fluctuation in the mean increase in the number of final roots, live roots, dead roots, and dead plantlets that were higher than the *Trichoderma* inoculation treatment. The results also showed that the best inoculation time on *Ceratorhiza* and *Trichoderma* was day 3 and 4. The adaptation process had the effect of increasing the number of dead roots in weeks 1 and 2. The adaptation process stopped at the beginning of week 4 with the number of new roots appearing a lot.

**Conclusion:** Orchid mycorrhizal *Ceratorhiza* shows the value of effectiveness test compared with *Trichoderma*. The results of this study are expected to be basic information in efforts to cultivate natural orchids in Indonesia.

**Keywords:** Orchid mycorrhizal; ceratorhiza; trichoderma; *Phalaenopsis amabilis* (L.) blume; Indonesia.

## 1. INTRODUCTION

*Phalaenopsis* is a genus of orchids, which some of its members have important role as parent crosses. Approximately 30 of the total 62 species are spread throughout Indonesia. The presence of this genus in its native habitat (nature) has been reported to have greatly diminished, even some of the members have been recorded to the IUCN red list version 2013.2 due to excessive exploration and forest degradation. Thus, it is very necessary to conserve the existence of native *Phalaenopsis* orchids in Indonesia through the efforts of preservation and protection of plants.

The presence of endophytic mycorrhizal as Orchid Mycorrhizal Fungi (OMF) in orchid plants is known to play an important role in stimulating orchid seed germination [1], supporting the supply of plantlet nutrition [2,3], helping the formation of more buds and flower buds [2] and control biological agents by

---

<sup>1</sup>Faculty of Mathematics and Natural Sciences, University of Lampung, Lampung, 35145, Indonesia.

\*Corresponding author: E-mail: mahfutkariem@yahoo.com, mahfut.mipa@fmipa.unila.ac.id;

inducing resistance to bacterial infection of *Erwinia chrysanthemi* [4] and inhibiting the replication of ORSV and CymMV [5]. But its presence in different plants can be as a disease-causing agent (pathogen).

Previous studies have identified the isolates of *Ceratorhiza* and *Trichoderma* isolated from the roots of the orchid *Phalaenopsis amabilis* L. (Blume) in Yogyakarta, Indonesia. Identification of the two is distinguished by observing morphological and molecular characters [6]. Molecular analysis was carried out based on the identification of the rDNA-ITS sequence, while morphological analysis was done by observing the surface colour, appearance and colony growth rate, hyphae colour and diameter, shape and size of monilioid cells, and the number of nuclei [7].

*Ceratorhiza* isolate in this study were reported Mahfud et al. [6] as one group of Ceratobasidium were isolated from the roots of orchid plants in Yogyakarta. The results of molecular analysis showed 600-750 bp of DNA products located on the ITS1-5.8S-ITS4 region. The sequenced products showed insertion and substitution occurrences, which may result in strain diversity and possible variation. Reconstruction of phylogenetic trees using *Maximum Parsimony* and *Bootstrap-1000* approach showed showed the Indonesian isolate is at the basal clade and already far apart from the other isolates.

Soelistijono et al. [8] reported that induction of *Ceratorhiza* and *Tricoherma mycorrhizal* are as a biofertilizer (organic fertilizer). Mycorrhiza works to improve the structure of the soil around plant roots by breaking down organic substances in the soil. The presence of organic substances in the soil is abundant but the shape and size that cannot be absorbed by plants. Besides saving costs, the use of mycorrhizal is very safe for the environment. In the sequel, the mycorrhizal application can accelerate the growth and development of orchid plantlets. Based on this, further research needs to be carried out on the effectiveness of both Indonesian isolate endophytic mycorrhizal and their role as biofertilizer. This research is expected to be important information in the cultivation and development potential of *Phalaenopsis amabilis* L. (Blume) in Indonesia.

## **2. MATERIALS AND METHODS**

### **2.1 Source of Orchid Mycorrhizal Isolates**

The isolates used were *Ceratorhiza* and *Trichoderma* isolate collected from the root of *P. amabilis* L. (Blume) in Yogyakarta. Pure isolates were bred on Potato Dextrose Agar (PDA) media. Inoculum incubation is carried out in a dark room at room temperature until the age of 7 days.

### **2.2 Source of Orchid Plantlet**

The orchid plantlet used was 12 months old *P. amabilis* L. (Blume) cultured from seeds on Murashige and Skoog (MS) media. Plantlets are removed from culture bottles and soaked in a fungicide solution (2 g/l water) for 20 minutes. Plantlet is then planted on sterile moss media. Orchids are grown properly for 1 month before treatment. Watering is done twice a week using a spray tool.

### **2.3 Orchid Mycorrhizal Inoculation**

Orchid mycorrhizal inoculation in this study used the method of Nuangmek et al. [9]. Orchid mycorrhizal are grown in Petri dishes 9 cm in diameter. Plantlet is placed in a petri dish containing orchid mycorrhizal for 1, 2, 3 and 4 days. Then the plantlets are regrown on sterile moss growing media and acclimatized in a greenhouse. This study consisted of 4 treatments. Each treatment was repeated 3 times, each repetition of 5 plantlets, so that the total plantlet used was 60.

### **2.4 Observation of the Orchid Mycorrhizal Effectiveness Test**

The effectiveness of orchid mycorrhizal on plantlets was carried out *in vivo*. Observation of each treatment is carried out every day for the next month. Observation variables include the number of initial and final roots, the number of live and dead roots and the number of living and dead plants.


### 3. RESULTS AND DISCUSSION

#### 3.1 Inoculation of Orchid Mycorrhizal

The results of orchid mycorrhizal inoculation showed hyphae would envelop the roots of the plantlet. The hyphae were getting thicker with the length of treatment (Fig. 1).

#### 3.2 Observation of Orchid Mycorrhizal Effectiveness Test

The effectiveness of orchid mycorrhizal was carried out through observations of growth and survival ability of post-inoculation plantlet shown in Tables 1-4.



**Fig. 1. Orchid mycorrhizal inoculation by placing plantlets on (A) *Ceratorhiza* and (B) *Trichoderma* isolates in a petri dish at 4 days treatment**

**Table 1. Mean effectiveness of orchid mycorrhizal test results at 1<sup>st</sup> week**

Orchid mycorrhizal isolates	Duration of incubation (Day)	Observation of effectiveness Test (1 <sup>st</sup> week)					
		∑ Root (Initial)	∑ Root (End)	∑ Root (Life)	∑ Root (Dead)	∑ Plantlet (Life)	∑ Plantlet (Dead)
Control	-	7	7	7	0	3	0
<i>Ceratorhiza</i>	1	11	7	7	4	3	0
	2	10	10	10	0	3	0
	3	10	7	7	3	3	0
	4	8	9	3	6	2	1
<i>Trichoderma</i>	1	8	8	8	0	3	0
	2	5	5	5	0	3	0
	3	11	10	10	1	3	0
	4	7	6	7	0	3	0

The test results showed that the *Ceratorhiza* inoculation treatment showed a fluctuation in the mean increase in the number of final roots, live roots, dead roots, and dead plantlets that were higher than the *Trichoderma* inoculation treatment. The highest observation of the highest final root number in *Ceratorhiza* inoculation was at incubation time of 1 day which was 3 at week 2 (Table 2) and 2 at week 4 (Table 4), whereas at *Trichoderma* inoculation was at incubation time at 3 day namely 4 at week 3 (Table 3) and -2 at week 4 (Table 4).

The highest number of life roots in *Ceratorhiza* inoculation was at 4 days incubation ie 1 at week 3 (Table 3) and -3 at week 4 (Table 4), whereas at *Trichoderma* inoculation was at 4 day incubation time at 4 at week 3 (Table 3) and 4 weeks (Table 4). The highest mean number of dead roots in *Ceratorhiza* inoculation was at incubation time of 1 day which was 3 at week 3 (Table 3) and 6 at week 4 (Table 4), whereas at *Trichoderma* inoculation was at incubation time of 2 days namely 3 at week 3 (Table 3) and 1 at week 4 (Table 4). Finally, the mean observation the highest number of dead plantlets in *Ceratorhiza* inoculation was at an incubation time of 1 day which is 1 at week 4 (Table 4), whereas in *Trichoderma* inoculation no dead plantlets were found.

**Table 2. Mean effectiveness of orchid mycorrhizal test results at 2<sup>nd</sup> week**

Orchid mycorrhizal isolates	Duration of incubation (Day)	Observation of effectiveness test (2 <sup>nd</sup> week)					
		∑ Root (Initial)	∑ Root (End)	∑ Root (Life)	∑ Root (Dead)	∑ Plantlet (Life)	∑ Plantlet (Dead)
Control	-	7	7	7	0	3	0
<i>Ceratorhiza</i>	1	7	7	7	0	3	0
	2	7	5	5	2	3	0
	3	8	9	9	1	3	0
	4	3	3	3	3	3	0
<i>Trichoderma</i>	1	8	7	7	1	3	0
	2	5	5	5	0	3	0
	3	10	10	9	3	3	0
	4	6	5	5	1	3	0

**Table 3. Mean effectiveness of orchid mycorrhizal test results at 3<sup>rd</sup> week**

Orchid mycorrhizal isolates	Duration of incubation (Day)	Observation of effectiveness test (3 <sup>rd</sup> week)					
		∑ Root (Initial)	∑ Root (End)	∑ Root (Life)	∑ Root (Dead)	∑ Plantlet (Life)	∑ Plantlet (Dead)
Control	-	7	7	7	0	3	0
<i>Ceratorhiza</i>	1	7	7	7	0	3	0
	2	8	8	8	0	3	0
	3	10	7	7	3	3	0
	4	3	3	3	0	3	0
<i>Trichoderma</i>	1	7	7	7	0	3	0
	2	5	5	5	0	3	0
	3	11	11	11	0	3	0
	4	5	5	5	0	3	0

**Table 4. Mean effectiveness of orchid mycorrhizal test results at 4<sup>th</sup> week**

Orchid mycorrhizal isolates	Duration of incubation (Day)	Observation of effectiveness test (4 <sup>th</sup> week)					
		∑ Root (Initial)	∑ Root (End)	∑ Root (Life)	∑ Root (Dead)	∑ Plantlet (Life)	∑ Plantlet (Dead)
Control	-	7	7	7	0	3	0
<i>Ceratorhiza</i>	1	7	8	8	0	3	0
	2	8	10	7	3	3	0
	3	7	9	9	1	3	0
	4	3	5	5	0	3	0
<i>Trichoderma</i>	1	7	5	5	2	3	0
	2	5	7	5	2	3	0
	3	9	9	9	3	3	0
	4	5	6	6	0	3	0

The results of this study showed that the inoculation of *Ceratorhiza* gave more effect on the number of dead roots in week 1 and 2 compared to *Trichoderma* inoculation. Although at weeks 3 and 4 a large number of new roots appear (Fig. 2). The results also showed that the best inoculation time for *Ceratorhiza* and *Trichoderma* was day 3 and 4. This adaptation process stopped at the beginning of week 4. As of week 4, the mean number of dead roots decreased and the number of root increases increased. Based on visual observations, the root undergoes decay as a process of adaptation to the orchid mycorrhizal inoculation treatment. This is due to the faster growth of *Ceratorhiza* hypha so that it can associate with plants faster. The faster the orchid mycorrhizal is associated with the host plant, the higher the capacity to absorb nutrients and cause increased growth. Muslim et al. [10] explained that the main principle of orchid mycorrhizal work is to infect the root system of the host plant and produce hyphae intensively so that it can increase the capacity of plants to absorb nutrients.



Fig. 2. Development of plantlet results from mycorrhizal inoculation per week; (1) week 1; (2) week 2; (3) week 3; (4) week 4. The arrows indicate the emergence of new roots

Orchid plants require orchid mycorrhizal infections to complete their life cycle. An important role of orchid mycorrhizal in plant growth is its ability to absorb nutrients both macro and micro. The treatment of orchid mycorrhizal inoculation on orchids is known to be able to increase the efficiency of inhibition of N nutrient absorption to increase plant growth, such as increasing length, width and the number of leaves and roots. Element N is a building material for amino acids/ proteins, enzymes, nucleic acids, nucleoproteins and alkaloids. N deficiency will limit cell division and distribution [11].

#### 4. CONCLUSION

The results of the effectiveness of orchid mycorrhizal isolates in Indonesia and its role as biofertilizer showed that the inoculation treatment of *Ceratorhiza* showed fluctuations in the average increase in the number of final roots, life roots, dead roots, and dead plantlets which were higher than those in the *Trichoderma* inoculation treatment. The results also showed that the best inoculation time on *Ceratorhiza* and *Trichoderma* was day 3 and 4. The adaptation process had the effect of increasing the number of dead roots in week 1 and 2. The adaptation process stopped at the beginning of week 4 with the number of new roots appearing a lot.

#### 3 COMPETING INTERESTS

Author has declared that no competing interests exist.

#### REFERENCES

1. Shao SC, Burgess KS, Sanders JMC, Liu Q, Fan XL, Huang H, Gao JY. Using *in situ* symbiotic seed germination to restore over-collected medicinal orchids in Southwest China. *Front Plant Sci.* 2017;8:888.
2. Silva JAT, Kerbauy GB, Zeng S, Chen Z, Duan J. *In vitro* flowering of orchids. *Reviews in Biotechnology.* 2013;34(1).
3. Wu PH, Huang DD, Chang DCN. Mycorrhizal symbiosis enhances *Phalaenopsis* orchid's growth and resistance to *Erwinia chrysanthemi*. *African Journal of Biotechnology.* 2011; 10(50):10095-10100.
4. Wu FH, Chan MT, Liao DC, Hsu CT, Lee YW, Daniell H, Duvall MR, Lin CS. Complete chloroplast genome of *Oncidium gower ramsey* and evaluation of molecular markers for identification and breeding in *Oncidiinae*. *BMC Plant Biology.* 2010;10(68):1-12.
5. Tong TG. Research Focus: Beneficial Endophytes in plants; 2019. Available:[http://abrc.sinica.edu.tw/2010E/view2/faculty\\_E1\\_Print.php?id=cgtong](http://abrc.sinica.edu.tw/2010E/view2/faculty_E1_Print.php?id=cgtong) (Accessed October 11<sup>th</sup> 2019)
6. Mahfut, Daryono BS, Indrianto A, Somowiyarjo S. *Malaysian Journal of Microbiology.* 2020; 16(1):68-72.
7. Gopireddy BM, Devi GU, Kumar KV, Babu TR, Naidu TCM. Cultural and morphological characterization of *Rhizoctonia solani* f. sp. *sasakii* isolates collected from different districts of Andhra Pradesh. *Int. J. Curr. Microbiol. App. Sci.* 2017;6(11):3457-3469.
8. Soelistijono Priyatmojo A, Semiarti E, Sumardiyono C. Characterization of pathogenic *Rhizoctonia* sp. and mycorrhizal *Rhizoctonia* isolates on terrestrial orchid plant *Spathoglottis plicata*. *Biota.* 2011;16(2):371-380.
9. Nuangmek W, Mckenzie EHC, Lumyong S. Endophytic fungi from wilt banana (*Musa accuminata colla*) work againts antracnose disease caused by *Colletotricum musae*. *Research Journal of Microbiology.* 2008;3(5):368-374.
10. Muslim A, Horinouchi H, Hyakumachi M. Biocontrol of *Fusarium* wilt of spinach by the plant growth promoting fungus *Fusarium equiseti* Gf18. *Journal of Plant Pathology.* 2010;92(1):146-254.
11. Alghamdi SA. Influence of mycorrhizal fungi on seed germination and growth in terrestrial and epiphytic orchids. *Saudi Journal of Biological Sciences.* 2019;26(3):495-502.



**Biography of author**



**Dr. Mahfut**

Faculty of Mathematics and Natural Sciences, University of Lampung, Lampung, 35145, Indonesia.

He was born on 9 September 1981 in Merbau Mataram, South Lampung. In 2006 he obtained his Bachelor of Biology degree from the Faculty of Mathematics and Natural Sciences, University of Lampung. In 2011 he obtained a Master's degree (Master of Science) and in 2018 obtained his doctorate degree from the Biology Postgraduate Program, Gadjah Mada University, Yogyakarta. The title of the thesis of the author is "The Effect of the Insecticide Made from Profenofros Active on Organ Reproduction and Reversal of Tomatoes (*Lycopersicum esculentum* Mill.)", Entitled Thesis "Detection and Characterization of Odontoglossum ringspot virus (ORSV) Isolates of Java and Bali, Indonesia", and titled Dissertation "Variation Genetic rbcL Genes and CP Genes, and Induction of Natural Phalaenopsis Resistance in Indonesia Against Odontoglossum ringspot virus with Orchid Mycorrhiza ". Currently the he is actively working as a lecturer in the Department of Biology, Faculty of Mathematics and Natural Sciences, University of Lampung since 2014.

---

© Copyright (2020): Authors. The licensee is the publisher (Book Publisher International).

**DISCLAIMER**

This chapter is an extended version of the article published by the same author(s) in the following journal.  
Journal of Annual Research & Review in Biology, 33(4): 1-7, 2019.

**Reviewers' Information**

- (1) Ann A. J. Mofunanya, University of Calabar, Nigeria.
- (2) R. Mahalakshmi, India.

## **London Tarakeswar**

### **Registered offices**

India: Guest House Road, Street no - 1/6, Hooghly, West Bengal, PIN-712410, India, Corp. Firm  
Registration Number: L77527, Tele: +91 8617752708, Email: [contact@sciencedomain.org](mailto:contact@sciencedomain.org),  
(Headquarters)

UK: Third Floor, 207 Regent Street, London, W1B 3HH, UK  
Fax: +44 20-3031-1429 Email: [contact@sciencedomain.org](mailto:contact@sciencedomain.org),  
(Branch office)

## ● 16% Overall Similarity

Top sources found in the following databases:

- 16% Internet database
- 5% Publications database
- Crossref database
- Crossref Posted Content database
- 3% Submitted Works database

### TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	<b>bookpi.org</b> Internet	4%
2	<b>mjm.usm.my</b> Internet	2%
3	<b>repository.udsm.ac.tz:8080</b> Internet	2%
4	<b>researchgate.net</b> Internet	1%
5	<b>journalajbge.com</b> Internet	<1%
6	<b>journalrepository.org</b> Internet	<1%
7	<b>Rehab M. El-Gharabawy, Amira S. Ahmed, Thara I. Al-Adhadh. "Amelior...</b> Crossref	<1%
8	<b>journaljalsi.com</b> Internet	<1%

9	<b>sciencedomain.org</b>	Internet	<1%
10	<b>journalmrji.com</b>	Internet	<1%
11	<b>2pressrelease.net</b>	Internet	<1%
12	<b>coursehero.com</b>	Internet	<1%
13	<b>E. Abba, M. Amina, Y. Lamogo, A. Rejoice, A. Jemimah, K. P. Yoriyo. "S...</b>	Crossref	<1%
14	<b>ikpress.org</b>	Internet	<1%
15	<b>timacad.ru</b>	Internet	<1%

## ● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)
- Manually excluded text blocks

---

### EXCLUDED SOURCES

<b>Mahfut ., Budi Setiadi Daryono, Ari Indrianto, Susamto Somowiyarjo. "Effectiv...</b>	<b>44%</b>
Crossref	
<b>bp.bookpi.org</b>	<b>17%</b>
Internet	
<b>repository.lppm.unila.ac.id</b>	<b>16%</b>
Internet	
<b>journalarrb.com</b>	<b>13%</b>
Internet	
<b>pesquisa.bvsalud.org</b>	<b>11%</b>
Internet	
<b>251news.com</b>	<b>8%</b>
Internet	
<b>sersc.org</b>	<b>6%</b>
Internet	
<b>ejournal.unp.ac.id</b>	<b>3%</b>
Internet	
<b>semantic scholar.org</b>	<b>3%</b>
Internet	

**ojs.uajy.ac.id**

Internet

**3%**

EXCLUDED TEXT BLOCKS

**ABSTRACT**Aim: The existence of Orchid Mycorrhizal Fungi (OMF) has a role to sti...

journalarrb.com

---

**1. INTRODUCTION**Phalaenopsis is a genus of orchids, which some of its members...

sersc.org

---

**Effectiveness Test of Orchid Mycorrhizal Isolate (Ceratorhiza and Trichoderma) In...**

sersc.org