

PAPER NAME

Enumerate the Number of Vertices Labeled Connected Graph of Order Seven Containing No Parallel Edges

AUTHOR

Muslim Ansori

WORD COUNT

5019 Words

CHARACTER COUNT

21912 Characters

PAGE COUNT

8 Pages

FILE SIZE

727.3KB

SUBMISSION DATE

Aug 19, 2022 9:52 PM GMT+7

REPORT DATE

Aug 19, 2022 9:53 PM GMT+7

● 7% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 2% Internet database
- 5% Publications database
- Crossref database
- Crossref Posted Content database
- 2% Submitted Works database

● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources
- Manually excluded text blocks

Enumerate the Number of Vertices Labeled Connected Graph of Order Seven Containing No Parallel Edges

Muslim Ansori¹, Wamiliana^{1*}, Fitriani¹, Yudi Antoni¹, Desiana Putri¹

¹Department of Mathematics, Faculty of Mathematics and Natural Science, Universitas Lampung, Bandar Lampung, 35145, Indonesia

*Corresponding author: wamiliana.1963@fmipa.unila.ac.id

Abstract

A graph that is connected $G(V,E)$ is a graph in which there is at least one path connecting every two vertices in G ; otherwise, it is called a disconnected graph. Labels or values can be assigned to the vertices or edges of a graph. A vertex-labeled graph is one in which only the vertices are labeled, and an edges-labeled graph is one in which only edges are assigned values or labels. If both vertices and edges are labeled, the graph is referred to as total labeling. If given n vertices and m edges, numerous graphs can be made, either connected or disconnected. This study will be discussed the number of disconnected vertices labeled graphs of order seven containing no parallel edges and may contain loops. The results show that the number of vertices labeled connected graph of order seven with no parallel edges is $N(G_{7,m,g})_l = 6,727 \times C_6^m$; while for $7 \leq g \leq 21$, $N(G_{7,m,g})_l = k_g C_{g-1}^{(m-(g-6))}$, where $k_7 = 30,160$, $k_8 = 30,765$, $k_9 = 21,000$, $k_{10} = 28,364$, $k_{11} = 26,880$, $k_{12} = 26,460$, $k_{13} = 20,790$, $k_{14} = 10,290$, $k_{15} = 8,022$, $k_{16} = 2,940$, $k_{17} = 4,417$, $k_{18} = 2,835$, $k_{19} = 210$, $k_{20} = 21$, $k_{21} = 1$.

Keywords

Vertex Labeled, Connected Graph, Order Seven, Loops

Received: 5 April 2022, Accepted: 13 July 2022

<https://doi.org/10.26554/sti.2022.7.3.392-399>

1. INTRODUCTION

Without any doubt, one of the most widely used fields of mathematics is graph theory, especially to represent a real-life problem because of the flexibility of drawing a graph. The date of birth of graph theory began with the publication of Euler's solution regarding the Konigsberg problem in 1736, one of the mathematical fields with specific birth date is graph theory (Vasudev, 2006). Given a set $V = \{v_1, v_2, \dots, v_n\}$ of vertices/nodes, $V \neq \emptyset$, and a set of edges $E = \{e_{ij} | i, j \in V\}$, a graph $G(V,E)$ is a structure that consists of ordered pair of V and E . In real-life problems, the cities, buildings, airports, and others can be portrayed by vertices, while the roads that connect the cities, the pipes that connect the buildings, the flight paths that connect the airports, and others can be portrayed by edges. In order to represent real-life problems, on the edges, we can assign nonstructural information such as distance, time, flow, cost, and others by setting a non-negative number $c_{ij} \geq 0$.

There is a lot of application of graph theory in real-life problems, such as in computer science, chemistry, biology, sociology, agriculture, and others. In computer science, the graph structure plays an important role, for example, in designing a database, software engineering, and network system (Singh, 2014). The database is used for interconnecting analysis, as a

storage system with index-free adjacency, as a tool for graph-like-query, and for other purposes. In network design, graph-based representation makes the problem easier to visualize and provides a more accurate definition. In agriculture, the dynamic closures of the accounting structure are represented by a directed graph (Álvarez and Ehnts, 2015). The structure of graphs, together with discrete mathematics, are applied in chemistry to model the biological and physical properties of chemical compounds (Burch, 2019). The theoretical graph concept also was used by Gramatica et al. (2014) to represent or describe the possible modes of action of a given pharmacological compound. In biology, a phylogenetic tree was represented by a leaf-labeled tree (Huson and Bryant, 2006; Brandes and Cornelsen, 2009), while Mathur and Adlakha (2016) represented DNA using a combined tree. Hsu and Lin (2008) presented many graph theoretical concepts in engineering and computer science, and Al Etaiwi (2014) used the concepts of a complete graph, cycle graph, and minimum spanning tree to generate a complex cipher text. Priyadarsini (2015) explored the use of graph theory concepts, expander, and extremal graphs, in the design of some ciphers, whereas Ni et al. (2021) created ciphers using corona and bipartite graphs. In agriculture, graph theory concepts were used to

group agricultural workers performing manual tasks (Kawakura and Shibasaki, 2018), while the concept of graph coloring to optimize a farmer's goal was used by Kannimuthu et al. (2020). The relationship and unification of graph theory and physical-chemical measures (such as boiling and melting point, covalent and ionic potentials, and electronic density) make molecular topology can describe molecular structure comprehensively. A weighted directed graph, connectivity matrix, and Dijkstra's algorithm were used by Holmes et al. (2021) in plasma chemical reaction engineering. The basic structure of a directed graph is mostly used for the visualization of the reactions. Moreover, they use Gephi, an open-source graph software for visualization.

In 1874, Cayley counted the number of hydrocarbon isomers C_nH_{2n+2} (Cayley, 1874), and this process is similar to enumerating the number of a binary tree. Bona (2007) discussed the method of enumerating trees and forests. Redfield and Pólya are two other mathematicians that worked independently with graphical enumeration, especially in graph coloring (Bogart, 2004), and in graph enumeration, a comprehensive explanation of Pólya's counting theorem is one of the most powerful tools.

The number of graphs that can be formed for labeled and unlabeled graphs is different if we are given n vertices and m edges. For example, given $n=3$ and $m=2$, the number of simple connected unlabeled graphs that can be constructed is only one, while if every vertex is assigned labeled, The maximum number of graphs that can be created is three. The higher the order of the graph, the more labeled graph are formed. Agnarsson and Greenlaw (2006) gave the formula to enumerate graphs. However, no formula for enumerating graphs with special properties such as planarity or connectivity was provided.

There are some studies that have been done concerning the enumeration of the vertex-labeled graph with connectivity properties. In 2017, Amanto et al. (2017) proposed the formula to count disconnected vertices labeled graphs of order maximal four. For order five, the number of labeled vertices in connected graphs with no loops and may contain maximal five parallel edges had been proposed by Wamiliana et al. (2019). Amanto et al. (2021) studied the relationship between the formula for the number of connected vertex labels with no loops in graphs of order five and order six. Wamiliana et al. (2020) also discussed the number of vertices labeled connected graphs of order six with no parallel edges and a maximum of ten loops, while Puri et al. (2021) gave the formula to compute the number of vertices labeled connected graphs of order six without loops, while Anso et al. (2021) proposed the number of vertices labeled connected graphs of order seven with no loops.

The article is organized as follows: Section I provides information about graphs, graph applications in various fields, and previous research related to this topic. Section II discusses Observation and Investigation, while Section III discusses Results and Discussion. Section IV contains the conclusion.

2. OBSERVATION AND INVESTIGATION

Suppose that we are given the number of vertices $n=7$, and the number of edges m . We will construct connected graphs $G(V,E)$ of order n . Since the graph must be connected, then $m \geq 6$. Moreover, every vertex is labeled. Let g as the number of non-loop edges, $g \geq n-1$.

We start firstly by constructing all basic patterns of connected graphs of order seven. Note that the basic patterns contain no loops. The basic pattern starts with $m=6$, and with $n=7$, $m=6$, and constructs all possible patterns. After all possible patterns for $m=6$ are already constructed, then we continue with $m=7$, and so on until $m=21$. When $m=21$, only one pattern can be constructed because parallel edges are not allowed. Figure 1 shows some examples of patterns for $m=6$, Figure 2 shows some patterns that are isomorphic with the first graph in the second row of the graphs in Figure 1, and Figure 3 shows when $m=21$.




Figure 1. Some Basic Patterns for $n=7$ and $m=6$

Note that all isomorphic graphs will be counted in the pattern. However, we do not need to construct isomorphic graphs. Figure 2 shows the patterns of isomorphic graphs of the pattern of the first graph in the second row of Figure 1.




Figure 2. Some Patterns are Isomorphic with the First Graph in the Second Row in Figure 1




Figure 3. The Basic Pattern for $n=7$ and $m=21$




Figure 4. The Procedure

After constructing the basic pattern, the enumeration step begins. It begins from the first pattern of $n=7$ dan $m=6$ by adding one loop so that $m=7$, calculating the number of graphs that are able to be formed, and then continuing with this pattern by increasing the number of loops ($m=9$), and so on. Continue with this similar manner until the last pattern. The procedure can be put in the following diagram:

3. RESULTS AND DISCUSSION

The first step, as given in Figure 4 is constructing all possible patterns. Because there are many patterns obtained and due to limitation of space, here we give some patterns and also the number of all possible graphs formed according to the patterns. The obtained graphs are grouped by m and g , for example, for $n=6$, $m=6$, and $g=6$, the patterns are:

The results for all patterns are shown in Table 1 below:

Please note that in the table the dash sign (-) means there is impossible to construct the graph, while the empty space on the table means that we are not calculate more because g is fixed in each column, adding more edges simply adds more loops, and the constructed graph already constitute a sequence of numbers. The number in each column is able to be written as multiplication of a fix number and a sequence of number so that Table 2 can be rewritten in Table 3 as follow:

From Table 3 we can see that for every $g=6, 7, \dots, 21$, the number of graphs obtained are bigger as m increases, and the number of graphs obtained are multiplication of a fix number. For example, for $g=6$, the fix number is 6,727, and the number of graphs increases follows a certain pattens of sequence which is 1, 7, 28, 84, 210, 462, 924, 1,716, 3,003, 5,005.

| | | | | | | | | | |
|----|----|----|-----|-----|-----|-----|------|------|------|
| 1 | 7 | 28 | 84 | 210 | 462 | 924 | 1716 | 3003 | 5005 |
| 6 | 21 | 56 | 126 | 252 | 462 | 792 | 1287 | 2002 | |
| 15 | 35 | 70 | 126 | 210 | 330 | 495 | 715 | | |

Table 1. The Pattern for $n=7$, $m=6$, and $g=6$

| The patterns | The number of isomorphic graphs |
|--------------|---|
| | $C_1^7 \cdot C_6^6 = 7$ |
| | $\frac{7!}{2} = 2,520$ |
| | $C_1^7 \cdot C_3^6 \cdot C_2^3 = 420$ |
| | $C_1^7 \cdot C_2^6 \cdot C_3^4 \cdot 1 = 420$ |
| | $C_1^7 \cdot C_3^6 \cdot 3! = 840$ |
| | $C_1^7 \cdot C_1^6 \cdot C_2^5 \cdot C_3^3 = 420$ |
| | $C_1^7 \cdot C_1^6 \cdot C_1^5 \cdot C_4^4 = 210$ |
| | $C_2^7 \cdot C_1^5 \cdot C_2^4 \cdot 2! = 1,260$ |
| | $C_2^7 \cdot C_1^5 \cdot C_2^4 \cdot C_2^2 = 630$ |
| Total | 6,727 |

Table 2. The Number of Vertices Labeled Connected Graph of Order Seven Containing No Parallel Edges

1 The number of vertices labeled connected order seven graphs with no parallel edges

| m | g | | | | | |
|-----|------------|-------------|-------------|-------------|-------------|-------------|
| | 6 | 7 | 8 | 9 | 10 | 11 |
| 6 | 6,727 | - | - | - | - | - |
| 7 | 47,089 | 30,160 | - | - | - | - |
| 8 | 188,356 | 211,120 | 30,765 | - | - | - |
| 9 | 565,068 | 844,480 | 215,355 | 21,000 | - | - |
| 10 | 1,412,670 | 2,533,440 | 861,420 | 147,000 | 28,364 | - |
| 11 | 3,107,874 | 6,333,600 | 2,584,260 | 588,000 | 198,548 | 26,880 |
| 12 | 6,215,748 | 13,933,920 | 6,460,650 | 1,764,000 | 794,192 | 188,160 |
| 13 | 11,543,532 | 27,867,840 | 14,213,430 | 4,410,000 | 2,382,576 | 752,640 |
| 14 | 20,201,181 | 51,754,560 | 28,426,860 | 9,702,000 | 5,956,440 | 2,257,920 |
| 15 | 33,668,635 | 90,570,480 | 52,792,740 | 19,404,000 | 13,104,168 | 5,644,800 |
| 16 | - | 150,950,800 | 92,387,295 | 36,036,000 | 26,208,336 | 12,418,560 |
| 17 | - | - | 153,978,825 | 63,063,000 | 48,672,624 | 24,837,120 |
| 18 | - | - | - | 105,105,000 | 85,177,092 | 46,126,080 |
| 19 | - | - | - | - | 141,961,820 | 80,720,640 |
| 20 | - | - | - | - | - | 134,534,400 |

1 The number of vertices labeled connected order seven graphs with no parallel edges

| m | g | | | | |
|-----|-------------|-------------|------------|------------|------------|
| | 12 | 13 | 14 | 15 | 16 |
| 12 | 26,460 | - | - | - | - |
| 13 | 185,220 | 20,790 | - | - | - |
| 14 | 740,880 | 145,530 | 10,290 | - | - |
| 15 | 2,222,640 | 582,120 | 72,030 | 8,022 | - |
| 16 | 5,556,600 | 1,746,360 | 288,120 | 56,154 | 2,940 |
| 17 | 12,224,520 | 4,365,900 | 864,360 | 224,616 | 20,580 |
| 18 | 24,449,040 | 9,604,980 | 2,160,900 | 673,848 | 82,320 |
| 19 | 45,405,360 | 19,209,960 | 4,753,980 | 1,684,620 | 246,960 |
| 20 | 79,459,380 | 35,675,640 | 9,507,960 | 3,706,164 | 617,400 |
| 21 | 132,432,300 | 62,432,370 | 17,657,640 | 7,412,328 | 1,358,280 |
| 22 | - | 104,053,950 | 30,900,870 | 13,765,752 | 2,716,560 |
| 23 | - | - | 51,501,450 | 24,090,066 | 5,045,040 |
| 24 | - | - | - | 40,150,110 | 8,828,820 |
| 25 | - | - | - | - | 14,714,700 |

1 The number of vertices labeled connected order seven graphs with no parallel edges

| m | g | | | | |
|-----|------------|------------|-----------|---------|-------|
| | 12 | 13 | 14 | 15 | 16 |
| 17 | 4,417 | - | - | - | - |
| 18 | 30,919 | 2,835 | - | - | - |
| 19 | 123,676 | 19,845 | 210 | - | - |
| 20 | 371,028 | 79,380 | 1,470 | 21 | - |
| 21 | 927,570 | 238,140 | 5,880 | 147 | 1 |
| 22 | 2,040,654 | 595,350 | 17,640 | 588 | 7 |
| 23 | 4,081,308 | 1,309,770 | 44,100 | 1,764 | 28 |
| 24 | 7,579,572 | 2,619,540 | 97,020 | 4,410 | 84 |
| 25 | 13,264,251 | 4,864,860 | 194,040 | 97,02 | 210 |
| 26 | 22,107,085 | 8,513,505 | 360,360 | 19,404 | 462 |
| 27 | - | 14,189,175 | 630,630 | 36,036 | 924 |
| 28 | - | - | 1,051,050 | 63,063 | 1,716 |
| 29 | - | - | - | 105,105 | 3,003 |
| 30 | - | - | - | - | 5,005 |

Table 3. Alternative form of Table 2

| The number of vertices labeled connected order seven graphs with no parallel edges | | | | | | |
|--|-------------|--------------|--------------|--------------|--------------|--------------|
| m | g | | | | | |
| | 6 | 7 | 8 | 9 | 10 | 11 |
| 6 | 1x6,727 | - | - | - | - | - |
| 7 | 7x6,727 | 1x30,160 | - | - | - | - |
| 8 | 28x6,727 | 7x30,160 | 1x30,765 | - | - | - |
| 9 | 84x6,727 | 28x30,160 | 7x30,765 | 1x21,000 | - | - |
| 10 | 210x6,727 | 84x30,160 | 28x30,765 | 7x21,000 | 1x28,364 | - |
| 11 | 462x6,727 | 210x30,160 | 84x30,765 | 28x21,000 | 7x28,364 | 1x26,880 |
| 12 | 924x6,727 | 462x30,160 | 210x30,765 | 84x21,000 | 28x28,364 | 7x26,880 |
| 13 | 1,716x6,727 | 924x30,160 | 462x30,765 | 210x21,000 | 84x28,364 | 28x26,880 |
| 14 | 3,003x6,727 | 1,716x30,160 | 924x30,765 | 462x21,000 | 210x28,364 | 84x26,880 |
| 15 | 5,005x6,727 | 3,003x30,160 | 1,716x30,765 | 924x21,000 | 462x28,364 | 210x26,880 |
| 16 | - | 5,005x30,160 | 3,003x30,765 | 1,716x21,000 | 924x28,364 | 462x26,880 |
| 17 | - | - | 5,005x30,765 | 3,003x21,000 | 1,716x28,364 | 924x26,880 |
| 18 | - | - | - | 5,005x21,000 | 3,003x28,364 | 1,716x26,880 |
| 19 | - | - | - | - | 5,005x28,364 | 3,003x26,880 |
| 20 | - | - | - | - | - | 5,005x26,880 |

| The number of vertices labeled connected order seven graphs with no parallel edges | | | | | |
|--|--------------|--------------|--------------|-------------|-------------|
| m | g | | | | |
| | 12 | 13 | 14 | 15 | 16 |
| 12 | 1x26,460 | - | - | - | - |
| 13 | 7x26,460 | 1x20,790 | - | - | - |
| 14 | 28x26,460 | 7x20,790 | 1x10,290 | - | - |
| 15 | 84x26,460 | 28x20,790 | 7x10,290 | 1x8022 | - |
| 16 | 210x26,460 | 84x20,790 | 18x10,290 | 7x8,022 | 1x2,940 |
| 17 | 462x26,460 | 210x20,790 | 84x10,290 | 28x8,022 | 7x2,940 |
| 18 | 924x26,460 | 462x20,790 | 210x10,290 | 84x8,022 | 28x2,940 |
| 19 | 1,716x26,460 | 924x20,790 | 462x10,290 | 210x8,022 | 84x2,940 |
| 20 | 3,003x26,460 | 1,716x20,790 | 924x10,290 | 462x8,022 | 210x2,940 |
| 21 | 5,005x26,460 | 3,003x20,790 | 1,716x10,290 | 924x8,022 | 462x2,940 |
| 22 | - | 5,005x20,790 | 3,003x10,290 | 1,716x8,022 | 924x2,940 |
| 23 | - | - | 5,005x10,290 | 3,003x8,022 | 1,716x2,940 |
| 24 | - | - | - | 5,005x8,022 | 3,003x2,940 |
| 25 | - | - | - | - | 5,005x2,940 |

| The number of vertices labeled connected order seven graphs with no parallel edges | | | | | |
|--|-------------|-------------|-----------|----------|---------|
| m | g | | | | |
| | 12 | 13 | 14 | 15 | 16 |
| 17 | 1x4,417 | - | - | - | - |
| 18 | 7x4,417 | 1x2,835 | - | - | - |
| 19 | 28x4,417 | 7x2,835 | 1x210 | - | - |
| 20 | 84x4,417 | 28x2,835 | 7x210 | 1x21 | - |
| 21 | 210x4,417 | 84x2,835 | 28x210 | 7x21 | 1x1 |
| 22 | 462x4,417 | 210x2,835 | 84x210 | 28x21 | 7x1 |
| 23 | 924x4,417 | 462x2,835 | 210x210 | 84x21 | 28x1 |
| 24 | 1,716x4,417 | 924x2,835 | 462x210 | 210x21 | 84x1 |
| 25 | 3,003x4,417 | 1,716x2,835 | 924x210 | 462x21 | 210x1 |
| 26 | 5,005x4,417 | 3,003x2,835 | 1,716x210 | 924x21 | 462x1 |
| 27 | - | 5,005x2,835 | 3,003x210 | 1,716x21 | 924x1 |
| 28 | - | - | 5,005x210 | 3,003x21 | 1,716x1 |
| 29 | - | - | - | 5,005x21 | 3,003x1 |
| 30 | - | - | - | - | 5,005x1 |

| | | | | | | |
|----|----|----|----|-----|-----|-----|
| 20 | 35 | 56 | 84 | 120 | 165 | 220 |
| 15 | 21 | 28 | 36 | 45 | 55 | |
| | 6 | 7 | 8 | 9 | 10 | |
| | | 1 | 1 | 1 | 1 | |

Notate $(G_{7,m,g})_l$ as the number of vertices labeled connected graphs of order seven containing no parallel edges (loops are allowable) with the number of edges is m and the number of non loop edges is g .

Result 1: Given $m > 6, g = 6$, the total number of vertices labeled connected graphs of order seven with no parallel edges is $N(G_{7,m,g})_l = 6,727 \times C_6^m$

Proof:

Consider the above sequence of numbers.

That sequence of numbers is able to be represented by polynomial of order six because the fixed

$$Q_5m = \alpha_6m^6 + \alpha_5m^5 + \alpha_4m^4 + \alpha_3m^3 + \alpha_2m^2 + \alpha_1m + \alpha_0$$

The following system of equations is obtained by substituting $m = 6, 7, 8, 9, 10, 11, 12$ to $Q_5(m)$.

- 6, 727 = 46, 656 α_6 + 7, 776 α_5 + 1, 296 α_4 + 216 α_3 + 36 α_2 + 6 α_1 + α_0 (1)
- 47, 089 = 117, 649 α_6 + 16, 807 α_5 + 2, 401 α_4 + 343 α_3 + 49 α_2 + 7 α_1 + α_0 (2)
- 188, 356 = 262, 144 α_6 + 32, 768 α_5 + 4, 096 α_4 + 512 α_3 + 64 α_2 + 8 α_1 + α_0 (3)
- 565, 068 = 531, 441 α_6 + 59, 049 α_5 + 6, 561 α_4 + 729 α_3 + 81 α_2 + 9 α_1 + α_0 (4)
- 1, 412, 670 = 1, 000, 000 α_6 + 100, 000 α_5 + 10, 000 α_4 + 1, 000 α_3 + 100 α_2 + 10 α_1 + α_0 (5)
- 3, 107, 874 = 1, 771, 561 α_6 + 161, 051 α_5 + 14, 641 α_4 + 1, 331 α_3 + 121 α_2 + 11 α_1 + α_0 (6)
- 6, 215, 748 = 2, 985, 984 α_6 + 248, 832 α_5 + 20, 736 α_4 + 1, 728 α_3 + 144 α_2 + 12 α_1 + α_0 (7)

These equations form a system of equations that can be transformed into a matrix $Ax = b$ as follow:

$$\begin{bmatrix} 46,656 & 7,776 & 1,296 & 216 & 36 & 6 & 1 \\ 117,649 & 16,807 & 2,401 & 343 & 49 & 7 & 1 \\ 262,144 & 32,768 & 4,096 & 512 & 64 & 8 & 1 \\ 531,441 & 59,049 & 6,561 & 729 & 81 & 9 & 1 \\ 1,000,000 & 100,000 & 10,000 & 1,000 & 100 & 10 & 1 \\ 1,771,561 & 161,051 & 14,641 & 1,331 & 121 & 11 & 1 \\ 2,985,984 & 248,832 & 20,736 & 1,728 & 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} \alpha_6 \\ \alpha_5 \\ \alpha_4 \\ \alpha_3 \\ \alpha_2 \\ \alpha_1 \\ \alpha_0 \end{bmatrix} = \begin{bmatrix} 6,727 \\ 47,089 \\ 188,356 \\ 565,068 \\ 1,412,670 \\ 3,107,874 \\ 6,215,748 \end{bmatrix}$$

By solving this system of equations we get: $\alpha_6 = \frac{6,727}{720}, \alpha_5 = \frac{100,905}{720}, \alpha_4 = \frac{57,175}{720}, \alpha_3 = \frac{151,575}{720}, \alpha_2 = \frac{1,843,198}{720}, \alpha_1 = \frac{807,240}{720}, \alpha_0 = 0$.

Thus

$$\begin{aligned} Q_5(m) &= \alpha_6m^6 + \alpha_5m^5 + \alpha_4m^4 + \alpha_3m^3 + \alpha_2m^2 + \alpha_1m + \alpha_0 \\ &= \frac{6,727}{720}m^6 - \frac{100,905}{720}m^5 + \frac{57,175}{720}m^4 - \frac{151,575}{720}m^3 \\ &\quad + \frac{1,843,198}{720}m^2 - \frac{807,240}{720}m \\ &= \frac{6,727}{720}(m^6 - 15m^5 + 85m^4 - 225m^3 + 274m^2 - 120m) \\ &= \frac{6,727 \cdot 3 - 1)(m-2)(m-3)(m-4)(m-5)(m-6)!}{6.5.4.3.2.1(m-6)!} \\ &= 6,727 \times C_6^m \end{aligned}$$

Please note that for every g (every column, the sequence of numbers is the same, except the multiplier). Thus, the

polynomial related to the sequence of numbers is the same. However, because the multipliers are different, it will cause different formulas.

Result 2: Given $m \geq 7, g = 7$, the total number of vertices labeled connected graphs of order seven with no parallel edges is $N(G_{7,m,g})_l = 30,160 \times C_6^{(m-1)}$

Proof:

The polynomial that represents the sequence is the same which is

$$Q_5m = \alpha_6m^6 + \alpha_5m^5 + \alpha_4m^4 + \alpha_3m^3 + \alpha_2m^2 + \alpha_1m + \alpha_0$$

However, for $m = 7$, the numbers of graphs are different. The following system of equations is obtained by substituting $m = 7, 8, 9, 10, 11, 12, 13$ to the equation.

- 30, 160 = 117, 649 α_6 + 16, 807 α_5 + 2, 401 α_4 + 343 α_3 + 49 α_2 + 7 α_1 + α_0 (8)
- 211, 120 = 262, 144 α_6 + 32, 768 α_5 + 4, 096 α_4 + 512 α_3 + 64 α_2 + 8 α_1 + α_0 (9)
- 844, 480 = 531, 441 α_6 + 59, 049 α_5 + 6, 561 α_4 + 729 α_3 + 81 α_2 + 9 α_1 + α_0 (10)
- 2, 533, 440 = 1, 000, 000 α_6 + 100, 000 α_5 + 10, 000 α_4 + 1, 000 α_3 + 100 α_2 + 10 α_1 + α_0 (11)
- 6, 333, 600 = 1, 771, 561 α_6 + 161, 051 α_5 + 14, 641 α_4 + 1, 331 α_3 + 121 α_2 + 11 α_1 + α_0 (12)
- 13, 933, 920 = 2, 985, 984 α_6 + 248, 832 α_5 + 20, 736 α_4 + 1, 728 α_3 + 144 α_2 + 12 α_1 + α_0 (13)
- 27, 867, 840 = 4, 826, 809 α_6 + 371, 293 α_5 + 28, 561 α_4 + 2, 197 α_3 + 169 α_2 + 13 α_1 + α_0 (14)

These equations form a system of equations that can be transformed into a matrix $Ax = b$ as follow:

$$\begin{bmatrix} 117,649 & 16,807 & 2,401 & 343 & 49 & 7 & 1 \\ 262,144 & 32,768 & 4,096 & 512 & 64 & 8 & 1 \\ 531,441 & 59,049 & 6,561 & 729 & 81 & 9 & 1 \\ 1,000,000 & 100,000 & 10,000 & 1,000 & 100 & 10 & 1 \\ 1,771,561 & 161,051 & 14,641 & 1,331 & 121 & 11 & 1 \\ 2,985,984 & 248,832 & 20,736 & 1,728 & 144 & 12 & 1 \\ 4,826,809 & 371,293 & 28,561 & 2,197 & 169 & 13 & 1 \end{bmatrix} \begin{bmatrix} \alpha_6 \\ \alpha_5 \\ \alpha_4 \\ \alpha_3 \\ \alpha_2 \\ \alpha_1 \\ \alpha_0 \end{bmatrix} = \begin{bmatrix} 30,160 \\ 211,120 \\ 844,480 \\ 2,533,440 \\ 6,333,600 \\ 13,933,920 \\ 27,867,840 \end{bmatrix}$$

By solving this system of equations we get: $\alpha_6 = \frac{30,160}{720}, \alpha_5 = \frac{633,360}{720}, \alpha_4 = \frac{5,278,000}{720}, \alpha_3 = \frac{22,167,600}{720}, \alpha_2 = \frac{48,979,840}{720}, \alpha_1 = \frac{53,202,240}{720}, \text{ and } \alpha_0 = \frac{21,715,200}{720}$.

Thus

$$\begin{aligned} Q_5(m) &= \alpha_6m^6 + \alpha_5m^5 + \alpha_4m^4 + \alpha_3m^3 + \alpha_2m^2 + \alpha_1m + \alpha_0 \\ &= \frac{30,160}{720}m^6 - \frac{633,360}{720}m^5 + \frac{5,278,000}{720}m^4 - \frac{22,167,600}{720}m^3 \\ &\quad + \frac{48,979,840}{720}m^2 - \frac{53,202,240}{720}m + \frac{21,715,200}{720} \\ &= \frac{30,160}{720}(m^6 - 21m^5 + 175m^4 - 735m^3 + 624m^2 - 1764m + 720) \\ &= \frac{30,160 \cdot 3 - 1)(m-2)(m-3)(m-4)(m-5)(m-6)}{720} \\ &= \frac{30,160 \cdot 3 - 1)(m-2)(m-3)(m-4)(m-5)(m-6)(m-7)!}{6.5.4.3.2.1(m-7)!} \\ &= 30,160 \times C_6^{(m-1)} \end{aligned}$$

The following results are obtained by doing the similar manner:

For $m \geq 8, g = 8$, is $N(G_{7,m,g})_l = 30,765 \times C_7^{(m-2)}$

For $m \geq 9, g = 9$, is $N(G_{7,m,g})_l = 21,000 \times C_8^{(m-3)}$

For $m \geq 10, g = 10$, is $N(G_{7,m,g})_l = 28,364 \times C_9^{(m-4)}$

For $m \geq 11, g = 11$, is $N(G_{7,m,g})_l = 26,880 \times C_{10}^{(m-5)}$

For $m \geq 12, g = 12$, is $N(G_{7,m,g})_l = 26,460 \times C_{11}^{(m-6)}$

For $m \geq 13, g = 13$, is $N(G_{7,m,g})_l = 20,790 \times C_{12}^{(m-7)}$

For $m \geq 14, g = 14$, is $N(G_{7,m,g})_l = 10,290 \times C_{13}^{(m-8)}$

For $m \geq 15$, $g=15$, is $N(G_{7,m,g})_l = 8,022 \times C_{14}^{(m-9)}$
 For $m \geq 16$, $g=16$, is $N(G_{7,m,g})_l = 2,940 \times C_{15}^{(m-10)}$
 For $m \geq 17$, $g=17$, is $N(G_{7,m,g})_l = 4,417 \times C_{16}^{(m-11)}$
 For $m \geq 18$, $g=18$, is $N(G_{7,m,g})_l = 2,835 \times C_{17}^{(m-12)}$
 For $m \geq 19$, $g=19$, is $N(G_{7,m,g})_l = 210 \times C_{18}^{(m-13)}$
 For $m \geq 20$, $g=20$, is $N(G_{7,m,g})_l = 21 \times C_{19}^{(m-14)}$
 For $m \geq 21$, $g=21$, is $N(G_{7,m,g})_l = C_{20}^{(m-15)}$

Note that the multiplier for $g=6$ is the same as the multiplier of $t=6$ in Ansori et al. (2021), as well as $g=7$ with $t=7$, and so on until $g=21$ with $t=21$. However, the formulas are different because in Ansori et al. (2021) the formula are for connected vertex labeled graph without loops while in this study is for connected vertices labeled graph without parallel edges. For example, for $g=8$, $N(G_{7,m,g})_l = 30,765 \times C_7^{(m-2)}$, while in Ansori et al. (2021), for $t=8$, $N(G_{7,m,8}) = 30,765 \times C_7^{(m-1)}$.

4. CONCLUSIONS

Based on the above reasoning, we may conclude that the number of vertices in a labeled connected graph of order seven with no parallel edges is $N(G_{7,m,g})_l = 6,727 \times C_6^m$ for $g=6$, while for $7 \leq g \leq 21$, $N(G_{7,m,g})_l = k_g C_{g-1}^{(m-(g-6))}$, where $k_7 = 30,160$, $k_8 = 30,765$, $k_9 = 21,000$, $k_{10} = 28,364$, $k_{11} = 26,880$, $k_{12} = 26,460$, $k_{13} = 20,790$, $k_{14} = 10,290$, $k_{15} = 8,022$, $k_{16} = 2,940$, $k_{17} = 4,417$, $k_{18} = 2,835$, $k_{19} = 210$, $k_{20} = 21$, $k_{21} = 1$.

5. ACKNOWLEDGMENT

The Research Center Universitas Lampung funded this study under Postgraduate research grant project No. 815/UN26.21/PN/2022, and the authors gratefully acknowledge the funding.

REFERENCES

- Agnarsson, G. and R. Greenlaw (2006). *Graph Theory: Modeling, Applications, and Algorithms*. Prentice-Hall, Inc.
- Al Etaiwi, W. M. (2014). Encryption Algorithm Using Graph Theory. *Journal of Scientific Research and Reports*, **3**(19); 2519–2527
- Álvarez, M. C. and D. Ehnts (2015). *The Roads Not Taken: Graph Theory and Macroeconomic Regimes in Stock-Flow Consistent Modeling*. Working Paper
- Amanto, A., N. Notiragayu, L. Zakaria, and W. Wamiliana (2021). The Relationship of the Formulas for the Number of Connected Vertices Labeled Graphs with Order Five and Order Six without Loops. *Desimal: Jurnal Matematika*, **4**(3); 357–364
- Amanto, A., W. Wamiliana, M. Usman, and R. Permatasari (2017). Counting the Number of Disconnected Vertex Labeled Graphs with Order Maximal Four. *Science International Lahore*, **29**(6); 1181–1186
- Ansori, M., W. Wamiliana, and F. Puri (2021). Determining the Number of Connected Vertex Labeled Graphs of Order Seven without Loops by Observing the Patterns of Formula for Lower Order Graphs with Similar Property. *Science and Technology Indonesia*, **6**(4); 328–336
- Bogart, K. P. (2004). *Combinatorics Through Guided Discovery*. LibreText
- Bona, M. (2007). *Introduction to Enumerative Combinatorics*. McGraw-Hill Science/Engineering/Math
- Brandes, U. and S. Cornelsen (2009). Phylogenetic Graph Models Beyond Trees. *Discrete Applied Mathematics*, **157**(10); 2361–2369
- Burch, K. J. (2019). Chemical Applications of Graph Theory. In *Mathematical Physics in Theoretical Chemistry*. Elsevier, pages 261–294
- Cayley, P. (1874). LVII. On the Mathematical Theory of Isomers. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, **47**(314); 444–447
- Gramatica, R., T. Di Matteo, S. Giorgetti, M. Barbiani, D. Bevec, and T. Aste (2014). Graph Theory Enables Drug Repurposing—How a Mathematical Model Can Drive the Discovery of Hidden Mechanisms of Action. *PloS One*, **9**(1); e84912
- Holmes, T. D., R. H. Rothman, and W. B. Zimmerman (2021). Graph Theory Applied to Plasma Chemical Reaction Engineering. *Plasma Chemistry and Plasma Processing*, **41**(2); 531–557
- Hsu, L. H. and C. K. Lin (2008). *Graph Theory and Interconnection Networks*. CRC Press
- Huson, D. H. and D. Bryant (2006). Application of Phylogenetic Networks in Evolutionary Studies. *Molecular Biology and Evolution*, **23**(2); 254–267
- Kannimuthu, S., D. Bhanu, and K. Bhuvaneshwari (2020). A Novel Approach for Agricultural Decision Making Using Graph Coloring. *SN Applied Sciences*, **2**(1); 1–6
- Kawakura, S. and R. Shibasaki (2018). Grouping Method Using Graph Theory for Agricultural Workers Engaging in Manual Tasks. *Journal of Advanced Agricultural Technologies*, **5**(3); 173–181
- Mathur, R. and N. Adlakha (2016). A Graph Theoretic Model for Prediction of Reticulation Events and Phylogenetic Networks for DNA Sequences. *Egyptian journal of Basic and Applied Sciences*, **3**(3); 263–271
- Ni, B., R. Qazi, S. U. Rehman, and G. Farid (2021). Some Graph-Based Encryption Schemes. *Journal of Mathematics*, **2021**; 1–8
- Priyadarsini, P. (2015). A Survey on Some Applications of Graph Theory in Cryptography. *Journal of Discrete Mathematical Sciences and Cryptography*, **18**(3); 209–217
- Puri, F., M. Usman, M. Ansori, and Y. Antoni (2021). The Formula to Count the Number of Vertices Labeled Order Six Connected Graphs with Maximum Thirty Edges without Loops. *Journal of Physics: Conference Series*, **1751**(1); 012023
- Singh, R. P. (2014). Application of Graph Theory in Computer Science and Engineering. *International Journal of Computer Applications*, **104**(1)
- Vasudev, C. (2006). *Graph Theory with Applications*. New Age

- International
Wamiliana, W., A. Amanto, M. Usman, M. Ansori, and F. C. Puri (2020). Enumerating the Number of Connected Vertices Labeled Graph of Order Six with Maximum Ten Loops and Containing No Parallel Edges. *Science and Technology Indonesia*, 5(4); 131–135
- Wamiliana, W., A. Nuryaman, A. Sutrisno, and N. Prayoga (2019). Determining the Number of Connected Vertices Labelled Graph of Order Five with Maximum Number of Parallel Edges is Five and Containing No Loops. *Journal of Physics: Conference Series*, **1338**; 012043

● 7% Overall Similarity

Top sources found in the following databases:

- 2% Internet database
- Crossref database
- 2% Submitted Works database
- 5% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

| | | |
|---|--|-----|
| 1 | F C Puri, Wamiliana, M Usman, Amanto, M Ansori, Y Antoni. "The For... | 2% |
| | Crossref | |
| 2 | Sriwijaya University on 2022-03-01 | 2% |
| | Submitted works | |
| 3 | Amanto, Notiragayu, F C Puri, Y Antoni, Wamiliana. "Counting the num... | 1% |
| | Crossref | |
| 4 | repository.lppm.unila.ac.id | <1% |
| | Internet | |
| 5 | Mohammed J. Zaki. "Predicting Protein Folding Pathways", Advanced I... | <1% |
| | Crossref | |
| 6 | discovery.researcher.life | <1% |
| | Internet | |
| 7 | RMIT University on 2021-09-18 | <1% |
| | Submitted works | |
| 8 | ejournal.radenintan.ac.id | <1% |
| | Internet | |

● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)
- Manually excluded text blocks

EXCLUDED SOURCES

sciencetechindonesia.com

Internet

54%

doaj.org

Internet

11%

sciencetechindonesia.com

Internet

4%

EXCLUDED TEXT BLOCKS

Science and Technology Indonesiae-ISSN:2580-4391 p-ISSN:2580-4405Vol

www.coursehero.com

of Mathematics, Faculty of Mathematics and Natural Science, Universitas Lampung

Wamiliana, A Nuryaman, Amanto, A Sutrisno, N A Prayoga. "Determining the Number of Connected Vertices..."