

PAPER NAME

Sub-exact sequence of rough groups

AUTHOR

Nevi Setyaningsih

WORD COUNT

3118 Words

CHARACTER COUNT

14325 Characters

PAGE COUNT

6 Pages

FILE SIZE

440.4KB

SUBMISSION DATE

Aug 19, 2022 10:09 PM GMT+7

REPORT DATE

Aug 19, 2022 10:09 PM GMT+7

● 15% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 12% Internet database
- 15% Publications database
- Crossref database
- Crossref Posted Content database
- 8% Submitted Works database

● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources
- Manually excluded text blocks



Sub-exact sequence of rough groups

Nevi Setyaningsih¹, Fitriani^{1*}, Ahmad Faisol¹

¹ Universitas Lampung, Indonesia

✉ fitriani.1984@fmipa.unila.ac.id *

Article Information

Submitted May 05, 2021

Revised August, 12 2021

Accepted August 15, 2021

Keywords

Exact Sequence;

Sub-exact Sequence;

Group;

Rough Set.

Abstract

Rough Set Theory (RST) is an essential mathematical tool to deal with imprecise, inconsistent, incomplete information and knowledge. Some algebraic structures, such as groups, rings, and modules, have been presented on rough set theory. The sub-exact sequence is a generalization of the exact sequence. In this paper, we introduce the notion of a sub-exact sequence of groups. Furthermore, we give some properties of the rough group and rough sub-exact sequence of groups.

INTRODUCTION

Pawlak first introduced the rough set theory in 1982 (Pawlak, 1991). This is an advanced theory of set theory, in which the subsets of the universal set are explained by a tuple of lower approximations and upper approximations. The basic concept of a rough set is the equivalence relation.

Let U is a finite non-empty set called universal set, and θ is an equivalence relation in U . The tuple (U, θ) is called the approximation space (Miao et al., 2005). The lower approximation of a set is a combination of all equivalence classes, the smallest subset. The upper approximation of a set X , denote by $Apr(X)$, combines all equivalence classes with the largest set. A tuple $(A, B) \in P(U) \times P(U)$ is called a rough set in (U, θ) if $(A, B) = Apr(X)$ for $X \in P(U)$, where $P(U)$ is the power set of (U) and $Apr(X)$ is the approximation of X .

In 1994, Biswas and Nanda (Biswas & Nanda, 1994) introduced the rough subgroup. Furthermore, Kuroki (Kuroki, 1997) introduced the idea of rough ideals in semigroups in 1997. In 2001, Han Suqing (Miao et al., 2005) investigated homomorphism and isomorphism on rough set. Subsequently, Davvaz (Davvaz, 2004) studied the relationship between the rough set and the ring theory by considering the ring as a universal set and introducing the abstraction ideal notation and the abstraction subring concerning the ideal of a ring. In 2006, Davvaz and Mahdavi-pour (Davvaz & Mahdavi-pour, 2006) investigated the rough module. Furthermore, rough semigroups on approximation space (Bagirmaz & Ozcan, 2015), homomorphism of rough groups (Jesmalar, 2017), rough G -modules (Isaac & Paul, 2017), and direct products of rough subgroups (Bagirmaz, 2019) are investigated. Besides that, Sinha and Prakash (Sinha & Prakash, 2016) studied the exact sequence of the rough module. They define the exact sequence of rough modules on the rough ring $Apr(R)$.

Let R be a ring and let $A \xrightarrow{f} B \xrightarrow{g} C$ is the exact sequence over R -module such that $\text{Im } f = \text{Ker } g (=g^{-1}(0))$. Davvaz & Parnian-Garamaleky (1999) introduced the concept of U -exact by replacing 0 with sub-module $U \subseteq C$. Motivated by Davvaz and Parnian-Garamaleky, who introduced the U -exact sequence concept. In 2016, Fitriani et al. gave new ideas from

How to cite

Setyaningsih, N., Fitriani, F., & Faisol, A. (2021). Sub-exact sequence of rough groups. *Al-Jabar: Jurnal Pendidikan Matematika*, 12(2), 267-272.

E-ISSN

2540-7562

Published by

Mathematics Education Department, UIN Raden Intan Lampung.

B with arbitrary submodule X of B (Fitriani et al., 2016). Besides that, motivated by the U-exact sequence, Elfiyanti et al. give an Abelian property of the category of U-complexes (Elfiyanti et al., 2016). In 2017, Aminizadeh et al. (Aminizadeh et al., 2018) introduced quasi exact sequence of S-acts. Furthermore, Fitriani et al. gave an X-sub-linearly independent module a new concept (Fitriani et al., 2017). In 2018, they introduced a U-generated module (Fitriani et al., 2018b). Furthermore, they established U -basis and U -free modules by using the concept of a sub-exact sequence of modules (Fitriani et al., 2018a). In addition, the sub-exact sequences can be used to determine the Noetherian property of the submodule of the generalized power series module (Faisol et al., 2021).

Let U', U, U'' be R -modules and X be a submodule of U . The triple (U', U, U'') is said to be an X -sub-exact on U if there is R -homomorphisms f and g so the sequence of R -modules $U' \xrightarrow{f} U \xrightarrow{g} U''$ is exact. The set of all submodule X in L such that the sequence $U' \xrightarrow{f} U \xrightarrow{g} U''$ is exact, is denoted by $\sigma(U', U, U'')$.

Based on the definition of sub-exact sequence, we introduce the notion of a sub-exact sequence of rough groups in this paper. Furthermore, we give some properties of the rough group and sub-exact sequence of rough groups. These properties can be used to determine all submodules X in U so that the triple (U', U, U'') is rough X -sub-exact.

METHODS

The research methods are based on literature, especially those related to the standard set, the upper and lower approximations space, the rough group, the exact sequence, and the sub-exact sequence. Our research steps are as follows. First, we define the rough sub-exact sequences of rough groups. Furthermore, we investigate the properties of the rough group and construct the example of the rough sub-exact sequence of groups using the finite set. Finally, we investigate the properties of the rough sub-exact sequences of rough groups.




Figure 1. Research stage diagram

RESULTS AND DISCUSSION

Before we construct the rough sub-exact of groups, we recall the definition of rough group, sub-exact sequence, and exact sequence of the rough module over the rough ring as follows.

Definition 1. (Miao et al., 2005) Let $K = (U, R)$ be an approximation space and $*$ be a binary operation defined on U . A subset $G (G \neq \emptyset)$ of universe U is called a rough group if $Apr(G) = (\underline{Apr}(G), \overline{Apr}(G))$ the following properties are satisfied:

- (1) for every $x * y \in \overline{Apr}(G), x * y \in G$;
- (2) association property holds in $\overline{Apr}(G)$;

- (3) there exists $e \in \overline{Apr}(G)$ such that $x \in G, x * e = e * x = x$; e is called the rough identity element of rough group G ;
- (4) for every $x \in G$, there exists $y \in G$, such that $x * y = y * x = e$; y is called the rough inverse element of x in G .

Definition 2. (Fitriani et al., 2016) Let K, L, M be R -modules and X be a submodule of L . Then the triple (K, L, M) is said to be X -sub-exact at L if there exist R -homomorphisms f and g such that the sequence of R -modules and R -homomorphisms $K \xrightarrow{f} X \xrightarrow{g} M$ is exact.

Definition 3. (Sinha & Prakash, 2016) A sequence $Apr(M') \xrightarrow{f} Apr(M) \xrightarrow{g} Apr(M'')$ of two homomorphism of a module over the ring (R) is said to be rough exact if $Im f = Ker g$. This happens if and only if $gf = 0$, and the relation $g(x)=0, x \in Apr(M)$, implies that $x = f(x')$ for some $x' \in Apr(M')$.

Based on the definition of the sub-exact sequence of modules, we define the exact sequence of rough groups as follows.

Definition 4. Let U', U, U'' be rough groups, and X be a rough subgroup of U . Then the triple (U', U, U'') is said to be rough X -sub-exact at U if there exist rough homomorphisms f and g such that the sequence of rough groups and rough group homomorphisms:

$$Apr(U') \xrightarrow{f} Apr(U) \xrightarrow{g} Apr(U'')$$

is exact.

Based on Definition 4, every rough exact sequence is a rough sub-exact sequence of groups. However, the converse need not be true. Therefore, before we give the properties of a rough sub-exact sequence, we provide the properties of the rough group as follows.

Proposition 5. Let (U, θ) be an approximation space, and let X_1, X_2 be subsets of U . If X_1 and X_2 are rough groups with $\overline{Apr}(X_1) = \overline{Apr}(X_2)$, then $X_1 \cup X_2$ is also the rough group.

Proof. Let $*$ be a binary operation defined on U . It is clear that $\overline{Apr}(X_1 \cup X_2) = \overline{Apr}(X_1) \cup \overline{Apr}(X_2)$. By assumption, $\overline{Apr}(X_1) = \overline{Apr}(X_2)$. Let $a \in \overline{Apr}(X_1 \cup X_2)$ and $b \in \overline{Apr}(X_1 \cup X_2)$. Then $a \in \overline{Apr}(X_1)$ and $b \in \overline{Apr}(X_1)$. Since X_1 is the rough group, we have $a * b \in \overline{Apr}(X_1)$. Hence $a * b \in \overline{Apr}(X_1 \cup X_2)$. For every $a, b, c \in X_1 \cup X_2$, the associative property holds in $\overline{Apr}(X_1 \cup X_2)$, i.e., $a * (b * c) = (a * b) * c$. Since X_1 is a rough group, we have the identity element $e \in \overline{Apr}(X_1) = \overline{Apr}(X_1 \cup X_2)$. Finally, we want to show that every element in $\overline{Apr}(X_1 \cup X_2)$ has an inverse. By assumption, $\overline{Apr}(X_1) = \overline{Apr}(X_2)$, so that $\overline{Apr}(X_1 \cup X_2) = \overline{Apr}(X_1)$. This implies that every element in the set $X_1 \cup X_2$ has an inverse in $\overline{Apr}(X_1 \cup X_2)$. Therefore, it proved $(X_1 \cup X_2)$ is a rough group. \square

By applying Proposition 5 to a finite number of subsets U , we have the following property of the rough group.

Corollary 6. Let (U, θ) be an approximation space, and let X_1, X_2, \dots, X_n be subsets of U . If X_1, X_2, \dots, X_n are rough groups with $\overline{Apr}(X_1) = \overline{Apr}(X_2) = \dots = \overline{Apr}(X_n)$, then $\cup_{i=1}^n X_i$ is also the rough group.

Example 1. Let $U = \{0,1,2,3,4, \dots, 149\}$. We define the relation θ in U , where $u \theta v$ if and only if $u - v = 13k$, for some $k \in \mathbb{Z}$. It is easy to show that θ is an equivalence relation on

U . From this equivalence relation, we have 13 equivalence classes as described in the following table.

Table 1. The Equivalence Classes of U

No	The Equivalence Class	The Elements
1	$E_1 = [1]$	{1,14,27,40,53,66,79,92,105,118,131,144}
2	$E_2 = [2]$	{2,15,28,41,54,67,80,93,106,119,132,145}
3	$E_3 = [3]$	{3,16,29,42,55,68,81,94,107,120,133,146}
4	$E_4 = [4]$	{4,17,30,43,56,69,82,95,108,121,134,147}
5	$E_5 = [5]$	{5,18,31,44,57,70,83,96,109,122,135,148}
6	$E_6 = [6]$	{6,19,32,45,58,71,84,97,110,123,136,149}
7	$E_7 = [7]$	{7,20,33,46,59,72,85,98,111,124,137}
8	$E_8 = [8]$	{8,21,34,47,60,73,86,99,112,125,138}
9	$E_9 = [9]$	{9,22,35,48,61,74,87,100,113,126,139}
10	$E_{10} = [10]$	{10,23,36,49,62,75,88,101,114,127,140}
11	$E_{11} = [11]$	{11,24,37,50,63,76,89,102,115,128,141}
12	$E_{12} = [12]$	{12,25,38,51,64,77,90,103,116,129,142}
13	$E_{13} = [0]$	{0,13,26,39,52,65,78,91,104,117,130,143}

The tuple (U, θ) is an approximate space. Furthermore, we give a subset $X = \{8,9,10,14,28,42,56,70,80,94,108,122,136,140,141,142\} \subseteq U$. Then the lower approximations of X is $\underline{Apr}(X) = \emptyset$, and the upper approximations of X is $\overline{Apr}(X) = E_1 \cup E_2 \cup E_3 \cup E_4 \cup E_5 \cup E_6 \cup E_7 \cup E_8 \cup E_9 \cup E_{10} \cup E_{11} \cup E_{12} \cup E_{13} = U$. The rough set $Apr(X)$ is the ordered pair of the lower and upper approximations written as $Apr(X) = (\underline{Apr}(X), \overline{Apr}(X)) = (\{\}, \{0,1,2,3,4,5,6,7,8, \dots, 147,148,149,149\})$. Next, we define the binary operation $+_{150}$ on rough set $Apr(X)$. We will show that X is a rough group.

- (1) For every $a, b \in X$, $(a +_{150} b) \in \overline{Apr}(X)$,
- (2) Association property holds in $\overline{Apr}(X)$;
- (3) There exists $0 \in \overline{Apr}(X)$ such that for every $x \in X$, $x +_{150} 0 = 0 +_{150} x = x$.
- (4) Table 2 shows that every element in X has an inverse in $\overline{Apr}(X)$.

Table 2. Inverse Element on X

No	1	2	3	4	5	6	7	8
$x \in X$	8	9	10	14	28	42	56	70
Inverse of x	142	141	140	136	122	108	94	80

It follows from Table 2 that every element of X has an inverse in $\overline{Apr}(X)$. Hence X is a rough group on U .

If we take $X_1 = \{8,9,10,14,28,31,42,56,70,80,94,108,119,122,136,140,141, 142\}$, and $X_2 = \{8,9,10,14,28,42,44,56,70,80,94,106,108,122, 136,140,141,142\}$. We can see that every element of X_1 has invers in X_1 . The same thing happened to the set X_2 . Next, the sets X_1 and X_2 satisfy all the properties of rough group. Hence, X_1 dan X_2 are rough groups. Furthermore, we will show that $X_1 \cup X_2$ is a rough group in approximation space (U, θ) .

We have $X_1 \cup X_2 = \{8,9,10,14,28,31,42,44,56,70,80,94,106,108,119,122, 136,140, 141,142\}$. This implies $\overline{Apr}(X_1 \cup X_2) = U$. If we take $+_{150}$ as a binary operation in U , we get $(a +_{150} b) \in \overline{Apr}(X_1 \cup X_2)$, for every $a, b \in X_1 \cup X_2$. Furthermore, the associative property holds in $\overline{Apr}(X_1 \cup X_2)$. It has an $e \in \overline{Apr}(X_1 \cup X_2)$ identity element, i.e. $0 \in \overline{Apr}(X_1 \cup X_2)$ so that for each $x \in X_1 \cup X_2$, $x +_{150} 0 = 0 +_{150} x = x$. Then 0 is an element of

identity in $X_1 \cup X_2$. Every element in the $X_1 \cup X_2$ has an inverse in $\overline{Apr}(X_1 \cup X_2) = U$. So, the set $(X_1 \cup X_2)$ is a rough group in approximation space (U, θ) . This shows that the union of two rough groups with the same upper approximation is also a rough group.

Next, we will give the properties of the sub-exact sequence of rough groups.

Proposition 7. Let (U, θ) be an approximation space, and let X_1, X_2 be rough groups of U , such that $\overline{Apr}(X_1) = \overline{Apr}(X_2)$. The triple (U', U, U'') is a rough X_1 -sub-exact sequence if and only if the triple (U', U, U'') is a rough X_2 -sub-exact sequence.

Proof. We assume that the triple (U', U, U'') is a rough X_1 -sub-exact sequence. Based on Definition 4, the following sequence:

$$Apr(U') \xrightarrow{f} Apr(X_1) \xrightarrow{g} Apr(U'')$$

is rough exact. It implies $\text{Im}(f) = \text{Ker}(g)$, where f is a rough homomorphism from $Apr(U')$ to $Apr(X_1)$, and g is a rough homomorphism from $Apr(X_1)$ to $Apr(U'')$. The rough homomorphism f maps every element of $\overline{Apr}(U')$ to $\overline{Apr}(X_1)$, and the rough homomorphism g maps every element of $\overline{Apr}(X_1)$ to $Apr(U'')$. By hypothesis, $\overline{Apr}(X_1) = \overline{Apr}(X_2)$ and hence the following sequence:

$$Apr(U') \xrightarrow{f} Apr(X_2) \xrightarrow{g} Apr(U'')$$

is rough exact. So, the triple (U', U, U'') is a rough X_2 -sub-exact sequence.

To prove the triple (U', U, U'') is a rough X_2 -sub-exact sequence implies the triple (U', U, U'') is a rough X_1 -sub-exact sequence can be shown in a similar way.

CONCLUSIONS

The sub-exact sequence of rough groups is a generalization of the exact sequence of rough groups. Furthermore, the union of finite rough groups which the same upper approximation is also a rough group. If (U, θ) is an approximation space, and X_1, X_2 are rough groups of U such that $\overline{Apr}(X_1) = \overline{Apr}(X_2)$, then the triple (U', U, U'') is an X_1 -sub-exact sequence if and only if the triple (U', U, U'') is an X_2 -sub-exact sequence.

ACKNOWLEDGMENTS

The authors want to thank the reviewers for their insightful suggestions and efforts towards improving our manuscript. Furthermore, the authors wish to thank the Research Institutions and Community Service of Universitas Lampung for this research support and funding under the Research Contract No: 1667/UN 26.21/PN/2021.

AUTHOR CONTRIBUTIONS STATEMENT

NS is the lead investigator in this study. She collected literature related to sub-exacts and group in algebra. FF and AF help in analyzing the development of theory about the group.

REFERENCES

- Aminizadeh, R., Rasouli, H., & Tehranian, A. (2018). [Quasi-exact sequences of s-acts](#). *Bull.Mallays. Math.Sci.Soc.*, 42(5), 2225–2235.
- Bagirmaz, N. (2019). [Direct products of rough subgroups](#). *Eskişehir Technical University Journal of Science and Technology A - Applied Sciences and Engineering*, 20(3), 307–316.

- Bagirmaz, N., & Ozcan, A. F. (2015). Rough semigroups on approximation spaces. *International Journal of Algebra*, 9(7), 339–350.
- Biswas, R., & Nanda, S. (1994). Rough groups and rough subring. *Bull. Polish Acad. Sci. Math.*, 42(3), 251–254.
- Davvaz, B. (2004). Roughness in rings. *Information Sciences*, 164(4), 147–163.
- Davvaz, B., & Mahdavi-pour, M. (2006). Roughness in modules. *Information Sciences*, 176(24), 3658–3674.
- Davvaz, B., & Parnian-Garamaleky, Y. A. (1999). A note on exact sequences. *Bulletin of The Malaysian Mathematical Society*, 22(1), 53–56.
- Elfiyanti, G., D. Nasution, I. M., & Amartiwi, U. (2016). Abelian property of the category of U-complexes chain of U-complexes. *Int. J. Math. Anal.*, 10(17), 849–853.
- Faisol, A., Fitriani, & Sifriyani. (2021). Determining the noetherian property of generalized power series modules by using X-sub-exact sequence. *J. Phys.: Conf. Ser.*, 1751(1), 012028.
- Fitriani, Surodjo, B., & Wijayanti, I. E. (2016). On sub-exact sequences. *Far East Journal of Mathematical Sciences*, 100(7), 1055–1065.
- Fitriani, Surodjo, B., & Wijayanti, I. E. (2017). On X-sub-linearly independent modules. *Journal of Physics: Conference Series*, 893(1), 1–6.
- Fitriani, Wijayanti, I. E., & Surodjo, B. (2018a). A generalization of basis and free modules relatives to a family of R-modules. *Journal of Physics: Conference Series*, 1097(1), 1–6.
- Fitriani, Wijayanti, I. E., & Surodjo, B. (2018b). Generalization of U-generator and M-subgenerator related to category $\sigma[M]$. *Journal of Mathematics Research*, 10(4), 101–106.
- Li, F., & Zhang, Z. (2014). The homomorphisms and operations of rough groups. *The Scientific World Journal*, 2014, 1–6.
- Isaac, P., & Paul, U. (2017). Rough G-modules and their properties. *Advances in Fuzzy Mathematics*, 12(1), 93–100.
- Jesmalar, L. (2017). Homomorphism and isomorphism of rough group. *International Journal of Advance Research, Ideas and Innovations in Technology*, 3(3), 1382–1387.
- Kuroki, N. (1997). Rough ideals in semigroups. *Inform. Sci.*, 100, 139–163.
- Miao, D., Han, S., Li, D., & Sun, L. (2005). Rough group, rough subgroup and their properties. In *Lecture Notes in Artificial Intelligence*, 3641, 104–113.
- Pawlak, Z. (1991). *Rough sets-theoretical aspects of reasoning about data*. Dordrecht, Kluwer.
- Sinha, & Prakash. (2016). Rough exact sequences of modules. *International Journal of Applied Engineering Research*, 11(4), 2513-2517.

● 15% Overall Similarity

Top sources found in the following databases:

- 12% Internet database
- Crossref database
- 8% Submitted Works database
- 15% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	tandfonline.com Internet	4%
2	B H S Utami, Fitriani, M Usman, Warsono, J I Daoud. "Sub-Exact Seque... Crossref	2%
3	Binus University International on 2018-08-06 Submitted works	2%
4	nozdr.ru Internet	2%
5	m-hikari.com Internet	1%
6	repository.lppm.unila.ac.id Internet	1%
7	Jiuping Xu, Rui Qiu, Zhimiao Tao. "Rough Approximation Operators in ... Crossref	<1%
8	Jawaharlal Nehru Technological University on 2014-08-22 Submitted works	<1%

9	yuhuaqian.com	Internet	<1%
10	trademarkelite.com	Internet	<1%
11	Lingyun Yang, Luoshan Xu. "Algebraic aspects of generalized approxi...	Crossref	<1%
12	irep.iium.edu.my	Internet	<1%

● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)
- Manually excluded text blocks

EXCLUDED SOURCES

ejournal.radenintan.ac.id	82%
Internet	
ejournal.radenintan.ac.id	69%
Internet	
pasca.unila.ac.id	43%
Internet	

EXCLUDED TEXT BLOCKS

<http://ejournal.radenintan.ac.id/index.php/al-jabar/index>
news.teknokrat.ac.id