

PAPER NAME

Atmospheric Hg Levels in Tree Barks Due to Artisanal.pdf

AUTHOR

Endang L Widiastuti

WORD COUNT

6575 Words

CHARACTER COUNT

41427 Characters

PAGE COUNT

14 Pages

FILE SIZE

6.9MB

SUBMISSION DATE

Sep 11, 2022 10:40 PM GMT+7

REPORT DATE

Sep 11, 2022 10:41 PM GMT+7**● 9% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 8% Internet database
- 7% Publications database
- Crossref database
- Crossref Posted Content database
- 4% Submitted Works database

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded sources
- Manually excluded text blocks

Article

Atmospheric Hg Levels in Tree Barks Due to Artisanal Small-Scale Gold Mining Activity in Bunut Seberang Village in Indonesia

Azhary Taufiq ¹, Melya Riniarti ^{1,*}, Endang Linirin Widiastuti ², Hendra Prasetya ^{3,4,*}, Slamet Budi Yuwono ¹, Ceng Asmarahman ¹ and Tedy Rendra ⁵

- ¹ Department of Forestry, Faculty of Agriculture, University of Lampung, Jl. Prof. Dr. Ir. Sumantri Brojonegoro, RW.No: 1, Gedong Meneng, Kec. Rajabasa, Kota Bandar Lampung, Lampung 35141, Indonesia; azharytaufiq.at@gmail.com (A.T.); slamet.budi@fp.unila.ac.id (S.B.Y.); ceng.asmarahman@fp.unila.ac.id (C.A.)
- ² Department of Biology, Faculty of Mathematics and Science, University of Lampung, Jl. Prof. Dr. Ir. Sumantri Brojonegoro, RW.No: 1, Gedong Meneng, Kec. Rajabasa, Kota Bandar Lampung, Lampung 35141, Indonesia; endang.linirin@fmipa.unila.ac.id
- ³ National Research and Innovation Agency (BRIN), Gedung B.J. Habibie, Jakarta Pusat 10340, Indonesia
- ⁴ Research Institute for Humanity and Nature, Kyoto 603-8047, Japan
- ⁵ Department of Environmental Science, Faculty of Multidisciplinary, University of Lampung, Jl. Prof. Dr. Ir. Sumantri Brojonegoro, RW.No: 1, Gedong Meneng, Kec. Rajabasa, Kota Bandar Lampung, Lampung 35141, Indonesia; dyrendra@gmail.com
- * Correspondence: ² melya.riniarti@fp.unila.ac.id (M.R.); hendra.prasetya@brin.go.id (H.P.)



Citation: Taufiq, A.; Riniarti, M.; Widiastuti, E.L.; Prasetya, H.; Yuwono, S.B.; Asmarahman, C.; Rendra, T. Atmospheric Hg Levels in Tree Barks Due to Artisanal Small-Scale Gold Mining Activity in Bunut Seberang Village in Indonesia. *Atmosphere* **2022**, *13*, 633. <https://doi.org/10.3390/atmos13040633>

Academic Editors: Young-Ji Han, Chris G. Tzanis and Artur Badyda

Received: 24 January 2022

Accepted: 12 April 2022

Published: 15 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Mercury (Hg) is a useful heavy metal; however, it is toxic to both humans and the environment. Tree bark is an excellent bioindicator, which has been proven to be effective in studying the level of atmospheric Hg contamination. This study aimed to determine the distribution of evaporated Hg using the total weight of Hg (THg) in tree barks in Indonesia at the artisanal and small-scale gold mining (ASGM) area of Bunut Seberang Village and Lampung University, respectively. Samples were taken using purposive sampling, based on the criteria of forestry trees at a height level of 1.3 m above ground as wide as 100 cm². The samples were analyzed by atomic absorption spectrometry and Scanning electron microscopy to determine the levels of THg and to investigate the bark structures. Results showed that the highest THg values were found in a *Magnolia champaca* sample (56.5 µg), followed by *Swietenia mahagoni* (45.8 µg) and *Swietenia mahagoni* (33.5 µg). All species studied showed THg levels in the tree barks at an elevation from 30 to 320 m above sea level. The Hg amounts found in the sampled barks indicated the dispersion of Hg throughout the ASGM area, which signified hazardous atmospheric conditions in the area.

Keywords: ASGM; atmospheric Hg; tree bark; forestry tree; atmospheric pollution

1. Introduction

Mercury (Hg) is a toxic heavy metal, which is well known for its harmful effects in the environment. The European Union Commission and World Health Organization have stated that Hg even at low concentrations could affect human health and has become a global environmental concern [1]. Reduction or elimination of Hg use [2]. Hg transport and disposal present serious risks to the environment [3]. In artisanal and small-scale gold mining (ASGM), Hg has been used to extract gold (Au) from an ore to form a stable Hg–Au amalgam. Then, the amalgam is heated to evaporate the Hg and isolate the Au. Hg amalgamation has been used for thousands of years to mine Au and Ag in this way [4].

An uncontrolled use of Hg in gold mining occurs due to low technical knowledge [5]. In Indonesia, ASGM has been the second most significant source of Hg pollution after coal burning [6]. Recognizing the problems associated with the use of Hg in mining, over 130 countries have signed the United Nation's Minamata Convention as an agreement to

5 significantly reduce the emission and usage of Hg [7]. During an entire ore amalgamation process, 20 g of Hg is typically used to produce 1 g of Au, where 19 and 1 g of the Hg pass or are released to the tailings and atmosphere, respectively [3]. Globally, Hg amalgamation activities in mining could lead to the annual release of up to 1000 tons of Hg to the environment. Specifically, Hg release from Indonesia has been estimated between 100 and 150 tons per year [8].


Various studies have focused on the determination of Hg concentrations in various bioindicators including tree-like foliage [9], leaf litter [10], tree bark, and roots [11]. In particular, tree barks have been used to assess the state of Hg contamination in the environment [12]. In one study, higher Hg was found in tree foliages and barks than in wood samples [13]. Foliage materials typically contain high Hg concentrations because foliage absorbs elemental Hg from the atmosphere through its stomata [14]. Although bark and wood are less exposed to atmospheric Hg than foliage, bark can capture and retain atmospheric Hg through surface sorption [13]. The enrichment of trace elements in a tree bark can be used to trace pollution sources [15]. This is because airborne particles are often trapped within the structure of a tree bark, where these particles accumulate over a period of several years [16].

6 This study aimed to determine the atmospheric pool of Hg in an ASGM area at Bunut Seberang Village, Lampung, Indonesia, using forestry tree barks as bioindicators.

2. Materials and Methods


2.1. Study Location

The test and control samples were collected in December 2020 in an ASGM area of Bunut Seberang Village, Pesawaran District, and in January 2021 in the University of Lampung, Lampung Province, respectively, as shown in Figure 1. The control samples were selected to compare the research main samples. The distance of main test samples obtained in the ASGM area ranged from 0 to 1.2 km. For the control samples, the location was 32 km from the ASGM location. Six forestry trees were found in the ASGM areas and three forestry trees were found in control areas, respectively. The trees were selected due to their rough bark characteristics, which could retain atmospheric pollutants such as Hg [13].



(a)

1 Figure 1. Cont.



(b)

Figure 1. The research sampling locations were (a) Bunut Seberang Village, Lampung Province, Pesawaran District, Indonesia (test samples), and (b) University of Lampung, Lampung Province, Bandar Lampung, Indonesia (control samples).

2.2. Sampling Methods

Evaporated elemental Hg has been mostly found in trees, which are located around 1.3 m from the ground [12]. The dimensions of tree bark samples obtained were 10 cm × 10 cm. The samples were taken at a height of 1.3 m to ensure homogeneity [13]. The collection of tree barks was conducted by selecting features that faced gold refining locations at the ASGM site. The trees selected for sampling have a minimum diameter of 20 cm. Bark samples were collected by purposive sampling based on rough stem morphology [17]. Based on the sampling used, there were six tree species (*C. pentandra* ($n = 1$), *M. champaca* ($n = 1$), *P. falcataria* ($n = 2$), *P. acerifolium* ($n = 2$), *S. mahagoni* ($n = 7$), and *T. grandis* ($n = 5$)), all with rough bark characteristics as shown in Figure 2. The total number of samples collected was 18 (n (total) = 18). The locations of the sampled trees were selected as shown in Figure 2. The collected rough barks were in form of hard chipped, broken scaled, and grooved barks. Trees that have such rough bark characteristics include *T. grandis*, *Swietenia* sp., and *Schima wallichii* [18]. The collected samples were kept in coded plastic sample bags.

The sample distances in the ASGM location were measured by conducting field observations at the location. The elevation data were obtained using the secondary data from the Digital Elevation Model and Rupa Bumi Indonesia maps of Lampung Province for 2017 with a scale of 1:10,000. The wind direction data for 2010–2020 were obtained from four stations of the Meteorology, Climatology, and Geophysical Agency, which are located in Bandar Lampung, Pesawaran, South Lampung, and North Lampung, Lampung, Indonesia.




Figure 2. The locations of the sampled trees at Bunut Seberang Village including *C. pentandra* ($n = 1$), *M. champaca* ($n = 1$), *P. falcata* ($n = 2$), *P. acerifolium* ($n = 2$), and *S. mahagoni* ($n = 7$) and at Bunut Seberang Village such as *T. grandis* ($n = 5$) (n (total sampled trees) = 18).

2.3. Atomic Absorption Spectrometry (AAS) and Scanning Electron Microscopy (SEM)

The mercury concentrations were determined by Atomic Absorption Spectrometry (AAS). The barks were cut to obtain the Fragment Dimension (FD), approximately 100 cm^2 ($10 \text{ cm} \times 10 \text{ cm}$) from the tree. Samples were dried in an oven at $300 \text{ }^\circ\text{C}$ for 48 h. A dried sample was weighed and the weight was counted as the dry weight (DW) (g). Each sample was analyzed by Image J from Rasband, W.S., ImageJ, U. S. National Institutes of Health, Bethesda (Maryland, USA) and cut as wide as 14 to 18 cm^2 to measure the Real Square (RS). Each sample was mashed using a Philips Plastic HR2115/00 mixer by Phillips Inc. (Eindhoven, Netherlands) speed of 13,000 rpm.

The AAS analyses were conducted using the AAS method described in the Indonesian National Standard 01-2896-1998. A 5 g sample was placed in a plastic tube, then dissolved in 25 mL H_2SO_4 18N, 20 mL HNO_3 7N, and 1 mL 2% of Na_2MoO_4 . The samples then were sent to PT Superintending Company of Indonesia (PT SUCOFINDO) for AAS analysis using an Agilent 240FS-VGA 7 from Agilent Technologies Inc. (Mulgrave, Victoria, Australia). AAS analysis was conducted using the AAS method described in the Indonesian National Standard 01-2896-1998. Hg results were reported as concentrated Hg (CHg) before conversion to total weight Hg (THg).

Scanning Electron Microscopy (SEM) was used to inspect and identify the anatomical features of the tree bark samples [19]. The SEM analysis uses ZEISS/EVO MA 10, The Carl Zeiss Foundation, Munich, Germany. SEM uses a procedure involving electrons to magnify the image of a sample, which allowed the inspection of the tree bark samples in detail [20].

The accumulated Hg was measured as THg (µg), which was calculated by Equation (1) [12,21,22].

$$THg = (DW \times CHg) \times \frac{(FD)}{(RS)} \tag{1}$$

where the THg is Total Weight of Mercury, CHg is Concentrated Hg, DW is Dry Weight, FD is Fragment Dimension, RS is Real Square.

2.4. Statistical Analysis

Statistical analyses were performed using an IBM SPSS Version 26 SPSS Inc. (Chicago, IL, USA). To determine whether the THg, CHg, bark thickness, elevation, and distance values were normally distributed, a Shapiro–Wilk measurement at the 5% level was used followed by a one-way analysis of variance at the 5% level of significance [21]. A bivariate correlation analysis was used to determine the correlation between bark thickness, elevation, distance, and THg.

3. Results

THg Levels in the Tree Bark Samples Based on Distance and Elevation

The THg levels found in the Bunut Seberang Village are presented in Table 1.

Table 1. THg Levels Found in Bunut Seberang Village.

No.	Name	CHg (µg/g)	THg (µg)	Bark Thickness (mm)	Elevation (m asl *)	Distance to Nearest ASGM (m)
1	<i>C. pentandra</i> 1	0.33	11.7	9.64	80	247
	Mean	0.33	11.7	9.64	80	247
2	<i>M. champaca</i> 1	1.01	56.5	10.4	30	47
	Mean	1.01	56.5	10.4	30	47
3	<i>P. falcataria</i> 1	0.34	2.19	5.66	310	580
4	<i>P. falcataria</i> 2	0.20	4.24	19.5	240	619
	Mean	0.27	3.21	12.6	275	600
5	<i>P. acerifolium</i> 1	0.37	23.2	9.74	200	223
6	<i>P. acerifolium</i> 2	0.37	19.2	10.5	320	604
	Mean	0.37	21.2	10.1	260	414
7	<i>S. mahagoni</i> 1	0.09	5.63	0.60	50	1215
8	<i>S. mahagoni</i> 2	0.68	32.2	0.76	80	894
9	<i>S. mahagoni</i> 3	0.14	5.97	0.56	110	982
10	<i>S. mahagoni</i> 4	0.35	17.9	0.50	50	1069
11	<i>S. mahagoni</i> 5	0.78	45.8	0.62	90	723
12	<i>S. mahagoni</i> 6	0.53	33.5	0.80	60	233
13	<i>S. mahagoni</i> 7	0.10	2.21	0.60	120	1250
	Mean	0.38	20.5	0.63	80	909
14	<i>T. grandis</i> 1	0.20	6.66	0.48	30	985
15	<i>T. grandis</i> 2	0.30	11.6	0.30	50	1111
16	<i>T. grandis</i> 3	0.17	4.41	0.49	120	186
17	<i>T. grandis</i> 4	0.48	17.7	0.48	80	607
18	<i>T. grandis</i> 5	0.42	3.50	0.17	150	713
	Mean	0.31	8.77	0.38	86	720

* asl, above sea level.

The results in Table 1 showed that all of the sampled forestry trees contained Hg. The highest THg amount was found in *M. champaca* (56.5 µg), which was followed by *S. mahagoni*, *P. acerifolium*, and *C. pentandra* (2.21–45.8 µg). Various THg amounts were found in *T. grandis* from 3.50 to 17.7 µg.

The bark thickness did not show any significant correlation to the THg. The thickest bark sample was from *P. falcataria* 2 (bark thickness = 19.5 mm), which contained a lower THg of 4.24 µg than the thin bark sample from *S. mahagoni* 5 (bark thickness = 0.62 mm) with

a measured THg of 45.8 µg. Another example was *C. pentandra* (bark thickness = 9.64 mm), which contained a lower THg of 11.7 µg than *S. mahagoni* 6 (bark thickness = 0.80 mm) with a measured THg of 33.5 µg.

Meanwhile, THg was affected by CHg found in the samples. *Magnolia champaca* gave the highest value for CHg of 1.01 µg/g. It also contained the highest amount Hg with a THg of 56.5 µg. The *M. champaca* and *S. mahagoni* 7 samples both gave the second largest CHg value of 0.78 µg/g. They also contained the second largest THg of 45.8 µg.

Based on the results, THg values found in the studied areas were not affected by elevation. Similar values for THg were generally found at any elevation level. *P. falcataria* 1 (THg = 5.66 µg) that was obtained at an elevation level of 310 m asl showed a similar THg with *T. grandis* 5 and *T. grandis* 1 (THg = 3.50 and 6.66 µg, respectively), which were obtained at an elevation level of 150 and 30 m asl, respectively. *C. pentandra* 1 which was sampled at an elevation level of 80 m asl contained a THg of 11.7 µg-DW. This was lower than the THg found in *S. mahagoni* 2 (32.2 µg), which was sampled at the same elevation level of 80 m asl.

The THg values found were also not affected by distance. *M. champaca* was the closest sample location (47 m from the ASGM location) providing the highest THg value (56.5 µg) compared to the other samples. In a further location, a high THg value (45.8 µg) was also found in *S. mahagoni* 5, which was sampled 723 m away from the ASGM location. These numbers showed the variation in atmospheric Hg, which was not dependent on distance.

The results from control samples are summarized in Table 2.

Table 2. THg values found in the control location.

No.	Name	CHg (µg/g)	THg (µg)	Bark Thickness (mm)
1	<i>M. champaca</i> 1	<0.0002	<0.02	3.20
2	<i>M. champaca</i> 2	<0.0002	<0.02	3.10
3	<i>M. champaca</i> 3	<0.0002	<0.02	2.73
4	<i>A. cadamba</i> 1	<0.0002	<0.02	4.38
5	<i>A. cadamba</i> 2	<0.0002	<0.02	4.69
6	<i>A. cadamba</i> 3	<0.0002	<0.02	4.05
7	<i>S. mahagoni</i> 1	<0.0002	<0.02	10.60
8	<i>S. mahagoni</i> 2	<0.0002	<0.02	8.89
9	<i>S. mahagoni</i> 3	<0.0002	<0.02	5.45
10	<i>S. mahagoni</i> 4	0.0120	0.26	12.40
11	<i>S. mahagoni</i> 5	<0.0002	<0.02	6.70

The control samples showed THg values of <1 µg. Plants have been classified as hyperaccumulators of Hg and toxic if the Hg concentration in a plant exceeds 10 [23] and 1 ppm [24], respectively. Therefore, the trees in the control area were not toxic.

Wind direction, elevation, and bark type have been found to determine the levels of heavy metal contamination in tree barks [25]. This was because Hg spreads in the atmosphere due to wind [26]. Wind also causes the dispersion of Hg over long distances [27].

As seen in Table 1, all the sampled trees in the ASGM area contained and were above the Hg detection limit (0.02 µg); therefore, the air near this area was most likely contaminated by Hg. Hence, *M. champaca*, *S. mahagoni*, *C. pentandra*, *P. acerifolium*, *P. falcataria*, and *T. grandis* samples in Table 1 were used to determine the atmospheric Hg contamination in the studied area. To determine the distribution rate of THg in the ASGM study area, the distance and elevation of each collected bark were defined. These are summarized in Table 2.

4. Discussion

4.1. THg in the Tree Bark Samples

Barks contain Hg (Table 1) and main organic matters of carbon and oxygen. Hg found in a bark sample was either from within or between the atmosphere in ASGM area and

1

stored in the tree bark, especially in the external layer of the barks [28]. The amount of Hg stored in tree bark is often defined by the factors of bark roughness and sample location [29]. The differences in Hg concentrations between tree species in the studied location could be explained by the variations in wind flow during 2010–2020 as shown in Figure 3 [30] and different rates of absorption in the six tree species considered [13].




Figure 3. Atmospheric Hg distribution in the Bunut Seberang Village between 2010 and 2020.

THg contamination in tree barks did not exhibit significant variation in this study, with the exception of *M. champaca* 1, *S. mahagoni* 2, *S. mahagoni* 5, and *S. mahagoni* 6 with THg values of >20 µg. The surface magnification of the samples is shown in Figure 4. The samples of *M. champaca* and *S. mahagoni* showed textures of clustered and fibrous-grouped, respectively. These samples have a rough-textured bark. The THg found in *P. acerifolium* was low, because its bark was less rough especially when compared with *M. champaca* [31] (see Figure 4). The SEM results showed that the Hg levels found in *M. champaca* ranged from 19.6 to 25.1%. The level was 19.6% in *S. mahagoni*. After considering the SEM data, the low roughness of a tree bark seemed to have minimized the possibility of accumulation of atmospheric Hg. In fact, rougher tree barks have been known to be better accumulators of heavy metals [32].




Figure 4. SEM micrographs of *M. champaca* at magnifications of (a) 1000, (b) 3000, and (c) 5000 \times . SEM micrographs of *S. mahagoni* at magnifications of (d) 1000, (e) 3000, and (f) 5000 \times . The colored box (□: □, □, □, or □) in a panel depicts the analyzed area.

We found that the tree bark thickness did not define the THg level found in bark (see Figure 5). The bark thickness only affected the THg value by 0.6%. The bark thickness varied from 0.17 to 19.5 mm as shown in Figure 5. The differences in the bark thickness can vary in shape and thickness, depending on the species and ecological factors [33]. In a study with bark thickness values from 0 to 0.6 mm of samples, the heavy metal adsorption capacity was found to be higher in the outer bark than in the inner bark [34]. This was because the uptake of heavy metals by the tree inner bark (via phloem) was lower than the outer bark (via rhytidome).




Figure 5. (a) Bark thickness (mm) versus THg found, (b) bark thickness versus THg found (control), (c) distance from ASGM (m) versus THg found, (d) distance from ASGM (m) versus THg found (control), (e) elevation versus THg found, and (f) elevation versus THg found (control).

4.2. Elevation, Distance, and Bark Thickness

A bark accumulates atmospheric Hg as it is continually exposed to air pollution for many years; therefore, it could provide precise information about the changes in the atmospheric condition of a certain ecosystem [35]. A study of the chemical constituents in barks is an appropriate method for tracing pollution sources [12,13,15,21,22]. The distances shown in Figure 4 showed a negative trend toward THg levels in tree barks. All tree barks contained Hg in both areas near and far from the ASGM location studied. The bark has been found to have the highest capacity for heavy metal absorption, followed by cones, needles, and wood [33]. Tolerant tree species have been used as an indicator for the accumulation of air pollutants [36]. Hg concentrations have been determined by the analyses of tree tissues including bark [15].

Hg spreads in the atmosphere [26] and disperses over long distances due to wind [27]. Tree barks have been known to retain and capture atmospheric Hg through surface sorption [13] even over long distances. This was observed in our work. A high THg value of 45.8 μg was found for the *S. mahagoni* 5, which was sampled at a long distance of 723 m from the ASGM location (see Figure 5).

Heavy metal adsorption on biomass such as bark is a physiochemical process [35]. In the case of a bark, heavy metals typically accumulate through the wet and dry deposition processes [37]. Briefly, a tree bark absorbs gaseous Hg at its surface. The absorbed Hg is then bound to thick containing molecules such as tannins [38]. The wet and dry depositions of Hg have been defined as the air-to-surface flux in the presence and absence of precipitation, respectively [39]. For these reasons, tree bark can be used as a heavy metal bioindicator where their outer portion consists of dead materials, which do not interfere with their growth cycles [35].

All the tested samples contained atmospheric Hg. THg values were observed in the samples obtained from 0 to 320 m asl. A previous study found that atmospheric Hg increased with altitude [40]. Conversely, our measurements indicated that atmospheric Hg did not increase with altitude as shown in Figures 4 and 5. The THg values were rather grouped and dispersed throughout all the altitudes studied. Additionally, shown in Figure 5, the group with an elevation from 0 to 100 m asl provided a maximum value of 56.5 μg , with median and minimum values of 17.8 and 5.63 μg , respectively. For the group with an elevation from 100 to 320 m asl, the maximum value was 23.2 μg , with median and minimum values of 4.32–2.19 μg , respectively.

P. falcataria 2 from an elevation of 240 m asl showed a lower THg than *S. mahagoni* 6 from an elevation of 60 m asl. In addition, the sources of Hg in the ASGM site were dispersed due to the wind factor. The dispersion of windblown atmospheric Hg in the 2010–2020 study period occurred from north to south. Therefore, wind largely controlled the movement of atmospheric Hg in our research sites [41] as shown in Figure 6.

4.3. Wind Direction

The major transport of evaporated Hg comes from various sources in the atmosphere [42] and occurs in the gaseous phase [43]. The movement of heavy metals depends on numerous factors including temperature, surface waters, air masses, and wind [41]; therefore, they are transported over long distances even at global scales [44]. Wind in the study area was recorded from 2010 to 2020. Wind was found to blow mostly from north to south as shown in Figure 6. Wind movement that carried Hg was also affected by a mountainous area (e.g., *P. falcataria* 1 and *P. falcataria* were taken at the elevation level of 310 and 240 m asl, respectively), which diverted the wind to a lower elevation. A previous study that has coastal and hilly areas stated that topography significantly influences the Hg [12]. Nevertheless, in this study that has mountainous areas did not significantly affect the Hg.




Figure 6. (a) Distance, (b) thickness, and (c) elevation. ■ = 2q box, □ = 3q box.

The mountainous area in this study was located in the south of the ASGM area. The presence of elevation and trees have been known to cause airflow fluctuations and lower wind speeds [45]. These affect the total Hg contamination. It has been found that wind influences the distribution of heavy metals in some areas brought by pollutant sources [46]. *M. champaca* provided the highest Hg contamination (56.5 µg) at a distance of 47 m from the ASGM location. *S. mahagoni* 6, which was the sample closest to the ASGM area (distance of 233 m), also exhibited an elevated THg (33.5 µg). *S. mahagoni* 2 (32.2 µg) and *S. mahagoni* 5 (45.8 µg) exhibited high Hg contamination, although their distances (894 and 723 m, respectively) from the studied area were greater than the other samples.

5. Conclusions

In this study, about 18 samples, with 6 different species, were found during the field sampling. The *M. champaca* was indicated as the highest THg (56.5 µg) among other samples then followed by *C. pentandra* 11.7 µg, *P. falcataria* 1 ranged from 2.19 µg to 4.24 µg, *P. acerifolium* ranged from 19.2 µg to 23.2 µg, *T. grandis* ranged from 3.50 µg to 17.7 µg, and *S. mahagoni* ranged from 2.21 µg to 45.8 µg. The highest elevation on this study is at 320 m asl (*P. acerifolium* 2) and the lowest elevation is at 30 m asl (*M. champaca* and *T. grandis* 1). The furthest sample location is *S. mahagoni* 7 at 1250 m, and the closest sample is *M. champaca* at 47 m from ASGM site. This study showed that the wind direction and the topography (closed mountain area) is the important factor to attach evaporated mercury in the tree bark. Meanwhile, the distance, elevation, and bark thickness showed no significant effect on the attachment of evaporated mercury.

Author Contributions: A.T. was the lead author who is still doing a postgraduate program at Lampung University. M.R., H.P. and E.L.W. were the supervisors. Furthermore, S.B.Y. and C.A. were advisers. T.R. was the assistant that helped the lead author finish this research. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by The Research Institute for Humanity and Nature (RIHN: a constituent member of NIHU), Project No.14200102.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: Azhary Taufiq, one of the authors of this study, would like to thank the University of Lampung for providing the master's degree scholarships and RIHN Project No.14200102 for funding this research, and the Pesawaran Regency's Forestry Department.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. WHO (World Health Organisation). Exposure to Mercury: A Major Public Health Concern. 2009. Available online: <http://www.who.int/phe/news/Mercury-flyer.pdf> (accessed on 4 January 2021).
2. Holmes, P.; James, K.A.F.; Levy, L.S. Is low level environmental mercury exposure of concern to human health? *Sci. Total Environ.* **2004**, *408*, 171–182. [CrossRef] [PubMed]
3. Telmer, K.H.; Stapper, D. *Evaluating and Monitoring Small Scale Gold Mining and Mercury Use: Building a Knowledge-Base with Satellite Imagery and Fieldwork*. UNDP/GEF/UNIDO Project EG/GLO/01/G34; Final Report to the United Nations Industrial Development Organization; UNIDO: Vienna, Austria, 2007.
4. De Lacerda, L.D.; Salomons, W. *Mercury from Gold and Silver Mining a Chemical Time Bomb?* Springer: Berlin, Germany, 1998; pp. 1–151.
5. Green, C.S.; Lewis, P.J.; Wozniak, J.L.; Drevnick, P.E.; Thies, M.L. A comparison of factors affecting the small-scale distribution of mercury from artisanal small-scale gold mining in a Zimbabwean stream system. *Sci. Total Environ.* **2019**, *647*, 400–410. [CrossRef] [PubMed]
6. Limbong, D.; Kumpampung, J.; Rimper, J.; Arai, T.; Miyazaki, N. Emissions and environmental implications of mercury from artisanal gold mining in North Sulawesi, Indonesia. *Sci. Total Environ.* **2020**, *302*, 227–236. [CrossRef]
7. Xu, J.; Garcia, A.B.; Lagerkvist, A.; Bertilsson, S.; Sjöblom, R.; Kumpiene, J. Sources and remediation techniques for mercury contaminated soil. *Environ. Int.* **2015**, *74*, 42–53. [CrossRef] [PubMed]
8. Veiga, M.M.; Maxson, P.A.; Hylander, L.D. Origin and consumption of mercury in small-scale gold mining. *J. Clean. Prod.* **2006**, *14*, 436–447. [CrossRef]
9. Siwik, E.L.; Campbell, L.M.; Mierle, G. Fine-scale mercury trends in temperate deciduous tree leaves from Ontario, Canada. *Sci. Total Environ.* **2009**, *407*, 6275–6279. [CrossRef]
10. Blackwell, B.D.; Driscoll, C.T.; Maxwell, J.A.; Holsen, T.M. Changing climate alters inputs and pathways of mercury deposition to forested ecosystems. *Biogeosciences* **2014**, *119*, 215–228. [CrossRef]
11. Gueguen, F.; Stille, P.; Geagea, M.L.; Boutin, R. Atmospheric pollution in an urban environment by tree bark biomonitoring-Part I: Trace element analysis. *Chemosphere* **2012**, *86*, 1013–1019. [CrossRef]
12. Prasetya, H.; Sakakibara, M.; Omori, K.; Laird, J.S.; Sera, K.; Idham, A.K. *Mangifera indica* as Bioindicator of Mercury Atmospheric Contamination in an ASGM Area in North Gorontalo Regency, Indonesia. *Geosciences* **2018**, *8*, 31. [CrossRef]

13. Yang, Y.; Yanai, R.D.; Driscoll, C.T.; Montesdeoca, M.; Smith, K.T. Concentrations and content of mercury in bark, wood, and leaves in hardwoods and conifers in four forested sites in the Northeastern USA. *PLoS ONE* **2018**, *13*, e0196293. [[CrossRef](#)]
14. Fleck, J.A.; Grigal, D.F.; Nater, E.A. Mercury uptake by trees: An observational experiment. *Water Air Soil Pollut.* **1999**, *115*, 513–523. [[CrossRef](#)]
15. Lahd Geagea, M.L.; Stiller, P.; Millet, M.; Perrone, T. Tree characteristics and Pb, Sr and Nd isotopic compositions of steel plant emissions. *Sci. Total Environ.* **2007**, *373*, 404–419. [[CrossRef](#)] [[PubMed](#)]
16. Catinon, M.; Ayrault, S.; Clocchiatti, R.; Boudouma, O.; Asta, J.; Tissut, M.; Ravel, P. The anthropogenic atmospheric elements fraction: A new interpretation of elemental deposits on tree barks. *Atmos. Environ.* **2009**, *43*, 1124–1130. [[CrossRef](#)]
17. Ismail, A.Y.; Hendrayana, Y.; Saputra, R.H. Inventarisasi dan identifikasi sebaran hutan rakyat di Kabupaten Majalengka. *Wanaraksa* **2016**, *10*, 31–40.
18. Indriyanto. *Dendrologi: Suatu Teori & Praktik Menyidik Pohon*; Plantaxia: Bandar Lampung, Indonesia, 2012; pp. 1–124.
19. Sawidis, T.; Breuste, J.; Mitrovic, M.; Pavlovic, P.; Tsigaridas, K. Trees as bioindicator of heavy metal pollution in three European cities. *Environ. Poll.* **2011**, *12*, 3560–3570. [[CrossRef](#)]
20. Ji, Z. Use of compositional and combinatorial nanomaterial libraries for biological studies. *Sci. Bull.* **2016**, *10*, 755–771. [[CrossRef](#)]
21. Prasetya, H.; Sakakibara, M.; Sera, K.; Laird, S.T. Evaluation of the total mercury weight exposure distribution using tree bark analysis in an artisanal and small-scale gold mining area, North Gorontalo Regency, Gorontalo Province, Indonesia. *Int. J. Environ. Res. Public Health* **2022**, *19*, 33. [[CrossRef](#)]
22. Rendra, T.; Riniarti, M.; Yuwono, S.B.; Prasetya, H.; Widiastuti, E.L.; Bakri, S.; Taufiq, A. Mapping atmospheric mercury in Lampung Province, Indonesia using bark of multipurpose tree species. *Atmosphere* **2022**, *13*, 2. [[CrossRef](#)]
23. Kosegeran, A.O.; Rondonuwu, S.; Simbala, H.; Rumondor, M. Kandungan merkuri pada tumbuhan paku (*Diplazium accedens* Blume) di daerah tambang emas Tatelu-Talawaan, Kabupaten Minahasa Utara. *J. Ilm. Sains* **2015**, *15*, 59–65. [[CrossRef](#)]
24. Kabata-Pendias, A.; Pendias, H. *Soils and Plants*, 3rd ed.; CRC Press: New York, NY, USA, 2001; pp. 1–403.
25. Odukoya, O.O.; Arowolo, T.T.; Bamgbose, O. Pb, Zn and Cu levels in tree barks as indicator of atmospheric pollution. *Environ. Int.* **2000**, *26*, 11–16. [[CrossRef](#)]
26. Javarabad, D.M.; Azadfar, D.; Arzanesh, H.M. The ability to filter heavy metals of lead, copper and zinc in some species of tree and shrub. *Int. J. Adv. Biol. Biomed. Res.* **2013**, *1*, 53–60.
27. Griggs, T.; Liu, L.; Talbot, R.W.; Torres, A.; Lan, X. Comparison of atmospheric mercury speciation at a coastal and an urban site in Southeastern Texas, USA. *Atmosphere* **2020**, *11*, 73. [[CrossRef](#)]
28. Świsłowski, P.; Kříž, J.; Rajfur, M. The Use of Bark in Biomonitoring Heavy Metal Pollution of Forest Areas on the Example of Selected Areas in Poland. *Ecol. Chem. Eng. S* **2020**, *27*, 195–210. [[CrossRef](#)]
29. Tye, A.M.; Hodgkinson, E.S.; Rawlins, B.G. Microscopic and chemical studies of metal particulates in attic dust: Evidence for historical atmospheric smelter emission, Humberside, UK. *J. Environ. Monit.* **2006**, *8*, 904–912. [[CrossRef](#)]
30. Han, Y.J.; Kim, P.R.; Lee, G.S.; Lee, J.I.; Noh, S.; Yu, S.M.; Park, K.S.; Seok, K.S.; Kim, H.; Kim, Y.H. Mercury concentrations in environmental media at a hazardous solid waste landfill site and mercury emissions from the site. *Environ. Earth Sci.* **2017**, *76*, 361. [[CrossRef](#)]
31. Krisnawati, H.; Kallio, M.; Kanninen, M. *Anthocephalus cadamba* Miq. *Ekologi, Silviculture dan Produktivitas*; CIFOR: Bogor, Indonesia, 2011; pp. 1–22.
32. Barnes, D.; Hammadah, M.A.; Ottaway, J.M. The lead, copper and zinc contents of tree rings and barks, a measurement of local pollution. *Sci. Total Environ.* **1976**, *5*, 63–67. [[CrossRef](#)]
33. Pásztor, Z.; Mohácsiné, I.; Gorbacheva, G.; Börcsök, Z. The Utilization of Tree Bark. *BioResources* **2016**, *11*, 7859–7888. [[CrossRef](#)]
34. Aoyama, M.; Kishino, M.; Jo, T.S. Biosorption of Cr (VI) on Japanese cedar bark. *Sep. Sci. Technol.* **2004**, *39*, 1149–1162. [[CrossRef](#)]
35. Patel, K.S.; Yadav, A.; Sahu, Y.K.; Lata, L.; Milosh, H.; Corns, W.T.; Martin-Ramos, P. Tree bark as a bioindicator for arsenic and heavy metal air pollution in Rajnandgaon District, Chhattisgarh, India. *Hazard Toxic Radioact. Waste* **2020**, *24*, 1–5. [[CrossRef](#)]
36. Nursal, F.B. Akumulasi timbal (Pb) pada Talus Lichenes di Kota Pekanbaru. *Biogenesis* **2005**, *1*, 47–50.
37. Tovar-Sanchez, E.; Surez-Rodrigues, R.; Ramires-Trujillo, A.; Valencia-Cavas, L.; Hernandez-Plata, I.; Mussali-Galante, P. The Use of Biosensors for Biomonitoring Environmental Metal Pollution. In *Biosensors for Environmental Monitoring*; Rinken, T., Kivirand, K., Eds.; IntechOpen: London, UK, 2019. [[CrossRef](#)]
38. Chiarantini, L.; Rimondi, V.; Bardelli, F.; Benvenuti, M.; Cosio, C.; Costagliola, P. Mercury speciation in *Pinus nigra* barks from monte Amiata (Italy): An X-ray absorption spectroscopy study. *Environ. Pollut.* **2017**, *227*, 83–88. [[CrossRef](#)] [[PubMed](#)]
39. Zhang, L.; Lyman, S.; Mao, H.; Lin, C.; Gay, D.A.; Wang, S.; Gustin, M.S.; Feng, X.; Wania, F. A synthesis of research needs for improving the understanding of atmospheric mercury cycling. *Atmos. Chem. Phys.* **2017**, *17*, 9133–9144. [[CrossRef](#)]
40. Bernhoft, A.R. Mercury Toxicity and Treatment: A Review of The Literature. *Environ. Public Health* **2011**, *2012*, 460508. [[CrossRef](#)] [[PubMed](#)]
41. Briffa, J.; Sinagra, E.; Blundell, R. Heavy metal pollution in the environment and their toxicological effects on humans. *Heliyon* **2020**, *6*, e04691. [[CrossRef](#)]
42. Sprovieri, F.; Pirrone, N.; Ebinghaus, R.; Kock, H.; Dommergue, A. A review of worldwide atmospheric mercury measurements. *Atmos. Chem. Phys.* **2010**, *10*, 8245–8265. [[CrossRef](#)]
43. Siddiqi, Z.M. Transport and Fate of Mercury (Hg) in the Environment: Need for Continuous Monitoring. In *Handbook of Environmental Materials Management*; Hussain, C., Ed.; Springer: Cham, Switzerland, 2018; pp. 1–20. [[CrossRef](#)]

44. Schuster, P.F.; Krabbenhoft, D.P.; Naftz, D.L.; Cecil, L.D.; Olson, M.L.; Dewild, J.F. Atmospheric mercury deposition during the last 270 years: A glacial ice core record of natural and anthropogenic sources. *Environ. Sci. Technol.* **2002**, *36*, 2303–2310. [[CrossRef](#)]
45. Zhang, M.; Bae, W.; Kim, J. The effects of the layouts of vegetation and wind flow in an apartment housing complex to mitigate outdoor microclimate air temperature. *Sustainability* **2019**, *11*, 3081. [[CrossRef](#)]
46. Boylan, H.M.; Cain, R.D.; Kingston, H.M. A new method to assess mercury emissions: A study of three coal-fired electric-generating power station configurations. *Air Waste Manag. Assoc.* **2003**, *53*, 1318–1325. [[CrossRef](#)]

● 9% Overall Similarity

Top sources found in the following databases:

- 8% Internet database
- Crossref database
- 4% Submitted Works database
- 7% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	mdpi-res.com Internet	2%
2	repository.lppm.unila.ac.id Internet	1%
3	Satomi Kimijima, Masayuki Sakakibara, Masahiko Nagai. "Investigation... Crossref	1%
4	businessdocbox.com Internet	<1%
5	link.springer.com Internet	<1%
6	mdpi.com Internet	<1%
7	Khageshwar Singh Patel, Ankit Yadav, Yaman Kumar Sahu, Lesia Lata, ... Crossref	<1%
8	Louisa J. Esdaile, Justin M. Chalker. "The Mercury Problem in Artisanal... Crossref	<1%

- 9 **ncbi.nlm.nih.gov** <1%
Internet
-
- 10 **Hamid Shirkhanloo, Mohammad Ali Fallah Mehrjerdi, Hamid Hassani. "I...** <1%
Crossref
-
- 11 **hdl.handle.net** <1%
Internet
-
- 12 **jpoll.ut.ac.ir** <1%
Internet
-
- 13 **Mark D. Loewen, Subodh Sharma, Gregg Tomy, Feiyue Wang, Paul Bull...** <1%
Crossref
-
- 14 **Muhammad Yousaf, Khakhathi L. Mandiwana, Khurram S. Baig, Julia L...** <1%
Crossref
-
- 15 **Ya-Chen Shih, Chen-Yi Ke, Cheng-Ju Yu, Chi-Yu Lu, Wei-Lung Tseng. " ...** <1%
Crossref
-
- 16 **Shuo Zhang, Lina Zhou, Lexin Jia, Jinsheng Li, Biying Liu, Yueming Yua...** <1%
Crossref

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded text blocks
- Manually excluded sources

EXCLUDED SOURCES

Azhary Taufiq, Melya Riniarti, Endang Linirin Widiastuti, Hendra Prasetya, Slam... 62%

Crossref

EXCLUDED TEXT BLOCKS

atmosphereArticleAtmospheric Hg Levels in Tree Barks Due to ArtisanalSmall-Sca...

Azhary Taufiq, Melya Riniarti, Endang Linirin Widiastuti, Hendra Prasetya, Slamet Budi Yuwono, Ceng Asmara...

Citation: Taufiq, A.; Riniarti, M.;Widiastuti, E.L.; Prasetya, H.; Yuwono,S.B.; Asmarah...

Azhary Taufiq, Melya Riniarti, Endang Linirin Widiastuti, Hendra Prasetya, Slamet Budi Yuwono, Ceng Asmara...

Abstract: Mercury (Hg) is a useful heavy metal; however, it is toxic to both humans...

www.mdpi.com