

PAPER NAME

Isolation and identification entomopatho
gen fungi as candidate.pdf

AUTHOR

Endang L Widiatuti

WORD COUNT

2173 Words

CHARACTER COUNT

12580 Characters

PAGE COUNT

6 Pages

FILE SIZE

529.2KB

SUBMISSION DATE

Oct 23, 2022 11:15 AM GMT+7

REPORT DATE

Oct 23, 2022 11:15 AM GMT+7

● 11% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 9% Internet database
- 8% Publications database
- Crossref database
- Crossref Posted Content database
- 7% Submitted Works database

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded sources
- Manually excluded text blocks

PAPER • OPEN ACCESS

Isolation and identification entomopathogen fungi as candidate of bioinsecticide from flies and cockroaches' (Insect vector's disease)

To cite this article: E Rosa *et al* 2021 *J. Phys.: Conf. Ser.* **1751** 012049

View the [article online](#) for updates and enhancements.

You may also like

- [Exploration of entomopathogenic fungi as potential biocontrol of corn earworm \(*Helicoverpa armigera* \(Hübner\)\)](#)
Amelia Sebayang, Andi Tenrirawe and Hishar Mirsam
- [Virulence of Entomopathogenic Fungi Isolates against Green Ladybug *Nezara viridula* L. \(Hemiptera: Pentatomidae\)](#)
Muhammad Agung Permadi, Rafiqah Amanda Lubis, Mukhlis et al.
- [Entomopathogenic fungi as biological agents in forest plant pest control: A systematic review](#)
T Alfina and N F Haneda



The Electrochemical Society
Advancing solid state & electrochemical science & technology

243rd ECS Meeting with SOFC-XVIII

More than 50 symposia are available!

Present your research and accelerate science

Boston, MA • May 28 – June 2, 2023

[Learn more and submit!](#)

2

Isolation and identification entomopathogen fungi as candidate of bioinsecticide from flies and cockroaches' (Insect vector's disease)

E Rosa¹, C.N Ekowati¹, T T Handayani¹, E L Widiastuti¹

¹ Faculty of Mathematics and Natural Sciences, University of Lampung

email: emantisrosa@gmail.com¹, ecolilacto@gmail.com², elwidi@yahoo.com⁴

Abstract: Chemical insecticide has been used for many years to eradicate insects as disease vector, in which causing negative effects not only to those of target insects but also to the environment both in the short and long term. Therefore, it is necessary to determine an alternative biological control of this insect by using natural insecticide (bioinsecticide), such as using fungi as entomopathogen. As microorganism, fungi, contain of bioactive compound with their toxicity could kill the target insect from larvae to adults, called entomopathogen fungi. This entomopathogen fungi are presumably eco-friendly and able to produce toxic compound which can kill target insect such as cyclopeptida destruxin A, B, C, D and desmethyldestruxin B. Two of the fungi which known to be entomopathogenic are *Metarhizium anisopliae* and *Beauveria bassiana*. Yet, there should be more others, therefore it is necessary to explore and isolate any potential bio-insecticide of entomopatogen fungi for disease vectors such as from flies (*Musca domestica*) and cockroach (*Periplaneta americana*). This study found three isolated fungi from flies (L1, L2, L3). They were *Geotrichum* sp., *Penicillium* sp., and *Aspergillus* sp. Three isolated fungi from cockroach (K1, K2, K3) were also found, they were identified as *Aspergillus* sp., except for K3 *Penicillium* sp.

Keyword: bioinsecticide, entomopathogen, *Aspergillus*, *Penicillium*, *Geotrichum*

1. Introduction

Many cases of vector-borne diseases in the community are transmitted by flies and cockroaches, which act as vectors for parasitic diseases such as ascariasis, diarrhea and dysentery. Chemical insecticides are widely used in controlling such vectors, however, negative externalities emerged from the usage of such insecticides are evident in recent studies, i.e. resistance to vectors, the death of non-target animals, and pollution [1,2]. Therefore, it is necessary to explore the potential of natural materials that are not only effective in eradicating insects that acts as a vector role, but also safe for both health and environment; such as entomopathogenic fungi.

Entomopathogenic fungi which are known to be able killing mosquito larvae are *Geotrichum candidum*, *Beauveria bassiana*, *Metarhizium anisopliae*. These fungi produce *destruxin B* and *beauvaricin* compounds which are considered effective in killing larvae (larvicides) [3]. This study by employing *Beauveria bassiana* against the Orthoptera Order, found that increasing in conidia concentration up to 108 conidia/ml caused 43.33% death [3]. Isolation of entomopathogenic fungi, *Metarhizium anisopliae*,

also indicated able to control flies (*Musca domestica* L.) (Diptera: Muscidae) [4], while this *Metarhizium* sp. isolate was also pathogenic to *S. litura* eggs [5]. Yet, the ability of fungi to kill target insects depends on the enzymatic activity of fungi itself [6]. These enzymes contribute in the process of fungal infection in target insects, starting from spore penetration into the host's body until attacking the insect's digestive tract. The types of fungi that are potential to act as bioinsecticides for insect disease vectors is still limited, therefore, this study thrives to explore the isolation and identification of entomopathogenic fungi originating from flies and cockroaches.

2. Methods

This research was conducted in April - November 2018 at the Microbiology Laboratory of the Department of Biology, Faculty of Maths and Sciences, The University of Lampung. The tools and materials used in this study were adjusted to meet the requirement of conducting this study, while the equipment followed the standards of isolation and identification for fungi by the Microbiology Laboratory. This research was conducted through two stages; the isolation stage and the identification stage. Insects used as entomopathogenic fungal agents are originated from *Musca domestica* (flies) and *Periplaneta americana* (cockroaches). The identification process was performed, both by macroscopic and microscopic, on the morphology of colony.

Entomopathogen Fungi Isolation

Insects used to determined for entomopathogenic fungi agents were cockroaches and flies. These insects were left to die, then placed in a petri dish covered with a damp sterile filter paper. The petri dishes were left at room temperature (28-30°C), for 10 days until the insect's body was covered with fungi. The fungi that grows on the insect body were isolated then planted on the PDA medium. Then, they were incubated at room temperature (28-30°C), for 14 days. As fungi started to emerge, they were left to be purified on a sloping medium for identification. Macroscopic and microscopic identifications were carried out by following other studies [7, 8]. The goal from macroscopic identification was to identify colony's color and shape, while microscopic identification aimed at identifying the hyphal structure, spore's shape and location, and the presence of special structures (vesicles, sterigata, vialid l) following other study [9]. The results of the observations were analyzed descriptively and presented in tables and figures

3. Results and Discussion

The study indicated that there were 6 fungi isolates collecting from houses flies and cockroaches. Their characters can be seen in Table 1 for macroscopic characters and Table 2 for microscopic characters.


Table 1. Macroscopic morphological characteristics of the colony of fungi isolates from flies and cockroaches

No	Animal sample	Isolate code	Color		Shape	Edge
			Surface	Base		
1	House flies	L1	White	Hyaline	Circular	Entire
2		L2	Green	Hyaline	Circular	Undulate
3		L3	Yellowish-white	Hyaline	Rhizoid	Filamentous
4	Cockroaches	K1	Grayish-green	White	Circular	Undulate
5		K2	Black	White	Circular	Undulate
6		K3	White	Hyaline	Myceloid	Filamentous


From the macroscopic characters then identification of entomopathogenic fungi was further investigated as well as their microscopic characters. Macroscopic characteristics of Isolate L1 (Table 1) consisted of colony with white and cotton-like filaments, they also had smooth and even edges. Isolate L1 was consisted of insulated hyaline hyphae which did not have conidiophores. Their isolate colony and morphology of entomopathogenic fungi from flies and cockroaches could be seen in Figure 1.

Table 2. Results of the microscopic character identification of entomopathogenic fungi isolates from flies and cockroaches

Isolate	Conidia Structure	Conidiophore	Hypha	Vesicle	Phialide	Foot-cell	Identification
L1	Short & cylindrical	-	Insulated	-	-	-	<i>Geotrichum sp.</i>
L2	Globose	Branched off	Insulated	-	single	-	<i>Penicillium sp.</i>
L3	Globose	Erect	Insulated	Oval	single	Yes	<i>Aspergillus sp.</i>
K1	Globose	Erect	Insulated	Oval	single	Yes	<i>Aspergillus sp.</i>
K2	Globose	Erect	Insulated	Oval	single	Yes	<i>Aspergillus sp.</i>
K3	Globose	Branched off	Insulated	-	single	-	<i>Penicillium sp.</i>



L1



L2, K3




Figure 1. Isolate colony and morphology of entomopathogenic fungi from flies and cockroaches

Conidia (arthospores) are short and cylindrical with insulated end that form chain-shaped, and at the size of $5\ \mu\text{m}$ (Figure 1, L1). This morphological characteristic gives well resemblance to the *Geotrichum* sp. genus.

The colony from L2 and K3 isolates were either green or white, with white edges. The result showed that L2 and K3 have a morphological structure like a brush. The conidia was round-shaped or cylindric-shaped with a diameter of $2.5\ \mu\text{m}$. It also resembled chain-shaped and was located at the end of a single phialide. The result also revealed branch that support the round-shaped phialide. The conidiophores were branched off. The length of the conidiophores was ranging from 22.5 to $27.5\ \mu\text{m}$.

Based on the morphological characteristics observed, L2 and K3 isolates had characteristics in accordance with findings by Kurasein [8], in which *Penicillium* sp. genus was characterized by its feature of hyphae with septa, conidia, sterigma, and conidioconidia (shown in Figure 1). In addition, other characteristics of *Penicillium* sp. were also noted, such as having branched mycelium, conidioconidia that appear on the surface of the conidia with clustered sterigma, and conidia form chains [9].

Isolates L3, K1 and K2 were characterized by insulated hyphae with a diameter of $7.5\ \mu\text{m}$. The conidiophore was at length $275\ \mu\text{m}$, while also having upright and simple structure. On the top of the conidiophore, single phialide was attached to the vesicle. The vesicles were oval-shaped with a diameter of $25 \times 20\ \mu\text{m}$. Its conidia was $5\ \mu\text{m}$ in length with a round-chain-shaped at the end of pumpkin-shaped phialide (Table 2). This morphological feature resembles to the genus *Aspergillus* sp (Figure 1, L3, K1, K2).

Microscopically of *Aspergillus* sp. with a bluish-green round-shaped conidia, while the head of the conidia (vesicle) is *clavate* or round-shaped, and as its colony grows, it would experience transformation into oval-shaped (columnar) and own conidia stalk (conidiophora) [10]. In addition, according to other study [11], *Aspergillus* sp. had sterigmata which seemingly cover the upper half part of the vesicles, it also had conidia with a serrated surface.

The size of conidia in *Aspergillus* might vary due to the differences in gene regulation that determine the establishment of conidia through the conidiation process. Such process leads to the difference in gene expression [12, 13].

The color and time needed for growth of fungi colonies might be affected by several factors; i.e. nutrition, pH, temperature and humidity. Different types of fungi require different nutrition. Therefore, even different types of fungi might be grown in the same media and environmental conditions, it was possible that they would respond differently. Each type of fungi had a different optimum requirement of pH, temperature, and humidity to support its growth, for instance, *Aspergillus flavus* fungi requires pH of 6, temperature of $25\ ^\circ\text{C}$ and humidity of 90; whereas *Penicillium chrysogenum* requires pH of 7, temperature of $30\ ^\circ\text{C}$ and humidity of 90 for its growth [14].

4. Conclusion

1. Three fungi isolates were obtained houseflies, they were L1, L2 and L3 and through macroscopic and microscopic identification they revealed to be *Geotrichum sp.*, *Penicillium sp.*, *Aspergillus sp.*
2. Three fungi isolates also found from cockroaches, they were K1, K2 and K3, which were identified to be *Aspergillus sp.* and *Penicillium sp.*

References

- [1] Nejati Jalil., Boostanbakhsh Atefeh. 2012. Cockroaches bacterial infections in wards of hospitals, Home dan city, west of Iran. *Asian Pacific Journal of Tropical Disease*. Vol. 2: 381-384
- [2] Rahman ,M.S dan Soviana ,L.2016. Perbedaan Kerentanan status nyamuk *Aedes aegypti* terhadap Malathion di Kabupaten Bantul Yogyakarta Jurnal Kesehatan Masyarakat. Volume 11 no 2.
- [3] Dewantara Neil, A. Wibowo N, Felicia J.2017. Efektivitas *Beauveria bassiana* (Bals.) Vuillemin Sebagai Pengendalian Hama Belalang Kayu (*Valanga nigricornis* Burm). 1-14. ISSN.
- [4] Yunizar, N., Rahmawati.,Kustiati. 2018. Patogenitas Isolat Jamur Entomopatogen *Metarhizium anisopliae* terhadap Lalat Rumah (*Musca domestica* L.) (Diptera: Muscidae). *Jurnal Protobiont*. Vol. 7(3): 77-82.
- [5] Trizelia, M.Y. Syahrawati, Dan A. Mardiah. 2011. Patogenisitas Beberapa Isolat Cendawan Entomopatogen *Metarhizium* spp. terhadap Telur *Spodoptera litura* Fabricius (Lepidoptera: Noctuidae). *J. Entomol. Indon.*, 8 (1): 45-54
- [6] Halimah, N. Imaningsih W. Mariana. 2018. Karakterisasi Morfologi Jamur Entomopatogen di Hutan Mandiangin Banjarbaru, Kalimantan Selatan. *Jurnal Mikologi Indonesia*. Vol. 2(1): 39-48
- [7] Humber, A.R. 2005. Entomopathogenic Fungal Identification. American Phytopathological Society and Entomological Society of America. Las Vegas, Nevada
- [8] Barnett dan B.B. Hunter. 1998. *Illustrated Genera of Imperfect Fungi*. 4th Edition. Macmillan Publishing Company. New York.
- [9] Ganjar, I., Wellyzar Sjamsuridzal dan ariyanti Oetari. 2006. Mikologi Dasar dan Terapan. Yayasan Obor Indonesia. Jakarta.
- [10] Kurasein, Riccardo AA Muzzarelli.2009. Chitin. *Elsevier*.
- [11] Fardiaz, S. 1989. *Penuntun Praktikum Mikrobiologi Pangan*. PAU Pangan dan Gizi IPB. Bogor.
- [12] Redig, P. 2005. Mycotic infections in birds I: Aspergillosis. *North American Veterinary Conference Proceedings*, Eastern States Veterinary Association. 1192-1194.
- [13] Hyuk, Jae and Yu. 2010. Regulation of Development in *Aspergillus nidulans* and *Aspergillus fumigatus*. *Journal of Mycobiology*. Vol. 38(4) : 229-237.
- [14] Praja, Ratih Novita., Yudhana, Aditya. 2017. Isolasi dan Identifikasi *Aspergillus* spp. Pada Paru-Paru Ayam Kampung yang dijual di Pasar Banyuwangi. *Jurnal Medik Veteriner*. Vol. 1 No. 1 : 6-11.

● 11% Overall Similarity

Top sources found in the following databases:

- 9% Internet database
- Crossref database
- 7% Submitted Works database
- 8% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	faculty.uobasrah.edu.iq	Internet	3%
2	irep.iium.edu.my	Internet	2%
3	eprints.umsida.ac.id	Internet	2%
4	O Lisa, S Hanum, S Samingan, M Bahi, A Alfizar, L Fitri, N Novita, S Sya...	Crossref	2%
5	University of Surrey on 2021-10-20	Submitted works	<1%
6	Higher Education Commission Pakistan on 2016-09-06	Submitted works	<1%
7	Sriwijaya University on 2021-08-16	Submitted works	<1%
8	Hafiz Fauzana, Febriliani Arda, Nelvia, Rusli Rustam, Fifi puspita. " Tes...	Crossref	<1%

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded text blocks
- Manually excluded sources

EXCLUDED SOURCES

E Rosa, C.N Ekowati, T T Handayani, E L Widiastuti. "Isolation and identificatio... 80%

Crossref

repository.lppm.unila.ac.id 79%

Internet

Victorian Institute of Technology on 2022-10-10 6%

Submitted works

iopscience.iop.org 5%

Internet

L Budiarti, N S P Nuryanti. "Entomopathogenic Fungus from Rice Rhizosphere ... 5%

Crossref

EXCLUDED TEXT BLOCKS

Journal of Physics: Conference SeriesPAPER • OPEN ACCESSYou may also like

Victorian Institute of Technology on 2022-10-10

To cite this article: E Rosa et al 2021 J. Phys.: Conf. Ser. 1751 012049

repository.lppm.unila.ac.id

View the article online for updates and enhancements

repository.lppm.unila.ac.id

Virulence of Entomopathogenic Fungisolates against Green Ladybug Nezaraviridu...

eprints.umsida.ac.id

Helicoverpa armigera (Hübner) Amelia Sebayang, Andi Tenrirawe and Hishar Mirsam

L Budiarti, N S P Nuryanti. "Entomopathogenic Fungus from Rice Rhizosphere Soil in Politeknik Negeri Lamp..."

2020 Journal of Physics: Conference Series IOP Publishing

eprints.umsida.ac.id

1 Faculty of Mathematics and Natural Sciences, University of Lampung email: ema...

irep.iium.edu.my

Exploration of entomopathogenic fungi as potential biocontrol of corn earworm

iopscience.iop.org