

Materials Science and Engineering
Conference Series

1173

Preface

It has been a tremendous experience to organize the 6th International Conference of Science, Technology, and Interdisciplinary Research (IC-STAR 2020). The annual international conference is held in conjunction with other conferences throughout series of event spanning from October to December 2020. The scope of IC-STAR 2020 includes chemical and biotechnology, civil and environmental, computer science and information technology, electrical and communications, materials sciences, industrial engineering, mechanical and aerospace, geodetic and geophysics, applied mathematics and physics sciences, and also the interdisciplinary research. IC-STAR conference series intend to integrate engineering and life sciences fields in order to provide a cross section roadmap from basic research, technological and developments, processes development, and their applications and integrity in the real-world usage.

The event's series started with a virtual event which is hosted at Universitas Lampung, in Bandar Lampung, Indonesia, on 20 – 21 October 2020. The conference featured Keynote Speakers list of Prof. Ahmad Saudi Samosir (Universitas Lampung, Indonesia), Prof. Rizalman Mamat (Universiti Malaysia Pahang, Malaysia), Prof. Sunny J. Kalayathankal (K. E. College, India), Prof. Naoya Kasai (Yokohama National University, Japan) and Prof. Gavin Brown (Auckland University, New Zealand). During the event, there are about 50 authors presenting their papers in oral presentation.

The series proceeded with another virtual event hosted at Institut Teknologi Bandung, in Bandung, Indonesia, on 25-26 November 2021 by ICASA-MATRIX. Keynote Speakers for the conference were Prof. Koji Shimoyama (Tohoku University, Japan), Prof. Erry YT Adesta (International Islamic University of Malaysia, Malaysia), Prof. Tatacipta Dirgantara (Institut Teknologi Bandung, Indonesia) and Prof. Bambang Kismono Hadi (Institut Teknologi Bandung, Indonesia). There were about 40 oral presentations during the event.

The series concluded with a virtual event jointly hosted and supported by Universiti Teknologi Malaysia, Curtin University Malaysia, Universiti Teknologi Brunei, and Universitas Lampung, on 8 – 9 December 2021. The Keynote Speakers were Prof. Hadi Nur (Universiti Teknologi Malaysia, Malaysia), Prof. Ramesh Singh (Universiti Teknologi Brunei, Brunei Darussalam), Dr. Rino Rakhmata Mukti (Institut Teknologi Bandung, Indonesia), Prof. Mohammed Rafiq Abdul Kadir (Universiti Teknologi Malaysia, Malaysia), Prof. M. Akbar Rhamdhani (Swinburne University, Australia) and Prof. Ahad Ali (Lawrence Technological University, USA). There were about 80 oral presentations in the event.

We appreciate all distinguished speakers and all authors who presented during the conference series. We also acknowledge the effort of all committees, including all reviewers, for their contribution to the conference series and in making this proceeding. Thank you and hope to see you in future events of the conference series.

Publication and Proceeding Editors,

Irza Sukmana (Universitas Lampung, Indonesia)

Khairuddin (Universitas Lampung, Indonesia)

Seri Rahayu Ya'akub (Universiti Teknologi Brunei, Brunei Darussalam)

Fethma M Nor (Curtin University, Malaysia)

Taufiq Mulyanto (Institut Teknologi Bandung, Indonesia)

Muhammad Hanif (Universitas Lampung, Indonesia)



Patron

- **Prof. Dr. Karomani, M.Si** – *Rector, Universitas Lampung, Indonesia*
- **Prof. Akio Tomiyama** – *Kobe University, Japan*
- **Prof. Hadi Nur** – *Universiti Teknologi Malaysia, Malaysia*
- **Prof. Toshiro Ohashi** – *Hokkaido University, Japan*
- **Prof. Gurumurthy Hegde** – *BMS College of Engineering, India*
- **Prof. Gamal Baroud** – *Universite de Sherbrooke, QC – Canada*

Steering Committee

- **Prof. Dr. Heryandi, S.H., M.S.** – *Vice Rector for Academic, Universitas Lampung, Indonesia*
- **Prof. Suharso, Ph.D.** – *Vice Rector for Cooperation, Universitas Lampung, Indonesia*
- **Dr. Eng. Helmy Fitriawan** – *Dean, Faculty of Engineering, Universitas Lampung, Indonesia*
- **Dr.rer.nat. Ir. R. M. Rustamaji, M.T., IPU** – *Head, Forum Teknik BKS PTN Barat, Indonesia*
- **Dr. Ir. Lusmeilia Afriani, D.E.A** – *Director of Research Institute and Community Service, Universitas Lampung, Indonesia*
- **Dr. M. Irsyad** – *Universitas Lampung, Indonesia*
- **Dr. Vera A. Noorhidana** – *University of Lampung, Indonesia*

International Steering Committee

- **Prof. Gunawarman** – *Universitas Andalas, Indonesia*
- **Dr. Denni Kurniawan** – *Universiti Teknologi Brunei, Brunei Darussalam*
- **Prof. Rafaqat Hussain** – *COMSATS Institute of Information Technology, Pakistan*
- **Dr. Basem Abu Izneid** – *University of Business & Technology, Saudi Arabia*
- **Dr. Ahmad Kafrawi Nasution** – *Muhammadiyah University of Riau, Indonesia*
- **Dr. Deddy H. B. Wicaksono** – *Swiss German University, Indonesia*
- **Dr. Fethma M Nor** – *Curtin University Sarawak, Malaysia*
- **Assoc. Prof. Taufiq Mulyanto** – *Institut Teknologi Bandung, Indonesia*
- **Dr. Uceu Suhuddin** – *Helmholtz-Zentrum Geesthacht, Germany*
- **Dr. Taha Abd Elrahman** – *Unviersite de Sherbrooke, QC, Canada*
- **Dr. Dandi Bachtiar** – *International Network on Science, Technology and Education Programme, INSTEP Network, Indonesia*
- **Dr. Yanuar Z. Afief** – *Universiti Malaysia Sarawak, Malaysia*
- **Dr. M. Fakhrol Ulum** – *IPB University, Indonesia*

Organizing Committee

- **Assoc. Prof. Irza Sukmana** – *Conference Chair*
- **Assoc. Prof. Sarkowi** – *Conference co-Chair*
- **Dr. Eng. Misfa Susanto** – *General Secretary*
- **A. Yudi Eka Risano, S.T., M.Eng.** – *Treasurer*

TPC Coordinators

- **Dr. Eng. Khairudin** – *Scientific Chair*
- **Gigih Forda Nama, S.T., M.T.** – *ICT*
- **Dr. Nandi Khairudin** – *Review and Publication Chair*
- **Martinus, S.T., M.Sc.** – *Virtual Meeting Chair*

Contact

All questions about submissions should be emailed to icstar@eng.unila.ac.id and for *WhatsApp*, please contact **+62 812 7964 334**

Table of contents

[Volume 1173](#)

2021

[Previous issue](#)[Next issue](#)

The 6th International Conference of Science, Technology, and Interdisciplinary Research 2020 (IC-STAR 2020) 20th-21st October 2020, Bandar Lampung, Indonesia

Accepted papers received: 29 July 2021

Published online: 01 September 2021

Papers

012001

OPEN ACCESS

Characteristics of medium-low rank coal blending on performance and efficiency steam power plant

Hendri, R Nurhasanah and A F Lubis

012002

OPEN ACCESS

Comparison of the winglet aerodynamic performance in unmanned aerial vehicle at low Reynolds number

S H S Putro, B J Pitoyo, N Pambudiyatno, Sutardi and W A Widodo

OPEN ACCESS

Structure from Motion (SfM) method to characterize fluvial sedimentology in Way Semaka river in Lampung province, Indonesia

R C Wibowo, B S Mulyatno, O Dewanto and M Sarkowi

012004

OPEN ACCESS

Optimization, modelling and simulation to scale-up the production of rice bran extract and defatted rice bran

G Fitriyano, N H Fithriyah, R A Nugrahani, A B Syamsudin and M Kosasih

012005

OPEN ACCESS

Optimization of shell and tube heat exchanger using the water cycle algorithm

M F I M Hanafi, A Bahreininejad and N Uddin

012006

OPEN ACCESS

Classification of *Civet* and *Canephora* coffee using Support-Vector Machines (SVM) algorithm based on order-1 feature extraction

R Z H Suyoto, M Komarudin, G F Nama and T Yulianti

012007

OPEN ACCESS

Geophysical approach for assessment of seawater intrusion in the coastal aquifer of Bandar Lampung, Indonesia

Rustadi, I G B Darmawan, N Haerudin, Suharno and A Setiawan

012010

OPEN ACCESS

Coal velocity and proximate analysis relationship using Multiple Linear Regression

S Erfani, F M S Siregar, A Zaenudin, Rustadi, I B S Yogi and R C Wibowo

Geophysical approach for assessment of seawater intrusion in the coastal aquifer of Bandar Lampung, Indonesia

Rustadi^{1,2,*}, I G B Darmawan², N Haerudin², Suharno², and A Setiawan¹

¹ Department of Environment Science, Universitas Lampung, Jalan Prof. Soemantri Brojonegoro No.1, Bandar Lampung 35145, Indonesia

² Department of Geophysical Engineering, Universitas Lampung, Jalan Prof. Soemantri Brojonegoro No.1, Bandar Lampung 35145, Indonesia

*Email: rustadi_2007@yahoo.com

Abstract. Overexploitation in 1980-2000 caused a lot of groundwater loss in the aquifer coastal of Bandar Lampung. Groundwater has been degraded with increasing salinity due to the intrusion of seawater. In 2013, the intrusion extended 400 m from the coastline. Subsurface geological mapping is carried out to examine alluvial formation's material constituents and the ability to transfer seawater to land. Three well explorations combined with three geoelectric tomography and 51 points of gravity. The gravity method is used to map structures that can connect coastal aquifers and land aquifers. Meanwhile, geoelectric tomography to map the presence of aquifers and low resistivity zones by seawater intrusion. The coastline has relatively shallow aquifers at a depth of 2 - 10 m in a sand layer with coral inserts, while the lower part is composed of silt and clay. The aquifer has experienced a decrease in resistivity of fewer than 10 ohms.m appears in the three circular patterns, caused by seawater. In addition to the permeable zone of shallow aquifers, there is a northwest-southeast trending structure that can channel seawater to land aquifers.

Keywords: Coastal aquifer, seawater intrusion, geophysics, Bandar Lampung

1. Introduction

Since the Dutch colonial period, the Bandar Lampung coast has been the gateway to commerce between regions. The VOC relied on Teluk Betung to collect and transport the spices obtained in Lampung Province. As an economic center for a long time, it has been able to trigger regional development. Developing the manufacturing industry, tourism, and settlements with a population density of 8050/km².

The main component of clean water sources on the coast of Bandar Lampung is groundwater. Overexploitation in 1980 - 2010 has caused degradation in coastal aquifers and increased sodium and chloride, caused by seawater's intrusion. In 2013, the distribution zone reached a radius of 500 m from the coastline [1]. Seawater intrusion in coastal areas is a process of infiltrating saltwater from the sea into groundwater on land. In natural conditions, groundwater will flow continuously into the sea. When the balance is disturbed, the intrusion could be extended inland. The problem of groundwater contamination by seawater has become a global problem [2-4].



Increasing groundwater salinity on the coast of Bandar Lampung requires regular monitoring and is supported by integrated data. However, there are various problems, and the research focuses on mapping aquifer layers and faults that can affect the interaction between seawater and freshwater on land. Geoelectric and gravity methods are used to obtain initial information on coastal aquifers [5] and structures [6]. Geoelectric mapping is often performed to delineate aquifers [7-9] and evaluate zones that have been exposed by seawater intrusion [10-12]. Meanwhile, the gravity method is quite useful in interpreting the thickness of the sediment layer and the possible presence of structures [13-15].

2. Coastal geology in Bandar Lampung

The geology of the coastal of Bandar Lampung is composed of sedimentation from land runoff and ocean wave circulation to produce coastal alluvial deposits (Qa). This formation has a limited distribution and forms inconsistent contacts with the Quaternary sedimentary rock formations, Young Mountain Rock Formation (Qhv) and the Lampung Formation (QTI) and Tertiary sedimentary rocks, the Campang Formation (TPOC), and the Tarahan Formation (TPOT). The geological setting of the Bandar Lampung coast and its surroundings is shown in Figure 1.

Sedimentary rock formations and volcanic products cover bedrock in the form of metamorphic rocks consisting of migmatite, schist, gneiss, alabaster, and quartzite, which are included in the Gunung Kasih Complex (Pzg). Gunung Kasih Complex is in pre-tertiary age, and in some places, it has a relatively shallow position [16]. The Bandar Lampung region has a complex geological setting that is influenced by tectonic pressure, which results in an increase in magmatic through the bedrock and forming intrusions and lenses of igneous rock. The tectonic process forms various faults in the direction of the subduction path. Various faults can act as permeable zones that allow interactions between the saline water to move to extend inland.




Figure 1. Geology map, well exploration, measure point of gravity, and line ERT of the study area.

3. Materials and methods

There are three exploration wells in coastal alluvial deposits (Qa) intending to determine soil mechanics' constituent lithology and characterization. Imaging to map the conductive zone related to aquifers' presence and seawater intrusion distribution is carried out through 3 lines of electrical resistivity tomography (ERT). Measurements using the ARES instrument, with the Wenner-Schlumberger configuration, electrode spacing 6 m. Other sediment formations and faults were

analyzed through gravity measurements to map the alluvial coastal sediment layer's thickness. Measurements using a LaCoste and Romberg gravity meter with points scattered randomly. Each data point from the well is represented by a red dot (W), the electrical resistivity tomography (ERT) is symbolized by green rectangular (G), and a thick black dot represents the gravity measuring point.

4. Results and discussions

4.1. Gravity

The anomaly Bouguer from gravity data is shown in Figure 2, and the anomaly value ranges from 32 - 60 mgal. Low anomalies correlate with young formations, especially coastal alluvials in the south. There are indications of basins in coastal alluvials with anomalies ranging from 34 - 38 mgal, as shown on Fig. 2.




Figure 2. Map of the Bouguer anomaly with contour intervals of 2 mgal.

The contour pattern of the Bouguer anomaly in the Lampung Bay area is in line with the existence of the fault structure in the area. The low anomaly in the south is oriented towards the northwest-southeast. The direction of this anomaly is also in line with the direction of the dominant structure. An incision profile was carried out on the A-A' path to identify this dominant structure's effect. Qualitative interpretation through 2.5 D modelling on slice A-A' to obtain subsurface images of the basin on the coast of Bandar Lampung.

The modelling results are shown in Figure 2 show that the two main structures are trending northwest-southeast. This structure is a normal fault structure with opposite slopes to form a sag basin. This structural pattern confirms the presence of the basin from the Bouguer anomaly contour. This basin area is interpreted as a zone of the potential groundwater aquifer.

4.2. Electrical resistivity tomography

The measurement data of three ERT lines in G-1, G-2, and G-3 is processed using Res2DInv software. The resulting image is shown in Figures 3, 4, and 5.

Tomographic inversion results is presented in Figure 3 show a low resistivity value at low depth. These results indicate that groundwater has been mixed with seawater in shallow aquifers. However, this pattern still does not show a seawater intrusion pattern, but rather a mixture of groundwater in coastal areas. This condition is reinforced by observations in the field of saltwater conditions in the coastal area of Lampung Bay. The majority of residents no longer use groundwater wells and instead use clean water by buying. Meanwhile, in the port area (Panjang Port), the resistivity tomography inversion results also showed the same pattern as the Lampung Bay (Figure 4).




Figure 3. Inverted electrical resistivity models for Wenner-Schlumberger measurements on ERT line 1.




Figure 4. Inverted electrical resistivity models for Wenner-Schlumberger measurements on ERT line 2.

The absence of a seawater intrusion resistivity pattern in this measurement indicates that seawater contamination only occurs in shallow groundwater aquifers, specifically in coastal areas. This condition is seen at low resistivity values with a thickness of up to 10 m at the beginning of the measurement path. The low resistivity value indicates the effect of mixed saltwater on groundwater. Two resistivity patterns appeared that formed a trench in the middle of the measurement path with a low resistivity value <10 Ohm.m. This pattern is interpreted as a trench or shallow aquifer pathway that may have been formed due to the Lampung-Panjang Fault's existence. This low resistivity pattern also indicates the presence of an accretion structure filled with saline water.

Although the ERT line 1 and line 2 measurement paths have not shown a seawater intrusion pattern, exciting results are shown by the third line. This route is located in the industrial area and the population, namely Srengsem Village. This area is the location for massive groundwater utilization. Several problems have emerged regarding the need for clean water in this area. The low resistivity pattern indicating the seawater intrusion zone is shown in Figure 5.




Figure 5. Inverted electrical resistivity models for Wenner-Schlumberger measurements on ERT line 3.

The seawater intrusion channel that appears in the middle of the track is composed of three circular patterns. This pattern has a low resistivity value of fewer than 10 ohms. The seawater path is estimated to be in the geological structure gap that develops in the region. This seawater intrusion pattern can occur due to the massive use of groundwater to disturb the aquifer's stability and the soil.

4.3. Well exploration

The constituent materials in the coastal alluvial formation produce variations in lithology and thickness in the three wells. The drilling results' interpretation is shown in Figure 6 and site class testing from the SPT compression test, the layers are still in the soft group up to a depth of 12 m in well-1, 26 m in well-2, and 32 m in well-3.

The first well (Well-1) layer is dominated by clay at a depth of 2 - 25 meters below sea level. Meanwhile, the sand layer with coral inserts is in the second layer with a thickness of 2 meters, overlain by a clay layer above it. The thickness of the clay layer in well-1 indicates an impermeable layer that holds the aquifer above it so that seawater intrusion cannot penetrate the clay layer. However, surface aquifers that are not too thick are likely to be contaminated by seawater. The sand layer's density with coral inserts shows that the shallow marine still influences forming this rock layer to deep marine depositional environment.




Figure 6. Lithology section on three explorations well

The location of well-2 and well-3 is in the same location, namely Srengsem Village, Panjang District. Well-3 is closer to the coast than well-2. Well-3 starts with a seawater depth of up to 14 m until it meets a silt layer with coral and clay insert at the bottom. This layer indicates a depositional environment in the deep sea to the shallow sea transition area. In line with well-2, the presence of a silt layer with coral and clay insertion at the bottom indicates a similar depositional environment in well-3. However, in well-2, there is an aquifer layer of groundwater in the form of coral sand with a thickness of approximately 6 m. This layer has the potential to act as a groundwater pathway that can be contaminated by seawater.

5. Conclusions

The results of the geophysical approach to assessing the intrusion zone of seawater in the Bandar Lampung coastal aquifer have shown a mixed pattern of groundwater and seawater. The gravity method has succeeded in identifying the groundwater aquifer's basin area in Lampung Bay controlled by two normal fault structures. One of the three ERT measurement paths successfully detected a seawater intrusion pattern in the Srengsem Village area, namely the ERT line 3. The resistivity value is relatively low (<10 ohms), appears in the three circular patterns, and is interpreted as a seawater intrusion zone. This finding is also reinforced from drill data, especially in well-2, which shows the groundwater aquifer layer is in a sand layer with coral inserts.

Acknowledgements

Acknowledgments were delivered to the Universitas Lampung, especially the Department of Geophysical Engineering and all colleges who have helped collect data, processing, and model, until completing this paper.

References

- [1] Syafridi, Zaenudin A, Kusumastuti D I, and Suharno 2013 Penggunaan metode geolistrik untuk pemodelan intrusi air laut di daerah pesisir Kota Bandar Lampung *Jurnal Teori dan Aplikasi Fisika* **01**(02)
- [2] Loáiciga H A, Pingel T J, Garcia E S 2012 Sea water intrusion by sea-level rise: Scenarios for the 21st century *Ground Water* **50**(1):37–47
- [3] Das S, Maity P K, Das R 2018 Remedial Measures for Saline Water Ingression in Coastal Aquifers of South West Bengal in India *MOJ Ecol Environ Sci* **3**(1)
- [4] Perera M D N D, Ranasinghe T K G P, Piyadasa R U K, Jayasinghe G Y 2018 Risk of seawater intrusion on coastal community of Bentota river basin Sri Lanka. *Procedia Eng* **212** 699–706 <https://doi.org/10.1016/j.proeng.2018.01.090>
- [5] Pratama W R 2019 Aplikasi Metode Geolistrik Resistivitas Konfigurasi Wenner-Schlumberger Untuk Mengidentifikasi Litologi Batuan Bawah Permukaan dan Fluida Panas Bumi Way Ratai di Area Manifestasi Padok di Kecamatan Padang Cermin Kabupaten Pesawaran Provinsi Lampung *J. Geofis. Eksplor.* **5** 30–44
- [6] Wibowo R C, Mulyatno B S, Romosi M and Zaenudin A 2018 Determination of slip surface area using geoelectric, MASW, and soil mechanics data in cimuncang village, west Java *EAGE-HAGI 1st Asia Pacific Meet. Near Surf. Geosci. Eng.* 0–4
- [7] El_Hameed A G A, El-Shayeb H M, El-Araby N A, Hegab M G 2017 Integrated geoelectrical and hydrogeological studies on Wadi Qena, Egypt *NRIAG J Astron Geophys* **6**(1):218–29 <http://dx.doi.org/10.1016/j.nrjag.2017.03.003>
- [8] Mohamaden M I I, Abuo Shagar S, Allah G A 2009 Geoelectrical survey for groundwater exploration at the Asyuit governorate, Nile Valley, Egypt *J King Abdulaziz Univ Mar Sci.* **20**(1):91–108
- [9] Sultan S A, Essa K S A T, Khalil M H, El-Nahry A E H, Galal A N H 2017 Evaluation of groundwater potentiality survey in south Ataq-northwestern part of Gulf of Suez by using resistivity data and site-selection modeling *NRIAG J Astron Geophys* **6**(1):230–43 <http://dx.doi.org/10.1016/j.nrjag.2017.02.002>
- [10] Bouderbala A, Remini B, Hamoudi A S 2016 Geoelectrical investigation of saline water intrusion into freshwater aquifers: A case study of Nador coastal aquifer, Tipaza, Algeria. *Geofis Int.* **55**(4):239–53
- [11] Alfarrak N, Walraevens K 2018 Groundwater overexploitation and seawater intrusion in coastal areas of arid and semi-arid regions *Water (Switzerland)* **10**(2)
- [12] Sathish S, Elango L, Rajesh R, Sarma V S 2011 Application of Three Dimensional Electrical Resistivity Tomography to Identify Seawater Intrusion **4**(1):21–8
- [13] Al Farajat M 2009 Characterization of a coastal aquifer basin using gravity and resistivity methods: A case study from Aqaba in Jordan *Acta Geophys* **57**(2):454–75
- [14] Pool D R 2008 The utility of gravity and water-level monitoring at alluvial aquifer wells in southern Arizona *Geophysics* **73**(6)
- [15] Delinom R M 2009 Structural geology controls on groundwater flow: Lembang fault case study, West Java, Indonesia *Hydrogeol J.* **17**(4):10 11–23
- [16] Darmawan I G B, Setijadji L D and Wintolo J 2015 Geology and Geothermal System in Rajabasa Volcano South Lampung Regency, Indonesia (Approach to Field Observations, Water Geochemistry and Magnetic Methods) *Proceedings World Geothermal Congress* (Melbourne, Australia).