

Liquid Organic Fertilizer from Plant Extracts Improves the Growth, Yield and Quality of Sweet Corn (*Zea mays* L. var. *saccharata*)

By Darwin Habinsaran Pangaribuan; Sarno;Kus Hendarto;Priyanto;Ajeng Kusuma Darma;Tika Aprillia

16
Liquid Organic Fertilizer from Plant Extracts Improves the Growth, Yield and Quality of Sweet Corn (*Zea mays* L. var. *saccharata*)

Darwin Habinsaran Pangaribuan^a, Sarno, Kus Hendarto, Priyanto, Ajeng Kusuma Darma and Tika Aprillia

Department of Agrotechnology, Faculty of Agriculture, University of Lampung, 35145 Bandar Lampung, Indonesia

11
ABSTRACT

The objective of this research was to study the effect of concentration and dose of liquid organic fertilizer (LOF) derived from an extract mixture derived from *lamtoro* leaves, banana humps, and coconut fibers on the growth, yield, quality, and nutrient uptake of sweet corn. Experiment 1 comprised six treatments, namely a control without LOF, recommended inorganic fertilizers, and LOF at a concentration of 15 ml l⁻¹, 30 ml l⁻¹, 45 ml l⁻¹, or 60 ml l⁻¹. Experiment 2 consisted of five treatments, namely recommended inorganic fertilizers, and LOF doses of 25 l ha⁻¹, 50 l ha⁻¹, 75 l ha⁻¹ or 100 l ha⁻¹. Results showed that LOF consistently increased the growth, yield and quality of sweet corn. Application of LOF with a concentration of 60 ml l⁻¹ or a dose of 100 l ha⁻¹ showed the highest yield compared to other treatments. The quality of sweet corn increased markedly. It was concluded that LOF could be applied as an additional supplement to inorganic fertilizers used for sweet corn organic farming in the tropics.

Keywords: Banana hump, coconut fiber, leucaena extract, nutrient uptake, organic farming

INTRODUCTION

The community prefers sweet corn (*Zea mays* L. var. *saccharata*) because of its relatively high sugar content. Organic sweet corn production needs can be improved by improving fertilization techniques. Fertilizers for sweet corn can be given in the solid or liquid form. Solid organic fertilizers tend to be taken up by the crop relatively slowly (Hartz et al., 2000; Johnson et al., 2012). Another alternative in organic tropical agriculture is the use of liquid organic fertilizers (LOF), which are taken up more quickly by crops.

The application of fertilizer into the plants through leaves was successfully carried out on corn plants (Amanullah et al., 2014). Similarly, the application of LOF was carried out on various vegetable crops such as sweet corn (Fahrurrozi et al., 2016) and tomatoes (Zhai et al., 2009). Hartz et al. (2010) reported that the application of LOF could function the same as conventional fertilization with solid fertilizers. Furthermore, the researchers stated that nitrogen derived from liquid organic fertilizers is more readily available and the nitrification process takes place faster.

One cultivation technique used to increase growth, yield, and quality of sweet corn grown organically are LOF made from the waste of plant materials available around the farmer's garden. There is still a lack of a scientific basis on the utilization of plant extracts as LOF. Plant wastes such as *lamtoro* (*Leucaena leucocephala*) leaves, banana humps, and coconut husks are widely available around farmers' gardens in tropical areas. *Lamtoro* leaves are often used as animal feed. *Lamtoro* leaves contain significant amounts of nitrogen, protein, vitamins, and minerals (Farinu et al., 1992; Meulen et al., 1979). Liquid organic fertilizer from the leaves of *lamtoro* is a source of nitrogenous nutrients for plants. Banana humps contain microbial decomposers of organic matter, such as *Bacillus* sp., *Aeromonas* sp. and *Aspergillus niger* (Suhastyo et al., 2013). These microbes were able to decompose organic matter. ¹⁵ The main content of coconut fibers is cellulose, hemi-cellulose, and lignin (Arsyad et al., 2015). If the coconut fiber is soaked and fermented with water, the potassium contained in coconut fiber will dissolve in soaking water so that the liquid fertilizer coconut husk is high in potassium.

¹⁰ The purpose of this study was to determine the effect of concentration and dose of liquid organic fertilizer made from an extract mixture of *lamtoro* leaves, banana humps and coconut fibers on the growth, production, quality, and nutrient uptake of sweet corn plants.

MATERIALS AND METHODS

This research was carried out in Kota Sepang experimental station, Bandar Lampung, Indonesia. The study area was classified as having Ultisol soil. A field study was set up in RCBD (Randomized Complete Block Design) and replicated three times. The experiment consisted of two sub-experiments. Experiment 1 (April to June 2016) consisted of six treatments, namely an unfertilized control, recommended inorganic fertilizer, and application of LOF with a concentration of 15 ml l⁻¹, 30 ml l⁻¹, 45 ml l⁻¹, or 60 ml l⁻¹. Experiment 2 (April to June 2017) consisted of five treatments, namely recommended inorganic fertilizer, and an application of LOF with a dose of 25 l ha⁻¹, 50 l ha⁻¹, 75 l ha⁻¹, or 100 l ha⁻¹.

The research began with making plant extract LOF following a modified procedure developed by Astuti et al. (2014). *Lamtoro* leaves, banana humps, and coconut fibers were cut into small pieces and then separately put into containers, into which brown sugar and rice washing water were added and stirred evenly. Then EM-4 was added, after which the container was closed and then fermented for 21 days and filtered. Liquid organic fertilizer from *lamtoro* leaves, banana humps, and coconut fibers was then mixed in a ratio of 1: 1: 1. The nutrient content of the plant extracts (Table 1) was analyzed in Soil Laboratory, Faculty of Agriculture, University of Lampung.

All plots were hand hoed before planting with sweet corn. The area of each plot was 3 m × 3 m, and the row spacing was 70 cm × 20 cm. The application of liquid fertilizer was carried out by spraying evenly on the upper and lower part of leaves. The application of LOF was carried out once a week from 2 to 7 weeks after planting (WAP). The inorganic recommended fertilizers were given included urea at 300 kg ha⁻¹, SP-36 at 150 kg ha⁻¹, and KCl at 100 kg ha⁻¹. Agronomic practices used in this study included watering, growing, control of weeds, pests, and diseases organically. Harvesting was conducted at 70 days after planting.

The parameters observed in this study were the number of leaves, stem diameter, plant height, leaf area index (LAI), SPAD value, the weight of husked ear, the weight of unhusked ear, ear length, ear diameter, shoot dry weight, °Brix value, yield, and N, P and K nutrient uptake. Statistical analysis of the data was conducted using MINITAB v. 16.0 software. Analysis of variance and least significant difference (LSD) tests was used to test overall treatment effects and between-treatment differences and was conducted at the 5% probability level.

RESULTS AND DISCUSSION

Results

The LOF application affected the vegetative parameters of the plants. Leaf number and stem diameter were greater in LOF-applied plant than the control (Table 2). The results of the analysis of variance showed that the application of LOF increased the number of leaves and stem diameter. The number of leaves and stem diameter treated with concentrations of 15 ml l⁻¹, 30 ml l⁻¹, 45 ml l⁻¹, and 60 ml l⁻¹ were not significantly different. The lowest number of leaves and the lowest stem diameter were recorded in the control treatment (Table 2).

The LOF application increased plant height, leaf area index, and the greenness of the leaves (indicated by SPAD value) (Table 3). The plant height, LAI, and SPAD value were greater in LOF-applied plants than with the inorganic fertilizer treatment. Plant height and LAI increased with increasing doses of LOF until 75 l ha⁻¹; above these levels, there was no significant additional increase. The greenness level of leaves (SPAD value) in the 100 l ha⁻¹ treatment was higher than in the other treatments, while doses of 50 l ha⁻¹ and 75 l ha⁻¹ were not significantly different from each other. The lowest plant height, LAI and SPAD value were recorded with the inorganic fertilizer treatment (Table 3).

LOF application also affected the generative parameters of plants. The results of the analysis of variance indicated that the LOF application increased the weight of unhusked ears and weight of husked ears (Tables 4 and 5). Weights of unhusked ears and weight of husked ears in the LOF treatment were significantly higher than the control treatment (Table 4). The lowest weights of unhusked ears, the weight of husked ears, ear length, and ear diameter were recorded in the control plants (Table 4).

The LOF application increased ear length (Tables 4 and 5). Ear lengths at concentrations of 30 ml l⁻¹, 45 ml l⁻¹, and 60 ml l⁻¹ were not significantly different from the treatment of inorganic fertilizer (Table 4), while ear length between doses of 25 l ha⁻¹ and inorganic fertilizer did not show any significant differences (Table 5). The ear length between doses of 50 l ha⁻¹, 75 l ha⁻¹, and 100 l ha⁻¹ did not show any significant differences (Table 5).

The application of LOF increased the ear diameter (Tables 4 and 5). The lowest ear diameter was recorded in control plants (Table 4). The diameter of the ear between

¹ concentrations of 30 ml l⁻¹, 45 ml l⁻¹, 60 ml l⁻¹ (Table 4) as well as between doses of 75 l ha⁻¹ and 100 l ha⁻¹ (Table 5) were not significantly different.

The application of LOF consistently increased sweet corn yields (Tables 6 and 7). The yield of sweet corn at all LOF concentrations (Table 6) was not significantly different. The lowest yield was recorded in control treatment. The highest yield was obtained at a concentration of 60 ml l⁻¹ (Table 6) and dosed 100 l ha⁻¹ (Table 7). The application of LOF increased the dry shoot weight (Tables 6 and 7). ¹⁸ The lowest shoot dry weight was noted in control treatment (Table 6).

Application of LOF increased plant nutrient uptake (Table 8). Nutrient uptake of P and K in the LOF treatment were significantly higher than those treated with inorganic fertilizer. Also, nutrient uptake of N, P, and K between doses of 25 l ha⁻¹ and 50 l ha⁻¹ as well as between doses 75 l ha⁻¹ and ¹³ 100 l ha⁻¹ did not show any significant differences (Table 8).

The LOF application increased the quality of sweet corn as indicated by sugar content (Tables 6 and 7) as indicated by °Brix value. The lowest sugar content was recorded in control treatment (Table 6). The sugar content between all concentrations of LOF (Table 6) and between doses of 75 l ha⁻¹ and 100 l ha⁻¹ (Table 7) were not significantly different.

Discussion

Vegetative growth of plants is represented by the number of the parameters of leaves, stem diameter, and plant height. The highest number of leaves and stem diameter were seen concentrations of 45 ml l⁻¹ and 60 ml l⁻¹ in experiment 1. At the beginning of growth, plants can absorb nutrients from LOF, which are used by plants to support growth. Nitrogen (Table 1) is needed for vegetative growth. Nitrogen is a key element for producing vegetative biomass on the early growth of maize (Massignam et al., 2009).

The fact that LOF increased the leaf area index might be due to an increase in the number of leaves (Table 3). ⁶ Fageria et al. (2009) stated that foliar application requires higher LAI for absorbing applied nutrient solution in sufficient amounts. Table 3 showed that the application of LOF at a dose of 100 l ha⁻¹ gave a higher value of leaf greenness compared to the other treatments. According to Nugroho (2015) nitrogen, in addition to stimulating plant growth, also gave the green color of the leaves. The darker the green color of leaves on corn plants, the higher the amount of nitrogen absorbed by the plants.

The application of LOF gave positive results in terms of the generative growth of plants. The results showed that the concentration of 60 ml l⁻¹ gave the best results in terms of the weight of unhusked ears and husked ears, and the treatment with a dose of 100 l ha⁻¹ gave the highest ear diameter and the heaviest weight of the ears. It can be inferred that the higher the dose or concentration of LOF given, the higher the generative growth of plants. According to Zafar et al. (2013), enlargement of ear diameter is related to the availability of phosphorus; phosphorus greatly influences the formation of ears.

Liquid organic fertilizer application increased the yield of sweet corn. This means that the contribution of nutrients from LOF can be complementary to the recommended solid fertilizer. The results of this study are supported in various other LOF studies. Minardi et al. (2015) found that banana corm extract increased the available P in soil. Aini et al. (2017) stated that the treatment of banana humps had a significant effect on the growth and yield of soybean. Working with *Gleichenia linearis* plant extract, Aulya et al. (2018) showed that the application of 100 mg l⁻¹ was the most effective concentration in increasing plant height and leaf area of maize compared to a control.

The results showed that the treatment of a dose of LOF 100 l ha⁻¹ gave a high degree of N, P, and K nutrient uptake, and this was attributed to the adequate supply of nutrients from LOF. This means that the higher the dose of LOF application, the higher the nutrient uptake of N, P, and K; in contrast to Muktamar et al. (2016) showed that increase in rates of LOF significantly raised nitrogen uptake by sweet corn but did not raise phosphorus and potassium uptake. Mukuralinda et al. (2010) also showed high P nutrient uptake due to the application of organic fertilizer compared to control.

The application of N nutrients in plants has a direct role in the formation of amino acids, proteins, nucleic acids, enzymes, nucleoproteins, and alkaloids (Mokhele et al., 2012), which are needed for the vegetative growth process of plants and increase the greenness of leaves. Leaf N nutrient uptake has a close relationship with the leaf greenness level, characterized by the SPAD value. Increasing nutrient uptake of P enhanced the metabolic processes in the plant to become more active. P is known as the forming element of ATP for energy sources. In addition to P, K also plays an important role in improving the quality of plant fruit. The application of liquid organic fertilizer containing an element of K from coconut fiber extract improved the quality of sweet corn, which is characterized by a higher Brix value. Jifon and Lester (2009) found that the application of K fertilizer through

leaves increased the sugar content of muskmelon plants. Sweet taste in sweet corn involves potassium nutrients, which play a role in the activation of many enzymes that has a role in metabolic processes in plants. Enzymes that play a role in sugar synthesis are activated by K (Prajapati & Modi, 2012).

The research showed that in both experiments, between the all concentration of the LOF and inorganic recommended fertilizers (experiment 1) as well as between the doses of the LOF and inorganic recommended fertilizers (experiment 2) gave the same effect on all parameters of growth and yield of sweet corn. Sweet corn plants fertilized with LOF had the same growth and yield with those of recommended fertilizers. Therefore, it implied that LOF could be a complementary fertilizer. Many researchers have also succeeded in applying the foliar application of LOF to sweet corn (Muktamar et al., 2017) and corn (Aulya et al., 2018).

From this experiment, it can be recommended to utilize local plant resources around farmers such as *lamtoro* leaves, banana humps, and coconut fibers to be used as LOF. The high content of N, P, K nutrients in LOF (Table 1) means that the application of liquid organic fertilizer is sufficient to meet the needs of N, P, and K nutrients for sweet corn plants. The mixture of plant residues contained the basic nutrient requirements for plants. *Lamtoro* leaves have a high content of protein, carbohydrate, and nitrogen (Devi et al., 2013). Banana humps contained P and K (Bahtiar et al., 2016). *Azospirillum*, *Azotobacter*, *Bacillus* sp., *Aeromonas* sp., and *Aspergillus niger* and the other phosphate solubilizing bacteria were identified in banana humps (Minardi, et al., 2015; Suhastyo et al., 2013). Banana humps naturally had bioactive compounds to be used as plant growth regulators (Ulfa et al., 2013), namely auxin, gibberellin and cytokinin. Coconut fiber had a high P and K (Abad et al., 2002). It can be said that the application of LOF supplied nutrients, microbes, and hormones leading to better growth, yield and quality of sweet corn.

The use of liquid organic fertilizer from plant materials can be used as a complementary fertilization technology in developing organic agriculture. However, the nutrient content of plant extract is dependent on the type of organic waste used, the fermentation period, and the storage of plant extract. In its application, liquid organic fertilizer needs to be mixed with a surfactant to increase the efficiency of uptake by plant leaves. The application of liquid organic fertilizer is an inexpensive and effective technology in organic farming. Therefore, local-based liquid organic fertilizers that contain extracts of *lamtoro*, banana hump, and coconut fiber have the potential to support organic

farming. Further research is still needed to determine the appropriate concentration and doses of liquid organic fertilizer for each horticultural and food crop.

CONCLUSIONS

Application of LOF with the concentration of 60 ml l⁻¹ or dose 100 l ha⁻¹ showed the best growth, yield, and quality of sweet corn compared to lower concentrations or lower doses. Application of LOF improved the agronomic quality of sweet corn. Sweet corn sprayed with LOF absorbed a higher amount of N, P, and K. It is recommended that LOF could be a complementary fertilization technology in sweet corn organic farming.

ACKNOWLEDGEMENT

This research was funded by the Research Grant 2017, Faculty of Agriculture, University of Lampung. The authors would like to thank to Rugayah and Y. C. Ginting for scientific discussions during manuscript preparation.

REFERENCES

- Abad, M., Noguera, P., Puchades, R., Maqueira, A., & Noguera, V. (2002). Physico-chemical and chemical properties of some coconut coir dusts for use as a peat substitute for containerised ornamental plants. *Bioresource Technology*, 82(3), 241-245. doi: 10.1016/S0960-8524(01)00189-4
- Aini, D. N., Sugiyanto, B., & Herlinawati. (2017). Aplikasi mikroorganisme lokal bonggol pisang dan pupuk kandang kambing terhadap produksi kedelai (*Glycine max* L. Merrill) varietas Baluran [Application of local microorganism derived from stem of banana and goat manure on the yield of soybean (*Glycine max* L. Merrill) var. Baluran]. *Agriprima, Journal of Applied Agricultural Sciences*, 1(1), 35-43. doi: 10.25047/agriprima.v1i1.13
- Amanullah, Kakar, K. M., Khan, A., Khan, I., Shah, Z., & Hussain, Z. (2014). Growth and yield response of maize (*Zea mays* L.) to foliar NPK-fertilizers under moisture stress condition. *Soil Environment*, 33(2), 116-123.
- Arsyad, M., Wardana, I. N. G., Pratikto, & Irawan, Y. S. (2015). The morphology of coconut fiber surface under chemical treatment. *Revista Materia*, 20(1), 169-177.
- Astuti, T., Amir, Y. S., Yelni, G., & Isyaturriyadhah (2014). The result of biotechnology by local microorganisms to banana peel on rumen fluid characteristics as ruminant feed. *Journal of Advanced Agricultural Technologies*, 1(1), 28-31. doi:10.12720/joaat.1.1.28-31
- Aulya, N. R., Noli, Z. A., Bakhtiar, A., & Mansyurdin. (2018). Effect of plant extracts on growth and yield of maize (*Zea mays* L.). *Pertanika Journal of Tropical Agricultural Sciences*, 41(3), 1193-1205.

- Bahtiar, S. A., Muayyad, A., Ulfaningtias, L., Anggara, J., Priscilla, C., & Miswar. (2016). Pemanfaatan kompos bonggol pisang (*Musa acuminata*) untuk meningkatkan pertumbuhan dan kandungan gula tanaman jagung manis (*Zea mays* L. *saccharata*) [Compost use banana weevil (*Musa acuminata*) to boost growth and content of sugar sweet corn (*Zea mays* L. *saccharata*)]. *Agritrop, Journal of Agricultural Science*, *14*(1), 18-22.
- Devi, M. V., Ariharan, V., & Prasad, P. N. (2013). Nutritive value and potential uses of *Leucaena leucocephala* as biofuel - A mini review. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, *4*(1), 515-521.
- Fageria, N. K., Filho, M. P. B., Moreira, A., & Guimarães, C. M. (2009). Foliar fertilization of crop plants. *Journal of Plant Nutrition*, *32*(6), 1044–1064. doi:10.1080/01904160902872826
- Fahrurrozi, Mukhtar, Z., Dwatmadji, Setyowati, N., Sudjtmiko, S., & Chozin, M. (2016). Growth and yield responses of three sweet corn (*Zea mays* L. var. *saccharata*) varieties to local-based liquid organic fertilizer. *International Journal on Advanced Science Engineering and Information Technology*, *6*(3), 319-323.
- Farinu, G. O., Ajiboye, S. O., & Ajao, S. (1992). Chemical composition and nutritive value of leaf protein concentrate from *Leucaena leucocephala*. *Journal of the Science of Food and Agriculture*, *59*(1), 127-129. doi: 10.1002/jsfa.2740590119
- Hartz, T. K., Mitchell, J. P., & Giannini, C. (2000). Nitrogen and carbon mineralization dynamics of manures and composts. *HortScience*, *35*(2), 209–212. doi: 10.21273/HORTSCI.35.2.209
- Hartz, T. K., Smith, R., & Gaskell, M. (2010). Nitrogen availability from liquid organic fertilizers. *HortTechnology*, *20*(1), 169-172. doi: 10.21273/HORTTECH.20.1.169
- Jifon, J. L., & Lester, G. E. (2009). Foliar potassium fertilization improves fruit quality of field-grown muskmelon on calcareous soils in south Texas. *Journal of the Science of Food and Agriculture*, *89*(14), 2452–2460. doi: 10.1002/jsfa.3745

- Johnson, H. J., Colquhoun, J. B., Bussan, A. J., & Laboski, C. A. M. (2012). Estimating nitrogen mineralization of composted poultry manure, organic fertilizers, and green manure crops for organic sweet corn production on a sandy soil under laboratory conditions. *HortTechnology*, 22(1), 37-43. doi: 10.21273/HORTTECH.22.1.37
- Massignam, A. M., Chapman, S. C., Hammer, G. L., & Fukai, S. (2009). Physiological determinants of maize and sunflower grain yield as affected by nitrogen supply. *Field Crops Research*, 113(3), 256-267. doi: 10.1016/j.fcr.2009.06.001
- Meulen, U., Struck, S., Schulke, E., & Harith, E. A. E. (1979). A review on the nutritive value and toxic aspects of *Leucaena leucocephala*. *Tropical Animal Production*, 4(2), 113-126.
- Minardi, S., Hartati, S., Widijanto, H., & Wulandari, D. (2015). The use of extract banana corm and phosphate rock to increase available-P in Alfisols. *Sains Tanah, Journal of Soil Science and Agroclimatology*, 12(2), 76-81. doi: 10.15608/stjssa
- Mokhele, B., Zhan, X., Yang, G., & Zhang, X. (2012). Nitrogen assimilation in crop plants and its affecting factors. *Canadian Journal of Plant Science*, 92(3), 399-405. doi: 10.4141/CJPS2011-135
- Muktamar, Z., Fahrurrozi, Dwatmadji, Setyowati, N., Sudjtmiko, S., & Chozin, M. (2016). Selected macronutrients uptake by sweet corn under different rates liquid organic fertilizer in closed agriculture system. *International Journal on Advanced Science Engineering and Information Technology*, 6(2), 258-261.
- Muktamar, Z., Sudjtmiko, S., Chozin, M., Setyowati, N., & Fahrurrozi. (2017). Sweet corn performance and its major nutrient uptake following application of vermicompost supplemented with liquid organic fertilizer. *International Journal on Advanced Science Engineering and Information Technology*, 7(2), 602- 608.
- Mukuralinda, A., Tenywa, J. S., Verchot, L., Obua, J., Nabahungu, N. L., & Chianu, J. N. (2010). Phosphorus uptake and maize response to organic and inorganic fertilizer

- inputs in Rubona, Southern Province of Rwanda. *Agroforestry Systems*, 80(2), 211-221. doi:10.1007/s10457-010-9324-9
- Nugroho, W. S. (2015). Penetapan standar warna daun sebagai upaya identifikasi status hara (N) tanaman jagung (*Zea mays* L.) pada tanah Regosol [Determination of a standard of green color on corn (*Zea mays* L.) leaves as the marker of nitrogen adequacy on Regosol soil]. *Planta Tropika, Journal of Agro Science*, 3(1), 8-15. doi: 10.18196/pt.2015.034.8-15
- Prajapati, K., & Modi, H. A. (2012). The importance of potassium in plant growth - A review. *Indian Journal of Plant Sciences*, 1(2), 177-186.
- Suhastyo, A. A., Anas, I., Santosa, D. A., & Lestari, Y. (2013). Studi mikrobiologi dan sifat kimia mikroorganisme lokal (MOL) yang digunakan pada budidaya padi metode sri (System of Rice Intensification) [Studies of microbiology and chemical properties of the local microorganisms (MOL) used in rice farming by SRI method (System of Rice Intensification)]. *Sainteks*, 10(2), 29-39.
- Ulfa, F., Sengin, E. L., Baharuddin, Syaiful, S. A., Sennang, N. R., Rafiuddin, ... Ifayanti. (2013). Potential of plant extracts as growth exogenous regulators of potato seeds. *International Journal of Agriculture Systems*, 1(2), 98-103.
- Zafar, M., Abbasi, M. K., & Khaliq, A. (2013). Effect of different phosphorus sources on the growth, yield, energy content and phosphorus utilization efficiency in maize at Rawalakot Azad Jammu and Kashmir Pakistan. *Journal of Plant Nutrition*, 36(12), 1915-1934. doi:10.1080/01904167.2013.819892
- Zhai, Z., Ehret, D. L., Forge, T., Helmer, T., Lin, W., Dorais, M., & Papadopoulos, A. P. (2009). Organic fertilizers for greenhouse tomatoes: Productivity and substrate microbiology. *HortScience*, 44(3), 800-809. doi: 10.21273/HORTSCI.44.3.800

Liquid Organic Fertilizer from Plant Extracts Improves the Growth, Yield and Quality of Sweet Corn (*Zea mays* L. var. *saccharata*)

ORIGINALITY REPORT

11%

SIMILARITY INDEX

PRIMARY SOURCES

1	jchr.org Internet	42 words — 1%
2	media.neliti.com Internet	34 words — 1%
3	doaj.org Internet	29 words — 1%
4	arhiva.nara.ac.rs Internet	28 words — 1%
5	Faysal Fayez Eliyan, Akram Alfantazi. "Electrochemical investigations on the corrosion behavior and corrosion natural inhibition of API-X100 pipeline steel in acetic acid and chloride-containing CO ₂ -saturated media", <i>Journal of Applied Electrochemistry</i> , 2012 Crossref	26 words — 1%
6	krshikosh.egranth.ac.in Internet	14 words — < 1%
7	agroteknologi.fp.uns.ac.id Internet	14 words — < 1%
8	Yang Zhang, Karel Ghyselbrecht, Boudewijn Meesschaert, Luc Pinoy, Bart Van der Bruggen. "Electrodialysis on RO concentrate to improve water recovery in wastewater reclamation", <i>Journal of Membrane Science</i> , 2011 Crossref	13 words — < 1%

-
- 9 C. C.G. St. Martin, R. A.I. Brathwaite. "Compost and compost tea: Principles and prospects as substrates and soil-borne disease management strategies in soil-less vegetable production", *Biological Agriculture & Horticulture*, 2012
Crossref 12 words — < 1%
-
- 10 L M Mooy, A Hasan, R Onsili. "Growth and yield of Tomato as influenced by the combination of liquid organic fertilizer concentration and branch pruning", *IOP Conference Series: Earth and Environmental Science*, 2019
Crossref 12 words — < 1%
-
- 11 chimie-biologie.ubm.ro
Internet 12 words — < 1%
-
- 12 ir.kagoshima-u.ac.jp
Internet 11 words — < 1%
-
- 13 Rafael Roehrig, Walter Boller, Carlos A. Forcelini, Amanda Chechi. "USE OF SURFACTANT WITH DIFFERENT VOLUMES OF FUNGICIDE APPLICATION IN SOYBEAN CULTURE", *Engenharia Agrícola*, 2018
Crossref 11 words — < 1%
-
- 14 pscipub.com
Internet 11 words — < 1%
-
- 15 Arsyad, Muhammad, I Nyoman Gede Wardana, Pratikto, and Yudy Surya Irawan. "The morphology of coconut fiber surface under chemical treatment", *Matéria (Rio de Janeiro)*, 2015.
Crossref 11 words — < 1%
-
- 16 bioinfopublication.org
Internet 11 words — < 1%
-
- 17 Pavel Zhalnerchyk, Andrzej A. Przybyła, Franciszka Jaumień. "Influence Of Chemicals Of Arbolin Group On Branching Of Maiden Trees Of Three Apple Cultivars", *Journal of Horticultural Research*, 2015
Crossref 10 words — < 1%

18 Mohammad Kazem Souri, Farzaneh Yaghoubi Sooraki. "Benefits of organic fertilizers spray on growth quality of chili pepper seedlings under cool temperature", Journal of Plant Nutrition, 2019
Crossref 10 words — < 1%

19 scholarsresearchlibrary.com
Internet 10 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF