

PAPER NAME

**2018EN_PROVISION OF LOW-DOSE-INT.
pdf**

WORD COUNT

4484 Words

CHARACTER COUNT

25034 Characters

PAGE COUNT

6 Pages

FILE SIZE

359.8KB

SUBMISSION DATE

Jul 28, 2022 3:51 PM GMT+7

REPORT DATE


Jul 28, 2022 3:53 PM GMT+7**● 16% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 15% Internet database
- 4% Publications database
- Crossref database
- Crossref Posted Content database
- 1% Submitted Works database

● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded text blocks



1 PROVISION OF LOW-DOSE PROPRANOLOL IN PATIENTS WITH ACUTE ISCHEMIC STROKE AGAINST THE LEVELS OF TNF-A, IL-10, AND THE RATIO OF TNF-A / IL-10 AT “ABDUL MOELOEK” GENERAL HOSPITAL, BANDAR LAMPUNG, INDONESIA

Roezwir Azhari¹, Ellyza Nasrul², Al Rasyid³, Adnil Edwin⁴ and Endang Nurcahyani*⁵

¹Department of Neurology, “Abdul Moeloek” General Hospital, Bandar Lampung, Indonesia.

²Department of Clinical Pathology, Faculty of Medicine, University of Andalas, Padang, Indonesia.

³Department of Neurology, Cipto Mangunkusumo, National General Hospital, Jakarta, Indonesia.

⁴Department of Psychiatry, Faculty of Medicine, University of Andalas, Padang, Indonesia.

⁵Dept. of Biology, Faculty of Mathematics and Natural of Science, University of Lampung, Bandar Lampung, Indonesia.

³*Corresponding Author: Endang Nurcahyani

Dept. of Biology, Faculty of Mathematics and Natural of Science, University of Lampung, Bandar Lampung, Indonesia.

⁴Article Received on 30/10/2017

Article Revised on 21/11/2017

Article Accepted on 12/12/2017

1 ABSTRACT

Brain and the immune system are linked through sympathetic, parasympathetic nervous system and HPA axis. Relationship between brain and immune system is thought to be through molecules secreted by the necrotic brain tissues. Immune system alterations in ischemic stroke can be seen with the presence of lymphopenia, immediate increment and followed by decrement of proinflammatory cytokines (TNF- α) and elevated anti-inflammatory cytokines (IL-10). According to animal study, this condition can be controlled by giving propranolol. The aim of this study was to demonstrate the effect of 2x10 mg propranolol on TNF- α , IL-10, and TNF- α /IL-10 ratio in acute ischemic stroke patients with lymphopenia. This is an experimental research with pretest-posttest match control group study design. Samples used in this study were 40 ischemic stroke patients with lymphopenia. The study was conducted in “Abdul Moeloek” General Hospital, Bandar Lampung, Indonesia from July, 2016 to February, 2017. Patients were divided into 2 groups; ischemic stroke group without propranolol, and ischemic stroke group with propranolol. TNF- α and IL-10 levels were examined by ELISA. The results showed that reduction of TNF- α levels in the group treated with propranolol was lower (p=0.221). IL-10 levels increased in the group without propranolol (p=0.547). TNF- α /IL-10 ratio in the group with propranolol was higher (p=0.053). Ischemic stroke affects the patient's immune system that show in alteration of TNF- α , IL-10 levels and ratio of TNF- α /IL-10. Provision of propranolol 2x10 mg increase TNF- α level, decrease IL-10 level and increase ratio TNF- α /IL-10, but not significant statistically.

KEYWORDS: Ischemic Stroke, Propranolol, IL-10, TNF- α , TNF- α /IL-10 Ratio.

INTRODUCTION

Stroke has become a global public health problem, killing millions people and causing major disabilities worldwide.^[1] Until now ischemic stroke therapy has not been satisfactory due to various factors and the complexity of pathophysiology.^[7] Food and Drug Administration (FDA) approved therapy is rTPA (recombinant Tissue Plasminogen Activator). However, this therapy is limited due to the short time for intervention, only up to 4.5 hours post-stroke onset. In addition, the strict criteria of inclusion and exclusion results in the few patients who are able to receive rTPA treatment even in the United States are less than 5% of patients meet the criteria and are able to get the treatment.^[2] Treatment with rTPA is quite effective in reopening the clogged blood vessels, but increasing the

risk of cerebral hemorrhage and reperfusion itself induces other pathological processes, because the entry of circulating immune cells into brain tissue exacerbates inflammation and damage.^[3]

In the early stroke phase there was an increase in Tumor Necrosis Factor- α (TNF- α) due to excessive monocyte activation followed by a decrease due to monocyte fatigue.^[4] TNF- α is a potent proinflammatory cytokine. Produced mainly by active macrophages. Other cells can also produce these cytokines such as CD4 + lymphocytes, NK cells, neutrophils, mast cells, eosinophils and neurons.^[5] TNF- α plays an important role during brain ischemia, disrupts endothelial function, induces thrombogenesis through elevated levels of Plasminogen Activator Inhibitor (PAI-1) and Platelet

Activating Factor (PAF) and inhibits endogenous TPA activity. Some apoptotic pathways and neuronal damage are also caused by elevated levels of NF- α [6]

Interleukin 10 (IL-10) is an anti-inflammatory cytokine. This substance decreases expression of Th1 cytokines, macrophages, increases B cell survival, proliferation, and antibody production. [42] In mice with B cell deficiency, the infarct volume increasingly spread, this suggests the role of IL-10 in the regulation of the post-stroke immune system. IFN- γ production by Th1, T cell proliferation and cytokine response, TNF- α production by monocytes and macrophages, are all inhibited by IL-10. [7] The TNF- α /IL-10 ratio reflects a balance between Th1 and Th2 proinflammation and cellular immunity versus anti-inflammatory and humoral immunity. In patients with ischemic stroke, there was a decrease in the ratio of TNF- α /IL-10. [8,9]

Inoculation 200 cfu *S. pneumoniae* to MCAO mice can cause severe pneumonia, whereas in control mice 200,000 cfu are required to obtain the same severity of infection. [10] It has been shown that there was a post-stroke immunosuppression in animals and cerebral arterial media occlusion cause spontaneous infection in mice. [11] Similar infections may also be observed in stroke patients. [12,13] The suppression of the immune system by the nervous system actually protects the brain from further inflammatory damage, but at the same time increases susceptibility to infection. [14] Infection is a complication of 30% of patients with acute stroke. [15] Post-stroke infection is directly correlated with the severity of stroke (high NIHSS value), and extent of infarction. [16] Post-stroke infections prolong the duration of treatment and 80% are due to pneumonia and urinary tract infections. [17] Pneumonia in stroke patients will increase 3 times mortality within 30 days and worse clinical outcomes. [18]

Propranolol is a beta blocker, used to treat hypertension, irregular heart rhythm, thyrotoxicosis, anxiety and tremor. These drugs include a class of non-cardioselective sympatholytic beta blockers that can penetrate the blood-brain barrier. As non-selective beta blockers, drugs work to inhibit epinephrine and norepinephrine at β 1- and β 2- receptors. [19] Propranolol 2x10 mg can overcome the symptoms of tachycardia, hyperhidrosis, hypertension and hyperthermia, caused by sympathetic hyperactivity in post-stroke patients. [20]

The production of TNF- α is regulated by the α 2- and β -adrenoreceptors present in the terminal adrenergic nerves and macrophages. Using α 2- and β -adrenergic drugs (clonidine, CH-38083, isoproterenol, propranolol), it was found that α 2- and β - adrenoreceptors were involved in the production of TNF- α induced by lipopolysaccharide (LPS). [21] In addition to inducing an anti-apoptotic pathway, propranolol therapy significantly increases the activity of TNF- α and interferon γ (IFN γ). [22] In macrophages, endogenous norepinephrine inhibits the

ability of phagocytosis to limit its immune system activity. In addition, norepinephrine decreases the cytotoxicity activity of natural killer (NK) cells, expansion of CD8 + T cells, and increases T helper 2 response. [23]

MATERIAL AND METHODS

This study was an experimental study with pretest-postes control group, conducted from July, 2016 to February, 2017. Subjects consisted of 40 ischemic stroke patients with chronic hypertension and lymphopenia who fulfilled the inclusion and exclusion criteria at "Abdul Moeloek" General Hospital, Bandar Lampung, Indonesia. Subjects are divided into two groups with the same characteristics (match control group). The first group consisted of 20 ischemic stroke patients not given propranolol. The second group consisted of 20 ischemic stroke patients given propranolol 2x10 mg. Diagnosis of ischemic stroke is confirmed through CT-Scan head examination performed maximum until the third day. The clinical weight of stroke was measured using NIHSS. Examination of TNF- α and IL-10 levels was performed on the fourth day using ELISA method using Quantikine® HS Human Immunoassay, R & D Systems, specifically for research by Prodia Laboratory, Jakarta, Indonesia.

This study has passed the ethical test from the ethics committee of The Faculty of Medicine at University of Andalas, Padang, Indonesia. Participants who participated in the study have signed an informed consent form. Propranolol 2x10 mg was inspired by a case report by [20] given after the first blood sample was taken. Seven days later re-sampled blood. The TNF- α /IL-10 ratio was obtained by comparing TNF- α and IL-10 levels. Statistical tests were performed by comparing the TNF- α , IL-10 deltas and the pretest-postest TNF- α /IL-10 delta ratios between groups using independent T tests. TNF- α and IL-10 were tested using the Mann-Whitney test. Differences in levels of TNF- α , IL-10, and TNF- α /IL-10 ratios were found to be significant when $p < 0.05$.

RESULTS AND DISCUSSION

Provision of Low Dose Propranolol In Acute Ischemic Stroke Patients Against TNF- α level, IL-10 level, and TNF- α /IL-10 Ratio are presented in Table 1, Table 2, and Table 3.

Table 1: TNF- α level.

	TNF- α without Propranolol (Mean \pm SD)	TNF- α with Propranolol (Mean \pm SD)	P
Pretest	8,300 \pm 6,070	6,769 \pm 5,920	
Post Test	5,380 \pm 3,740	5,953 \pm 3,900	
Δ TNF- α	2,920 \pm 4,870	0,816 \pm 7,370	0,221

Table 1 shows that a decrease in TNF- α levels of more groups not given propranolol 2x10 mg but, the difference **Table 2: IL-10 level.**

	IL-10 without Propranolol (Mean \pm SD)	IL-10 with Propranolol (Mean \pm SD)	P
Pretest	0,702 \pm 0,60	0,761 \pm 0,39	
Post Test	0,884 \pm 1,22	0,663 \pm 0,62	
Δ IL-10	0,182 \pm 1,25	0,098 \pm 0,64	0,547

Based on Table 2., in groups not given propranolol, an increase in IL-10 levels was found. The group given propranolol 2x10 mg was decreased. However, the

in TNF- α levels decreased in both groups was not statistically significant ($p > 0.05$).

difference in IL-10 between groups was not statistically significant ($p > 0.05$).

Table 3: TNF- α /IL-10 Ratio.

	TNF-α/IL-10 Rratio Without Propranolol (Mean \pm SD)	TNF-α/IL-10Ratio with Propranolol (Mean \pm SD)	P
Pretest	14,79 \pm 1,24	9,73 \pm 6,56	
Posttest	10,81 \pm 8,13	19,97 \pm 3,06	
Δ TNF α /IL10	-3,98 \pm 1,24	+10,24 \pm 2,92	0,053

Table 3 presented that The TNF- α / IL-10 ratio in the group without propranolol decreased, whereas propranolol was elevated, indicating the effect of propranolol in improving anti-inflammatory conditions in ischemic stroke patients. However, the difference in TNF- α / IL-10 ratio between groups was not statistically significant ($p > 0.05$).

Increased proinflammatory cytokines (TNF- α and IL-1 β) post-stroke causes sympathetic nerve activation and HPA axis,^[24] further increased norepinephrine release from sympathetic nerve endings and glucocorticoids from the adrenal cortex and increased epinephrine and norepinephrine release from medulla adrenal.^[12] All mentioned above cause immunosuppression conditions characterized by decreased levels of TNF- α and elevated levels of IL-10^[25,26,12] In addition,^[27] have described clearly that T cells in the spleen (T-ChAT cells) expressing β 2AR will be activated by norepinephrine released at the splenic nerve to produce acetylcholine. Acetylcholine binds to the α 7nACh receptor, consequently the production of proinflammatory cytokines by macrophages is inhibited. In the liver, norepinephrine (NE) released by the sympathetic nerve endings changes the iNCT cell response that initially produces IFN- γ (Th1) changes producing IL-10 (Th2). It is clear that NE can change the production of cytokines from Th1 to Th2.^[27] Another mechanism occurs through the bonding of acetylcholine with the α 7nACh receptor which inhibits I κ B phosphorylation so that NF κ B does not undergo translocation to the nucleus, consequently the production of proinflammatory cytokines is inhibited. While other pathways are suspected through JAK2 pathways that activate STAT3 monomer changes to STAT3 dimer and into NF κ B binding nuclei so that NF κ B is inactivated.^[28]

Propranolol is a non-selective adrenergic beta blocker, which can decrease sympathetic nervous activity in both β 1 and ad adrenergic receptors^[19] The β -adrenergic receptor can regulate the production of TNF- α through catecholamines. Propranolol therapy causes blockade of NE bond with β AR, TNF- α production is not suppressed, and IL-10 is reduced so that systemic immunosuppression that occurs after stroke is reduced.^[11,10] The results of the research are not significant, the explanation is as follows. The path is inhibited only the sympathetic nerve pathway, while the HPA axis path is not. Glucocorticoids may inhibit the expression of chemoattractant and pro-inflammatory cytokines such as IL-1 β , IL-6 and TNF- α .^[29] In this study, the first TNF- α examination was performed on day 4, and the second day-to-11 post stroke examination. In this phase there has been a decrease in TNF- α levels due to post-stroke biphasic response.^[30] Within 1 week of therapy, the mean TNF- α level was lowered from 6.769 \pm 5.92 (pretest) to 5.953 \pm 3.9 (posttest). This shows that the immunosuppression process continues. At a dose of 2x10 mg, propranolol can overcome the symptoms of tachycardia, hyperhidrosis, and hyperthermia, caused by sympathetic hyperactivity in post stroke patients,^[20] but can not overcome post stroke immunosuppression.

Overcoming post stroke immunosuppression in experimental animals, determined by the dose of propranolol administered. The higher the dose of propranolol, the stronger inhibitory effect on immunosuppression.^[11] At a dose of 80 mg, BEST found poor patient outcomes due to decreased cerebral blood flow. However, the dose was administered from the first day of stroke.^[30,31] used a dose of 160 mg but his research was stopped without a clear explanation. Even though there is no definite dose of propranolol to treat

immunosuppression in stroke patients, it appears that a dose of 2×10 mg is not sufficiently strong to overcome severe immunosuppression. While high doses increase the risk of death especially when given in the early stroke phase.^[30]

The results were not significant in this study may also be due to differences in age range between research subjects too far (44 years-87 years).^[32] examined the effect of age with TNF levels. In the study, high levels of TNF α were found in 81-year-old subjects compared to young and healthy controls. However, in the study of^[43] stated that in older rats the production of TNF α in macrophages decreased compared with young rats. In another study,^[33] suggested that postmenopausal women had higher levels of TNF α than did younger controls.^[34] also stated the same thing. TNF α in menopausal women is higher than in women of reproductive age. It is influenced by estrogen hormones that decrease at the age of menopause. The hormone estrogen can suppress the production of inflammatory cytokines such as IL-1 β , IL-6 and TNF α . In addition, estrogen affects the Th1/Th2 balance with stimulation dominance on Th2. Meanwhile, according to^[35] levels of TNF α also showed an increase in men with decreased testosterone, and decreased testosterone occurred in elderly men. Thus, it can be concluded that TNF α levels are equally high in women and older men.

Based on the above discussion, the results of this study are supported by many existing literatures, in which TNF- α changes are not significant although in the propranolol therapy group. Other risk factors that contribute to changes in the immune system are not examined further. A considerable age range among study subjects may also be influential. In the group not given propranolol there was an increase in IL-10 levels. In the group given propranolol there was a decrease in IL-10 levels. Differences in IL-10 levels in both groups were not statistically significant ($P > 0.05$).

Patients at risk for stroke-related infection were patients with high IL-10 levels (≥ 14.5 pg / mL) with different KIT ELISA, low TNF- α levels and low TNF- α / IL-10 ratios.^[9] Because propranolol can inhibit the NE bond with β AR, propranolol therapy may reduce IL-10 levels as an anti-inflammatory cytokine.^[25,12] In this study, one patient died of severe pneumonia, where the level of IL-10 in second examination increased more than 17x compared to the first examination (data not shown). It can not be concluded whether infection is caused by high levels of IL-10 or an infection that causes elevated levels of IL-10.

Many studies support the theory above. Some of them are as follows. Oxprenolol therapy (β receptor antagonists) *in vivo* in LPS-induced rats, significantly reducing serum IL-10 levels.^[36] Propranolol may decrease elevated levels of IL-10 due to norepinephrine stimulation in THP-1 cells.^[37] High levels of IL-10 were

found in stroke-related infections.^[12,9] The results of this study indicate that in the group whom not given propranolol still increased in levels of IL-10 means still decrease immune function. While in the second group, the level decreased in the second examination within 1 week interval. Although not significant, the increase of IL-10 in the first group was consistent with existing theories, that during the immunosuppression period there was an increase in IL-10 levels,^[25,26,12] and decreased levels of IL-10 after provision of propranolol according to the theory of,^[28,27] blockade of norepinephrine bonds with β -adrenergic receptors by propranolol will also decrease the immunosuppression response through decreased levels of IL-10 due to suppression of Treg cells.^[37]

Research about the association levels of IL-10 and age is still inconclusive. In the study of,^[43] IL-10 production in rat macrophages was not affected by age. These results are in contrast to the study of^[44] which suggests that in women of menopausal age the levels of IL-10 increase.^[38] also stated the same thing. Levels of IL-10 in menopausal women increase, but in men of the same age did not increase.

It is recalled that the HPA pathway is not inhibited where glucocorticoids may increase the expression of anti-inflammatory cytokines such as IL-10 and TGF- β .^[29] Other risk factors that contribute to changes in the immune system are not examined further. A considerable age range among study subjects may also be influential. The TNF- α / IL-10 ratio in the first group decreased while the second group increased. The ratio difference was not statistically significant ($p > 0.05$).

Under normal circumstances, there is a balance between pro- and anti-inflammatory cytokines. This balance will be impaired when there is a pathological process in the brain, and potentially determine the ability of subsequent immunological responses.^[9,26] This TNF- α /IL-10 ratio, being the balance reference between Th1 and Th2 (proinflammatory and cellular immunity vs. antiinflammatory and humoral immunity). In stroke patients this ratio changed, even this decrease in ratio was already seen 1 day after onset of ischemic stroke.^[39]

In this study, it was found that TNF- α / IL-10 in the group without propranolol decreased, whereas propranolol was increased. Although not significant in both groups, the TNF- α / IL-10 ratio decrease in the first group occurred due to immunosuppression due to sympathetic system activation.^[25,40,12] The increase in TNF- α / IL-10 ratio in the second group (propranolol therapy) was due to propranolol therapy inhibiting norepinephrine bond with β AR.^[37,28,27]

The study (retrospectively) by^[45] showed that beta-blockers in the acute phase of ischemic stroke showed a decrease in the incidence of infection and a 3-month mortality rate compared with ischemic stroke patients

who did not receive beta-blockers.^[45] investigated the administration of propranolol in the infarct stroke model and found that propranolol may protect experimental animals from immunosuppression and life-threatening infections, despite an increase in specific autoreactivity of brain antigens, but did not exacerbate functional outcomes.^[41]

CONCLUSION

Low dose propranolol may improve the clinical symptom of sympathetic nerves hyperactivity but have not significantly improved immune system disorders in patients with ischemic stroke. Ischemic stroke affects the patient's immune system that show in alteration of TNF- α , IL-10 levels and ratio of TNF α /IL-10. Provision of propranolol 2x10 mg increase TNF- α level, decrease IL-10 level and increase ratio TNF α /IL-10, but not significant statistically.

ACKNOWLEDGEMENT

Thanks the author to Prodia Laboratory, Bandar Lampung, Indonesia and Prodia Laboratory, Jakarta, Indonesia for blood samples analyze.

BIBLIOGRAPHY

1. Roger VL, Go AS, Lyold-Jones DM. Heart disease and stroke statistics-2011 updates. *Circulation*, 2011; 123(4): e18-e209.
2. Roth JM. Recombinant tissue plasminogen activator for the treatment of acute ischemic stroke. *Proc (Bayl Univ Med Cent)*, 2011; 24(3): 257–259.
3. Nour, M, Scalzo F, Liebeskind DS. Ischemia-reperfusion injury in stroke. *Intervent Neurol*, 2013; 1: 185–199.
4. Liesz A, Dalpke A, Mracsko E. AMP signaling is a key pathway inducing immune modulation after brain injury. *J Neur*, 2015; 35(2): 583–598.
5. Tarkowski E. Intrathecal release of pro- and anti-inflammatory cytokines during stroke. *Clin Exp Immunol*, 1997; 110(3): 492-9.
6. Tuttolomondo A, Pecoraro R, Pinto A. Studies of selective TNF Inhibitors in the treatment of brain injury from stroke and trauma: a review of the evidence to date. *Drug Design, Development and therapy*, 2014; 8: 2221-2239.
7. Shim R and Wong CHY. Ischemia, immunosupresion and infection-tackling the predicaments of post-stroke complications. *Int J Mol Sco*, 2016; 17(64): 1-18.
8. Vogelgesang A, May VEL, Grunwald U, Bakkeboe M, Langner S. Functional Status of Peripheral Blood T-Cells in Ischemic Stroke Patients. *PLoS ONE*, 2010; 5(1): e8718.
9. Ashour WMR, Al-Anwar AD, Kamel AE, Aidaros MA. Predictors of early infection in cerebral ischemic stroke. *Journal of Medicine and Life*, 2016; 9(2): 163-169
10. Prass K, Braun JS, Dirnagl U, Meisel C, Meisel A. Stroke propagates bacterial infection aspiration to pneumonia in amodel of cerebral ischemia. *Stroke*, 2006; 37: 2607-2612.
11. Prass K, Meisel C, Hoflich C, Braun J, Halle E, Wolf T, Ruscher K, Victorov IV, Priller J, Dirnagl U, Volk HD, Meisel A. Stroke-induced immunodeficiency promotes spontaneous bacterial infections and is mediated by sympathetic activation reversal by post stroke T helper cell type 1-like immunostimulation. *J Exp Med*, 2003; 198: 725-36.
12. Chamorro A, Amaro S, Vargas M. Interleukin 10, monocytes and increased risk of early infection in ischemic stroke. *Journal of Neurology, neurosurgery, and psychiatry*, 2006; 77: 1279-1281.
13. Vermeij FHAL, Scholte op Reimer WJ, de Man HAL. Stroke-associated infection is an independent risk factor for poor outcome after acute ischemic stroke: data from the Netherlands Stroke Survey. *Cerebrovasc Dis*, 2009; 27, 465–471.
14. Kraft P, Dreschler C, Schuhmann MK. Characterization of peripheral immune cell subsets in patients with acute and chronic cerebrovascular disease: a case-control study. *Int J Mol Sci*, 2015; 16: 25433-25449
15. Westendorp WF, Nederkoorn HALJ., Vermeij JD, Dijkgraaf MG, van de Beek D. Post-stroke infection: a systematic review and meta-analysis. *BMC Neurol*, 2011; 11: 1-7.
16. Hug A, Dalpke A, Wieczorek N, Giese T, Lorenz A, Auffarth G, Liesz A, Veltkamp R. Infarct volume is a major determiner of post-stroke immune cell function and susceptibility to infection. *Stroke*, 2009; 40: 3226–3232.
17. George AJ, Boehme AK, Siegler JE. Hospital-Acquired Infection Underlies Poor Functional Outcome in Patients with Prolonged Length of Stay. *Hindawi*, 2013; 1-5.
18. Katzan IL, Cebul RD, Husak SH, Dawson NV, Baker DW. The effect of pneumonia on mortality among patients hospitalized for acute stroke. *Neurology*, 2003; 60: 620–5.
19. Steenen SA, van Wijk AJ, van der Heijden GJ. Propranolol for the treatment of anxiety disorders: Systematic review and meta-analysis. *Journal of Psychopharmacology*, 2015; 30(2): 128–139.
20. Siefferman JW dan Lai G. Propranolol for paroxysmal sympathetic hyperactivity with lateralizing hyperhidrosis after stroke. *Hindawi*, 2015; 1-3.
21. Szelényi J, Kiss JP, Vizi ES. Differential involvement of sympathetic nervous system and immune system in the modulation of TNF- α production by α 2- and β -adrenoceptors in mice. *Neuroimmunology*, 2000; 103(1): 34–40.
22. Jina JY, Zia AK. Propranolol inhibits growth of hemangioma-initiating cells but does not induce apoptosis. *Pediatr Res*, 2014; 75(3): 381–388.
23. Wrobel LJ, Bod L, Lengagne R. Propranolol induces a favourable shift of anti-tumor immunity in a murine spontaneous model of melanoma. *Oncotarget*, 2016; 7(47): 77825–77837.

24. A, Perrotta M, Lembo G, dan Carvenale D. The spleen: a hub connecting nervous and immune systems in cardiovascular and metabolic disease. *Int. J Mol Sci*, 2017; 18: 1-12.
25. Gill D dan Veltkamp R. Dynamics of T cell responses after stroke. *Current Opinion in Pharmacology*, 2016; 26: 26–32.
26. Urra X, Cervera A, Obach V, Climent N, Chamorro A. Monocytes are major players in the prognosis and risk of infection after acute stroke. *Stroke*, 2009a; 40: 1262-8.
27. Pavlov VA dan Tracey KJ (2017): Neural regulation of immunity: molecular mechanisms and clinical translation. *Nature Neuroscience*, 2017; 20(2): 156-166.
28. Cuoco J, Fennie C, dan Cheriyan GK. The cholinergic anti-inflammatory pathway: a novel paradigm for translational research in neuroimmunology. *iMedPub Journals*, 2016; 7(2): 1-7.
29. Sorrels SF dan Sapolsky RM. An inflammatory of glucocorticoid actions in CNS. *Brain Behav Immun*, 2007; 21(3): 259-72.
30. Mullen MT, McKinney JS, Kasner SE. Blood pressure management in acute stroke. *Journal of Human Hypertension*, 2009; 23: 559-569.
31. Haverkamp W. 2015. *Beta blocker in acute ischemic stroke (BIAS)*. <https://www.clinicaltrials.gov> (Downloaded at February 27th 2016).
32. Brunsgaard H, Pedersen AN, Schroll M. TNF- α , leptin, and lymphocyte function in human aging. *Life Sci*, 2000; 67(22): 2721-31.
33. Malutan AM, Dan M, Nicolae C, Carmen M. Proinflammatory and anti-inflammatory cytokine changes related to menopause. *Prz Menopauzalny*, 2014; 13(3): 162–168.
34. Gonzalez DA, Diaz BB, Rodriguez Perez Mdel C, Hernandez AG, Chico BN, de Leon AC. Sex hormones and autoimmunity. *Immunol. Lett.*, 2010; 133: 6-13.
35. Bouman A, Heineman MJ, Faas MM. Sex hormones and the immune response in humans. *Hum. Reprod. Updat*, 2005; 11: 411–423.
36. Suberville S, Bellocq A, Fouqueray B. Regulation of interleukin-10 production by beta-adrenergic agonists. *Eur J Immunol*, 1996; 26(11): 2601-5.
37. Wang H, Yan FL, Cunningham M. Potential specific immunological indicators for stroke associated infection are partly modulated by sympathetic pathway activation. *Oncotarget*, 2016; 7(32): 52404-52415.
38. Pietschmann P, Gollob E, Brosch S, Hahn P, Kudlacek S, Willheim M, Woloszczuk W, Peterlik M, Tragl KHAL. The effect of age and gender on cytokine production by human peripheral blood mononuclear cells and markers of bone metabolism. *Gerontol*, 2003; 38: 1119–1127.
39. Haeusler KG, Schmidt, WU, Fohring F. Cellular immunodepression preceding infectious complications after acute ischemic stroke in humans. *Cerebrovasc Dis*, 2008; 25: 50–58.
40. Urra X, Obach V, Climent N, Chamorro A (). Stroke induced immunodepression syndrome: from bench to bedside. *Current Molecular Medicine*, 2009b; 9: 195-202.
41. Römer C, Engel O, Winek K. Blocking stroke-induced immunodeficiency increases CNS antigen-specific autoreactivity but does not worsen functional outcome after experimental stroke. *J Neurosci*, 2015; 35(20): 7777-94.
42. Eskdale J, Kube D, Tesch H, Gallagher G. Mapping of the human IL10 gene and further characterization of the 5' flanking sequence. *Immunogenetics*, 1997; 46 (2): 120–8.
43. Dimitrijevic M, Stanojevic S, Vujic V, *et al.* 2014. Aging oppositely affects TNF- α and IL-10 production by macrophages from different rat strains *Biogerontology* (2014) 15:475–486
44. Deguchi K, Kamada M, Irahara M, Maegawa M, Yamamoto S, *et al.* 2001. Postmenopausal changes in production of type 1 and type 2 cytokines and the effects of hormone replacement therapy. *Menopause.*; 8:266–273.
45. Sykora M, Siarnik P, Diedler J. 2015. β -blockers, pneumonia, and outcome after ischemic stroke evidence from virtual international stroke trials archive. *Stroke*. 46:1269-1274.

● 16% Overall Similarity

Top sources found in the following databases:

- 15% Internet database
- Crossref database
- 1% Submitted Works database
- 4% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	repository.lppm.unila.ac.id	Internet	12%
2	scholar.unand.ac.id	Internet	<1%
3	sciencepublishinggroup.com	Internet	<1%
4	Federal University of Technology on 2019-06-20	Submitted works	<1%
5	"APASL Abstracts, Bali 2005", Liver International, 12/2005	Crossref	<1%
6	molecularbrain.biomedcentral.com	Internet	<1%
7	A. Arac. "Systemic augmentation of B-crystallin provides therapeutic b...	Crossref	<1%
8	pfxbxzz.paperopen.com	Internet	<1%

9	Edward E. Winger, Jane L. Reed, Sherif Ashoush, Tarek El-Toukhy, Sap...	<1%
	Crossref	
10	annals.org	<1%
	Internet	
11	coursehero.com	<1%
	Internet	
12	openaccessjournals.com	<1%
	Internet	
13	science.gov	<1%
	Internet	

● **Excluded from Similarity Report**

- Bibliographic material
 - Cited material
 - Manually excluded text blocks
 - Quoted material
 - Small Matches (Less than 10 words)
-

EXCLUDED TEXT BLOCKS

wjpls, 2018, Vol. 4, Issue 1

Federal University of Technology on 2019-06-20

et al

Universitas Indonesia on 2018-11-02

of Biology, Faculty of Mathematics and Natural of Science, University of Lampung, ...

repository.lppm.unila.ac.id