

PAPER NAME

2. Isolation and Characterization of Anoxygenic Photosynthetic Bacteria for Reducing Ammonia and Pr

AUTHOR

Sumardi Sumardi

WORD COUNT

4906 Words

CHARACTER COUNT

25697 Characters

PAGE COUNT

8 Pages

FILE SIZE

1.9MB

SUBMISSION DATE

Jan 28, 2022 12:56 PM GMT+7

REPORT DATE

Jan 28, 2022 12:57 PM GMT+7

● 5% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 4% Internet database
- Crossref database
- 2% Submitted Works database
- 1% Publications database
- Crossref Posted Content database

● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)
- Manually excluded text blocks



Asian Journal of Scientific Research

ISSN 1992-1454

science
alert
<http://www.scialert.net>

ANSI*net*
an open access publisher
<http://ansinet.com>



Research Article

Isolation and Characterization of Anoxygenic Photosynthetic Bacteria for Reducing Ammonia and Probiotics Candidate

Sumardi, Rochmah Agustrina, Bambang Irawan, Tugiyono, Desfika Ardia Putri and Achmad Arifiyanto

Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Lampung, Jl Prof. Soemantri Brodjonegoro, No. 1 Bandar Lampung, 35145, Indonesia

Abstract

Background and Objective: The accumulation of organic matter in an aquaculture pond system might increase the accumulation of ammonia compounds which made toxic in aquatic organisms. Increased ammonia concentrations affected enzymatic mechanisms in animals, increased oxygen consumption and interfered with oxygen transport. This research was aimed to reduce ammonia accumulation by increasing the rate of ammonia reduction by Anoxygenic Photosynthetic Bacteria (APB) which was also a potential candidate for probiotics. **Materials and Methods:** Bacteria were isolated from mangrove forests in the Hanura beach of the Pesawaran district in Lampung province. Strains were grown on SWC media at anoxygenic-photosynthetic condition, morphologically characterized, tested for pH resistance, salinity, antibiotic endurance, ability to compete with *Vibrio* sp. and ability to reduce ammonia. **Results:** A total of 7 isolates grew well at pH 7 and 10 and 3 and 6% of NaCl concentrations. L1 strain reduced ammonia by 62%. L2 strain was the most resistant to nalidixic acid, streptomycin, chloramphenicol, ampicillin and trimethoprim. **Conclusion:** Bacterial strain B has the highest competitive ability against *Vibrio* sp., which was able to survive up to 2.6 log_Σ.cell. Two probiotic candidates were photosynthetic bacteria. They were able to inhibit of *Vibrio* sp. growth, reduce the ammonia content by 22-33% and resistant into antibiotics.

Key words: *Vibrio*, ammonia, anoxygenic, photosynthetic bacteria, probiotic, vibriosis, aquaculture

Citation: Sumardi, R. Agustrina, B. Irawan, Tugiyono, D.A. Putri and A. Arifiyanto, 2021. Isolation and characterization of anoxygenic photosynthetic bacteria for reducing ammonia and probiotics candidate. Asian J. Sci. Res., 14: 6-12.

Corresponding Author: Sumardi, Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Lampung, Jl Prof. Soemantri Brodjonegoro No. 1 Bandar Lampung, 35145, Indonesia

Copyright: © 2021 Sumardi *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

There were several problems for aquaculture farms, either infectious or non-infectious. *Vibrio* sp. was one of infectious sources, namely vibriosis. Antibiotic utility was applied to treat the diseases. Unfortunately, antibiotics usage was lead to pathogen resistance¹. Ammonia accumulation was non-infectious problems. Ammonia was accumulated during feed digestion and excreted through their gills and in their feces by fish and shrimp. It was created an anaerobic condition at the pond².

Anaerobic Photosynthetic Bacteria (APB) were bacteria carried out decomposition activities even though oxygen levels in water or sediment were very few or even none at all. This was because they were able to utilize hydrogen donors from compounds that are reduced more strongly (H_2S , ammonia, nitrites and organic compounds) as a substitute for H_2O ³. According to Canfield *et al.*⁴, the presence of anoxygenic photosynthetic bacteria was able to reduce H_2S toxic compounds. Besides, they had good potential as a biocontrol agent⁵, as a heavy metal-lowering agent⁶ and an effective fishmeal replacement in aquaculture⁷. The effect of Fe ions exerts an increase in the absorption of light in photosynthetic APB at wavelengths⁸ of 300-400 nm.

Lampung province, Indonesia, had high production of shrimp hatchery that grew at mangrove habitat. Vibriosis and ammonia accumulation was serious threaten for shrimp aquaculture. Regarding the above reason the objective of the study was to isolate and characterize APB isolates from mangrove forests in the Hanura waters of Lampung as a candidate bioremediation agent in reducing ammonia compounds and candidates probiotic agents in competing against pathogenic bacteria *Vibrio* sp.

MATERIALS AND METHODS

Study area: This research project was conducted at the Department of Biology, Microbiology Lab, University of Lampung, Indonesia from December 2018 to June 2019.

Sample preparation: Anoxygenic photosynthetic bacterial isolates were isolated from several sample points in the Hanura Lampung mangrove forest including sea mud, mangrove mud, roots, leaf litter, flowers, snails and crabs. Each bacterial isolate was inoculated in liquid SWC media with a composition 2 g of bacto peptone, 1 g of yeast extract, 3 mL of glycerol, 250 mL of distilled water, 750 mL of sea-water and 15 g of agar) liquid media without agar. It incubated for 6-7 days in front of the lamp tungsten at a distance of

30-40 cm. A single loop of the anoxygenic photosynthetic bacterial isolate was taken and then scratched using the quadrant streak method⁹, on SWC agar (SWCA) and re-incubated for 5-6 days in an anaerobic jar in front of a 40-watt accent lamp with a distance of 30-40 cm. Furthermore, the growing bacterial isolates were transferred to the SWCA sloping media to obtain pure isolates¹⁰. Bacteria characterized through Gram staining, microscopic morphology, colony color and shape of the colony.

pH and salt-tolerant assay: Bacteria were purified at Petri dish containing SWCA media which was given a pH difference treatment of 4, 7 and 10. Then they incubated for 4-5 days by the anoxygenic photosynthetic condition. Bacteria were cultured into modified SWCA media with the addition of salt concentrations of 0, 3 and 6%. It was incubated for 4-5 days by anoxygenic photosynthetic. Their growth parameters observed according to Triyanto *et al.*¹¹ methods.

Ammonia reducing assay: A single loop of bacteria was grown on a screw tube containing liquid SWC media (5 mL). Each tube was then homogeneous with the vortex. The liquid SWC was added until it was full. The threaded tubes are then incubated at room temperature for 5-6 days in anaerobic photosynthetic conditions. They centrifuged for 10 min at 1000 rpm and 4°C and the supernatant was taken¹². The amount of 6 ml of bacteria culture supernatant was put into a test tube and add 1 drop of Seignette salt (15 g in 30 mL of distilled water) then homogenate and let stand for 2 min. Nessler (K_2HgI_4) as much as 0.25 mL was added and homogenized again for 15 min until a clear purple solution occurred. Ammonia reduction test was carried out using a UV-Vis Shimadzu spectrophotometer at a wavelength¹² of 420 nm.

Antibiotic assay: Bacteria were cultured at SWC liquid media in photosynthetic conditions for 3×24 hours at room temperature¹². Antibiotic assay conducted using the diffusion method. Bacteria in 0.1 mL suspension was poured on the surface of the SWCA media and placing antibiotic discs contained Chloramphenicol, Nalidixic Acid, Trimethoprim, Ampicillin and Streptomycin. It was incubated for 3×24 hours by the photosynthetic anaerobic condition. A clear zone around the antibiotic disk paper that indicated the inhibition zone was measured.

Anaerobic photosynthetic-bacteria against Vibriosp. Assay: Bacteria incubated at 14 mL of SWC liquid 3 days in photosynthetic threaded test tubes. A total of 10 mL isolates

of *Vibrio* sp. incubated in Nutrient Broth (NB) media (1×24 hours). They were calculated microscopically together with Gram staining to determine the population density of anoxygenic photosynthetic bacterial and *Vibrio* sp.

Both anoxygenic photosynthetic bacterial and *Vibrio* sp. were taken 1 mL and put into a threaded test tube containing NB liquid media at a density of 10^4 and 10^2 respectively (incubated for 2×24 hours by anoxygenic photosynthetic at room temperature). The population detected on the media used anaerobic photosynthetic pour method, while to detect the growth of *Vibrio* sp. incubated on TCBSA media on Petri dishes at room temperature¹³ for 24 hours.

RESULTS

Sample characterization: There were 7 strains of anoxygenic photosynthetic bacteria cultured from 12 location points. They were most Gram-negative isolates in the form of stem colonies while the colonies' color was beige, white, turbid, red, yellow and brown. Thereby they are taken which will be tested further presented in Table 1.

Salt tolerant and pH assay: Anoxygenic photosynthetic bacteria was tested in different pH and NaCl stresses, it was aimed to know the susceptibility of salinity and acidity level. The pH test was carried out using pH 4, 7 and 10 while the NaCl test with concentrations of 0, 3 and 6%. The result was obtained in Table 2 for salt-tolerant, due to all bacteria strains were taken from the mangrove ecosystem it was proven that they were salt tolerant. They were more appropriate to grow well at 3-6% salt concentration or above. Even though some of them kept tolerate the level of freshwater without salt content in the media.

Based on Table 3, the pH test results obtained showed that all isolates were unable to grow at pH 4. It was shown that the APB obtained from the isolation results could not grow

Table 1: Anoxygenic photosynthetic bacterial that succeeded culture

Sources of strain	Strain codes	Gram staining	Cell shape
Root	AS	Negative	Rods
Root	AM	Negative	Rods
Mud	L1	Negative	Rods
Mud	L2	Negative	Rods
Flower	B	Negative	Rods
Leaves	D	Negative	Rods
Leaves	B2DM	Negative	Rods

AS: Strain was isolated from fibrous root of mangrove, AM: Strain was isolated from root of mangrove, L1: Strain was isolated from mud of mangrove location 1, L2: Strain was isolated from mud of mangrove location 2, B: Strain was isolated from litter of flowers of mangrove, D: Strain was isolated from litter of leave of mangrove, B2DM: Red color-Strain was isolated from litter of flowers of mangrove

under acidic conditions. Whereas in alkaline conditions (pH 10), bacterial growth was inhibited in isolates L1.

Ammonia reducing assay: The results of a decrease in ammonia concentration tests are shown in Fig. 1, that anoxygenic photosynthetic bacterial isolates showed the different abilities in reducing ammonia concentration. Based on the results of the effectiveness test, the greatest effectiveness value is 62% L1 isolate and the smallest is B2DM 12%. Thus the L1 isolate was the best in suppressing the concentration of ammonia compared to other isolates. From this result, we knew that bacteria strain that taken from mud sediment had a high-ability to reduce ammonia. It was due to that they were allegedly doing that ecologically as long as they live at ammonia accumulation inside mud sediment of mangrove.

Antibiotic assay: Antibiotic susceptibility assay was presented in Table 4. The most resistance reactions were streptomycin and ampicillin. There were four isolates namely AS, B, D and

Table 2: Salt tolerant assay toward the anoxygenic photosynthetic bacterial culture

APB isolate	Concentration NaCl in medium (%)		
	0	3	6
AS	+	++	++
AM	+	++	++
L1	++	++	++
L2	++	++	++
B	+	++	++
D	+	++	++
B2DM	-	+	+
K1	++	++	++
B3DK	+	++	++
B4LM	++	+	+

-: No bacterial growth, +: Diameter of bacterial colony less than 0.5 cm, ++: Diameter of bacterial colony between 0.5-1.0 cm

Table 3: pH tolerant assay toward the anoxygenic photosynthetic bacterial culture

APB Isolate	Medium pH		
	4	7	10
AS	-	++	++
AM	-	++	++
L1	-	++	++
L2	-	+	-
B	-	+	+
D	-	+	+
B2DM	-	-	-
K1	-	+	+
B3DK	-	++	+
B4LM	-	++	++

-: No, +: Diameter of bacterial colony less than 0.5 cm, ++: Diameter of bacterial colony between 0.5-1.0 cm




Fig. 1: Percentage of reducing ammonia in anoxic photosynthetic bacteria compared to sources of strain isolation habitat

Table 4: Bacterial susceptibility toward antibiotic assay

APB isolate	Antibiotic assay				
	NA	S	C	A	T
AS	8.1	R	21.5	15.3	R
AM	11.7	10	16.4	R	18.1
L1	R	8,8	7.9	R	9.3
L2	17.5	5.2	25.2	6.2	7.1
B	12.4	R	R	R	9
D	9	R	19.1	10	10
B2DM	9.6	R	26.2	14.4	R

NA: Nalidixic acid, S: Streptomycin, C: Chloramphenicol, A: Ampicillin, T: Trimethoprim, R: Resistance

Table 5: Competition test results between APB and *Vibrio* sp.

No.	APB versus <i>Vibrio</i> sp.	Log Σ of initial cell		Log Σ of final cell	
		APB	<i>Vibrio</i> sp.	APB	<i>Vibrio</i> sp.
1	- <i>Vibrio</i> sp.	-	2.0	-	2.9
2	APB-B2DM -	4.0	-	2.2	-
3	APB-D -	4.0	-	2.5	-
4	APB-B -	4.0	-	2.3	-
5	APB-AS -	4.0	-	2.3	-
6	APB-AM -	4.0	-	1.9	-
7	APB-L1 -	4.0	-	2.1	-
8	APB-L2 -	4.0	-	2.2	-
9	APB-B2DM <i>Vibrio</i> sp.	4.0	2.0	2.4	2.2
10	APB-D <i>Vibrio</i> sp.	4.0	2.0	2.2	2.2
11	APB-B <i>Vibrio</i> sp.	4.0	2.0	1.9	0
12	APB-AS <i>Vibrio</i> sp.	4.0	2.0	2.2	1.4
13	APB-AM <i>Vibrio</i> sp.	4.0	2.0	2.6	0
14	APB-L1 <i>Vibrio</i> sp.	4.0	2.0	1.6	1.6
15	APB-L2 <i>Vibrio</i> sp.	4.0	2.0	2.4	2.4

1: Bacteria of *Vibrio* sp. 2: APB- B2DM 3: APB- D
 4: APB- B 5: APB- AS 6: APB- AM
 7: APB- L1 8: APB- L2 9: BFA B2DM X *Vibrio* sp.
 10: APB- D X *Vibrio* sp. 11: APB- B X *Vibrio* sp. 12: APB- AS X *Vibrio* sp.
 13: APB- AM X *Vibrio* sp. 14: APB- L1 X *Vibrio* sp. 15: APB- L2 X *Vibrio* sp.

B2DM which are resistant to streptomycin. Whereas AM, L1 and B isolates were resistant to ampicillin antibiotics.

Anaerobic photosynthetic-bacteria against *Vibrio* sp.

Assay: The results of the competition test between APB and *Vibrio* sp. shows the competitive ability of each of the different APB isolates. The biggest competition ability of

APB isolates was isolate B from mangrove litter samples at 2.6 and AM isolate at 1.9 Log Σ cell. The results of the competition test between APB and *Vibrio* sp. can be seen in Table 5.

The results of the competition between APB and *Vibrio* sp. this occurs because isolate B from APB comes from mangrove litter and AM isolates from mangrove roots are deep in the mud, where the aquatic environment is a dirty environment so many decaying bacteria also live in the environment. Therefore, APB B and AM isolates are accustomed to competing with decomposing microorganisms, one of which is *Vibrio* sp. Also, APB is thought to produce several antibacterial compounds.

DISCUSSIONS

Bacteria have genes for resistance that are obtained from the environment for a very long time or even after exposure to antibiotic selection. Bacteria can reproduce to propagate resistance genes both vertically and horizontally and have a long-term impact. Some bacteria have a natural ability to be resistant to antibiotics. This can occur because bacteria have enzymes that can damage drugs¹⁴. Also, streptomycin and ampicillin antibiotics are many types of antibiotics that are often used in aquatic environments. This is supported by research Kusmarwati *et al.*¹⁵ that 100% of *Vibrio parahaemolyticus* bacteria resistant to streptomycin and 90% of ampicillin in the cultivation of Vaname shrimp on the north coast of Java.

Furthermore, as shown in Table 5, the types of antibiotics of nalidixic acid, chloramphenicol and trimethoprim have sensitive reactions to APB isolates. This is presumably because all three types of antibacterials are still rarely used in an antibiotic environment. According to research Kusmarwati *et al.*¹⁵ informed that sensitive bacterium *Vibrio parahaemolyticus* on chloramphenicol antibiotics by 18%.

Makori *et al.*¹⁶ and Agustiyani *et al.*¹⁷ were described that the optimum aquaculture environment (ponds) requires the pH of soil conditions ranging from 6.34-8.10. The degree of acidity (pH) is one of the conditions for growing a microorganism. The relationship between the pH value and bacterial growth greatly affects the work system of bacterial metabolism itself to produce a substance called an enzyme. Enzyme activity will be maximum if the pH conditions are also optimum. Rice *et al.*¹⁸, was explained that the acidic pH of the bacterial cell membrane will become saturated by hydrogen ions thereby limiting membrane transport. This causes the acid substance that does not decompose to seep into the cell so that the ionization of the cell's pH changes. This change causes the process of sending amino acids from RNA to be inhibited so that it inhibits growth and can even kill the bacteria itself.

Then, the results of the APB isolate growth test with media containing different concentrations of NaCl showed that 8 isolates could grow at concentrations of 3% and 6% NaCl except for isolates K and B3DK. According to Suprayogi *et al.*¹⁹ that if the water contains 0.5-3% salt content, then the water can be called brackish water whereas if >3% can be said to be seawater.

Sodium chloride (NaCl) stress inhibited the growth of bacteria because it increased the osmotic pressure of the substrate which can cause water withdrawal from the microorganism cells so that the cell will lose water and shrink and cause slowing of microorganism activity¹⁸. This is supported by Membre research²⁰ where NaCl can cause denatured microbial protein besides NaCl also has high toxicity to microbes, salt ionization will produce Cl-ions that are toxic to microorganisms and can block the respiration system of the microorganism itself. Rubiano *et al.*²¹ salt were capable of reducing the solubility of O₂ in water, cause cells to be more sensitive to CO₂ and interfere with the work of proteolytic enzymes in microbial cells that function to break down proteins into amino acids.

In the study of antibiotic tests on APB isolates, the largest clear zone was found in B2DM isolates tested on chloramphenicol antibiotics by 26.2 mm and the smallest clear zone was L2 isolates on ampicillin antibiotics by 6.2 mm, Table 4. According to Li *et al.*²², the size of the clear zone contained in the antibiotic test is influenced by several factors including 1) the type and age of bacteria, 2) the concentration of antibiotic substances and the amount of inoculum or bacterial density, 3) levels of active substances or functional groups of antibiotic substances. Photosynthetic bacteria are known to have a central role in the nitrogen element cycle.

The addition of photosynthetic bacteria can also eliminate organic matter or other harmful substances in improving water quality and is included in the type of probiotics that provide nutrients and enzymatic contributions to fight pathogenic bacteria²³⁻²⁵.

In anaerobic sediments, some microbes break down organic matter with fermentation reactions that are capable of producing alcohol, ketones and other compounds. According to Madigan *et al.*²⁶ microorganisms can also use oxygen from nitrates, nitrites, sulfates and CO₂ to decompose organic material, but these microorganisms also emit nitrogen gas, ammonia, ferrous, manganous manganese, H₂S and methane as a result of metabolism.

Sea Water Complete (SWC) media used in the study contained organic compounds in the form of nitrogen from peptones which when incubated under anaerobic conditions for approximately 5 days will form ammonia compounds. According to Jeong *et al.*²⁷ an ammonia test can be carried out using a basic solution of mercury (II) iodide in Potassium Tetraiodomercurate (K₂HgI₄) or it can also be called a Nessler reagent to measure ammonia compounds calorimetric. In the working principle of the Nessler method, Nessler reagents when reacted with ammonia in the form of an alkaline solution will form a brownish-yellow colloidal dispersion.

As reported by Tsang *et al.*²⁸, ammonia can be reduced with the help of indigenous bacteria through metabolic processes. Ammonia degradation mechanism is found in aerobic conditions by nitrifying bacteria groups, namely Ammonia-Oxidizing Bacteria (AOB) such as Nitrosomonas and Nitrobacter. But apart from that also the anaerobic ammonia condition can be degraded by anaerobic ammonia-oxidizing (Anammox). According to Feng *et al.*²⁹, that the anoxygenic photosynthetic bacteria Rhodospirillum rubrum can be used as a bioremediation agent because it can reduce ammonia compounds by 56%.

Competition is a negative interaction that occurs between two or many microorganisms when the supply of food sources, oxygen, light and space needed is limited to survive and reproduce³⁰. APB which is anaerobic bacteria can decompose organic matter with fermentation reactions that can produce alcohol, ketones and other compounds. This is supported by research by Feng *et al.*²⁹, which states that anaerobic microorganisms can use oxygen from nitrates, nitrites, sulfates and CO₂ to decompose organic material, but emit nitrogen, ammonia, ferrous, manganous manganese, H₂S and methane as a result of metabolism. It suspected to be toxic to *Vibrio* sp.

According to Yang *et al.*³¹ the antimicrobial effect produced by bacteria is caused by several factors, namely the production of antibiotics, bacteriocin, siderophore, lysozyme, proteases, or hydrogen peroxidase and changes in pH values through the production of organic acids. The antimicrobial in question is bacteriocin, which is an extracellular component in the form of an antimicrobial protein compound that shows the opposite response to the target bacteria³². The bacteriocin compound that is currently widely used in fishery products is Nisin produced from *Lactococcus lactis*³³. Antimicrobial compounds which are bacteriocin will cause the target bacterial cells to become lysis. Besides, there was a struggle for nutrition and space between APB and *Vibrio* sp. as stated by Sumardi *et al.*³⁴ when a challenge test was conducted between Bacillus and *Vibrio* sp. Also, the difference in oxygen demand of each bacterium in which APB is a facultative anaerobic is that it can live in conditions of very little oxygen content until there is no oxygen at all while *Vibrio* sp. is a type of facultative aerobic bacteria wherein.

CONCLUSION

The result informed that bacteria were successfully cultured anaerobic photosynthetic-bacteria. L1 strain reduced ammonia by 62%. L2 strain was the most resistant to nalidixic acid, streptomycin, chloramphenicol, ampicillin and trimethoprim. Bacterial strain B has the highest competitive ability against *Vibrio* sp., which was able to survive up to 2.6 log.Σ.cell. L1 was potentially developed to bioremediation agent but has a low ability to pH and salinity stress it also happened to strain L2. Regarding the result of reducing ammonia, antibiotic susceptibility, pH and salinity stress, anaerobic photosynthetic-bacteria strain B was more appropriate selected as a candidate for bioremediation and prob

SIGNIFICANCE STATEMENT

This study found 2 probiotic candidates, APB-B and APB-AM strains, that were unique photosynthetic-bacteria. They were able to inhibit the growth of *Vibrio* sp. (*in vitro* conditions), able to reduce the ammonia content by 22-33% and resistant into antibiotics. This study will help researchers to uncover resistance antibiotics in BFA bacteria that occur in mangroves habitat. The resistance of these bacterial antibiotics has not been explored by many researchers. It was potentially developed as a probiotic candidate to overcome the shrimp disease.

ACKNOWLEDGMENT

This research is part of the Research Team Dr. Sumardi, M.Si based on Unila Postgraduate Research assignment letter No. Contract: 1572 / UN26.21 / PP / 2018, July 9 2018.

REFERENCES

1. Aguilera-Rivera, D., A. Prieto-Davó, G. Rodríguez-Fuentes, K.S. Escalante-Herrera and G. Gaxiola, 2019. A vibriosis outbreak in the Pacific white shrimp, *Litopenaeus vannamei* reared in biofloc and clear seawater. J. Invertebr. Pathol., 10.1016/j.jip.2019.107246
2. Durborow, R.M., D.M. Crosby and M.W. Brunson, 1997. Ammonia in fish ponds. Srac 1–28.
3. Hanada, S., 2005. Anoxygenic photosynthesis-A photochemical reaction that does not contribute to oxygen reproduction. Microb. Environ., 10.1264/jsme2.ME3101rh
4. Canfield, D.E., E. Kristensen and B. Thamdrup, 2005. The sulfur cycle. Adv. Mar. Biol., 48: 313-381.
5. Yuhana, M., 2010. Biocontrol agents in aquaculture: Production and their application. J. Akuakultur Indonesia, 9: 16-20.
6. Giotta, L., A. Agostiano, F. Italiano, F. Milano and M. Trotta, 2006. Heavy metal ion influence on the photosynthetic growth of *Rhodobacter sphaeroides*. Chemosphere, 62: 1490-1499.
7. Delamare-Deboutteville, J., D.J. Batstone, M. Kawasaki, S. Stegman and M. Salini *et al.*, 2019. Mixed culture purple phototrophic bacteria is an effective fishmeal replacement in aquaculture. Water Res. X., 10.1016/j.wroa.2019.100031
8. Sumardi, S. Farisi, R. Agustina and Yunita, 2019. Effects of magnetic fields and metal ions (Cu, Pb, Al and Fe) on growth of Anoxygenic Photosynthetic Bacteria (APB). Biospecies, 12: 42-50.
9. Cox, N.A., L.J. Richardson, D.E. Cosby, M.E. Berrang, J.L. Wilson and M.A. Harrison, 2017. A four-quadrant sequential streak technique to evaluate *Campylobacter* selective broths for suppressing background flora in broiler carcass rinses. J. Food Safety, 10.1111/jfs.12311
10. Higuchi-Takeuchi, M., K. Morisaki and K. Numata, 2016. A screening method for the isolation of polyhydroxyalkanoate-producing purple non-sulfur photosynthetic bacteria from natural seawater. Front. Microbiol., 10.3389/fmicb.2016.01509
11. Triyanto, A. Isnansetyo, I.D. Prijambada, J. Widada and Tarmiwati, 2009. An Isolation, characterization and proteolytic bacterial infection test from mud forest area mud. J. Perikan., 11: 13-18.

12. Huang, X., J. Ni, C. Yang, M. Feng, Z. Li and D. Xie, 2018. Efficient ammonium removal by bacteria rhodospseudomonas isolated from natural landscape water: China case study. *Water*, Vol. 10. 10.3390/w10081107
13. Yeung, M. and T. Thorsen, 2016. Development of a more sensitive and specific chromogenic agar medium for the detection of *Vibrio parahaemolyticus* and other *Vibrio* species. *J. Vis. Exp.*, 10.3791/54493
14. Egorov, A.M., M.M. Ulyashova and M.Y. Rubtsova, 2018. Bacterial enzymes and antibiotic resistance. *Acta Naturae*, 10: 33-48.
15. Kusmarwati, A., Y. Yenni and N. Indriati, 2017. Antibiotic Resistance in *Vibrio parahaemolyticus* from Vannamei Shrimp Originated from Northern Coast of Java for Export Market. *J. Pascapanen Bioteknologi Kelaut. Perikanan*, 12: 91-105.
16. Makori, A.J., P.O. Abuom, R. Kapiyo, D.N. Anyona and G.O. Dida, 2017. Effects of water physico-chemical parameters on tilapia (*Oreochromis niloticus*) growth in earthen ponds in Teso North Sub-County, Busia County. *Fish. Aqua. Sci.*, 20: 1-10.
17. Agustiyani, D., 2004. Effect of pH and organic substrate on growth and activities of ammonia-oxidizing bacteria. *Biodiversitas*, 5: 43-47.
18. Rice, K.C. and K.W. Bayles, 2008. Molecular control of bacterial death and lysis. *Microbiol. Mol. Biol. Rev.*, 72: 85-109.
19. Suprayogi, I., A. Nadjndji, Dijatnol and I. Muhammad, 2006. The phenomenon intrusion phenomenon in estrusi due to the effects of high tides on the river stream using the physical model approach. *J. Parifikasi*, 7: 133-138.
20. Membre, J.M. and P.M. Burlot, 1994. Effects of temperature, pH and NaCl on growth and pectinolytic activity of *Pseudomonas marginalis*. *Applied Environ. Microbiol.*, 60: 2017-2022.
21. Rubiano-Labrador, C., C. Bland, G. Miotello, J. Armengaud and S. Baena, 2015. Salt stress induced changes in the exoproteome of the halotolerant bacterium *Tistlia consotensis* deciphered by proteogenomics. *PLoS ONE*, 10.1371/journal.pone.0135065
22. Li, J., S. Xie, S. Ahmed, F. Wang and Y. Gu *et al.*, 2017. Antimicrobial activity and resistance: influencing factors. *Front. Pharmacol.* Vol. 8. 10.3389/fphar.2017.00364
23. Qi, Z., X.H. Zhang, N. Boon and P. Bossier, 2009. Probiotics in aquaculture of China-current state, problems and prospect. *Aquaculture*, 290: 15-21.
24. Verschuere, L., G. Rombaut, P. Sorgeloos and W. Verstraete, 2000. Probiotic bacteria as biological control agents in aquaculture. *Microbiol. Mol. Biol. Rev.*, 64: 655-671.
25. Cruz, P.M., A.L. Ibanez, O.A.M. Hermosillo and H.C.R. Saad, 2012. Use of probiotics in aquaculture. *ISRN Microbiol.*, Vol. 2012. 10.5402/2012/916845
26. Madigan, M., K. Bender, D. Buckley, W. Sattley and Stahl, 2019. *Brock Biology of Microorganisms*. 15th Edn., Pearson, Hudson Street, NY, USA, pp: 118-120.
27. Jeong, H., J. Park and H. Kim, 2013. Determination of NH⁺ in environmental water with interfering substances using the modified nessler method. *J. Chem.*, 10.1155/2013/359217
28. Sang, Y.F., Y.N. Wang, H. Wang, Y. Yang, Y. Zhang and H. Chua, 2017. Biodegradation of ammonia in biofiltration systems: changes of metabolic products and microbial communities. *Intechopen i.*, Vol. 13. 10.5772/intechopen.68155
29. Feng, Y., Y. Yu, Y. Wang and X. Lin, 2007. Biosorption and bioreduction of trivalent aurum by photosynthetic bacteria *Rhodobacter capsulatus*. *Curr. Microbiol.*, 55: 402-408.
30. Hibbing, M.E., C. Fuqua, M.R. Parsek and S.B. Peterson, 2010. Bacterial competition: surviving and thriving in the microbial jungle. *Nat. Rev. Microbiol.*, 8: 15-25.
31. Yang, S.C., C.H. Lin, C.T. Sung and J.Y. Fang, 2014. Antibacterial activities of bacteriocins: Application in foods and pharmaceuticals. *Front. Microbiol.*, Vol. 5. 10.3389/fmicb.2014.00241
32. Drider, D., G. Fimland, Y. Hechard, L.M. McMullen and H. Prevost, 2006. The continuing story of class IIa bacteriocins. *Microbiol. Mol. Biol. Rev.*, 70: 564-582.
33. Da Costa, R.J., F.L.S. Voloski, R.G. Mondadori, E.H. Duval and Â.M. Fiorentini, 2019. Preservation of meat products with bacteriocins produced by lactic acid bacteria isolated from meat. *J. Food Quality*, 10.1155/2019/4726510
- Sumardi, S. Farisi, C. Ekowati and R. Oktavia, 2019. Challenging test of bacillus bacteria in vitro probiotic candidates for vibrio harvey bacteria causing shrimp disease. *J. Biol. Papua*, 11: 57-63.

● 5% Overall Similarity

Top sources found in the following databases:

- 4% Internet database
- Crossref database
- 2% Submitted Works database
- 1% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	scialert.net Internet	2%
2	Rahmi, Jamaluddin Jompa, Akbar Tahir. "Transmission of Desulfovibri... Crossref	<1%
3	Multimedia University on 2017-10-02 Submitted works	<1%
4	oraas0.ied.edu.hk Internet	<1%
5	link.springer.com Internet	<1%
6	University of Sheffield on 2018-07-24 Submitted works	<1%
7	journal.uinjkt.ac.id Internet	<1%
8	repository.ung.ac.id Internet	<1%

● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)
- Manually excluded text blocks

EXCLUDED SOURCES

repository.lppm.unila.ac.id

Internet

88%

Sumardi ., Rochmah Agustrina, Bambang Irawan, Tugiyono ., Desfika Ardia Pu... **79%**

Crossref

mail.scialert.net

Internet

78%

EXCLUDED TEXT BLOCKS

OPEN ACCESSAsian Journal of Scientific ResearchISSN 1992-1454DOI: 10.3923/a...

repository.lppm.unila.ac.id