

PAPER NAME

**The growth performance of the Pacific white shrimp (*Litopenaeus vannamei*) cultured at various salini**

AUTHOR

**Supono Supono**

WORD COUNT

**3205 Words**

CHARACTER COUNT

**16661 Characters**

PAGE COUNT

**6 Pages**

FILE SIZE

**409.0KB**

SUBMISSION DATE

**Jan 11, 2023 2:48 PM GMT+7**

REPORT DATE

**Jan 11, 2023 2:48 PM GMT+7**


### ● 13% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 13% Internet database
- 10% Publications database
- Crossref database
- Crossref Posted Content database
- 8% Submitted Works database

### ● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources



# The growth performance of the Pacific white shrimp (*Litopenaeus vannamei*) cultured at various salinity conditions using single step acclimation

<sup>1,2</sup>Supono, <sup>1</sup>Rehulina T. Pinem, <sup>2</sup>Munti Sarida

<sup>4</sup>Department of Coastal and Marine Zone Management, Postgraduate Program, Lampung University, Indonesia; <sup>5</sup>Department of Aquaculture, Faculty of Agriculture, Lampung University, Indonesia. Corresponding author: Supono, [supono\\_unila@yahoo.com](mailto:supono_unila@yahoo.com)

**Abstract.** The Pacific white shrimp (*Litopenaeus vannamei*) is a euryhaline species that can be reared in aquatic environments with a wide range of salinity, from 0.5 to 40 ppt. The ability of Pacific white shrimps to adapt to extreme saline conditions allows the expansion of shrimp farming in inland aquatic environments, away from the coast. Therefore, this study aimed to analyze the specific growth rate (SGR), survival rate (SR), and feed conversion ratio (FCR) of Pacific white shrimp reared at various salinity levels using a single-step acclimation method. The study employed a completely randomized design with five treatments and three replications, namely T1, T2, T3, T4, and T5 at 5 ppt, 10 ppt, 15 ppt, 20 ppt, and 25 ppt, respectively. Each treatment unit was incorporated with Pacific white shrimp postlarvae (PL) 15 at a density of 2000 PL m<sup>-3</sup>. Subsequently, the shrimps were fed commercial pellets three times daily through a blind feeding program during the 40 days of rearing. The results showed that different salinities of culture media had an effect on SGR, SR, and FCR of Pacific white shrimp. The most desirable SGR and FCR were observed in the 15 ppt culture media at 12.3% and 1.5, respectively, while the best survival rate was observed in the 20 ppt salinity at 79.3%. Generally, Pacific white shrimp performed best in the culture media with a salinity of 15 ppt.

**Key Words:** euryhaline, growth, *L. vannamei*, survival rate.

**Introduction.** The Pacific white shrimp (*Litopenaeus vannamei*) is a euryhaline species that can tolerate a wide range of salinity between 0.5 to 45 ppt (Wyban & Sweeny 1991), allowing for high-yield rearing away from the coast. The rearing of Pacific white shrimp at low salinity is one of the best options to optimize the use of low-salinity ponds located far from seawater sources, due to the high salinity tolerance range of this shrimp (Chong-Robles et al 2014). Furthermore, this is important since there are numerous brackish water aquaculture centers located inland in Lampung Province, Indonesia that can be used for Pacific white shrimp farming (Supono 2021).

Although the Pacific white shrimp can adapt to a wide range of salinities, it grows best in iso-osmotic media at a salinity equal to the osmotic tolerance range of common shrimps (Anggoro et al 2018). The rate of osmosis is a measure of the processes that occur in aquatic animals during times of environmental stress (Cheng et al 2006). Differences in media salinity and fluid osmolarity in shrimp (hemolymph) can lead to increased energy requirements for adaptation, retardation of shrimp growth and metabolism (Gao et al 2016). Meanwhile, shrimps can only survive by utilizing available energy during conditions of low salinity, where an extremely high osmotic load reduces shrimp survival (Jayasankar et al 2009; Susilowati et al 2014). The salinity of the media can be adjusted based on the osmotic pressure of the media to increase the growth of cultured species (Lukas et al 2017). Acclimatization to temperature and salinity is an important factor that affects the success of low-salinity shrimp farming. Acclimatization systems for stocking shrimp juveniles are classified into one-step acclimation and gradual acclimation (Jayasankar et al 2009). Each technique has its advantages and disadvantages. The single-step acclimation is easy to implement, but sometimes has a

low survival rate. Gradual acclimation is difficult to implement in the field. Therefore, this study aimed to evaluate the growth rate, survival rate, and feed conversion ratio of Pacific white shrimp reared under various salinity conditions.

#### Material and Method

**Experimental design and setup.** The study was conducted from July to September 2021 at the Laboratory of Aquaculture, Lampung University, Indonesia. This study used a completely randomized design (CRD), which consisted of 5 treatments and 3 replications. The treatments include the following: T1 - culture media at salinity of 5 ppt; T2 - culture media at salinity of 10 ppt; T3 - culture media at salinity of 15 ppt; T4 - culture media at salinity of 20 ppt; T5 - culture media at salinity of 25 ppt.

The study made use of 15 plastic containers filled with 70 L of water, which was sterilized with chlorine at a concentration of 30 mg L<sup>-1</sup>. Each container was filled with 70 L of water at varying salinity based on the treatment. Furthermore, the aeration settings were performed in each maintenance container, with up to four aeration stones per container serving as an oxygen supply to the culture container. Each treatment unit was stocked with 140 Pacific white shrimp (PL 15), or the equivalent of 2000 PL m<sup>-3</sup>. The shrimp were obtained from a hatchery in Kalianda City, South Lampung. Acclimatization was conducted using a single-step acclimation system, while feeding was carried out using a blind feeding program three times daily at 6.30, 12.00, and 19.00, throughout the 45-day culture period. The feed used was commercial feed containing protein fat, fiber, ash, and moisture at 41%, 7%, 3%, 13%, and 10%, respectively. Sodium (Na), magnesium (Mg), calcium (Ca), and potassium (K) concentrations were analyzed using the U.S. EPA 200.7 method (U.S. EPA 1994), while pH, temperature, and dissolved oxygen were measured using a water quality checker.

**Statistical analysis.** Pacific white shrimp was evaluated based on specific growth rate (SGR), survival rate (SR), biomass, and feed conversion ratio (FCR). The data were analyzed using ANOVA at a 95% confidence level (p<0.05) using the SPSS 23.0 software. Further testing was conducted using the Duncan test when there were significant differences between treatments.

**Results and Discussion.** Table 1 shows the data from observations made during the study on SGR, SR, biomass, and FCR of Pacific white shrimp. The ANOVA statistical test results showed that the rearing of white shrimps in various saline media has a significant effect on their SGR, SR, biomass, and FCR (p<0.05).

Table 1

Growth performance of Pacific white shrimp

Parameter	T1 (5 ppt)	T2 (10 ppt)	T3 (15 ppt)	T4 (20 ppt)	T5 (25 ppt)
Growth rate (g)	0.6±0.1 <sup>a</sup>	1.4±0.2 <sup>c</sup>	2.8±0.1 <sup>d</sup>	0.8±0.2 <sup>ab</sup>	1.2±0.1 <sup>bc</sup>
SGR (%)	8.4±0.7 <sup>a</sup>	10.4±0.3 <sup>c</sup>	12.3±0.1 <sup>d</sup>	9.4±0.8 <sup>ab</sup>	10.3±0.2 <sup>bc</sup>
Biomass (g)	40.6±17.5 <sup>a</sup>	114±19.4 <sup>b</sup>	251.3±33.3 <sup>c</sup>	98.3±29.3 <sup>b</sup>	106±11.5 <sup>b</sup>
SR (%)	47±6 <sup>a</sup>	56±2 <sup>ab</sup>	63±11 <sup>b</sup>	79±6 <sup>c</sup>	61±2 <sup>b</sup>
FCR	1.9±0.2 <sup>b</sup>	1.8±0.1 <sup>b</sup>	1.5±0.1 <sup>a</sup>	1.6±0.1 <sup>a</sup>	1.8±0.2 <sup>b</sup>

Note. SGR - specific growth rate; SR - survival rate; FCR - feed conversion ratio, different superscripts indicate significant differences (p<0.05).

Table 2 shows the water quality data collected during the study, which included sodium, magnesium, calcium, potassium, pH, temperature, and dissolved oxygen. The water quality in the culture media is still in accordance with the standards for Pacific white shrimp culture.

Table 2

## Water quality parameters during culture

<i>N</i> <sub>o</sub>	Parameters	Unit	5 ppt	10 ppt	15 ppt	20 ppt	25 ppt	Standards
1	Sodium	mg L <sup>-1</sup>	1195	2384.5	3585	4774.6	5964.1	1522-7609 (Boyd 2018)
2	Magnesium	mg L <sup>-1</sup>	130.6	258.5	387.5	515.3	643.2	196-976 (Boyd 2018)
3	Calcium	mg L <sup>-1</sup>	73	142.2	212	281.2	350.4	60-290 (Boyd2018)
4	Potassium	mg L <sup>-1</sup>	65.1	128.8	193	256.6	320.3	55-275 (Boyd 2018)
5	pH		7.2-7.9	7.2-7.9	7.3-7.9	7.3-7.9	7.2-7.9	7.4-8.9 (Wyban & Sweeney 1991)
6	Temperature	°C	28-29	28-29	28-29	28-29	28-29	23-30 (Briggs et al 2004)
7	Dissolved oxygen	mg L <sup>-1</sup>	4.6-6.2	4.6-6.2	4.9-6.4	4.9-6.4	4.8-6.5	4-5 (Wyban & Sweeney 1991)

The use of single-step acclimation reduced SR of Pacific white shrimp. The results show that the SR of Pacific white shrimp ranged between 47 and 79%. According to Jayasankar et al (2009), the single-step acclimation of Pacific white shrimp in a media of 5 ppt salinity led to a SR of 53%. A sudden drop in salinity over a wide range makes osmoregulation difficult for aquatic animals, which can lead to death (Laramore et al 2001).

The highest SR was observed in the culture media with a salinity of 20 ppt, 79±6%. The shrimp had little difficulty in adapting to a salinity of 20 ppt, which was similar to the salinity of the previous culture media. Salinity differences have a significant impact on the SR of Pacific white shrimp. Meanwhile, shrimp mortality during rearing could be caused by a large difference in the osmotic pressure between the shrimp's body and its environment. Physiological processes can be disrupted when the osmotic value between the body fluids and the environment is too high, leading to stress and cell damage (Brocker et al 2012). Additionally, the SR of shrimps can be affected by the frequency of shrimp molting, cannibalism, and water quality. The low salinity treatment led to the incomplete molting of vannamei shrimps since the exoskeleton was not hard enough to trigger the release of the carapace. Therefore, this reduced the SR of Pacific white shrimp, provoking their gradual death.

The Pacific white shrimp in the culture media with a salinity of 15 ppt produced the best growth (SGR=12.3±0.1%), which was accompanied by a drop in the value of FCR. According to Wyban & Sweeney (1991) and Davis et al (2002), Pacific white shrimp grow very well at low salinities around 10-15 ppt, where there is an iso-osmotic equilibrium between the environment and the hemolymph of the shrimp. According to Jaffer et al (2020), the isosmotic point occurs at a salinity of 21.1 ppt. A previous study by Rahman et al (2016) obtained results consistent with those of this study, discovering that a salinity treatment of 15 ppt is the most effective for Pacific white shrimp growth. According to Bückle et al (2006), *L. vannamei* hyperregulates between salinities of 10 to 20 ppt and hyporegulates between 20 to 40 ppt.

The osmotic pressure of the culture medium has an effect on the osmoregulation of aquatic organisms, such as shrimp. According to Anggoro et al (2018), shrimps lose much water through the diffusion process at high salinity (hypertonic). Furthermore, they adapt to this condition by drinking much water and avoiding excess salt through various mechanisms, such as the hardening of the exoskeleton, which inhibits the molting process. The regulation of ion concentration in the blood of crustaceans is very important, since the osmotic pressure of the blood is higher than the external pressure of the environment at low salinity, and vice versa at high salinity. Additionally, the differences in osmotic pressure affect the bioenergetics of aquatic organisms (Rivera-Ingraham & Lignot 2017). According to Anggoro et al (2018), the salinity of iso-osmotic media has an effect on the level of osmotic workload required to maintain osmolarity pressure balance (media and blood) or electrolyte content balance; a higher osmotic work rate means more energy wasted on osmotic performance, which leads to a reduction in shrimp growth. According to Rachmawati et al (2012), the growth of white vannamei shrimps can also be inhibited when more energy is consumed during osmoregulation, since this reduces the amount of energy available for growth. Therefore,

aquatic organisms will grow optimally if the culture medium is iso-osmotic with their body fluids (Ferraris et al 1987).

Minerals are important for the growth and survival of cultured shrimps, especially potassium (Kaligis 2010), calcium (Kaligis 2015), and magnesium (Roy et al 2010). The salinity of the water can give a measure of its mineral content, which has an influence on the growth performance of shrimps, even though salinity itself plays a role in SGR and SR (Boyd 2018). Shrimp cultures often experience mineral deficiencies in low salinity, such as calcium and potassium deficiency. Table 3 shows the calcium and potassium concentrations in the culture media, which were still within the standards for Pacific white shrimp farming.

Table 3

Calcium and potassium concentrations in culture media and standards

Salinity (ppt)	Current study		Standards (Boyd 2018)	
	Calcium (mg L <sup>-1</sup> )	Potassium (mg L <sup>-1</sup> )	Calcium (mg L <sup>-1</sup> )	Potassium (mg L <sup>-1</sup> )
5	73	65	58	55
10	142.2	129	116	110
15	212	193	174	165
20	281.2	257	232	220
25	350.4	320	290	275

**Conclusions.** The salinity of the culture medium had an effect on SGR, SR, and FCR of Pacific white shrimp. The optimal SGR, biomass, and FCR occurred at 15 ppt salinity, while the optimal SR occurred at 20 ppt salinity. Therefore, it can be concluded that a media salinity of 15 ppt provided the best performance for Pacific white shrimp rearing.

**Acknowledgements.** This study was funded by Lampung University through the Postgraduate Research program in 2021.

**Conflict of Interest.** The authors declare that there is no conflict of interest.

## References

- Anggoro S. D., Suprpto D., Purwanti F., 2018 Osmoregulation pattern of fingerling vannamee shrimp (*Litopenaeus vannamei*) rearing in three molt stage iso-osmotic media. Ilmu Kelautan 23(3):119-122.
- Boyd C. E., 2018 Revisiting ionic imbalance in low-salinity shrimp aquaculture. Global Aquaculture Advocate, 4 p.
- Briggs M., Funge-Smith S., Subasinghe R., Phillips M., 2004 Introductions and movement of *Penaeus vannamei* and *Penaeus stylirostris* in Asia and the Pacific. RAP publication 2004/10. FAO Regional Office for Asia and the Pacific, Bangkok, Thailand, 1-92.
- Brocker C., Thompson D. C., Vasiliou V., 2012 The role of hyperosmotic stress in inflammation and disease. Biomolecular Concepts 3(4):345-364.
- Bückle L. F., Barón B., Hernández M., 2006 Osmoregulatory capacity of the shrimp *Litopenaeus vannamei* at different temperatures and salinities, and optimal culture environment. Revista de Biología Tropical 54(3):745-753.
- Cheng K. M., Hu C. Q., Liu Y. N., Zheng S. X., Qi X. J., 2006 Effects of dietary calcium, phosphorus, and calcium/phosphorus ratio on the growth and tissue mineralization of *Litopenaeus vannamei* reared in low-salinity water. Aquaculture 251(2-4):472-483.
- Chong-Robles J., Charmantier G., Boulo V., Lizárraga-Valdéz J., Enríquez-Paredes L. M., Giffard-Mena I., 2014 Osmoregulation pattern and salinity tolerance of the white shrimp *Litopenaeus vannamei* (Boone, 1931) during post-embryonic development. Aquaculture 422-423:261-267.

- Davis D. A., Saoud I. P., McGraw W. J., Rouse D. B., 2002 Consideration for *Litopenaeus vannamei* reared in inland low salinity waters. *Avances en nutricion acuicola VI memories del VI Simposium Internacional de Nutricion Acuicola 3 al 6 de September del 2002*, Cancun, Quintanaa Roo, pp. 73-90.
- Ferraris R. P., Estepa F. D. P., De Jesus E. G., Ladja J. M., 1987 Osmoregulation in *Penaeus monodon* effects of molting and external salinity. *The First Asean Fisheries Forum*, Manila, pp. 637-640.
- Gao W., Tian L., Huang T., Yao M., Hu W., Xu Q., 2016 Effect of salinity on the growth performance, osmolarity and metabolism-related gene expression in white shrimp *Litopenaeus vannamei*. *Aquaculture Reports* 4:125-129.
- Jaffer Y. D., Saraswathy R., Ishfaq M., Antony J., Bundela D. S., Sharma P. C., 2020 Effect of low salinity on the growth and survival of juvenile Pacific white shrimp, *Penaeus vannamei*: A revival. *Aquaculture* 515:734561.
- Jayasankar V., Jasmani S., Nomura T., Nohara S., Huong D. T. T., Wilder M. N., 2009 Low salinity rearing of the Pacific white shrimp *Litopenaeus vannamei*: Acclimation, survival and growth of postlarvae and juveniles. *JARQ* 43(4):345-350.
- Kaligis E. Y., 2010 [Growth rate, feed utilization efficiency, body potassium content, and postlarvae vaname (*Litopenaeus vannamei*, Boone) osmotic gradient on different potassium media]. *Jurnal Perikanan dan Kelautan Tropis* 6(2):92-97. [In Indonesian].
- Kaligis E. Y., 2015 [Growth response of white shrimp (*Litopenaeus vannamei*) reared in low salinity medium, fed different protein and calcium levels]. *Jurnal Ilmu dan Teknologi Kelautan Tropis* 7(1):225-234. [In Indonesian].
- Laramore S., Laramore C. R., Scarpa J., 2001 Effect of low salinity on growth and survival of postlarvae and juvenile *Litopenaeus vannamei*. *Journal of the World Aquaculture Society* 32(4):385-392.
- Lukas A. Y. H., Djokosetiyanto D., Budiardi T., Sudrajat A. O., Affandi R., 2017 Optimization of salinity and calcium on Indonesian shortfin eel *Anguilla bicolor* maintenance. *AAFL Bioflux* 10(4):951-961.
- Rachmawati D., Hutabarat J., Anggoro S., 2012 [Effect of different media salinity on the growth of tiger snail (*Babylonia spirata* L.) in the domestication process]. *Jurnal Ilmu Kelautan* 2(3):249-257. [In Indonesian].
- Rahman F., Rusliadi, Putra I., 2016 [Growth and survival rate of western white prawns (*Litopenaeus vannamei*) on different salinity]. *Journal Online Mahasiswa* 3(1). [In Indonesian].
- Rivera-Ingraham G., Lignot J. V., 2017 Osmoregulation, bioenergetics and oxidative stress in coastal marine invertebrates: Raising the questions for future research. *The Journal of Experimental Biology* 220(10):1749-1760.
- Roy L. A., Davis D. A., Saoud I. P., Boyd C. A., Pine H. J., Boyd C. E., 2010 Shrimp culture in inland low salinity waters. *Reviews in Aquaculture* 2(4):191-208.
- Supono, 2021 Current status of technical and economic analysis of inland shrimp culture in Lampung Province, Indonesia. *AAFL Bioflux* 14(1):218-216.
- Susilowati T., Hutabarat J., Anggoro S., Zainuri M., 2014 The improvement of the survival, growth and production of vaname shrimp (*Litopenaeus vannamei*) and seaweed (*Gracilaria verucosa*) based on polyculture cultivation. *International Journal of Marine and Aquatic Resource Conservation and Co-existence* 1(1):6-11.
- Wyban J. A., Sweeney J. N., 1991 Intensive shrimp production technology. *The Institute*, 158 p.
- \*\*\* U.S. EPA, 1994 Method 200.7: Determination of metals and trace elements in water and wastes by inductively coupled plasma-atomic emission spectrometry. Cincinnati, Ohio.

Received: 23 December 2021. Accepted: 26 February 2022. Published online: 30 April 2022.

Authors:

Supono, Department of Coastal and Marine Zone Management, Postgraduate Program, Lampung University, 1 Sumantri Brojonegoro St., 35141 Bandar Lampung, Indonesia, e-mail: supono\_unila@yahoo.com

Rehulina Tresia Pinem, Department of Coastal and Marine Zone Management, Postgraduate Program, Lampung University, 1 Sumantri Brojonegoro St., 35141 Bandar Lampung, Indonesia, e-mail: tresiarehulina@gmail.com

Munti Sarida, Department of Aquaculture, Faculty of Agriculture, Lampung University, 1 Sumantri Brojonegoro St., 35141 Bandar Lampung, Indonesia, e-mail: munti.sarida@fp.unila.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Supono, Pinem R. T., Sarida M., 2022 The growth performance of the Pacific white shrimp (*Litopenaeus vannamei*) cultured at various salinity conditions using single step acclimation. AACL Bioflux 15(2):1061-1066.

## ● 13% Overall Similarity

Top sources found in the following databases:

- 13% Internet database
- 10% Publications database
- Crossref database
- Crossref Posted Content database
- 8% Submitted Works database

### TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	<b>Universitas Muhammadiyah Purwokerto on 2022-11-02</b>	2%
	Submitted works	
2	<b>banglajol.info</b>	1%
	Internet	
3	<b>researchgate.net</b>	1%
	Internet	
4	<b>bioflux.com.ro</b>	<1%
	Internet	
5	<b>scribd.com</b>	<1%
	Internet	
6	<b>Supono Supono, Destriana Puspitasari, Munti Sarida. "Pengaruh Pena...</b>	<1%
	Crossref	
7	<b>Cinquanta, L.. "Physical pre-treatment of plums (Prunus domestica). P...</b>	<1%
	Crossref	
8	<b>garuda.ristekbrin.go.id</b>	<1%
	Internet	



9	<b>scilit.net</b>	Internet	<1%
10	<b>Jesus T. Ponce-Palafox, Ángel Alcalá Pavia, Dalia G. Mendoza López, ...</b>	Crossref	<1%
11	<b>Lilongwe University of Agriculture and Natural Resources on 2021-08-28</b>	Submitted works	<1%
12	<b>Luis Daniel Moreno-Figueroa, José Naranjo-Páramo, Alfredo Hernández...</b>	Crossref	<1%
13	<b>digilib.unila.ac.id</b>	Internet	<1%
14	<b>link.springer.com</b>	Internet	<1%
15	<b>Universitas Diponegoro on 2020-02-10</b>	Submitted works	<1%
16	<b>cfile205.uf.daum.net</b>	Internet	<1%
17	<b>etd.aau.edu.et</b>	Internet	<1%
18	<b>journal.trunojoyo.ac.id</b>	Internet	<1%
19	<b>mail.scialert.net</b>	Internet	<1%

**● Excluded from Similarity Report**

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)

## EXCLUDED SOURCES

---

<b>bioflux.com.ro</b>	<b>78%</b>
Internet	
<hr/>	
<b>repository.lppm.unila.ac.id</b>	<b>16%</b>
Internet	
<hr/>	
<b>University of Stirling on 2022-12-04</b>	<b>&lt;1%</b>
Submitted works	
<hr/>	
<b>University of Stirling on 2022-12-04</b>	<b>&lt;1%</b>
Submitted works	