PAPER NAME Prosiding IOP (earthworm).pdf	author Sri Y
WORD COUNT	CHARACTER COUNT
3761 Words	21176 Characters
PAGE COUNT	FILE SIZE
18 Pages	1020.3KB
SUBMISSION DATE	REPORT DATE
Jan 18, 2023 1:05 PM GMT+7	Jan 18, 2023 1:06 PM GMT+7

• 7% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

• 7% Publications database

• Excluded from Similarity Report

- Internet database
- Crossref Posted Content database
- Bibliographic material

- Crossref database
- Submitted Works database

PAPER • OPEN ACCESS

Earthworm population under different soil tillage and herbicide application at integrated field laboratory agriculture faculty, University of Lampung

To cite this article: S Yusnaini et al 2018 IOP Conf. Ser.: Earth Environ. Sci. 215 012015

View the article online for updates and enhancements.



IOP ebooks[™]

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

PAPER • OPEN ACCESS

International Conference on Organic Agriculture in the Tropics: State of the Art, Challenges and Opportunities

To cite this article: 2018 IOP Conf. Ser.: Earth Environ. Sci. 215 011001

View the article online for updates and enhancements.

International Conference on Organic Agriculture in the Tropics: State-of-the-Art, Challenges and Opportunities

August 20-24th, 2017

Content from 2 is work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1 IOP Conf. Series: Earth and Environmental Science **215** (2018) 011001

doi:10.1088/1755-1315/215/1/011001

EDITORS

Stefaan De Neve University of Ghent, Belgium stefaan.deneve@ugent.be

Benito Heru Purwanto Universitas Gadjah Mada, Indonesia benito@ugm.ac.id

Alim Isnansetyo Universitas Gadjah Mada, Indonesia isnansetyo@ugm.ac.id

Sri Nuryani Hidayah Utami UniversitasGadjahMada, Indonesia nuryani@ugm.ac.id

Junun Sartohadi Universitas Gadjah Mada, Indonesia junun@ugm.ac.id

Nur Ainun H.J. Pulungan Universitas Gadjah Mada, Indonesia nurainun.pulungan@ugm.ac.id

REVIEWERS

- 1. Alim Isnansetyo(Universitas Gadjah Mada, Indonesia)
- 2. Azwar Ma'as(Universitas Gadjah Mada, Indonesia)
- 3. Benito Heru Purwanto(Universitas Gadjah Mada, Indonesia)
- 4. Bram Moeskops (IFOAM-EU Group, Senior Scientific Coordinator TP Organics)
- 5. Clara Nicholls (University of California, Berkeley, US President of SOCLA (Latin American Scientific Society for Agro-ecology)
- 6. Didik Indradewa (Universitas Gadjah Mada, Indonesia)
- 7. Edhi Martono (Universitas Gadjah Mada, Indonesia)
- 8. Eko Hanudin (Universitas Gadjah Mada, Indonesia)
- 9. Gerold Rahmann (Thünen Institute of Organic Farming, Director, Braunschweig, Germany ISOFAR President)
- 10. Irham (Universitas Gadjah Mada, Indonesia)
- 11. Junun Sartohadi (Universitas Gadjah Mada, Indonesia)
- 12. Mette Olaf Nielsen (University of Copenhagen, Copenhagen, Denmark)
- 13. Michael Böhme (Humboldt Universität zu Berlin, Berlin, Germany)
- 14. Mohammad Nurcholis (Universitas Pembangunan Nasional "Veteran" Yogyakarta, Indonesia)
- 15. Reza Ardakani (Azad University, Karaj, Iran Trent University, Peterborough, Canada)
- 16. Sri Nuryani Hidayah Utami (Universitas Gadjah Mada, Indonesia)
- 17. Tomohide Sugino (Japan International Research Center for Agricultural Sciences (JIRCAS))
- 18. Wenliang Wu (China Agricultural University, Beijing, China)

TABLE OF CONTENT

	Authors	Paper Title
Soil Management and Soil Quality	Waluyati	Demand of organic fertilizer at rice cultivation in Klaten Regency
	Fitriatin	The Effect of phosphate solubilizing bacteria and organic fertilizer on phosphatase, available P, P uptake and growth sweet corn in andisols
	Hamzah	Biochar: Effects on crop growth
	Dermiyati	Soil fauna population during the maize (Zea mays I.) growth with the addition of organonitrophos, inorganic fertilizer and biochar
	Suryatmana	The potential of the mushroom log waste (MLW) and Azotobacter vinelandii to improve hydrocarbon biodegradation for rehabilitation of petroleum contaminated soil
	Sijabat	Labile carbon fraction, humic acid, and fulvic acid on organic and conventional farming of rice field in Imogiri and Berbah
	Setiawati	Utilization Azolla pinnata as substitution of manure to improve organic rice yield and paddy soil health
	Afany	The impact of horticulture farming practices in andisols forest on soils organic-p content in Tawangmangu, Central Java
	Widowaty	The soil organic dynamics from types biochar-organic fertilizers and soil
	Damarmoyo	Soil physical properties and abundance of soil fauna in conventional and organic rice field
	Putra	Influence of goat manure and Azolla on soil properties, nitrogen use efficiency, growth and yield of organic rice farming in Indonesia
Natural Plant Protection	Mora	Evaluation of botanical

		insecticides in controlling the
		population of fall armyworms
		(Spodoptera frugiperda Smith)
		present on corn crops (<i>Zea mays</i>)
		located in Santa Cruz,
		Guanacaste
	Chozin	The potential use of plant
		extracts for controlling northern
		leaf blight on organic sweet corn
		production
	Sudiatmiko	Changes in chemical properties
	, ,	of soil in an organic agriculture
		system
	Yusnaini	Farthworm population under
		different soil tillage and
		herbicide application at
		integrated field laboratory
		agriculture faculty University of
		Lampung
Organic Livestock Production	Realon-Garcia	Effects of diet composition on
		the energy balance of free-
		ranging organic broilers
Food Quality and Food Savety	Vunindanova	The analysis of endogenous
Tood Quality and Tood Savety	Turintuanova	auxin of shallot and its effect on
		the germination and the growth
		of organically cultivated molon
		(Cucumis malo)
	Kristamtini	(Cocumis melo)
	KIIStallitilli	of 20 local rice genetic resources
		of Vorwakarta Indonesia
	Kusmanadhi	Dreduction and quality of come
	Kusinanaum	adamama variation as affected
		by residual offect of worm
		sompost application
Feenemic Analysis of Organic Ferming		Maize forming performance in
Economic Analysis of Organic Farming		dry land with biochar and
	Acnah	manura in Kalitangah Villaga
	ASIIdii	Danggungraio District Plitar
		Paragonev Indonesia
		Productivity, profitability, and
		marketing of vegetable organic
	Irawan	naducto losson losent from
	lidwall	Pangkit Marbabu farmar's group
		in Somerong, Control Jove
		Organia on field recorded
		organic on-field research
		complined with information
	c	technology for last uptake,
	Sivakumar	sustainable, and profitable
		nuture using both macro and
		micro analysis of the system as a
		whole

Soil Bio-technology Effect of cow manure, neem compost and straw compost towards N uptake and soil fauna abundance in inceptisol paddy field, Berbah, Sleman Dwinta The effect of cow manure and neem compost toward NPK uptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, Indonesia Ardiantika Effect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, Kalitirto Panjaitan The effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.) Purwaningsih Paddy straw compost application to increase nitrogen fixation on soybean cultivars
Soil Bio-technology Effect of cow manure, neem compost and straw compost towards N uptake and soil fauna abundance in inceptisol paddy field, Berbah, Sleman Dwinta The effect of cow manure and neem compost toward NPK uptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, Indonesia Ardiantika Effect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, Kalitirto Panjaitan The effect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, Kalitirto Panjaitan The effect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, Kalitirto Panjaitan The effect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, Kalitirto Panjaitan The effect of organic fertilizer on nitrogen uptake and yield of red chili (Capsicum annum L.) Purwaningsih Paddy straw compost application on sovbean cultivars
DefinitionDefinitionDefinitionDefinitionDwintaDwintaCompost and straw compost towards N uptake and soil fauna abundance in inceptisol paddy field, Berbah, SlemanUtamiThe effect of cow manure and neem compost toward NPK uptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, IndonesiaArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (Capsicum annum L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
Dwintatowards N uptake and soil fauna abundance in inceptisol paddy field, Berbah, SlemanUtamiThe effect of cow manure and neem compost toward NPK uptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, IndonesiaArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (Capsicum annum L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
abundance in inceptisol paddy field, Berbah, SlemanUtamiThe effect of cow manure and neem compost toward NPK uptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, IndonesiaArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.)PurwaningsihPady straw compost application to increase nitrogen fixation on soybean cultivars
field, Berbah, SlemanUtamiThe effect of cow manure and neem compost toward NPK uptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, IndonesiaArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
UtamiThe effect of cow manure and neem compost toward NPK uptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, IndonesiaArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
Utamineem compost toward NPK uptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, IndonesiaArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
Utamiuptake, soil respiration, and rice production in organic paddy field in Imogiri Bantul, IndonesiaArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
Panjaitanproduction in organic paddy field in Imogiri Bantul, IndonesiaPanjaitanEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
ArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
ArdiantikaEffect of organic fertilizer on nitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
Ardiantikanitrogen uptake and yield of two different rice varieties in inceptisol, KalitirtoPanjaitanThe effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (Capsicum annum L.)PurwaningsihPaddy straw compost application to increase nitrogen fixation on soybean cultivars
Panjaitan Interent filter varieties in inceptisol, Kalitirto Panjaitan The effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.) Purwaningsih Paddy straw compost application to increase nitrogen fixation on soybean cultivars
Panjaitan The effect of organic fertilizer and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.) Purwaningsih Paddy straw compost application to increase nitrogen fixation on soybean cultivars
Panjaitan and compost of volcanic ash on growth and yield of red chili (<i>Capsicum annum</i> L.) Purwaningsih Paddy straw compost application to increase nitrogen fixation on soybean cultivars
Panjaitan growth and yield of red chili (Capsicum annum L.) Paddy straw compost application Purwaningsih to increase nitrogen fixation on soybean cultivars soybean cultivars
Capsicum annum L.)Paddy straw compost applicationPurwaningsihto increase nitrogen fixation onsoybean cultivars
Paddy straw compost application Purwaningsih to increase nitrogen fixation on soybean cultivars
Purwaningsih to increase nitrogen fixation on soybean cultivars
soybean cultivars
,
Compost derived from local
Riwandi organic materials as source of
plant nutrients
Effect of organic fertilizer on
Himawati
rice uptake in inceptisois Kalitirta Slaman
The effect of organic fertilizer
hased on Azolla microphylla
biomass and plant spacing to N
Utami and P uptake, soil compaction
and the yield of Pandanwangi
rice
Other Related Themes Effect of flooding duration on
nitrous oxide emission from
Purwanto organic and conventional rice
cultivation system in Central
Java, Indonesia
OFC Long-term organic mulching and
no-tillage practice increase
Niswati population and biomass of
eartnworm in sugarcane

INDEX

Α

Abundance of soil fauna Aerobic fermentation Anaerobic fermentation Andisols Andisols forest Antifungal activity Auxin Azolla green manure *Azolla microphylla Azolla pinnata Azotobacter vinelandii*

В

Bagasse Bangkit Merbabu farmer group Biochar Biodegradation Biophysic Botanical fungicide Botanical insecticides

С

Carbon Cassava Chicken excreta Compost Compost of volcanic dust Consumer preference Conventional practice Conventional rice field Corn cob Cow manure Crop performance Crop yield Curricula

D

Demand of organic fertilizer Disease intensity Dry land

Ε

Earthworm

Edamame Education Energy requirements Extension **F** Fixation Fulvic acid **IOP** Publishing

G Genetic resources Goat feces Goat manure Green soybean Growth analysis

H Health Herbicide Horticultural farming Humic acid

I Image processing Indonesia

J Jengkok

K Kinetics Klaten Regency

L

Labile carbon fractions Liquid bio-fertilizer Litosol Local rice

Μ

Macro Maize Maize farming Mediteran Melon

ICOAT 2017

IOP Conf. Series: Earth and Environmental Science 215 (2018) 011001 doi:10.1088/1755-1315/215/1/011001

Menthik Susu Mulching

Ν

N₂O emissions N use efficiency Neem cake Neem compost Nitrogen Nothern leaf blight Nutrient

0

On-field research Organic Organic agriculture Organic farming Organic fertilizer Organic mulching Organic practice Organic rice Organic rice Organic rice field Organic vegetable Organic-P Outdoor run

Ρ

Paddy soil Paddy straw biochar Parasitism Performance Petroleum waste pH Phosphatase Phosphate solubilizing Plant extracts Plant spacing Potassium

R

Red chilli Regosol Release Restricted feeding *Rhizobium japonicum* Rice Rice husk Rice husk biochar Rice production Rice productivity

S

Shallot Socio economic Soil amendment Soil chemical properties Soil compaction Soil fauna abundance Soil flooding Soil physical properties Soil properties Soil rehabilitation Soil respiration Soybean Spodoptera frugiperda Students Sugar cane Sustainability Sweet corn

IOP Publishing

T Tawangmangu Tillage Tropical acid soil

U Universities Uptake

W Worm compost

Y Yogyakarta



PAPER • OPEN ACCESS

Peer review statement

To cite this article: 2018 IOP Conf. Ser.: Earth Environ. Sci. 215 011002

View the article online for updates and enhancements.

Peer review statement

All papers published in this volume of *IOP Conference Series: Earth and Environmental Science* have been peer reviewed through processes administered by the proceedings Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

Content from 2 as work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

IOP Publishing

Earthworm population under different soil tillage and herbicide application at integrated field laboratory agriculture faculty, University of Lampung

S Yusnaini^{1*}, A Niswati¹, M A S Arif¹, Y Komalasari¹ and N Kaneko^{2*}

¹Faculty of Agriculture, University of Lampung, Jl. Prof. Sumantri Brojonegoro No.1 Bandar Lampung 35145, Lampung, Indonesia ²Soil Ecology Research Group, Graduate School of Environment and Information Sciences, YNU,79-7 Toki Wadai, Yokohama 240-8501, Japan

*Email: sri.yusnaini@fp.unila.ac.id and kanekono@ynu.ac.jp

Abstract. This study were aimed to determine effect of soil tillage and herbicide on earthworm populations. The experiment was designed by using completely randomize block design with two factors, i.e.; soil tillage systems consists of full tillage (T_1) and minimum tillage (T_0) and herbicide application (H) consists of herbicide (H₁) and non-herbicide (H₀). Herbicides with isopropylamine glyphosate + 2.4D active ingredient was application as treatment at level dosage 160 mL per 16 L of water. Soil sampling by using monolith and earthworm were collect by hand sorting methods. The data were analyzed by ANOVA and LSD test. The results showed that the soil tillage has no effect on the population and biomass of earthworms, but application of herbicide affected on earthworm population at one day after herbicide application. Earthworm population were higher in plot with no herbicide application (86 ind \cdot m⁻²) than that plot with herbicide application (33 ind \cdot m⁻²). However, at all combination treatment, earthworm population were increase followed by cassava growth. Population and biomass of earthworms found to be higher in the soil layer 0-10 cm than that 10-20 cm and 20-30 cm soil layers. Furthermore, soil water content had correlation with earthworm population.

1. Introduction

Earthworm is a soil macro fauna that presence in the soil is very important role in many soil functions such as nutrient availability, soil structure and organic matter dynamics [1]. However, earthworm distribution and activity are affected by soil moisture, organic matter, texture and pH [2]. Soil management likes soil tillage in agricultural soils can affect soil biota through changes in habitat [3],loss of organic matter [4], moisture and temperature dynamics [2]. Larthworm population change due to soil tillage depends on tillage intensity [2, 5]. Furthermore, leduced tillage intensity increased earthworm abundances and species diversity [3]. However, in reduce tillage systems, weeds become a problem [6], especially grasses and perennial weeds [7] and application of chemical herbicide have become the weed control strategy. Herbicide applications may reduce or maintain weed diversity [8, 9], but they can become a toxicological risk for earthworm. Earthworm are sensitive to the presence of chemical in the soil due to chemoreceptor distributed on their body surface [10]. Herbicides affect the feeding behavior of earthworms, which was reflected in the weight loss and reproductive capacity [11] and reduce cocoon production [12].

Content fron ² as work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd

ICOAT 2017	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 215 (2018) 012015	doi:10.1088/1755-1315/215/1/012015

The mortality of earthworms in soils and excessive use of herbicide is still vague. Some herbicides are directly toxic to earthworm while others have virtually no effects [13]. Based on this condition, it's necessary to study the effect of soil tillage and chemical herbicide application on the existence of earthworm in cassava cultivation.

2. Materials and Methods

⁵he research was conducted at the Integrated Field Laboratory, Agriculture Faculty University of Lampung (50°22'11.38" S, 1050°14'25.96" E) on May 2014. The experiment was designed by using completely randomize block design (RCBD) with two factors, i.e.; soil tillage systems consists of full tillage (T1) and minimum tillage (T0) and herbicide application (H) consists of with herbicide (H1) and without herbicide (H0). The Herbicides with active ingredient isopropylamine glyphosate + 2.4D was application as treatment at level dosage 160 mL per tank (16 L of water). Earthworm population sampling during cassava growth was done at 0, 3, 6, 11 month after planting cassava by using soil monolith and earthworm were collect by hand sorting methods. The data were analyzed by ANOVA and LSD test at 5% significant level and correlation between earthworm population and biomass with soil organic-C, soil pH, soil water content and soil temperature.

3. Results and Discussion

Based on statistical analysis lifect of tillage and herbicides on the earthworm population and biomass could be seen in table 1. Table 1 shown that the tillage systems has no effect on the earthworm populations and biomass in each month of observation, whereas herbicide application affect on earthworm populations only at 0 MAP.

Table 1. Analysis of variant effect of soil tillage and herbicideapplication on earthworm populations and biomass.

	_			Mo	nth			
Treatment	0 MAP		3MAP		6 MAP		11 MAP	
	[©] P	FB	EP	EB	EP	EB	EP	EB
Tillage (T)	ns		ns	ns	ns	ns	ns	ns
Herbicide (H)	*	ns	ns	ns	*	ns	ns	ns
ТхН	ns	ns	ns	ns	ns	ns	ns	ns

Note: MAP= Months after planting; EP= earthworms population; EB= earthworm biomass; ns= non-significant; *= significant at 5% level.

3.1. Earthworm populations and biomass

Based on statistical analysis, tillage systems has no effect on earthworm populations and biomass. It is might be caused that the duration of experiment is too short. Many atterature search revealed few studies that have assessed the effects of tillage systems on earthworms over the short and medium timescales simultaneously in both conventional (full) tillage and minimum tillage. However, it can be seen in figure 1, earthworm abundance on minimum tillage relatively higher than that earthworm population on full tillage due organic material content. Organic materials on minimum tillageare came from weeds and litter at surface layers, whereas on full tillage weeds and litter are cleared off from surface layer. Furthermore, earthworm distribution along soil depth was different. Earthwormpopulationper layer shows that neither the population nor the earthworm biomass was higher in the soil top layers (0–10 cm). This is caused by rich of organic material at top soil.Organic matter is one of the deciding factors that affect of macrofauna distribution on the below and above ground as well as a source of energy for soil fauna and microorganisms[14, 15].



Figure 1. Earthworm population at sampling time 0, 3, 6, and 11 MAP (MAP: Months after planting); T0: minimum tillage; T1: full tillage; H0: without herbicides; H1: herbicide application).

Figure 1 shows that earthworm population at 11 MAP was higher compared to others observation and the highest number of earthworm is in minimum tillage without herbicide application (T_0H_0) 190 ind·m⁻². The highest number of earthworm at 11 MAP might be caused by the full canopy of cassava make the soil microclimate, such us soil temperature and moisture are comfortable to earthworm living [2].

Based on ANOVA at table 1, herbicide application had significant effect on earthworm populations only at 0 MAP (before planting). Based on LSD test, weed control using herbicides has an earthworm population lower than that without application of herbicides (table 2), however at 3, 6 and 11 MAP herbicide application had no effect on earthworm number. The active ingredient glyphosate was eaten by earthworms can have an effect on earthworm breeding [16].

Table 2. The effect of herbicides	s application on	i earthworm po	pulation at 0 MAP.
------------------------------------------	------------------	----------------	--------------------

Treatment	Population (ind m ⁻²)
Without Herbicides (H ₀)	86 a
Herbicide Applications (H ₁)	33 b
LSD 0.05	37

3.2. The relationship between earthworm's population and biomass with some soil properties

Other factors that have an effect on earthworm populations i.e. soil properties such as soil organic carbon, soil pH, soil temperature and soil moisture content (humidity). The relationship between earthworm's population and biomass with some soil properties were shown in table 3. Based table 3, soil organic-C, and soil pH, have no correlation with earthworm populations and biomass. However, soil water content had positively correlated with earthworms population, it is caused by earthworms are extremely sensitive with soil humidity. Soil water content is needed by earthworms to keep his skin in order to function normally i.e. for respiration. Earthworm's biomass consists of 75–90% water, so the humidity is very important to earthworm life cycle. When the soil water content is too high, earthworms will move to find a suitable and then their skins could be normally functioning. The

optimum range of soil humidity for earthworm's growth is 15–50% [17]. Furthermore, temperature had positive correlation with earthworm populations. Earthworm is soil biota that very sensitive to the temperature, however the earthworms can still tolerate in temperatures above 30°C with high humidity. A temperature that is too low is also not preferred by earthworms, as it will affect hatching cocoon earthworms.

Table	3. Correlation	between	earthworms	population	$(ind n^{-2})$	and	earthworm
biomas	ss $(g \cdot m^{-2})$ with	soil prope	rties.				

	The correlation coefficient (r)			
Variable/Sampling time	The number of	Farthworm biomass		
	earthworms	Lattiworm biomass		
Soil analysis Before				
Cassava Planting 0 MAP				
C-organic	2 39 ^{ns}	0.49 ^{ns}		
soil pH	0.21^{ns}	0.51 ^{ns}		
Water content of soil	0.18 ^{ns}	0.31 ^{ns}		
Soil temperature	0.36^{ns}	0.01 ^{ns}		
Soil Analysis at 3 MAP				
C-organic	0.25 ^{ns}	0.03 ^{ns}		
soil pH	0.19 ^{ns}	0.21 ^{ns}		
Water content of soil	0.46^{ns}	0.15 ^{ns}		
Soil temperature	0.29 ^{ns}	0.01 ^{ns}		
Soil Analysis at 6 MAP				
C-organic	0.02^{ns}	0.26 ^{ns}		
soil pH	$0.37^{\rm ns}$	0.24^{ns}		
Water content of soil	0.79^{*}	0.41 ^{ns}		
Soil temperature	0.51^{*}	0.41 ^{ns}		
Soil Analysis at 11 MAP				
C-organic	31 ^{ns}	0.16 ^{ns}		
soil pH	0.28^{ns}	0.49 ^{ns}		
Water content of soil	0.03^{ns}	-0.64*		
Soil temperature	-0.31^{ns}	0.21^{ns}		

Note: note ns: not significant; *: significant.

4. Conclusion

Soil tillage system both full tillage or minimum tillage does not affect the population and biomass earthworms on the growing cassava season. While, herbicide application decreased earthworms population only at one month after herbicide application (0 MAP), but has no effect on earthworm populations and biomass at sampling time 3, 6 and 11 MAP. The earthworm population and biomass was higher in the topsoil (0–10 cm) compared with the deep soil layers (10–20 cm and 20–30cm). The soil moisture content and soil temperature had positive correlation with earthworm population.

References

- [1] Edwards C A and Arancon 2004Interactions among Organic Matter, Earthworms and Microorganism in Promoting Plant Growth(New York: CRC Press) pp 328–329
- [2] Curry J P 2004 Earthworm Ecology 91–113 DOI: 10.1201/9781420039719.pt3
- [3] Van Capelle C, SchraderS and Brunotte J 2012 European Journal of Soil Biology50 165–181
- [4] Hendrix PF, Mueller B R and Bruce RR 1992 Soil Biology and Biochemistry24(12) 1357–1361
- [5] Chan K Y 2001 Soil Tillage Research 57 179–191
- [6] Zarea MJ, Ghalavand A, Mohammadi Goltapeh E and Rejali F 2010*Journal of Plant Protection Research* **50** 463–469

- [7] Plesant JM, McCollum R E and Coble H D 1990*Agronomy Journal*82 102–112
- [8] Derkensen DA, Thomas AG, Lafond GP, Loeppky H A and Swanton CJ 1995 Weed Research**35** 311–320
- [9] Fryer JD and Chancellor R J 1970*Botanical Society of the British Isles Report***11**105–118
- [10] Rienecke S A and Rienecke A J 2007 Ecotoxicology and Environmental Safety 66 244-251
- [11] Venter J M and Reinecke A J 1988 Sublethal Ecotoxicological Effects of Dielrin on the Earthworm E. foetida (oligochaeta)In: Edwards CA Neuhauser EF (Eds) Earthworms in Waste and Environmental Management (Netherlands: SPB Academic Publishing) pp 337– 353
- [12] Muthukaruppan G, Janardhanan S and Vijayalakshmi GS 2005 Journal of Soils Sediments 5 82– 86
- [13] Brown AWA1978 Ecology of Pesticides(New York: John Wiley and Sons) 525 pp
- [14] Sugiyarto, Manan E, Edwl M, Yogi S, Eko H and Lily A 2007 Jurnal Biodiversitas7 96-100
- [15] Subowo 2010 Jurnal Sumberdaya Lahan 413–25
- [16] Madani R G 2013 Jurnal Mahasiswa Pendidikan Biologi2 175-182
- [17] Rukmana R 1999 Budi Daya Cacing Tanah(Yogyakarta: Kanisius)

• 7% Overall Similarity

Top sources found in the following databases:

• 7% Publications database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Crittenden, Stephen J "Biophysical Soil Quality of Tillage Systems in Publication	3%
2	Lindsey, Alex J "Evaluation of Humic Fertilizers on Turfgrass", Iowa St Publication	2%
3	Eduard, Wijnand. "Assessment of Mould Spore Exposure and Relations Publication	1%
4	Scanlon, Kate. "Play and Development of Social Skills in Infant Rhesus	<1%
5	Nzeyimana, Innocent. "Optimizing Arabica Coffee Production Systems	<1%
6	Wei, Yunan. "An advisory system for the development of unconvention Publication	<1%