

PAPER NAME

1. Short Communication.pdf

AUTHOR

S Bakri

WORD COUNT

6847 Words

CHARACTER COUNT

36780 Characters

PAGE COUNT

9 Pages

FILE SIZE

983.5KB

SUBMISSION DATE

Jan 16, 2023 7:52 AM GMT+7

REPORT DATE

Jan 16, 2023 7:53 AM GMT+7**● 20% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 19% Internet database
- 4% Publications database
- Crossref database
- Crossref Posted Content database
- 8% Submitted Works database

● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Cited material
- Manually excluded sources

1 Short Communication:

Assessing the relationship of Sumatran elephant's movement (*Elephas maximus sumatranus*) with vegetation intensity in Kotaagung Utara, Lampung Province, Indonesia using NDVI method

SAMSUL BAKRI, DELLA TIARA MONIK[✉], AGUS SETIAWAN, GUNARDI DJOKO WINARNO

Department of Environmental Science, Postgraduate Program, Universitas Lampung. Jl. Prof. Dr. Sumantri Brojonegoro No. 1, Bandar Lampung 35145, Lampung, Indonesia. Tel./fax. +62-271-783682, ✉email: dellatiaraaa@gmail.com

Manuscript received: 1 March 2022. Revision accepted: 21 March 2022.

Abstract. Bakri S, Monik DT, Setiawan A, Winarno GD. 2022. Short Communication: Assessing the relationship of Sumatran elephant's movement (*Elephas maximus sumatranus*) with vegetation intensity in Kotaagung Utara, Lampung Province, Indonesia using NDVI method. *Biodiversitas* 23: 1920-1928. The Indonesian Elephant Conservation Forum stated that the population of Sumatran elephants (*Elephas maximus sumatranus*) had decreased dramatically up to 70% in the last 20-30 years due to illegal hunting, land conversion, and encroachment, so the availability of elephant food in the wild is gradually inadequate. With the decline of the elephant population in Indonesia, several parties such as the government, NGO, and the public are beginning to monitor the Sumatran elephant. The monitoring is carried out by observing and studying the movements of elephants using a *GPS Collar*. The research aims to find out the consistency of elephant movement and its relationship to the availability of feed as indicated by the intensity of vegetation in KPH Kotaagung Utara, Lampung Province, Indonesia. This research used GIS technology (Correlation Citra Landsat 8 OLI and BIG Demnas Data), elephant movement data using *GPS Collar* in 2020, and land use data by BPKH Lampung. The results showed that the monthly movement pattern of elephants in 2020 was consistently monitored. From January-July, elephants are in the North area, while in August-December are in the South. The intensity of elephant movement in the Mixed Shrub Dryland Agriculture area is higher than in the other areas, at 107 points or about 90% of all the areas in KPH Kotaagung Utara. Meanwhile, in the Dryland and Shrub Agriculture areas, there are 6 points each, or about 5% of all the areas. The vegetation classification in KPH Kotaagung Utara is *dense* with an NDVI value range of 0.63-0.85. Furthermore, the regression results prove that NDVI and the monthly season affect the movement of elephants with a p-value of <0.001.

Keywords: KPH Kotaagung Utara, NDVI, Sumatran elephant (*Elephas maximus sumatranus*)

Abbreviations: API: Application Programming Interface; EEC: Earth Engine Computing; GPS: Global Positioning System; H: Hypothesis; IUCN: International Union for Conservation of Nature and Natural Resources; IVI: Important Value Index; KHL: Kawasan Hutan Lindung; KPH: Kesatuan Pengelolaan Hutan; NDVI: Normalized Difference Vegetation Index; NGO: Non-Governmental Organization; NIR: Near Infra-Red; R-Sq: R-Square (Coefficient of Determination); R-Sq (Adj) is the Adjusted R-Square (Coefficient of Determination); SEE: Standard Error of Estimate; WWF: World Wide Fund for Nature

INTRODUCTION

The Sumatran elephant (*Elephas maximus sumatranus*) is the largest herbivore in Sumatra (Indonesia) with the classification *Proboscidea* (Imtiyaz et al. 2021). According to the International Union for Conservation of Nature and Natural Resources (IUCN) 2022, Sumatran elephants are included in the *Red List Data Book* in the critically endangered category. According to the Directorate of Biodiversity Conservation, Directorate General of Natural Resources and Ecosystem Conservation, Ministry of Environment and Forestry of the Republic of Indonesia Jakarta (2020), in 2017, the Sumatran elephant population was estimated at 1,694-2,038 individuals found in 36 habitat pockets and spread over seven provinces, including Aceh, North Sumatra, Riau, Jambi, South Sumatra, Bengkulu and Lampung. The Government Regulation Number P.20/MenLHK/Setjen/

Kum.1/6/2018 about Protected Types of Plants and Animals in Indonesia strengthens the protection of Sumatran elephants, with the second amendment of Regulation of the Minister of Environment and Forestry Number P.106/MenLHK/Setjen/Kum.1/12/2018, however, the conversion of forest functions to agricultural and plantation areas, resulting in depression in elephant's natural habitats as a refuge, and become one of the factors's causing the decline of elephants population (Kumar et al. 2020).

In general, Sumatran elephants can be found in stable habitats, such as tropical rainforests in some parts of Indonesia (Moßbrucker et al. 2016). In line with Shaffer et al. (2019), elephants live in groups by traveling long distances out of their roaming area to fulfill their food needs and water, find social partners, and reproduce. The availability of adequate food and water will indirectly affect the activities and behavior of the elephants.

According to Wall et al. (2021), elephants often choose to be in protected areas, with security better than in other areas, aside from the abundance of food and water. However, as time goes by, anthropogenic activities, forest encroachment, and land clearing in protected forest areas are increasing, and the home range and food chain of the elephants that inhabit the rest of the forest are interrupted (Calabrese et al. 2017). These negative activities will destroy the elephant's natural habitat and trigger conflicts between elephants and humans. Thus, monitoring is needed to determine the elephant's roaming area. Analysis of land use maps was carried out using satellite imagery to see the greenery level of plants in the observation area (Prasetyo et al. 2017). Doing mapping and observation is one of the methods used to find out the relation between the roaming area and the wild elephant feed in KPH Kotaagung Utara. The first thing to do is see the intensity of elephant movements based on the land use map. According to Nedd et al. (2021) land use and land cover are important to be understood because such data refer to the purpose and the land function, such as wild animal habitat, land change, and development, as well as the physical appearance of the earth's surface such as bodies of water, rocks, built-up land, and others. Moreover, these appearances are valuable information for nurturing elephant programs that live naturally or domestically (Meytasari et al. 2014). The development of technology nowadays, like GPS (Global Positioning System) and non-invasive molecular technic, helps increase our understanding of observing and studying the wildlife ecology, especially wild elephants (Wong et al. 2018). After that, analyzing the vegetation intensity damage at KPH Kotaagung Utara using the vegetation

indexes. Index vegetation is needed to identify pixels covered by the size of vegetation proportion. According to Xue and Su (2017), the vegetation index is obtained from the greenness of vegetation or canopy based on digital signal intensity, or what is known as a fairly simple algorithm. There are many vegetation indexes with different algorithms, one of them is the vegetation index Normalized Difference Vegetation Index (NDVI). NDVI is an index that is often used to calculate the greenery scale, counted by the multispectral information as the normalization ratio between red-bow and closer infra-red this classification can be done with a good technique using remote sensing because of the special vegetation characteristic towards electromagnetic waves (Woźniak et al. 2020). Finally, do the correlation of NDVI value with the presentation of the elephant's move. Based on the background given, this observation aims to see the consistency of the elephant's movement and the relation with the food supply availability indicated by the vegetation's intensity in KPH Kotaagung Utara, Lampung Province, Indonesia.

MATERIALS AND METHODS

Research location

The research was conducted at a protected area, namely Kawasan Hutan Lindung (KHL), located inside KPH Kotaagung Utara, Tanggamus District, Lampung Province, Indonesia (Figure 1). Data analysis was carried out using a laptop and ArcGIS software in January 2022.




Figure 1. Research location KPH Kotaagung Utara, Tanggamus District, Lampung Province, Indonesia

Procedures

Types and data source

The data were collected from secondary data obtained from the internet, newspaper, or journal (Ajayi 2017). The data collected were GPS Collar data in 2020 kindly provided by WWF, land use data in KPH Kotaagung Utara consisting of BPKH data, a constellation of Citra Lansat 8 OLI, and BIG Demnas, as well as NDVI, which were obtained by searching using Google Earth Engine.

Data collecting

Secondary data were obtained from the results of mapping the consistency of the Sumatran elephant roaming area using the GPS Collar (data WWF) in 2020, consisting of land use data, a constellation of Citra Lansat 8 OLI and BIG Demnas data, and NDVI results using Google Earth Engine. Currently, there are 24 elephants in the province of Tanggamus, which are divided into 2 groups. GPS Collars were installed on each Bunga and Citra elephant group leader. According to Nazaruddin, chairman of the Mahout Forum, the elephant fitted with the GPS Collar is a female elephant aged 25-30 years, may weigh up to 2,500 kg (5511.56 lb.) and stand about 2.19 m (7.1 ft.). The data generated from the GPS Collar is in the form of recording the points of elephant movement per day every year. The data was obtained from WWF and has been granted permission to be processed. The goal of the GPS Collar is to track the movement patterns of animals in great detail (Stabach et al. 2020).

Data analysis

Distribution of coordinate points based on land cover

The elephants wearing GPS Collars were moved across different land covers. The data obtained were then processed and analyzed using the formula:

$$\text{Types of Land Cover} = \frac{(\text{The Number of Points in Land Cover})}{(\text{The Total of All Points})} \times 100\% \dots\dots(1)$$

Normalized Difference Vegetation Index (NDVI) analysis based on Earth Engine Computing (EEC)

Vegetation intensity analysis was carried out by NDVI using the red band and Near Infra-Red (NIR) in remote sensing to determine the vegetation index from the satellite (Shafizadeh-moghadam et al. 2021). Viewing the NDVI value of an area can be done by using Earth Engine Computing. Earth Engine is a platform developed by Google with various open-source data available in the data catalog and ready for analysis in multi-petabyte sizes processed through computing services with high-performance capacities (Gorelick et al. 2017). GEE offers substantial advantages over conventional processing, including easy-to-develop and open algorithms and the ability to do various computing tasks efficiently (Xia et al. 2020). This is because the Application Programming Interface (API) can be accessed through the related website (IDE), allowing for rapid generation of prototypes and visualizations in a large cloud computing chart (Gorelick et al. 2017; Sidhu et al. 2018; Chen et al. 2021) An NDVI score near 1 indicates that the area is becoming denser in

terms of vegetation cover; and vice versa, a value near 0 indicates that the area's vegetation is becoming rarer (Pahlefi et al. 2021). Ahmed and Akter (2017) describe the NDVI algorithm as follows:

$$NDVI = \frac{(NIR-R)}{(NIR+R)} \dots\dots\dots(2)$$

Where, NDVI is the *Normalized Difference Vegetation Index*; NIR is the spectral reflectance value in the near-infrared band; R is the spectral reflectance value in the red band. The vegetation density level is classified into five categories (Table 1): cloud and water, non-vegetation, sparse vegetation, moderate vegetation, and dense vegetation (Sunaryo and Iqmi, 2015; Hardianto et al. 2021). Correlations will be made between the data and data from image processing using NDVI at the same coordinates. Through statistical correlation analysis, data validation will demonstrate the similarity of image processing data to actual conditions. The higher the correlation coefficient, the more comparable the two compared outcomes are.

Multiple linear regression

The main motivation for developing the linear model in this research is to obtain a tool that can be used to accurately predict the proportion of an elephant herd or elephant area density (EAD) using two predictor variables (independent variables) in the form of NDVI data and the distribution of elephant movement variables according to time. While the NDVI variable reflects the environmental and ecological conditions of land use at the observation point in 2020, it shows the movement of elephants according to time (t), where (t) is January to December. To make a comparison between one month and another in influencing EAD, one month must be used as a reference. This means that the predictor variable by month that affects EAD needs to be treated as a dummy variable (Artaya 2019). In this study, December is used as a reference, meaning that in this modeling analysis, December must be given a score of 0. Meanwhile, other months are always compared to December in influencing EAD at every point. Mathematically, this description can be expressed by the postulates of the linear model as follows:

$$[EAD]_{it} = \beta_0 + \beta_1[NDVI]_{it} + \beta_2[JAN]_{it} + \beta_3 [FEB]_{it} + \beta_4[MAR]_{it} + \beta_5[APR]_{it} + \beta_6[MAY]_{it} + \beta_7[JUN]_{it} + \beta_8[JUL]_{it} + \beta_9[AUG]_{it} + \beta_{10}[SEP]_{it} + \beta_{11}[OCT]_{it} + \beta_{12}[NOV]_{it} + \xi_{it}$$

In the model, β_0 to β_{12} are the parameters found using the least-squares deviation method (OLS = Ordinary Least Square) using software, namely Minitab 16. This is an error or deviation originating from various variables that are not considered in the prediction model or errors in the measurement of both EAD and NDVI data. In this regard, the hypothesis tested is:

Hypothesis:

H0: $\beta_1 = \beta_2 = \beta_3 = \beta_4 = \beta_5 = \beta_6 = \beta_7 = \beta_8 = \beta_9 = \beta_{10} = \beta_{11} = 0$

H1: $\beta_1 \neq \beta_2 \neq \beta_3 \neq \beta_4 \neq \beta_5 \neq \beta_6 \neq \beta_7 \neq \beta_8 \neq \beta_9 \neq \beta_{10} \neq \beta_{11} \neq 0$

Where, H0 means that the EAD Value is not significantly different according to NDVI or according to the month of observation. And then, H1 means that the EAD Value is significantly different according to NDVI or the month of observation.

As for variables, symbols in models, units, data sources of variable response, and predictors are presented in Table 2.

Hypothesis testing

Hypothesis testing is a subset of inferential statistics that aims to draw conclusions using data from a population sample. Simultaneously, the F method was conducted to test the independent variable effect on the dependent variable. To determine if the independent variable had a partial effect on the dependent variable, the t-test was performed. The significance level used in this study was 10%. The coefficient of determination test is intended to determine the percentage or proportion of total variance explained by the dependent variable by the independent variable. Because the equation is multiple regression, the value used is R². Testing the multiple linear regression equation parameters was carried out using Minitab 16.

Table 1. Classification of vegetation density on NDVI value

Classification of vegetation density	NDVI value
Cloud and Water	-2.0-0
Non-Vegetation	0-0.21
Sparse	0.21-0.42
Moderate	0.42-0.63
Dense	0.63-0.85

Data Source: Sunaryo and Iqmi (2015); Hardianto et al. (2021)

Table 2. Variables, symbols in models, units and scores, data sources

Variable	Symbol	Data source
Elephant movement (Elephant Area Density)	[EAD] _{it}	World Wide Fund for Nature
NDVI	[NDVI] _{it}	Interpretation of Landsat Imagery
January	[JAN] _{it}	Research results in 2022
February	[FEB] _{it}	Research results in 2022
March	[MAR] _{it}	Research results in 2022
April	[APR] _{it}	Research results in 2022
May	[MAY] _{it}	Research results in 2022
June	[JUN] _{it}	Research results in 2022
July	[JUL] _{it}	Research results in 2022
August	[AUG] _{it}	Research results in 2022
September	[SEP] _{it}	Research results in 2022
October	[OKT] _{it}	Research results in 2022
November	[NOV] _{it}	Research results in 2022

RESULTS AND DISCUSSIONS

Movement patterns of Sumatran elephants

Wild animal behavior is the movement of animals to fulfill the stimuli in their bodies by utilizing the stimuli obtained from their environment. Movement behavior is carried out to find food, breed, or escape from predators and other disturbances. The behavior of wild animals is determined by two elements: primary and secondary factors (Gill et al. 2019). Primary factors encourage animals to move to meet their physiological demands, such as hunger, thirst, and sexual desire. As a result, it is believed that the distribution of feed, water, and mating is the primary predictor of a place's utilization. Additionally, secondary factors affect how space is used and may include variances in a place's microclimate, terrain conditions, risk of encountering predators, and risk of disease. Animals prioritize time and space, as these factors limit and scale animal behavior patterns (Torres et al. 2017).

In Tanggamus District, 24 Sumatran elephants (*Elephas maximus sumatranus*) were observed, divided into two groups: the Bunga group (16 elephants) and the Citra group (8 elephants). This is related to Williams et al. (2019), which state that elephants live with *matriarchal* patterns or large and complex social groups in the wild. Elephants move in a wide-roaming area, thus, they utilize a variety of habitat types, including swamp forest, peat swamp forest, lowland forest, and low montane rainforest (Sukumar 1999).

Elephants, as herbivores, require a greater amount of food than other herbivores due to their big body size. Elephant feed is made from plants, including leaves, fruit, stems, bark, tubers, gums, and roots of various plants. According to Riba'i et al. (2013), in general elephants consume five different plant parts: leaves, stems, fruit, roots, and skin. The quantity and diversity of food sources, which vary considerably between areas, are the primary reasons the wild elephant group in the KPH Kotaagung Utara has such a large roaming area. The findings of this study indicate that the Sumatran Elephant's movement in the KPH Kotaagung Utara from 2020 tends to be steady. Between January and July, elephants go to the North, whereas between August and December, elephants tend to travel to the Southern area of the KPH Kotaagung Utara (Figure 2).

Coordinate points distribution based on land cover

The movement of each group is influenced by the season and the condition of its habitat resources, especially feed and the available home range. During the dry season, groups of elephants typically wander from highland to lowland forest in search of feed, but during the rainy season, elephants do the opposite. Basically, elephants prefer areas with cold weather and will stop feeding activities when the weather is hot. During the dry season, groups of elephants typically wander from highland to lowland forest in search of food, but during the rainy season, elephants do the opposite. In general, elephants prefer chilly climates and will cease feeding activities in hot weather. This is aligned with Williams's (2009)

statement that elephants are warm-blooded animals (thermal cover), which means that in hot situations, the elephant will seek shelter and adjust its body temperature to adapt to its environment. Elephants also avoid sunburn by covering their body in soil or mud to keep cool (Imtiyaz et al. 2021). Elephant populations will thrive in habitats with a balanced blend of physical and biotic environments. Mineral salts, water availability, and slopes contribute to these physical circumstances. While biological variables in the form of vegetation composition or structure, vegetation profile, and feed availability (Shaffer et al. 2019). Elephants equipped with GPS collars migrate or travel in a variety of land use and cover types. In KPH Kotaagung Utara, there are 119 elephant movement coordinates distributed throughout 9 different land uses: Primary Dryland Forest, Secondary Dryland Forest, Settlement, Mining, Dryland Agriculture, Shrub Mixed Dryland Agriculture, Rice Fields, Scrub Bush, and Open Land are the different forms of land use. Elephant activity is more intense in the Mixed Shrub Dryland Agriculture area than in other places, with 107 points accounting for approximately 90% of all areas in KPH Kotaagung Utara. Meanwhile, the Dryland and Scrub Agriculture sectors

each have six points or approximately 5% of all areas. This is demonstrated in Table 3 and Figure 3.

Table 3. Distribution of the coordinates of Sumatran elephants fitted with GPS Collars on various types of land use in KPH Kotaagung Utara, Tanggamus District, Lampung Province, Indonesia

Land use	Number of points	Percentage	Wide (ha)
Primary Dryland Forest	0	0	206.3090
Secondary Dryland Forest	0	0	5,641.1765
Settlement	0	0	125.9465
Mining	0	0	96.2618
Dryland Farming	6	5%	2,002.1857
Mixed Shrub Dryland Agriculture	107	90%	45,772.5237
Rice Field	0	0%	851.3812
Shrubs	6	5%	1,485.7736
Open Land	0	0%	76.2225
Grand Total	119	100%	56,257.7808




Figure 2. Map of Sumatran elephant movement in KPH Kotaagung Utara, Tanggamus District, Lampung Province, Indonesia for 2020




Figure 3. Map of Sumatran elephant movement in KPH Kotaagung Utara Tanggamus District, Lampung Province, Indonesia based on land use 2020

Based on the elephant's land uses and roaming, Mixed Shrub Dryland Agriculture receives the most points, 107 out of 119. Due to the predominance of Mixed Shrub Dryland Agriculture usage by elephants in KPH Kotaagung Utara, such area is look like very ideal for Sumatran elephants who live around. The reasons are clear, why they really like those Mixed Shrub Dryland Agriculture. That are because the elephants still found plenty of food, and the ecosystem has a reasonably dense canopy cover, which protects elephants from direct sunburn. The second type of land use that elephant crossed the most is Dryland and Shrub Farm, which received six points out of 119 points. This is reinforced by Putri's research (2021) which states that the vegetation in the northern area is dominated by woody plant species such as harendong trees (*Melastoma affine*), masi-masian (*Mimusops elengi*), bendo (*Artocarpus elasticus*), and cocoa (*Theobroma cacao*), while the vegetation that dominates the southern area is non-woody plant species, such as reed grass (*Saccharum spontaneum*), tepus (*Phrynium pubinerve*), resam fern (*Dicranopteris linearis*), jerumo manok (*Leea indica*), Bandotan (*Ageratum conyzoides*), Kasapan (*Clidemia hirta*), reeds (*Imperata cylindrica*), bone grass (*Eleusine indica*), teki grass (*Cyperus rotundus*), and ganyongan (*Canna discolor*). Additionally, gamal tree (*Gliricidia sepium*) with an IVI of 111, and dadap (*Erythrina variegata*) with an IVI of 105 dominated the plant species of dryland agricultural ecosystem type in KPH Kotaagung Utara during the tree phase. The vegetation in an area has a positive influence on the elephant's habitat, which varies depending on the

structure and composition of vegetation that grows in the area (Farhan et al. 2019). Additionally, there was no evidence of Sumatran elephant points in six other types of land cover in the KPH Kotaagung Utara, namely: Settlement, Mining, Primary Dryland Forest, Secondary Dryland Forest, Rice Fields, and Open Land. This is because the land area is so small. Each NDVI's value and area can be viewed in Table 4.

Types of land use in the classification of vegetation density in the form of Non-Vegetation and Sparse, the Sumatran elephant is barely traversed. Apart from having a smaller land area compared to others, elephants are also one of the animals with a high level of sensitivity, making them minimize their activity on human territory. According to Abdullah et al. (2019), elephants can detect sounds in the frequency range from 1-20,000 Hz, with a hearing distance of roughly 10 km. Therefore, increased human activities around elephant habitats, such as mining, agriculture, or urbanization, will generate noise effects that decrease the quality and quantity of elephant habitats, which can impact population decline.

NDVI value based on elephant point distribution

Based on the data analysis, the results of the classification of elephant distribution points with NDVI variations in the KPH Kotaagung Utara for 2020 were classified into Cloud and Water, Non-Vegetation, Sparse Moderate, and Dense, as presented in Table 4. The NDVI value describes the vegetation cover above the ground surface with different brightness values based on the

difference in digital number values through a calculation of visible light and near-infrared. The classification can be seen in Figure 4.

According to Figure 4, the vegetation condition in KPH Kotaagung Utara is classified as dense. This is because KPH Kotaagung Utara has a relatively high vegetation density, with an average NDVI of 0.63-0.85 and 107 elephant distribution spots out of a total of 119. This is aligned with studies (Huang et al. 2020) indicating that the theoretical range of the NDVI value is -1 to 1 (the closer to 1, the more vegetation), with values less than 0 representing sea (water) or unoccupied land. The amount of land cover categories also influences the density classification, specifically Secondary Dryland Forest, Primary Dryland Forest, and Mixed Shrub Dryland Agriculture, which dominate in the KPH Kotaagung Utara, covering 51,620.0092 Ha of the total 56,020 Ha. Furthermore, Moderate's NDVI classification which has 12 elephant distribution points. Meanwhile, the Sparse, Cloud and Water, and Non-Vegetation classifications do not have elephant distribution points. A low NDVI value indicates that the vegetation is experiencing moisture stress, and a higher value indicates a higher density of green vegetation. Furthermore, according to Sugiyanto et al. (2017), elephants will choose fresh grass in the rainy season because it contains carbohydrates and low fiber content (*Lignohemicellulose*). Meanwhile, elephants prefer leaves containing high protein content in the dry season, like (8-10% *Malvaceae* and 10 to 20% Leguminosae). A low NDVI value indicates that the vegetation is under water stress, whereas a high value implies a dense canopy of greenery.

Multiple linear regression T-Test against dependent variables

The T-test, also known as the partial test, was employed to determine the effect of each independent variable of vegetation intensity (NDVI) on the dependent variable of elephant migration in this study (elephant area density). The T-test results on multiple linear regression can be seen in Table 5.

Table 5. Monthly distribution results and Vegetation Intensity (NDVI) on elephant movement (elephant area density) in KPH Kotaagung Utara, Tanggamus District, Lampung Province, Indonesia

Predictor	Coef	SE coef	T-value	P-value
Constant	1.688	0.118	14.27	0.000
[NDVI] _{it}	-2.220	0.282	-7.87	0.000
[JAN] _{it}	0.976	0.121	8.09	0.000
[FEB] _{it}	0.761	0.108	7.07	0.000
[MAR] _{it}	0.802	0.099	8.07	0.000
[APR] _{it}	0.759	0.106	7.15	0.000
[MAY] _{it}	0.755	0.118	6.42	0.000
[JUN] _{it}	0.736	0.093	7.88	0.000
[JUL] _{it}	0.423	0.074	5.56	0.000
[AUG] _{it}	0.335	0.065	5.15	0.000
[SEP] _{it}	0.014	0.060	2.33	0.022
[OCT] _{it}	0.215	0.731	2.94	0.004
[NOV] _{it}	0.201	0.072	2.76	0.007

S = 0.0726422; R-Sq = 60.29%; R-Sq(adj) = 55.80%

Note: S is the Standard Error of Estimate (SEE); R-Sq is the R-Square (Coefficient of Determination); R-Sq (Adj) is the Adjusted R-Square (Coefficient of Determination)

Table 4. Value and area of each NDVI classification in KPH Kotaagung Utara, Tanggamus District, Lampung Province, Indonesia for 2020

Classification of vegetation density	NDVI value	Type of land use	Total land use (ha)
Cloud and Water	-2.0-0	-	-
Non-Vegetation	0-0.21	Open Land, Settlement, and Mining	298.4309
Sparse	0.21-0.42	Rice Field	851.3812
Moderate	0.42-0.63	Dryland Farming, and Shrubs	3,487.9593
Dense	0.63-0.85	Secondary Dryland Forest, Primary Dryland Forest, and Mixed Shrub Dryland Agriculture	51,620.0092




Figure 4. NDVI classification on elephant point distribution in January-December 2020

8 The relationship between the dependent variable (EAD) and the independent variable (NDVI and the monthly distribution) in the regression:

$$\begin{aligned} \text{EAD} = & 1.688 - 2.220 \text{NDVI} + 0.976 [\text{JAN}]_{it} + \\ & 0.761 [\text{FEB}]_{it} + 0.802 [\text{MAR}]_{it} + 0.759 [\text{APR}]_{it} + \\ & 0.755 [\text{MAY}]_{it} + 0.736 [\text{JUN}]_{it} + 0.423 [\text{JUL}]_{it} + \\ & 0.335 [\text{AUG}]_{it} + 0.141 [\text{SEP}]_{it} + 0.215 [\text{OKT}]_{it} + \\ & 0.201 [\text{NOV}]_{it} \end{aligned}$$

The R-square value, or the resulting determinant coefficient, is 60.29% based on the data processing in Table 3. This value indicates that the diversity of about 60.29% of the vegetation intensity in the KPH Kotaagung Utara affects the presence of elephants. However, the other 39.71% is influenced by other aspects, like salticks. The thing that elephants do to keep their metabolism balanced is by salticks (Sitienei et al. 2012). According to Sach et al. (2019), mineral salts such as calcium, magnesium, and potassium contained in the soil or cliffs taken by the elephant's trunk are deemed helpful for elephants to help strengthen their bones, teeth, and tusks. Other factors that may influence elephant movement, such as the need for safe locations to carry out daily activities: rest, communicating, and dressing up, are elephant behaviors to defend themselves from threats (Setiawan et al. 2021). Additionally, some variables substantially affect elephant mobility (elephant area density) in KPH Kotaagung Utara, specifically monthly seasonal effects in January, February, April, May, June, July, August, October, and November. Sugiyanto et al. (2017) also revealed that elephants have foraging strategies when consuming grass and leave in the dry and wet seasons. Elephants will choose fresh grass in the wet season because it contains carbohydrates and low fiber content (*Lignoheemicellulose*). Meanwhile, elephants prefer leaves containing high protein content in the dry season, like (8-10% *Malvacea* and 10 to 20% *Leguminosae*) (Singh 2019). 3 The T-test and coefficient of the determination indicate that NDVI has a volume of 0.000, which corresponds to a 99% confidence interval. The NDVI coefficient value is -2.220, which means that if the value of vegetation intensity (NDVI) decreases by 1%, it will reduce the movement of elephants by 2.200 from the total movement of elephants in 56,257 hectares of the KPH Kotaagung Utara area. It demonstrates that the high NDVI represents land use and cover in the KPH Kotaagung Utara, which will affect Sumatran elephant distribution. This is consistent with the findings of Jim and Campos-Arceiz (2022), which indicated that the closed-canopy primary rainforest is a marginal habitat for elephants and humans alike and if the forest is harmed, the elephant's habitat will be disrupted. Essentially, grass, shrubs, or denser canopy biophysical characteristics enable satellite sensors to detect their reflectance as cover (Stagakis et al. 2010). However, according to (Hidayati et al. 2018), the vegetation index does not always accurately reflect the field conditions at the NDVI value.

According to this study, the intensity of dense vegetation is directly proportional to the distribution of Sumatran elephant points in KPH Kotaagung Utara. 2 The

intensity of elephant movement in the Mixed Shrub Dryland Agriculture area is higher than in the other areas, namely 107 points, or about 90% of all areas in KPH Kotaagung Utara. Meanwhile, in the dry and shrub agriculture areas, there are 6 points each, or about 5% of all the areas. Therefore, the vegetation classification in KPH Kotaagung Utara is categorized as *dense* with an NDVI value range of 0.63-0.85. Furthermore, the regression results prove that NDVI and the monthly season will affect the movement of elephants with a p-value of <0.001. Thus, the government, non-governmental organizations, and the community can accurately monitor the state of vegetation and elephant movement, thereby minimizing elephant-human conflicts resulting from land-use changes. Suppose it was determined that the status of the Sumatran elephant's habitat is deteriorating due to encroachment. In that case, it is hoped that specific parties can collaborate to restore the elephant's natural habitat to its original state.

ACKNOWLEDGEMENTS

The author would like to thank the KPH Kotaagung Utara, Lampung, Indonesia and the World Wide Fund for Nature Indonesia Southern Sumatra Program for allowing and assisting in data collection and all who have been involved in the preparation of this study. This study was funded by the World Wide Fund for Nature Indonesia Southern Sumatra Program.

REFERENCES

- Abdullah, Shabrina A, Khairil, Saputri M, Syafranti D, Ala N. 2019. The effect of anthropogenic noise on Sumatran elephant's anti-predator behavior in the Elephant Conservation Center. *J Phys Conf Ser* 1460 (1): 012067. DOI: 10.1088/1742-6596/1460/1/012067.
- Ahmed KR, Akter S. 2017. Analysis of landcover change in Southwest Bengal delta due to floods by NDVI, NDWI and K-means cluster with Landsat multi-spectral surface reflectance satellite data. *Remote Sens Appl Soc Environ* 8 (1): 168-181. DOI: 10.1016/j.rsase.2017.08.010.
- Ajayi, OV. 2017. Primary Sources of Data and Secondary Sources of Data. [Dissertation]. Benue State University, Makurdi. [Nigeria]. DOI: 10.13140/RG.2.2.24292.68481.
- Artaya IP. 2019. Regresi Linier Berganda Metode Dummy. *Qualitative Research Analysis Method*. DOI: 10.13140/RG.2.2.30936.75526. [Indonesian]
- Calabrese A, Calabrese JM, Songer M, Wegmann M, Hedges S, Rose R, Leimgruber P. 2017. Conservation status of Asian elephants: The influence of habitat and governance. *Biodivers Conserv* 26 (1): 2067-2081. DOI: 10.1007/s10531-017-1345-5.
- Chen S, Woodcock CE, Bullock ELAP, Service USF, Mountain R. 2021. Remote sensing of environment monitoring temperate forest degradation on google earth engine using landsat time series analysis. *Remote Sens Environ* 265: 1-10. DOI: 10.1016/j.rse.2021.112648.
- Directorate of Biodiversity Conservation, Directorate General of Natural Resources and Ecosystem Conservation, Ministry of Environment and Forestry of the Republic of Indonesia Jakarta. 2020. Sumatran elephants. www.mongabay.co.id/2020/08/12/rusaknya-habitat-ancaman-utama-kehidupan-gajah-sumatera/. [Indonesian]
- Farhan MR, Adawiyah RMK, Aisyah N, Nasrullah M, Triastuti A, Lestari S, Hasriati. 2019. Analisis Vegetasi Tumbuhan di Resort Pattunang-Karaenta Taman Nasional Batimurung Bulusaraung. Universitas Negeri Makassar, Makassar. [Indonesian]
- Gill M, Reveron AT, Ramirez AC, Maldonado O, Vandenberg JL, Erasquin GA. 2019. Influence of biological sex on social behavior,

- individual recognition, and non-associative learning in the adult gray short-tailed opossum (*Monodelphis domestica*). *Physiol Behav* 1 (21): 1-10. DOI: 10.1016/J.Physbeh.2019.112659.
- Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. 2017. Remote sensing of environment google earth engine: Planetary-scale geospatial analysis for everyone. *Remote Sens Environ* 202: 18-27. DOI: 10.1016/j.rse.2017.06.031.
- Hardianto A, Dewi PU, Feriansyah T, Sari NFS, Rifiana NS. 2021. Pemanfaatan citra lansat 8 dalam mengidentifikasi nilai indeks kerapatan vegetasi (NDVI) tahun 2013 dan 2019 (area studi: Kota Bandar Lampung). *J Geosains dan Remote Sensing (JGRS)* 2 (1): 8-15. DOI: 10.23960/jgrs.2021.v2i1.38. [Indonesian]
- Hidayati IN, Suharyadi R, Danoedoro P. 2018. Developing an extraction method of urban built-up area based on remote sensing imagery transformation index. *Forum Geografi* 32 (1): 1-10. DOI: 10.23917/forgeo.v32i1.5907.
- Huang S, Tang H, Hupy JP, Wang Y, Shao G. 2020. A commentary review on the use of normalized difference vegetation index (NDVI) in the era of popular remote sensing. *J For Res* 1 (32): 1-6. DOI: DOI: 10.1007/s11676-020-01155-1.
- Imtiyaz CD, Syarifuddin A, Triwanto J, Al-Hakim RR. 2021. Analysis of welfare levels, ecology, and animal management in Seblat Elephant Training Center, Bengkulu-Indonesia. *J Biotechnol Nat Sci* 1 (2): 64-74. [Indonesian].
- Kumar MA, Vijaykrishnan S, Singh M. 2020. Whose habitat is it anyway? Role of natural and anthropogenic habitats in conservation of charismatic species. *Trop Conserv Sci* 11 (1): 1-10. DOI: 10.1177/1940082918788451.
- Lim T, Campos-Arceiz A. 2022. A review of human-elephant ecological relations in the Malay Peninsula: adaptations for coexistence. *MDPI* 14 (1): 36-46. DOI: 10.3390/d14010036.
- Meytasari P, Bakri S, Herwanti S. 2014. Penyusunan kriteria domestikasi dan evaluasi praktek penguasaan gajah: Studi di Taman Nasional Way Kambas Kabupaten Lampung Timur. *J Sylva Lestari* 2 (2): 78-88. DOI: 10.23960/jsl2279-88. [Indonesian]
- Moßbrucker AM, Imron MA, Pudyatmoko S, Pratje PH, Sumardi. 2016. Modeling the fate of sumatran elephants in Bukit Tigapuluh, Indonesia: Research needs and implications for population management. *J Ilmu Kehutanan* 10 (1): 5-18. DOI: 10.22146/jik.12622.
- Nedd R, Light K, Owens M, James N, Johnson E, Anandhi A. 2021. A synthesis of land use/land cover studies: Definitions, classification systems, meta-studies, challenges and knowledge gaps on a global landscape. *MDPI* 10 (9): 984-994. DOI: 10.3390/land10090994.
- Pahlefi MR, Danoedoro P, Kamal M. 2021. The utilisation of sentinel-2A images and google earth engine for monitoring tropical Savannah grassland. *Geocarto Intl* 2: 1-15. DOI: 10.1080/10106049.2021.1914749.
- Prasetyo NN, Sasmito B, Prasetyo Y. 2017. Analisis perubahan kerapatan hutan menggunakan metode NDVI dan EVI pada citra satelit lansat 8 tahun 2013 dan 2016. *J Geodesi UNDIP* 6 (3): 21-27. [Indonesian]
- Putri AP. 2021. Analisis Ketersediaan jadi Jenis Tumbuhan Pakan dan Tempat Berlingdung Gajah Sumatera (*Elephas maximus sumatranus*) Pada Daerah Jelajah Gajah di KPHL Kotaagung Utara. [Undergraduate Thesis]. Universitas Lampung, Lampung. [Indonesian]
- Regulation of The Minister of Environment and Forestry of The Republic of Indonesia. 2018. Protected Types of Plants and Animals in Indonesia. P.20/MenLHK/Setjen/Kum.1/6/2018 and P.106/MenLHK/Setjen/Kum.1/12/2018. [Indonesian]
- Riba'i, Setiawan A, Darmawan A. 2013. Perilaku makan gajah Sumatera (*Elephas maximus sumatranus*) di Pusat Konservasi Gajah Taman Nasional Way Kambas. *Media Konservasi* 18 (2): 89-95. [Indonesian]
- Sach F, Dierenfeld ES, Evans SCL, Watts MJ, Yon L. 2019. African savanna elephants (*Loxodonta africana*) as an example of a herbivore making movement choices based on nutritional needs. *PeerJ* 7: 1-27. DOI: 10.7717/peerj.6260.
- Setiawan A, Monik DT, Charles Y, Rustiati EL, Ruciyansah Y, Anwar Z. 2021. An assessment of Sumatran elephant presences (*Elephas maximus sumatranus*) in Kotaagung Utara (Lampung, Indonesia) as a potential tourism attraction. *Biodiversitas* 22 (12): 5397-5407. DOI: 10.13057/biodiv/d221222.
- Shaffer LJ, Khadka KK, Hoek JVD, Naithani KJ. 2019. Human-elephant conflict: A review of current management strategies and future directions. *Front Ecol Evol* 1: 1-10. DOI: 10.3389/fevo.2018.00235.
- Shafizadeh-Moghadam H, Khazaei M, Alavipanah SK, Weng Q. 2021. Google earth engine for large-scale land use and land cover mapping: An object-based classification approach using spectral, textural and topographical factors. *GISci Remote Sens* 58 (6): 914-928. DOI: 10.1080/15481603.2021.1947623.
- Sidhu N, Pebesma E, Câmara G. 2018. Using google earth engine to detect land cover change: Singapore as a use case. *Eur J Remote Sens* 51 (1): 486-500. DOI: 10.1080/22797254.2018.1451782.
- Singh R. 2019. Potential Roles of Tree Leaves in Ruminant Nutrition. Pashudhan Praharee, India.
- Sitenei AJ, Jiwen G, Ngene SM, Paix MJDL, Waweru FK. 2012. Analysis in the concentration, determination and comparison of some mineral elements in the natural salt-licks utilized by elephants: Mt. Elgon National Park Case Study. *Open J Adv Mater Res* 2: 356-360. 1796-1800. DOI: 10.4028/www.scientific.net/AMR.356-360.1796.
- Stabach JA, Cunningham SA, Connette G, Mota JL, Reed D, Byron M, Songer M, Wachter T, Mertes K, Brown JL, Comizzoli P, Newby J, Monfort S, Leimgruber P. 2020. Short-term effects of GPS collars on the activity, behavior, and adrenal response of scimitar-horned oryx (*Oryx dammah*). *PLoS One* 15 (2): 1-22. DOI: 10.1371/journal.pone.0221843.
- Stagakis S, Markos N, Sykioti O, Kyparissis A. 2010. Monitoring canopy biophysical and biochemical parameters in ecosystem scale using satellite hyperspectral imagery: An application on a *Phlomis fruticosa* mediterranean ecosystem using multiangular CHRIS/PROBA observations. *Remote Sens Environ* 1 (114): 977-994. DOI: 10.1016/j.rse.2009.12.006.
- Sugiyanto EEL, Erianto, Prayogo H. 2017. Ketersediaan pakan gajah Sumatera [*Elephas maximus sumatranus* (Temminck, 1847)] di Resort Air Hitam Taman Nasional Tesso Nilo Kabupaten Pelalawan Provinsi Riau. *J Hutan Lestari* 5 (1): 147-155. DOI: 10.26418/jhl.v5i1.18836. [Indonesian]
- Sukumar R. 1999. The Asian Elephant: Ecology and Management. Cambridge University Press, Cambridge, UK.
- Sunaryo DK, Iqmi MZ. 2015. Pemanfaatan citra landsat 8 untuk pendeteksian dan mengetahui hubungan kerapatan vegetasi terhadap suhu permukaan. *J Spectra* 13 (25): 55-72. [Indonesian]
- Torres LG, Orben RA, Tolkova I, Thompson DR. 2017. Classification of animal movement behavior through residence in space and time. *PLoS One* 12 (1): 1-18. DOI: 10.1371/journal.pone.0168513.
- Wall J, Wittemyer G, Klinkenberg B, LeMay V, Blake S, Strindberg S, Henley M, Vollrath F, Maisels F, Ferwerda J, Hamilton ID. 2021. Human footprint and protected areas shape elephant range across Africa. *Curr Biol* 31 (11): 2437-2445. DOI: 10.1016/j.cub.2021.03.042.
- Williams E, Carter A, Hall C, Harrison SB. 2019. Social Interactions in Zoo-Housed Elephants: Factors affecting social relationships. *Animals (Basel)* 9 (10): 747. DOI: 10.3390/ani9100747.
- Williams TM. 2009. Heat transfer in elephants: Thermal partitioning based on skin temperature profiles. *J Zool* 222 (2): 235-245. DOI: 10.1111/j.1469-7998.1990.tb05674.x.
- Wong EP, Yon L, Walker SL, Mena AS, Wade J, Othman N, Saaban S, Arceiz AC. 2018. The elephant who finally crossed the road-Significant life events reflected in faecal hormone metabolites of a wild asian elephant. *Gajah* 48 (1): 4-11.
- Wóźniak E, Gabryszewski R, Dziob D. 2020. Remote sensing and electromagnetic wave behaviour to measure vegetation phenology with physics. *Phys Educ* 55 (4): 1-13. DOI: 10.1088/1361-6552/ab80ff.
- Xia Z, Guo X, Chen R. 2020. Automatic extraction of aquaculture ponds based on Google Earth Engine. *Ocean Coast Manag* 198: 105348. DOI: 10.1016/j.ocecoaman.2020.105348.
- Xue J, Su B. 2017. Significant remote sensing vegetation indices: A review of developments and applications. *J Sensors* 1 (1): 1-17. DOI: 10.1155/2017/1353691.

● 20% Overall Similarity

Top sources found in the following databases:

- 19% Internet database
- 4% Publications database
- Crossref database
- Crossref Posted Content database
- 8% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	doaj.org Internet	5%
2	researchgate.net Internet	5%
3	nveo.org Internet	3%
4	Universitas Jenderal Soedirman on 2022-04-14 Submitted works	1%
5	biodiversitas.mipa.uns.ac.id Internet	<1%
6	tandfonline.com Internet	<1%
7	ccsds.cosmos.ru Internet	<1%
8	hvtc.edu.vn Internet	<1%

9	repository.uia.ac.id	Internet	<1%
10	Institut Pertanian Bogor on 2022-01-13	Submitted works	<1%
11	Fajar Yulianto, Gatot Nugroho, Galdita Aruba Chulafak, Suwarsono Suw...	Crossref	<1%
12	A P Suka, Surati, D R Kurniasari, A Justianto. "Sources of income from ...	Crossref	<1%
13	Monash University on 2022-11-03	Submitted works	<1%
14	ijrrjournal.com	Internet	<1%
15	University of Edinburgh on 2022-11-03	Submitted works	<1%
16	pdfcoffee.com	Internet	<1%
17	jthornhillphotography.com	Internet	<1%
18	mdpi.com	Internet	<1%
19	A Zaitunah, Samsuri, N Hidayat. "Vegetation distribution analysis of gr...	Crossref	<1%
20	University of Queensland on 2018-08-20	Submitted works	<1%

- 21 **gssrr.org** <1%
Internet
-
- 22 **Yashar Jamei, Mehdi Seyedmahmoudian, Elmira Jamei, Ben Horan, Sa...** <1%
Crossref
-
- 23 **Abdullah, A Shabrina, Khairil, M Saputri, D Syafrianti, N Ala. "The effect...** <1%
Crossref
-
- 24 **Universidad Nacional de Colombia on 2021-09-24** <1%
Submitted works
-
- 25 **University of Leeds on 2022-08-29** <1%
Submitted works
-
- 26 **University of Melbourne on 2022-10-25** <1%
Submitted works

● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Quoted material
- Manually excluded sources

EXCLUDED SOURCES

smujo.id **66%**
Internet

smujo.id **10%**
Internet