

PAPER NAME

Joko Prasetyo_Ginting_Dkk_2021_Biodiversitas_Scopus.pdf

AUTHOR

Cipta Ginting

WORD COUNT

4982 Words

CHARACTER COUNT

27208 Characters

PAGE COUNT

6 Pages

FILE SIZE

722.6KB

SUBMISSION DATE

Feb 7, 2023 10:33 AM GMT+7

REPORT DATE

Feb 7, 2023 10:34 AM GMT+7

● 14% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 7% Internet database
- Crossref database
- 9% Submitted Works database
- 11% Publications database
- Crossref Posted Content database

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded sources
- Quoted material
- Manually excluded text blocks

The effect of biological agent and botanical fungicides on maize downy mildew

JOKO PRASETYO^{1,*}, CIPTA GINTING¹, HASRIADI MAT AKIN¹, RADIX SUHARJO¹, AININ NISWATI²,
AULIANA AFANDI¹, REZA ADIWIJAYA¹, SUDIONO¹, MUHAMMAD NURDIN¹

¹Department of Plant Protection, Faculty of Agriculture, Universitas Lampung. Jl. Prof. Soemantri Brojonegoro No. 1, Bandar Lampung 35141, Lampung, Indonesia. Tel.: +62-721-787029, *email: joko.prasetyo@fp.unila.ac.id, radix.suharjo@fp.unila.ac.id

²Department of Soil Science, Faculty of Agriculture, Universitas Lampung. Jl. Prof. Sumantri Brojonegoro No. 1, Bandar Lampung 35141, Lampung, Indonesia

Manuscript received: 4 February 2021. Revision accepted: 7 April 2021.

Abstract. Prasetyo J, Ginting C, Akin HM, Suharjo R, Niswati A, Afandi A, Adiwijaya R, Sudiono, Nurdin M. 2021. The effect of biological agent and botanical fungicides on maize downy mildew. *Biodiversitas* 22: 1652-1657. This study was conducted to evaluate the effect of the combination of *Trichoderma asperellum* and mycorrhizae with botanical fungicides made from betel leaf extract or turmeric extract against maize downy mildew. The study used a randomized complete block (RCB) design with three replications. The treatments applied were: single applications of *T. asperellum* (Ta); arbuscular mycorrhizal fungi (AMF); turmeric extract (T); betel extract (B); the combination of *T. asperellum* and AMF (TaM); turmeric extract and *T. asperellum* (TTa); turmeric extract and AMF (TM); turmeric extract + *T. asperellum* + AMF (TTaM); betel extract + *T. asperellum* (BTa); betel extract + AMF (BM); betel + *T. asperellum* + AMF (BTaM); and Control (C). The observed variables were disease incidence, incubation period, and shoot dry weight. The data were analyzed using ANOVA. Differences between treatments were tested by the LSD test at 5% significance level. The results showed that the combination of botanical fungicides with biological agents could reduce the incidence of downy mildew, extend the incubation period, and increase the dry weight of corn shoots. The best combination treatment was achieved from the combination of *T. asperellum* with betel extract. There was a synergistic effect between botanical fungicides and biological agents in reducing the disease incidence.

Keywords: Biological agents, botanical fungicides, downy mildew

INTRODUCTION

Corn is one of the most important carbohydrate sources in the world. Corn has an important role in fulfilling national and international market demand for food after rice and wheat (Tanklevska et al. 2020). The market demand for corn from year to year continues to increase along with the increasing population, improvement of community welfare, the development of livestock businesses, and the food industry's advancement. Corn is widely used by the food, beverage, chemical, and pharmaceutical industries (Ranum et al. 2014).

Based on its chemical composition and nutrient content, corn has prospects as a food and industrial raw material (Jabran et al. 2007; Ranum et al. 2014; Shah et al. 2016; Kazerooni et al. 2019). The use of corn as an industrial raw material provides added value for corn. Corn is used as a raw material in the feed and food industry and staple food in several regions in Indonesia (Bantacut et al. 2015). Corn kernels can be processed into corn flour, corn rice, and snacks. Corn can also be processed into cooking oil, margarine, and formula foods. Domestic production of corn in Indonesia has not been able to keep up with demand growth up to 2015 (Bantacut et al. 2015).

According to the Indonesia Central Bureau of Statistics (2016), dry shelled corn production in Lampung, Indonesia experienced fluctuations in production from 2010-2015,

i.e., 2,126,571 tons, 1,817,906 tons, 1,760,275 tons, 1,760,278 tons, 1,719,386 tons and 1,502,800 tons, respectively. One of the main factors responsible for decreasing corn production was downy mildew caused by *Peronosclerospora* spp.

Three *Peronosclerospora* species have been reported as the causative agent of downy mildew of corn in Indonesia, namely *P. maydis*, *P. philippinensis*, and *P. sorghi* (Muis et al. 2013; Rustiani et al. 2015; Muis et al. 2016). *P. maydis* (syn. *P. australiensis*) is the causative agent of downy mildew in several corn production areas in Lampung (Suharjo et al. 2020). Production loss in corn infected with this pathogen reaches 80%-100% due to the inability to produce kernels (Soenartiningih and Talanca 2010).

A metalaxyl fungicide is the primary choice of farmers to control downy mildew until now. However, the use of metalaxyl continuously in the long term has triggered resistance to the pathogen (Gisi and Sierotzki 2008; Burhanuddin 2009). Therefore, it is necessary to find another alternatives method to control the disease, such as the application of biological agents, i.e., *Trichoderma* (Puyam et al. 2016; Sood et al. 2020; Ginting et al. 2020) and mycorrhizal fungi (Akhtar and Siddiqui 2008; Tahar et al. 2010; Cameron et al. 2013; Pérez-de-Luque et al. 2017), and botanical fungicides, like turmeric (Ginting 2006; Rahman et al. 2016; Mamarabadi et al. 2018) and betel (Nalina and Rahim 2006; Rahman et al. 2016). The use of

biological agents and botanical fungicides is also safe for consumers and the environment. Biological agents and botanical fungicides are easily degraded and do not leave chemical residues on agricultural products (Yoon et al. 2013; Nega 2014; Bardin et al. 2015; Zaker 2016). This study was carried out to evaluate the effect of biological agents and botanical fungicides and their combination on downy mildew of corn. Botanical fungicides and biological agents were expected to have a synergistic effect in suppressing downy mildew.

MATERIALS AND METHODS

Plant material and planting media preparation

This study used the P27 variety (Dupont Indonesia) of corn seed. The seeds were planted in 36 plastic polybags (35 cm x 40 cm), each polybag containing 10 seeds. Plant maintenance includes watering and weeding.

The planting media was soil taken from around the Integrated Laboratory, Faculty of Agriculture, University of Lampung. Firstly, planting media (soil) was sterilized by steaming for 2 hours. Ten kg of sterilized soil was put into a polybag.

Preparation of biological agents

Two biocontrol agents were used in this study, namely *T. asperellum* and a consortium of three genera of Arbuscular Mycorrhizal Fungi (AMF), i.e. *Entrophospora* sp., *Gigaspora* sp., and *Glomus* sp. formulated in powder, was bought from Estate Crops Laboratory, Faculty of Agriculture, University of Lampung. The spore density of the AMF was 100 spore/g. The isolate of *T. asperellum* was a collection of the Plant Clinics, University of Lampung. *T. asperellum* was cultured on a sterile petri dish containing 10 mL of Potato Dextrose Agar (PDA). Potato Dextrose Agar (PDA) consists of 200g potato, 20g Dextrose, 20 g Agar, and 1000 mL distilled water. Conidia of *T. asperellum* were harvested by flooding plates containing 7 days old culture with 10 mL of sterile distilled water and grabbed gently using Drigalski spatula. The conidial suspension was transferred into Erlenmeyer 100 mL and diluted to the density of $\sim 10^5$ conidia/mL.

Preparation of botanical fungicides

Turmeric rhizome and betel leaves were weighed 500 g each, cleaned with sterile water, cut into small pieces, and air-dried. Subsequently, samples were dried in the oven at 50 °C for 36 hours. After drying, samples were ground using a pestle into a fine powder (600 mesh). Botanical fungicide was made by dissolving 10 g of fine powder into 100 mL of sterile distilled water and filtered using filter paper (Whatman no. 42). The suspension was centrifuged for 10 minutes at 300 rpm. The supernatant was collected for further use (Ayu et al. 2013).

Isolation of *Peronosclerospora* sp.

Peronosclerospora used in this study was collected from symptomatic corn plants taken from a cornfield located in Natar, South Lampung. The collection was

carried out at 03.00 am. Conidia from infected leaves were harvested using a small brush (no. 2). The brush was moistened before being used by dipping it in 20 mL sterile distilled water in a plastic bowl (400 mL of volume, 11 cm of lid diameter). The brush containing the conidia was immersed in 20 mL of distilled water previously used to moisten the brush. Conidia harvesting was continued until the water in the Petri dish was very cloudy due to the high conidia density. The conidial suspension was transferred into a graduated glass containing 500 mL sterile distilled water (Figure 1). Suspension of conidia was brought to the laboratory, and the conidial density was determined.

Peronosclerospora sp. inoculation and botanical fungicide application

The conidia suspension of *Peronosclerospora* ($\sim 10^5$ conidia/mL) was mixed with botanical fungicide suspension (1:9) and left for 1 hour. Inoculation was conducted by dropping 1 mL of the mixture of conidia suspension and botanical fungicide to the shoots of corn at 4-5 am, on the same day as conidia collection. As a control treatment, the plants were inoculated with a suspension of conidia (the same volume and the same conidia density) without applying biological control agents and botanical fungicides. Inoculation was carried out twice at the age of 7 and 10 days after planting.

Application of biological agents

Ten mL of *T. asperellum* conidia suspension ($\sim 10^6$ conidia/mL) was poured into each planting hole. One g of AMF powder was applied to each plant (Manila and Nelson 2014). The spore density of AMF powder was 100 spore/g.



Figure 1. Harvesting method of *Peronosclerospora* sp. conidia

Table 1. The scale of disease severity

Score category	Symptom depiction
0	No symptom
1	<10% of the total area of the leaves were infected
2	10-25% of the total area of the leaves were infected
3	25-50% of the total area of the leaves were infected
4	>50% of the total area of the leaves were infected

Observation and data collection

Observations were done daily for four weeks. The variables observed were disease incidence, disease severity, incubation period, and shoot dry weight. Downy mildew incidence was observed in the morning and was based on chlorosis symptoms in corn leaves. Disease incidence (DI) was calculated by dividing the number of infected plants (n) with the total plants observed (N). The incubation period is the time required for the appearance of disease symptoms calculated from the inoculation of downy mildew pathogen to the formation of symptoms. The disease severity was estimated on the scale based on the symptoms (Table 1.)

The DSI (Disease severity index) was calculated with the following formula: $\frac{\sum(n \times v)}{(N \times V)} \times 100\%$, whereas [n] is the total of infected leaves with a specific score, [v] is the score category of the symptom, [N] is total leaves observed, and [Z] is the highest score used. At the end of the observation, the corn plant was removed from the planting medium and then cleaned from the attached dirt. The shoot was cut into pieces, put in the envelope, and dried in the oven at 80 °C for 5 days to the constant weight.

RESULTS AND DISCUSSION

Two biological control agents (*T. asperellum* and AMF) and two botanical fungicides (turmeric extract and betel extract) were evaluated for their capability to suppress the development of downy mildew disease in corn. These biological control agents and botanical fungicides were applied in individual or combination applications.

The result showed that all the treatments, either single or in combination application were able to suppress disease incidence of corn downy mildew, compared to the control treatment. The symptoms in the control treatment were more severe than in the treatment groups (Figure 2).

Individual application of *T. asperellum* (Ta), Arbuscular Mycorrhizal Fungi (AMF) (M), turmeric (T), or betel (B) significantly reduced the incidence of disease compared to control (C). However, the incubation period and disease severity in these treatments were not significantly different from those in control plants. The treatments of AMF, turmeric, or betel extract produced a significantly higher dry shoot weight than control. Meanwhile, the dry shoot weight of the corn plants in *T. asperellum* treatment was not significantly different from that in control plants.




Figure 2. Disease symptoms on the inoculated plants. A. Control, B. Betel + *T. asperellum* (BTa), C. Betel + *T. asperellum* + AMF (BTaM), D. Turmeric + *T. asperellum* (TTa)




Figure 3. The downy mildew (DM) incidence of maize (%) in various treatments. Ta: *T. asperellum*, M: Arbuscular Mycorrhizal Fungi (AMF), T: turmeric extract, B: betel extract, TaM: *T. asperellum* + AMF, TTa: Turmeric + *T. asperellum*, TM: turmeric + AMF, TTaM: turmeric + *T. asperellum* + AMF, BTa: betel + *T. asperellum*, BM: betel + AMF, BTaM: betel + *T. asperellum* + AMF, C: Control.

Table 3. Average downy mildew incidence, incubation period, and dry shoot weight of corn on treatment combination of *Trichoderma asperellum*, Arbuscular Mycorrhizal Fungi (AMF), and botanical fungicides

Treatment	Downy mildew incidence (%)*	Disease severity index*	The incubation period (days)*	Weight of dry shoot (g)*
Ta	61.11b	66.70a	7.00c	20.73fg
M	66.67b	59.70a	6.00c	41.97bcd
T	66.67b	58.30a	5.33c	45.80bc
B	61.11b	73.60a	6.33c	28.60def
TaM	55.56b	79.20a	5.00c	24.07efg
TTa	5.56d	5.60bc	14.47b	41.30bcd
TM	27.78c	22.20bc	14.00b	33.97cde
TTaM	5.56d	5.60bc	18.70b	54.00b
BTa	0.00d	0.00c	25.00a	77.20a
BM	5.56d	5.60bc	19.30ab	36.77cde
BTaM	5.56d	5.60bc	18.30b	50.93bc
C	100.00a	97.20a	5.33c	14.60g

Note: *Numbers in the same column followed by the same letter were not significantly different based on the Least Significantly Difference (LSD) test at 5% of significant level. Ta: *T. asperellum*, M: Arbuscular Mycorrhizal Fungi (AMF), T: turmeric extract, B: betel extract, TaM: *T. asperellum* + AMF, TTa: Turmeric + *T. asperellum*, TM: turmeric + AMF, TTaM: turmeric + *T. asperellum* + AMF, BTa: betel + *T. asperellum*, BM: betel + AMF, BTaM: betel + *T. asperellum* + AMF, C: Control.

The treatment combination of botanical fungicides and biological agents have a synergistic effect in suppressing disease development and increasing plant performance compared to the individual treatment. The treatments' effects were consistent from the beginning to the end of the observation (Figure 3).

Application of *T. asperellum* + AMF (TaM), Tumeric + *T. asperellum* (TTa), Tumeric + AMF (TM), Tumeric + *T. Asperelum* + AMF (TTaM), betel + *T. asperellum* (BTa), betel + AMF (BM), betel + *T. asperellum* + AMF (BTaM) reduced significantly disease incidence than control (C). All the treatments significantly reduce disease severity, longer incubation period, and higher dry shoot weight than control, except treatment of *T. asperellum* + Mycorrhiza (TaM). The disease severity, incubation period, and dry shoot weight of the *T. asperellum* + Mycorrhiza (TaM) treatment group were not significantly different from those of the control (Table 3). The combined application of betel + *T. asperellum* (BTa) produced the best inhibitory activity to suppress disease development of corn downy mildew and the capability to increase dry shoot weight (Table 3).

All combinations of biological agents and botanical fungicides suppressed the disease development of corn downy mildew. *T. asperellum* induced the plant to be resistant to various diseases (Silva et al. 2011; Herrera-Téllez et al. 2019; Ramírez-Olier et al. 2019; Sood et al. 2020). It includes resistance to downy mildew in maize (Ginting et al. 2020). *Trichoderma* produced a variety of compounds that induce plant resistance locally and systemically against plant disease. It also induced plant resistance to survive in unfavorable environmental conditions. Moreover, it can promote plant growth by producing phytohormones (Shafawati and Siddiquee 2013; Mukherjee et al. 2013; Błaszczyk et al. 2014; Singh et al. 2002; Sood et al. 2020). *Trichoderma* spp. produce indole Acetic Acid (IAA) as growth hormone (Yudhanegara et al. 2016; Contreras-Cornejo et al. 2009) that increase lateral root

growth, multiply shoots and increase the biomass of shoots (Contreras-Cornejo et al. 2009; Haryuni 2013)

The Arbuscular Mycorrhizal Fungi (AMF) application affected plant growth and health due to more efficient absorption of nutrients because hyphae are bridging the gap between the root and soil biome (Whipps 2004; Tahat et al. 2010). Mycorrhiza increased the uptake of nitrogen (N), potassium (K), and phosphate (P) (Azcón-Aguilar and Barea 1996; Kabirun 2002; Hasanudin 2003; Musfal 2010), and increases the absorption of Cu and Zn (Liu et al. 2000; Watts-Williams et al. 2015; Diagne et al. 2020). Parapasan and Gusta (2014) reported that the mycorrhizal fungi applied to coffee seeds develop hyphae to infect plant roots. The hyphae increased nutrient absorption to support maximal plant growth (Azcón-Aguilar and Barea 1996; Tahat et al. 2010). Mycorrhiza enhance plant health and activate plant defense mechanism against pathogens (Azcón-Aguilar and Barea 1996; Whipps 2004; Tahat et al. 2010; Kamal et al. 2015; Diagne et al. 2020)

Botanical fungicides kill fungal pathogen due to their antifungal compounds. Sesquiterpene, essential ketone oil, turmeron, and artumeron in turmeric have biological activities as antibacterial, antioxidant, and antihepatotoxic (Mallmann et al. 2017; Amalraj et al. 2017). Turmeric has antifungal activity against several types of plant pathogenic fungi, including *Fusarium udum*, *Colletotrichum falcatum* Went, *Fusarium moniliforme* J. Sheld (Singh et al. 2002), *Xanthomonas axonopodis* pv. *manihotis* (Kuhn et al. 2006) and *Alternaria solani* (Balbi-Peña et al. 2006). Secondary metabolites in turmeric inhibit the growth of fungal mycelium (Hu et al. 2017).

Betel leaf extract has antifungal and antibacterial activity (Hertiana and Purwati 2002; Nguyen et al. 2020), decreasing the growth and formation of conidial fungi (Nalina and Rahim 2006). Chemical compounds in the essential oils of betel leaf were chavicol, enavibetol (betel phenol), allylprocatechol (hydroxychavikol), allylpyrocatechol-mono, diacetate, carvacrol, eugenol, p.cymene,

cineole, caryole, cadeneophyl, esragol, terpenes, sesquiterpene, phenyl propane, tannin, diastase, carotene, thiamine, riboflavin, nicotinic acid, vitamin C, sugar, starch and amino acids (Hertiana and Purwati 2002; Nguyen et al. 2020). Chavikol is a fragrance component in betel and has antibacterial properties with antibacterial activity five times stronger than ordinary phenols. The antifungal activity was contributed by chemical compounds of isoeugenol, limonene, and caryophyllene (Hertiana and Purwati 2002).

Trichoderma asperellum and betel leaf extract (BTa) is the best treatment combination in inhibiting corn downy mildew. BTa treatment increases the dry shoot weight. The BTa treatment can be recommended to control corn downy mildew.

ACKNOWLEDGEMENTS

Acknowledgments are conveyed to all those who helped this research. Thanks to the University of Lampung, Indonesia, especially LPPM for the financial support through the Institutional Research Grant scheme and the Faculty of Agriculture for permitting the use of the research facilities during this study.

REFERENCES

- Akhtar MS, Siddiqui ZA. 2008. Arbuscular mycorrhizal fungi as potential bioprotectants against plant pathogens. In: Siddiqui ZA, Akhtar MS, Futai K (eds.) *Mycorrhizae: Sustainable Agriculture and Forestry*. Springer, New York. DOI: 10.1007/978-1-4020-8770-7_3.
- Amalraj A, Pius A, Gopi S, Gopi S. 2017. Biological activities of curcuminoids, other biomolecules from turmeric and their derivatives-A review. *J Tradit Compl Med* 7 (2): 205-233. DOI: 10.1016/j.jtcme.2016.05.005.
- Ayu SR, Prasetyo J, Maryono T. 2013. The influence of some botanical fungicides against downy mildew in corns (*Zea mays* saccharata). *J Agrotek Tropika* 1 (1): 98-101. [Indonesian]
- Azcón-Aguilar C, Barea JM. 1996. Arbuscular mycorrhizas and biological control of soil-borne plant pathogens-an overview of the mechanisms involved. *Mycorrhiza* 6: 457-464. DOI: 10.1007/s005720050147.
- Balbi-Peña MI, Becker A, Stangarlin JR, Franzener G, Lopes MC, Schwan-Estrada KRF. 2006. Control of *Alternaria solani* in tomato by *Curcuma longa* extracts and curcumin-I in vitro evaluation. *Fitopatol Bras* 31 (3): 310-314. DOI: 10.1590/S0100-41582006000400012.
- Bantacut T, Akbar MT, Firdaus YR. 2015. Corn development for food security, industry and economy. *Pangan* 24 (2): 135-148. DOI: 10.33964/jp.v24i2.29. [Indonesian]
- Bardin M, Ajouz S, Comby M, Lopez-Ferber M, Grailot B, Siegwart M, Nicot PC. 2015. Is the efficacy of biological control against plant disease likely to be more durable than that of chemical pesticides?. *Front Plant Sci* 6: 566. DOI: 10.3389/fpls.2015.00566.
- Błaszczczyk L, Siwulski M, Sobieralski K, Lisiecka J, Jedryczka M. 2014. *Trichoderma* spp.-application and prospects for use in organic farming and industry. *J Plant Prot Res* 54 (4): 309-317. DOI: 10.2478/jppr-2014-0047.
- Central Bureau of Statistics. 2016. Production of Maize by Province (tons), 1993-2015. <https://www.bps.go.id> [June 6, 2017]
- Burhanuddin. 2009. Metalaxyl fungicide is not effective in downy mildew (*Peronosclerospora maydis*) in West Kalimantan and alternative control. *Prosiding Seminar Nasional Serealia*. Badan Penelitian dan Pengembangan Pertanian, Maros, 29 Juli 2009. [Indonesian]
- Cameron DD, Neal AL, van Wees SCM, Ton J. 2013. Mycorrhiza-induced resistance: more than the sum of its parts?. *Trends Plant Sci* 18 (10): 539-545. DOI: 10.1016/j.tplants.2013.06.004.
- Contreras-Cornejo HA, Macías-Rodríguez L, Cortés-Penagos C, López-Bucio J. 2009. *Trichoderma virens*, a plant beneficial fungus, enhances biomass production and promotes lateral root growth through an auxin-dependent mechanism in arabidopsis. *Plant Physiol* 149: 1579-1592. DOI: 10.1104/pp.108.130369.
- Diagne N, Ngom M, Djighlay PI, Fall D, Hocher V, Svistoonoff. 2020. Roles of arbuscular mycorrhizal fungi on plant growth and performance: importance in biotic and abiotic stressed regulation. *Diversity* 12 (10): 370. DOI: 10.3390/d12100370.
- Ginting C. 2006. Germination of *Hemileia vastatrix* uredospores on crude water extracts of zinger and turmeric rhizome and clove and Piper betle leaves. *Jurnal Hama dan Penyakit Tumbuhan Tropika* 6 (1): 32-40. DOI: 10.23960/j.hptt.1652-58.
- Ginting C, Prasetyo J, Dirmawati SR, Ivayani, Timotiwu PB, Maryono T, Widyastuti, Chafiska DIR, Asyifa A, Setyowati E, Pasaribu AHZ. 2020. Identification of maize downy mildew pathogen in Lampung. *Ann Res Rev Biol* 35 (7): 23-35. DOI: 10.9734/ARRB/2020/v35i730244.
- Gisi U, Sierotzki H. 2008. Fungicides mode of action and resistance in downy mildews. *Eur J Plant Pathol* 122: 157-167. DOI: 10.1007/978-1-4020-8973-2_12.
- Hasanudin H. 2003. Increasing of the nutrient and uptake availability of N and P and through corn yield of inoculation of mycorrhiza, Azotobacter and on ultisol organic matter. *J Agric Sci Indon* 5 (1): 83-89.
- Haryuni. 2013. Improvement of growth and yield of stevia (*Stevia rebaudiana* Bertoni M) through the application of *Trichoderma* sp. *Biosaintifika* 5 (2): 82-87.
- Herrera-Téllez VI, Cruz-Olmedo AK, Plasencia J, Gavilanez-Ruiz M, Arce-Cervantes O, Hernandez-Leon S, Saucedo-García M. 2019. The protective effect of *Trichoderma asperellum* on tomato plants against *Fusarium oxysporum* and *Botrytis cinerea* diseases involves inhibition of reactive oxygen species production. *Intl J Mol Sci* 20 (8): 1-13. DOI: 10.3390/ijms20082007.
- Hertiana T, Purwati. 2002. Essential oil distilled from ethanol extract of betel leaf (*Piper betle* L.) in several regions in Yogyakarta. *Gadjah Mada University Press, Yogyakarta*. [Indonesian]
- Hu Y, Zhang J, Kong W, Zhao G, Yang M. 2017. Mechanisms of antifungal and anti-aflatoxigenic properties of essential oil derived from turmeric (*Curcuma longa* L.) on *Aspergillus flavus*. *Food Chem* 220: 1-8. DOI: 10.1016/j.foodchem.2016.09.179.
- Jabran K, Ata Z, Farooq M. 2007. Maize: cereal with variety of uses. <https://www.dawn.com/news/236949/maize-cereal-with-a-variety-of-uses> [June 6, 2017].
- Kabirun S. 2002. Padi gogo's response to arbuscular mycorrhiza mushroom inoculation and phosphate fertilization in entisol. *J Ilmu Tanah & Lingkungan* 3 (2): 45-96. [Indonesian]
- Kamal R, Gusain YS, Sharma IP, Sharma S, Sharma AK. 2015. Impact of arbuscular mycorrhizal fungus, *Glomus intraradices*, *Streptomyces* and *Pseudomonas* spp. strain on finger millet (*Eleusine coracana* L.) cv Korchara under water deficit condition. *Afr J Biotechnol* 14 (48): 3219-3227. DOI: 10.5897/AJB2015.14479.
- Kazerooni EG, Sharif A, Nawaz H, Rehman R, Nisar S. 2019. Maize (Corn)-A Useful Source of human nutrition and health: a critical review. *Intl J Chem Biochem Sci* 15 (2019): 35-41.
- Kuhn OJ, Portz RL, Stangarlin JR, Del Águila M, Schwan-Estrada KRF, Franzener G. 2006. Effect of aqueous extract from turmeric (*Curcuma longa*) on *Xanthomonas axonopodis* pv. manihotis. *Semina: Ciênc Agrár* 27 (1): 13. DOI: 10.5433/1679-0359.2006v27n1p13.
- Liu A, Hamel C, Hamilton RI, Ma BL, Smith DL. 2000. Acquisition of Cu, Zn, Mn and Fe by mycorrhizal maize (*Zea mays* L.) grown in soil at different P and micronutrient levels. *Mycorrhiza* 9: 331-336. DOI: 10.1007/s005720050277.
- Mamarabadi M, Tanhaeian A, Ramezany Y. 2018. Antifungal activity of recombinant thanathin in comparison with two plant extracts and a chemical mixture to control fungal plant pathogens. *AMB Expr* 8: 180. DOI: 10.1186/s13568-018-0710-4.
- Mallmann CA et al. 2017. *Curcuma longa* L. Essential oil composition, antioxidant effect, and effect on *Fusarium verticillioides* and fumonisin production. *Food Control* 73: 806-813. DOI: 10.1016/j.foodcont.2016.09.032.
- Manila S, Nelson R. 2014. Biochemical changes induced in tomato as a result of arbuscular mycorrhizal fungal colonization and tomato wilt pathogen Infection. *Asian J Plant Sci Res* 4 (1): 62-68.

- Muis A, Pabendon MB, Nonci N, Waskito WBS. 2013. Genetic variability of downy mildew pathogens based on SSR Marker Analysis. *Penelitian Pertanian Tanaman Pangan* 32: 139-147. [Indonesian]
- Muis A, Nonci N, Pabendon MB. 2016. Geographical distribution of *Peronosclerospora* spp., the causal organism of maize downy mildew, in Indonesia. *Adv Agric Bot* 8: 143-155.
- Mukherjee PK, Horwitz BA, Herrera-Estrella A, Schmoll M, Kenerley CM. 2013. *Trichoderma* research in the genome era. *Ann Rev Phytopathol* 51: 105-129. DOI: 10.1146/annurev-phyto-082712-102353.
- Musfal. 2010. Potential of mycorrhiza arbuscular fungus to increase corn crop yield. *Jurnal Litbang Pertanian* 29 (4): 154-158. DOI: 10.25181/jppt.v14i3.161. [Indonesian]
- Nalina T, Rahim ZHA. 2006. Effect of *Piper betle* L. leaf extract on the virulence activity of *Streptococcus mutans*-An in vitro study. *Pak J Biol Sci* 9 (8): 1470-1475. DOI: 10.3923/pjbs.2006.1470.1475.
- Nega A. 2014. Review on concepts in biological control of plant pathogens. *J Biol Agric Health* 4 (27): 33-54.
- Nguyen LTT, Nguyen TT, Nguyen HN, Bui QTP. 2020. Simultaneous determination of active compounds in *Piper betle* Linn. leaf extract and effect of extracting solvents on bioactivity. *Eng Rep* 2 (10): 1-8. DOI: 10.1002/eng2.12246.
- Parapasan Y, Gusta AR. 2014. Waktu dan cara aplikasi cendawan mikoriza arbuskular (CMA) pada pertumbuhan bibit tanaman kopi. *Jurnal Penelitian Pertanian Terapan* 13 (3): 203-208. [Indonesian]
- Pérez-de-Luque A, Tille S, Johnson I, Pascual-Pardo D, Ton J, Cameron DD. 2017. The interactive effects of arbuscular mycorrhiza and plant growth-promoting rhizobacteria synergistically enhance host plant defenses against pathogens. *Sci Rep* 7: 16409. DOI: 10.1038/s41598-017-16697-4.
- Puyam A. 2016. Advent of *Trichoderma* as a bio-control agent-A review. *J Appl Nat Sci* 8 (2): 1100-1109. DOI: 10.31018/jans.v8i2.927.
- Rahman S, Biswas SK, Barman NC, Ferdous T. 2016. Plant extract as selective pesticide for integrated pest management. *Biotec Res J* 2 (1): 6-10.
- Ramírez-Olier J, Trujillo-Salazar J, Osorio-Echeverri V, Jaramillo-Ciro MM, Botero-Botero L. 2019. In vitro antagonism of *Trichoderma asperellum* against *Colletotrichum gloeosporioides*, *Curvularia lunata*, and *Fusarium oxysporum*. *Revista UIS Ingenierías* 18 (2): 159-166. DOI: 10.18273/revuin.v18n2-2019015.
- Ranum P, Pena-Rosas JP, Garcia-Casal MN. 2014. Global maize production, utilization, and consumption. *Ann N Y Acad Sci* 1312 (2014): 105-112. DOI:10.1111/nyas.12396.
- Rustiani US, Sinaga MS, Hidayat SH, Wiyono S. 2015. Tiga spesies *Peronosclerospora* penyebab penyakit bulai jagung di Indonesia. *Berita Biologi* 14: 29-37. DOI: 10.14203/beritabiologi.v14i1.1860. [Indonesian]
- Shafawati SN, Siddiquee S. 2013. Composting of oil palm fibres and *Trichoderma* spp. as the biological control agent: A review. *Int Biodeterior Biodegr* 85: 243-253. DOI: 10.1016/j.ibiod.2013.08.005.
- Shah TR, Prasad K, Kumar P. 2016. Maize—A potential source of human nutrition and health: A review. *Cogent Food Agric* 2 (1): 1-9. DOI: 10.1080/23311932.2016.1166995.
- Silva F, Ferreira S, Duarte A, Mendonça DI, Domingues FC. 2011. Antifungal activity of *Coriandrum sativum* essential oil, its mode of action against *Candida* species and potential synergism with amphotericin B. *Phytomedicine* 19 (1): 42-47. DOI: 10.1016/j.phymed.2011.06.033.
- Singh R, Singh BK, Upadhyay RS, Rai B, Youn-Su L. 2002. Biological control of *Fusarium* wilt disease of pigeon pea. *Plant Pathol J* 18 (5): 279-283. DOI: 1410-ECN-0101-2009-481-017566509.
- Soenartingsih, Talanca A. 2010. The spread of downy mildew (*Peronosclerospora maydis*) in corn in Kediri district In: Saenong S, Daud D, Amin N (eds). *Prosiding Seminar Ilmiah Dan Pertemuan Tahunan XX PEI-PFI Komda Sulsel*. Balai Penelitian Tanaman Serealia Maros. Maros, 5 November 2008. [Indonesian]
- Sood M, Kapoor D, Kumar V, Sheteiwiy MSS, Ramakrishnan M, Landi M, Aranita F, Sharma A. 2020. *Trichoderma*: The "Secrets" of a multitasking biocontrol agent. *Plants* 9 (6): 1-25. DOI: 10.3390/plants9060762.
- Suharjo R, Swibawa IG, Prasetyo J, Fitriana Y, Danaatmadja Y, Budiawan A, Roerts S, Noorhajati N, Ahmad M, Thines M. 2020. *Peronosclerospora australiensis* is a synonym of *P. maydis*, which is widespread on Sumatra, and distinct from the most prevalent java maize downy mildew pathogen. *Mycol Prog* 19 (11): 1309-1315. DOI: 10.1007/s11557-020-01628-x.
- Tahat MM, Kamaruzaman, Sijam, Othman R. 2010. Mycorrhizal fungi as biocontrol agent. *Plant Pathol J* 9 (4): 198-207.
- Tanklevska N, Petrenko V, Karnaushenko A, Melnykova K. 2020. World corn market: analysis, trends and prospects of its deep processing. *Agric Resour Econ* 6 (3): 96-111. DOI: 10.22004/ag.econ.305555.
- Watts-Williams SJ, Smith FA, McLaughlin MJ, Patti AF, Cavagnaro TR. 2015. How important is the mycorrhizal pathway for plant Zn uptake?. *Plant Soil* 390 (1): 157-166. DOI: 10.1007/s11104-014-2374-4.
- Whipps JM. 2004. Prospects and limitations for mycorrhizas in biocontrol of root pathogens. *Can J Bot* 82: 1198-1227. DOI: 10.1139/b04-082.
- Yoon MY, Cha B, Kim JC. 2013. Recent trends in studies on botanical fungicides in agriculture. *Plant Pathol J* 29 (1): 1-9. DOI: 10.5423/PPJ.RW.05.2012.0072.
- Yudha MK, Soesanto L, Mugiastuti E. 2016. The utilization of four *Trichoderma* sp. isolates for controlling clubroot disease in Chinese cabbage. *Jurnal Kultivasi* 15 (3): 143-149. [Indonesian]
- Zaker M. 2016. Natural plant products as eco-friendly fungicides for plant diseases control-A review. *Agriculturalist* 14 (1): 134-141. DOI: 10.3329/agric.v14i1.29111.

● 14% Overall Similarity

Top sources found in the following databases:

- 7% Internet database
- Crossref database
- 9% Submitted Works database
- 11% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Binus University International on 2019-07-08	2%
	Submitted works	
2	link.springer.com	<1%
	Internet	
3	frontiersin.org	<1%
	Internet	
4	intechopen.com	<1%
	Internet	
5	"Advances in Trichoderma Biology for Agricultural Applications", Sprin...	<1%
	Crossref	
6	themodern.farm	<1%
	Internet	
7	Mohammad Nasir Uddin, Zainul Abedin Siddique, Jabunnisa Akter, Md...	<1%
	Crossref	
8	pjp.pakps.com	<1%
	Internet	

- 9 "Plant-Microbe Interactions in Agro-Ecological Perspectives", Springer ... <1%
Crossref
- 10 "Plant Biotechnology: Recent Advancements and Developments", Sprin... <1%
Crossref
- 11 jurnal.permi.or.id <1%
Internet
- 12 mafiadoc.com <1%
Internet
- 13 olemiss.edu <1%
Internet
- 14 Biological Management of Diseases of Crops, 2013. <1%
Crossref
- 15 Free University of Bolzano on 2020-09-09 <1%
Submitted works
- 16 Hage-Ahmed, Karin, Johannes Krammer, and Siegrid Steinkellner. "The... <1%
Crossref
- 17 ijpba.info <1%
Internet
- 18 biorxiv.org <1%
Internet
- 19 Cranfield University on 2012-10-15 <1%
Submitted works
- 20 University of Sheffield on 2021-09-29 <1%
Submitted works

21	digilib.unila.ac.id	Internet	<1%
22	Associatie K.U.Leuven on 2022-01-12	Submitted works	<1%
23	Cristine Marie B. Brown, Robert A. Nepomuceno, Julieta A. Anarna, Ma...	Crossref	<1%
24	Joko Prasetyo, Sudiono ., Cipta Ginting, Yulia Citra Permatasari. "The E...	Crossref	<1%
25	Mansoura University on 2018-11-21	Submitted works	<1%
26	Papua New Guinea University of Technology on 2017-06-20	Submitted works	<1%
27	repository.ub.ac.id	Internet	<1%
28	"Mycorrhizosphere and Pedogenesis", Springer Science and Business ...	Crossref	<1%
29	Agriculture & Forestry University on 2022-09-02	Submitted works	<1%
30	Bioformulations for Sustainable Agriculture, 2016.	Crossref	<1%
31	F Yelli, J Prasetyo, T Maryono, M Fransiska. "Effect of Oxathiapiproline...	Crossref	<1%
32	Higher Education Commission Pakistan on 2012-09-17	Submitted works	<1%

- 33 **University of Stellenbosch, South Africa on 2021-10-04** <1%
Submitted works
-
- 34 **"Mycorrhiza - Nutrient Uptake, Biocontrol, Eco restoration", Springer Sci...** <1%
Crossref
-
- 35 **A.P. Formighieri, J.R. Stangarlin, C.C. Meinerz, G. Franzener, K.R.F. Sch...** <1%
Crossref
-
- 36 **Kenyatta University on 2020-09-07** <1%
Submitted works
-
- 37 **"Recent Trends in Mycological Research", Springer Science and Busine...** <1%
Crossref
-
- 38 **"Resources Use Efficiency in Agriculture", Springer Science and Busine...** <1%
Crossref
-
- 39 **"Symbiotic Soil Microorganisms", Springer Science and Business Medi...** <1%
Crossref
-
- 40 **Associatie K.U.Leuven on 2019-05-14** <1%
Submitted works
-
- 41 **Saifullah Omar Nasif, Abu Bakar Siddique, Abu Bakar Siddique, Md. Ma...** <1%
Crossref

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded sources
- Quoted material
- Manually excluded text blocks

EXCLUDED SOURCES

smujo.id	74%
Internet	
biodiversitas.mipa.uns.ac.id	74%
Internet	
smujo.id	4%
Internet	
repository.lppm.unila.ac.id	4%
Internet	
dokumen.pub	3%
Internet	
pfi.or.id	3%
Internet	
bibliotecavirtual.dgb.umich.mx:8083	1%
Internet	
The Friends School on 2017-11-01	<1%
Submitted works	
Junliang Zhao, Aihua Liang, Zhen Zhu, Yixiong Tang. "Regeneration of Chines...	<1%
Crossref	
Wenbo Chang, Xue Wang, Ying Lan, Meini Wang, Bin Wen, Lan Li. "Effects of P...	<1%
Crossref	

The British School - RJ on 2022-04-18

Submitted works

<1%

The British School - RJ on 2022-04-18

Submitted works

<1%

Universiti Malaysia Terengganu UMT on 2019-10-21

Submitted works

<1%

Universiti Malaysia Terengganu UMT on 2019-10-20

Submitted works

<1%

EXCLUDED TEXT BLOCKS

B I O D I V E R S I T A S I S S N : 1 4 1 2 - 0 3 3 XVolume 22, Number 4, April 2021E-ISSN: 2...

Universitas Jember on 2022-09-20

The effect

pfi.or.id

biological agent and botanical fungicides on maize downy mildew. Biodiversitas 2...

pfi.or.id

downy mildew

Kasetsart University on 2023-01-28

B I O D I V E R S I T A S 2 2 (4

Universitas Jember on 2022-09-20

A. Control, B

Kasetsart University on 2022-03-25

T. asperellum

www.smujo.id

T. asperellum + AMF

www.smujo.id

downy mildew

Kasetsart University on 2023-01-28

incubation period, and dry

www.smujo.id

BIODIVERSITAS 22 (4

Universitas Jember on 2022-09-20

Contreras-Cornejo HA, Macías-Rodríguez L, Cortés-Penagos C, López

link.springer.com

Bucio J. 2009. Trichoderma virens, a plant beneficial fungus

mafiadoc.com

enhances biomass production and promotes lateral root growth

vdoc.pub

through an auxin-dependent mechanism in arabidopsis. Plant Physiol149: 1579-15...

academica-e.unavarra.es

Diagne N, Ngom M

Mohamed Magdy Fahim Mansour, Manal Mohamed Emam, Karima Hamid Ali Salama, Amal Ahmed Morsy. ...

Roles of arbuscular mycorrhizal fungi on plant growth and

www.intechopen.com

performance: importance in biotic and abiotic stressed regulation

www.intechopen.com

Diversity 12 (10): 370. DOI: 10.3390/d12100370

Sevil Mohammadzadeh, Alireza Pirzad. "Biochemical responses of mycorrhizal-inoculated Lamiaceae (Lave...

Jurnal Hama dan Penyakit Tumbuhan Tropika 6 (1

ejournal.radenintan.ac.id

Ginting C, Prasetyo J

F Yelli, J Prasetyo, T Maryono, M Fransiska. "Effect of Oxathiapiproline-Famoxadone fungicide and Trichode...

2020. Identification of maize

F Yelli, J Prasetyo, T Maryono, M Fransiska. "Effect of Oxathiapiproline-Famoxadone fungicide and Trichode...

23-35.DOI:10.9734/ARRB/2020/v35i730244

insightsociety.org

Hasanudin H. 2003. Increasing of the nutrient and uptake availability of N

journals.areeo.ac.ir

and through corn yield of inoculation of mycorrhiza, Azotobacter and on ultisol org...

M Bahrani, M Davoodi, S Kazemeini. "Wheat Yield as Influenced by Nitrogen Rates, Sources and Tillage Syst...

Herrera-Téllez VI, Cruz-Olmedo AK, Plasencia J

link.springer.com

Arce-Cervantes O, Hernandez-Leon S, Saucedo-Garcia M. 2019. The protective effe...

Naglaa Taha, Said Kamel, Tamer Elsakhawy, Yousry Bayoumi, Alaa El-Dein Omara, Hassan El-Ramady. "Sust...

Numbers in the same column followed by the same letter were not significantly dif...

www.gssrr.org
