

PAPER NAME

**7167-Article Text-34071-2-10-20210105.
pdf**

WORD COUNT

5642 Words

CHARACTER COUNT

28636 Characters

PAGE COUNT

6 Pages

FILE SIZE

359.0KB

SUBMISSION DATE

Mar 3, 2023 2:21 PM GMT+7

REPORT DATE

Mar 3, 2023 2:22 PM GMT+7**● 20% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 13% Internet database
- 15% Publications database
- Crossref database
- Crossref Posted Content database
- 11% Submitted Works database

● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Manually excluded sources
- Manually excluded text blocks

Microbiological, physical and chemical properties of joruk (fermented fish product) with different levels of salt concentration

DYAH KOESOEMAWARDANI^{1,*}, LULU ULYA AFIFAH¹, NOVITA HERDIANA¹, A.S. SUHARYONO¹,
ESA GHANIM FADHALLAH¹, MAHRUS ALI²

¹Department of Agricultural Product Technology, Faculty of Agriculture, Universitas Lampung. Jl. Prof. Dr. Soemantri Brojonegoro No. 1, Bandar Lampung 35145, Lampung, Indonesia. Tel.: +62-721-701609. Fax.: +62-721-702767, *email: dyah.koesoemawardani@fp.unila.ac.id

²Department of Fisheries and Marine Sciences, Faculty of Agriculture, Universitas Lampung. Jl. Prof. Dr. Soemantri Brojonegoro No. 1, Bandar Lampung 35145, Lampung, Indonesia

Manuscript received: 18 November 2020. Revision accepted: 14 December 2020.

Abstract. *Koesoemawardani D, Afifah LU, Herdiana N, Suharyo AS, Fadhallah EG, Ali M. 2021. Microbiological, physical, and chemical properties of joruk (fermented fish product) with different levels of salt concentration. Biodiversitas 22: 132-136.* Joruk is a fermented fish product originated in Ogan Komering Ulu Timur, South Sumatra, Indonesia. The aim of this study was to determine the effect of salt on the microbiological, physical and chemical properties of joruk. This research was arranged in a Randomized Complete Block Design (RCBD), with treatment of salt concentrations at 5%, 10%, 15%, 20%, 25%, and 30% (w/w). The follow-up test used the Least Significance Difference (LSD) at 5%. The result of this study showed that the addition of 10% salt produced the best joruk with the best microbiological and chemical properties as follows: total LAB of 8.75 log cfu/g, total microbes of 13.25 log cfu/g, and total mold of 4.27 log cfu/g, pH of 5.85, total lactic acid of 2.97%, Total Volatile Base (TVB) of 153.05 mgN /100g, and water content of 59.33%. Based on this study, it is concluded that the addition of salt at different concentrations significantly affects the pH, total lactic acid, total lactic acid bacteria (LAB), and water content of joruk.

Keywords: Chemical, joruk, microbiological, physical

INTRODUCTION

Joruk is a fermented food that originated from Ogan Komering Ulu Timur, South Sumatra, Indonesia. Joruk is made from a mixture of Wader fish (*Rasbora* spp.), salt, palm sugar, and cooked rice that is fermented for one to two weeks. Joruk is similar to rusip and bekasam. Rusip is made without cooked rice and bekasam is made without palm sugars, besides that joruk and bekasam are made from freshwater fish, while rusip is made from saltwater fish (Puspita et al. 2017; Melia et al. 2019). Those ingredients are the success factors in the making of fermented fish products, so the development of joruk as one of fermented products requires proper process improvement. Anihouvi et al. (2006) and Kakati and Goswami et al. (2013) stated that fermented fish products are generally processed spontaneously and fermentation goes uncontrolled. This condition resulting in various quality of product and the product is susceptible to damage. Koesoemawardani et al. (2016) reported some of joruk were slimy which leads to spoilage. This condition indicating that the fermentation goes uncontrolled due to improper proportion usage of ingredients that can support the growth of the desired bacteria during fermentation such as lactic acid bacteria (LAB). According to Koesomawardani et al. (2016), Yuktika et al. (2017), and Koesoemawardani et al. (2019), the dominant microbes in fermented fish products are lactic acid bacteria.

The usage of salt is presumed to play an important role in controlling microbes during fermentation in the joruk's production, thus allowing the death of some bacteria that cannot stand in the presence of salt. Salt consists of NaCl will be dissociated to its constituent molecules, Na⁺ and Cl⁻. The Na⁺ ion is needed by the lactic acid bacteria as one of the factors supporting its growth, while the Cl⁻ ion binds the free water which lowering the availability of water in material, so the free water for microbes growth is decreased (Hadiwiyoto, 1993; Yuktika et al. 2017). Salt can select the unwanted bacteria, which is pathogenic and spoilage bacteria during fermentation, so the lactic acid bacteria are expected will dominate at the beginning of fermentation (Koesoemawardani et al. 2013). The activity of lactic acid bacteria which starts due to the decrease of water activity (Aw), water content, and acidic conditions, has become a critical point for inhibiting the growth of pathogen and spoilage bacteria. In addition, the salt used will also affect the characteristics of fermented fish products including physical, chemical, microbiology, and sensory properties (Kim et al. 2014; Fatimah et al. 2017). El Seikha et al. (2014) stated that the fish fermentation process that carried out without using salt will result in uncontrolled fermented products. Therefore, the aim of this study was to determine the effect of salt addition on the microbiological, physical, and chemical properties of joruk.

MATERIALS AND METHODS

Materials

The main materials used in this study were freshwater fish locally called Wader (*Rasbora spp.*), coarse salt (Tani Makmur), palm sugar obtained from the traditional producer, and cooked rice (rice variety of IR 64).

Preparation of materials

The raw materials used are wader fish (whole fish), rice from IR 64 varieties, palm sugar, and salt. The amount of rice and palm sugar expressed as percent per weight of fish in each treatment. The amount of rice is 10% (w/w) and palm sugar is 20% (w/w) used in the formulation of joruk. The first step was to prepare all materials and tools needed. Wader fish are washed with flowing water and then drained. Each sample (treatment) used 100 g of wader fish, and 10 g of rice. The palm sugar used in this research was in liquid form. The liquid palm sugar was prepared by heating the sugar and water (3:1) until it melts at 100 °C for 5 minutes, and then cooled down. The amount of palm sugar added is 20 ml. The salt is weighed per weight of fish according to the treatments, there are 5 g (5%), 10 g (10%), 15 g (15%), 20 g (20%), 25 g (25%), and 30 g (30%).

Joruk production

The production of joruk was carried out according to Koesoemawardani et al. (2019). The prepared wader fish were put in a 150 ml of jar, and then the liquid palm sugar, rice, and salt were added. The mix then stirred until blended, but the rice kept to not completely crumbling (still intact). After all the ingredients are evenly mixed, then the jars were closed tightly by wrapping duct tape on the lid of the jar. Joruk was ready to be analyzed after being stored for seven days in an anaerobic state. Anaerobic condition was created by putting a small jar (150 ml) per sample (treatment) into a large jar with the candle inside in lit condition. The jar is closed tightly by wrapping duct tape on the lid of the jar, and the anaerobic process started after the candle off.

Observation

Measurement of pH

The pH samples were measured using a digital pH-meter (Lovibond), after calibration using standard buffer solution of pH 4 and pH 7.

Water content

Determination of water content was carried out using gravimetric method following the AOAC (2005). Porcelain dishes were dried in the oven for 30 minutes, then cooled in a desiccator and weighed. A total of 5 g of the sample was weighed then put into a porcelain dish and dried in an oven at 105 °C for 3-5 hours. After that, it is cooled down in a desiccator for 15 minutes then weighed. After obtaining the results of the first weighing, then the dish containing the sample was dried again for 30 minutes after which it was cooled in a desiccator for 15 minutes. The measured water content is expressed in percent (%) and calculated using the following formula:

$$\text{Water content (\%)} = \frac{(W+W2)-W1}{W2} \times 100\%$$

Where:

W : Weight of porcelain dish (g)

W1 : Weight of porcelain dish and sample after drying (g)

W2 : Weight of sample (g)

Total lactic acid

The total acid measurement was carried out following the AOAC (2005). A sample of 10 grams was crushed with a Waring blender, then put into a 250 ml Erlenmeyer. Aquadest is added to the Erlenmeyer flask until it reaches the mark. The mixture is shaken out until homogeneous and then filtered. As much as 25 ml of filtrate was added 3 drops of phenolphthalein indicator, then titrated with 0.1 N NaOH solution to form a pink color. Every ml of 0.1 M NaOH was considered equivalent to 90.08 mg of lactic acid. Total lactic acid is calculated using the following formula:

$$\text{Total titratable acidity of lactic acid (mg/ml)} = \frac{\text{ml NaOH} \times \text{N NaOH} \times \text{M.E}}{\text{Volume of sample used}}$$

Where:

ml NaOH : Volume of NaOH used

N NaOH : Molarity of NaOH

M.E : Equivalent factor (90.08 mg)

Total volatile basic nitrogen (TVB-N)

The measurement of TVB-N was conducted following Gaseem (2019). Sample weighed 10 g and then was blended with 50 ml of fresh tap water. The blender was washed with 250 ml of fresh tap water into the distillation flask and 1-2 g of MgO was added to the mixture. TVB-N was released by boiling the mixture with magnesium oxide, which prevented volatile acids from distilling over into the boric acid. The distillate of volatile nitrogen was received in 25 ml of boric acid 2.0% then titrated by 0.1 N sulfuric acid. The TVB-N value was expressed in mg N/100 g and calculated as follow:

$$\text{TVB-N (mg N/100 g)} = \text{Titration (ml of 0.1 N acid)} \times 14$$

Total lactic acid bacteria

The measurement of total lactic acid bacteria was conducted following Khedil et al. (2009) and Karparvar et al. (2019). A sample of 1 g was put into a test tube containing 9 ml of sterile 0.85% physiological salt as a diluent to obtain a sample suspension with a 10⁻¹ dilution to a 10⁻¹⁰ dilution. A total of 1 ml of sample each 10⁻⁸, 10⁻⁹, and 10⁻¹⁰ dilutions were piped and put into each sterile petri dish, then poured ±15 ml of sterile MRS media agar (done in Duplo for each dilution) and shake evenly on the table or follow the pattern of number 8. After the agar media solidified, the plates were wrapped in paper and then incubated in upside-down position at 37 °C ± 1 °C for 48 ± 2 hours. The total number of lactic acid bacteria was counted (on a 30-300 colony scale) and expressed in cfu/g. The total lactic acid bacteria were calculated using the following formula:

$$\text{Total lactic acid bacteria} = \text{The number of colonies counted} \times (1/\text{dilution factor})$$

Total microbial count

The measurement of total microbial count follows the procedure of SNI 2333.3: 2015. A total of 1 g of sample was diluted with 9 ml of sterilized physiological saline (NaCl 0.85%). This dilution is calculated as a 10^{-1} dilution. Further dilution is carried out by dissolving 1 ml of the 10^{-1} dilution solution with 9 ml of physiological saline and counting it as a 10^{-2} dilution and so on up to 10^{-12} dilutions. A total of 1 ml of sample from each dilution 10^{-10} , 10^{-11} , and 10^{-12} were taken and put into each sterile petri dish, then poured \pm 15 ml of sterile PCA (done in Duplo for each dilution) and shake evenly on the table or follow the pattern of number 8. After the agar media solidified, the plates were wrapped in paper and then incubated in upside-down position at $35 \text{ }^{\circ}\text{C} \pm 1 \text{ }^{\circ}\text{C}$ for 48 ± 2 hours. The total number of microbes was counted and expressed in cfu/g.

Total mold

The measurement of total mold follows the procedure of SNI 2332.7: 2015. A total of 1 g of sample was diluted with 9 ml of sterilized physiological saline (NaCl 0.85%). This dilution is calculated as dilution 10^1 . Further dilution is carried out by dissolving 1 ml of the 10^{-1} dilution solution with 9 ml of physiological saline and counting it as a 10^{-2} dilution and so on up to 10^{-6} dilutions. A total of 1 ml of sample from each dilution of 10^{-4} , 10^{-5} , and 10^{-6} , poured plates were made on Potato Dextrose agar (PDA) medium. The incubation condition was at $25 \text{ }^{\circ}\text{C}$ for 3-5 days in a non-inverted position. The calculation was done using the Standard Plate Count (SPC) method, which is using a colony counter with the condition that the number of colonies per Petri dish is between 10-150 colonies, and there is no colony whose size covers more than half of the Petri dish.

Data analysis

This study used non-factorial Randomized Complete Block Design (RCBD) with four replications. The data analyzed by variance to determine whether there were differences among treatments. The similarity of data tested by the Tuckey test and following by the Smallest Real Difference (SRD) test at the level of 5% with SPSS 18.0 for Windows. Treatments given to each replication are salt concentrations (G) of 5% (G1), 10% (G2), 15% (G3), 20% (G4), 25% (G5), and 30% (G6) (w/w).

RESULTS AND DISCUSSION

Total Lactic Acid Bacteria (LAB)

The results of analysis of variance showed that salt concentration level had significant effect on total lactic acid bacteria of joruk during fermentation as shown in Table 1. The control treatment (0% of salt) was not carried out in this study because if the joruk was made without using salt it would result in uncontrolled fermentation products (unsuccessful). Total lactic acid bacteria in joruk ranged from 8.49-8.75 log cfu/g with the highest from 5%-10% salt concentration, 8.73-8.75 log cfu/g, and the lowest from 15%-30% salt concentration. The addition of high salt

concentration decreases the number of lactic acid bacteria, this phenomenon was inversely proportional to the observed pH value. This occurs because the addition of high salt causes the growth inhibition of lactic acid bacteria, so the lactic acid bacteria did not optimally produce lactic acid. Salt consists of its constituent molecules, Na^+ and Cl^- , where Na^+ is much needed by the lactic acid bacteria to support their growth factor, and Cl^- will binds the free water that causes the decrease of availability of water so that the free water for lactic acid bacteria's growth also reduced (Hadiwiyoto 1993; Yuktika et al. 2017). Due to the osmotic pressure of the solution, it causes plasmolysis which is followed by dehydration and then followed by the death of microorganisms. Salt also affects the water activity of the ingredients, so that it can control the growth of microorganisms. Therefore, the addition of high salt concentration causes lower water activity (A_w) value, results in the decreasing rate of lactic acid bacterial growth and increases the pH value (Aristyan, et al. 2014; Ahillah, et al. 2017). The growth of lactic acid bacteria was inhibited with salt concentration ranging from 15% to 30%. In this study, lactic acid bacteria growth was stunted due to high salt concentration starting from 15% to 30%. Total lactic acid bacteria results in this study were close to the results in Koesoemawardani et al. (2019), but they were lower than the results in Koesoemawardani et al. (2016) with 10.19 log cfu/g.

Total microbes

Total microbes in joruk with addition of different salt concentrations ranged from 13,39-13,18 log cfu/g. The result of analysis of variance showed that the addition of salt concentration had no significant effect on total microbes on joruk during fermentation (Table 1).

The same result also reported by Koesoemawardani et al. (2019) and Koesoemawardani et al. (2016). Furthermore, Desniar et al. (2012) reported the total microbes were not significantly different due to the total number of microbes counted are the microbes that involved in fermentation process, such as acid-forming bacteria (lactic, propionate, acetate), and several types of yeast and mold, while Widjajanti et al. (1996) reported that ingredient composition, including the salt used, also did not significantly affect the total number of microbes in bekasam because the available nutrients were sufficient for the growth of lactic acid bacteria, mold, and yeast, and their growth inhibitor product had not been formed.

Table 1. Microbiological properties of joruk

Treatment	Total LAB	Total microbes (log cfu/g)	Total mold
G1 (5%)	8.75 \pm 0.09 ^a	13.39 \pm 0.08	4.32 \pm 0.03
G2 (10%)	8.73 \pm 0.14 ^a	13.17 \pm 0.17	4.27 \pm 0.07
G3(15%)	8.57 \pm 0.10 ^b	13.24 \pm 0.19	4.29 \pm 0.08
G4 (20%)	8.50 \pm 0.25 ^b	13.24 \pm 0.18	4.36 \pm 0.06
G5 (25%)	8.49 \pm 0.15 ^b	13.20 \pm 0.14	4.30 \pm 0.06
G6 (30%)	8.49 \pm 0.07 ^b	13.24 \pm 0.20	4.27 \pm 0.08

Note: different letter in the same column indicated significant differences between treatments ($p < 0,05$)

The fermentation type of joruk is spontaneous fermentation, while the dominant total microbes are lactic acid bacteria (Koesoemawardani et al. 2016; Yuktika et al. 2017; Koesoemawardani et al. 2019). According to Paludan-Müller et al. (2002), the addition of high salt concentration causes certain types of microorganisms that are resistant to high salt levels will grow, such as halophilic bacteria. The total microbes measured in this study were also affected by the number of lactic acid bacteria that grow. Meanwhile, lactic acid bacteria can still grow in the range of 5% to 35% salt added (Zuidar et al. 2016; Ahillah et al., 2017).

Total mold

Total mold of joruk with addition of different salt concentrations ranged from 4.37-4.27 log cfu/g. The result of analysis of variance showed that the addition of salt concentration had no significant effect on total mold of joruk during fermentation (Table 1). This is because the available nutrients are still sufficient for mold and yeast, and also their growth inhibitor product had not been formed, so the salt addition had no significant effect on the mold's growth (Widjajanti et al. 1996). The physiological properties of mold were still within the limits of their growth requirements. According to Querol and Fleet (2006) and Feldmann (2012), the physiological properties of molds are including the requirements of water, growth temperature, oxygen requirement, pH, nutrients, and inhibitor compound. The observation results of water content in this study ranged from 44.74-61.78%, while the water content requirements for biochemical reactions in molds ranged on 90%, so it still supports mold growth. The optimum temperature for mold growth ranged from 25-30 °C or more, and in this study presumed that the storage condition still in room temperature (25-30 °C). Mold grows well at pH 2-8.5, while the pH in this study ranged from 4.65-6.11, so the mold still grows well in this condition. Meanwhile, the availability of oxygen still supports mold growth because the conditions of joruk's container in this study are facultatively anaerobic, so the molds still have oxygen to grow even in small amounts. Mold nutrients are in the form of carbon sources, nitrogen sources, energy sources, and growth factors (minerals and vitamins). The source of these nutrients in joruk was added in the same amount, while the treatment addition of salt was added on different concentration in this study only had an effect on lactic acid bacteria as the growth support. Therefore, the total mold in this research was not significantly different on each treatment. The inhibiting component was not present in joruk. According to Feldmann (2012), the inhibiting components of mold growth are including penicillium, clavasin, sorbic, propionic, and acetic acid. The growth of mold in this research was not different from Koesoemawardani et al. (2019).

pH value

The results of analysis of variance showed that the salt addition had significant effect on pH (Table 2). The pH value ranged from 4.65-6.11, while the pH value before fermentation on each treatment was not significantly

different ranged from 6.53-6.68. The highest pH value of joruk ranged from 5.96-6.11 obtained in the treatment salt addition of 15%-30%, while the lowest pH value of joruk is 4.65 from the treatment salt addition of 5%. The increase in pH value was due to the addition of the salt concentration used. The higher salt concentration used can inhibit and decrease the growth of lactic acid bacteria. This will reduce the ability of lactic acid bacteria to produce lactic acid. Therefore, the high concentration of salt addition was not effective in reducing pH (Hadiyanti dan Wikandari, 2013; Kusmawarti et al., 2011; Thariq et al. 2014).

The increase in pH value could be occurred due to increased activity of decomposing bacteria in non-protein nitrogen compound that produces volatile bases. In addition, the increase in the addition of salt concentration results in the fermentation process was not occurred perfectly, because the fish meat has been damaged by salt so the ability of lactic acid bacteria in fermentation is reduced (Maliki et al. 2019). The pH value of joruk in previous study reached 5.43-5.78 (Koesoemawardani et al. 2016) and 4.92-5.26 (Koesoemawardani et al. 2019). The pH values in those studies are lower due to the low concentration of salt used, 10%, while in this study use the highest concentration was 30%. Gaseem (2019) stated that pH value of each fermented fish product has different value due to processing methods, variations in the ingredients used and the variation of amount (material formulation) added. The pH is one of the quality parameters of fermented fish products, even though there is no standard. Fermented fish product with pH value of 6.5 or higher, it is considered as a bad quality (Anihouvi et al. 2012). The quality of joruk in this study is still suitable for consumption because the pH value of joruk ranged from 4.65-6.11.

Total lactic acid

The results of analysis of variance showed that the addition of salt significantly affected the total lactic acid during fermentation (Table 2). The total amount of lactic acid ranges from 1.98-3.06%. Salt addition supports the growth of lactic acid bacteria so that the fermentation process can run perfectly. Lactic acid bacteria will produce enzymes that hydrolyze complex carbohydrates into glucose and then will be converted into organic acids, especially lactic acid (Abdel-Rahman et al. 2013; Huang et al. 2020; Lestari et al., 2018; Rizal et al. 2020; Villalobos et al. 2020). However, if the salt added in higher concentration, the growth of lactic acid bacteria will be reduced and the production of lactic acid will be decreased. This result was in line with the total lactic acid bacteria observed in this study that the higher addition of salt can reduce the number of lactic acid bacteria. This is followed by a decrease in the formation of total lactic acid. The use of different proportions of ingredients in fermented fish products results in different total acids (Widjajanti et al. 1996). This could be causing the difference of total lactic acid produced in this study compared with Koesoemawardani et al. (2016) and Koesoemawardani et al. (2019), respectively, the total lactic acid formed was 0.71% and 6.21%-6.92%.

Table 2. Chemical properties of joruk

Treatment	pH Value	Total lactic acid (%)	Water content (%)
G1 (5%)	4.64±0.27 ^c	3.00±0.36 ^d	61.83±2.98 ^a
G2 (10%)	5.84±0.24 ^b	2.97±0.18 ^{cd}	59.33±2.53 ^{ab}
G3 (15%)	5.96±0.09 ^a	2.88±0.00 ^{bc}	55.76±4.70 ^{abc}
G4 (20%)	6.01±0.07 ^a	2.52±0.29 ^{ab}	51.49±5.20 ^{abcd}
G5 (25%)	6.01±0.15 ^a	2.34±0.20 ^a	44.73±8.08 ^d
G6 (30%)	6.11±0.08 ^a	1.98±0.20 ^a	49.90±8.76 ^{cd}

Note: different letter in the same column indicated significant differences between treatments (p<0,05)

Water content

The results of analysis of variance showed that the addition of salt concentration had a significant effect on the water content of joruk during fermentation (Table 2). The water content of joruk produced in study ranged from 44.73 to 61.87%. The higher addition of salt concentration causes a decrease in water content of joruk. Salt plays a role in attracting the water from fish body through an osmosis process, so the water content of joruk will decrease. The decrease of water content causes the water activity (A_w) value to decrease as the concentration of salt used increases. This result indicates that the higher salt concentration, the smaller water activity. Water activity affects microbial growth because it is a medium for growth and reproduction for bacteria (Aristyan et al. 2014; Majid, et al. 2014; Thariq et al. 2014; Fatimah et al. 2017; Puspita et al. 2019; Koesemawardani et al. 2020). The water content of joruk in this study was similar to the result reported by Koesemawardani et al. (2016) that is 61.04%, while Koesemawardani et al. (2019) reported joruk with a higher water content of 63.30%. This happens due to difference in ingredient proportion added, especially the usage of salt concentrations.

Best treatment selection

Joruk is one of fermented products with potential to be functional food because of the presence of lactic acid bacteria that is good for gut health. In fermented products, low total mold is desirable (Indonesian National Standard of fermented fish, SNI 2716: 2016). Based on these requirements, the treatment that produces joruk with high total lactic acid bacteria and low total mold will be selected as the best treatment. In this study, the treatment with the highest total lactic acid bacteria result was found in joruk with 10% salt concentration (w/w).

In conclusion, based on the results from this study, it can be concluded that salt concentration had a significant effect on total LAB, pH value, total lactic acid, and water content in joruk, however, it had no significant effect on total microbes and total mold in joruk. The addition of 10% salt produced the best joruk with the best microbiological, physical, and chemical properties as follows: total LAB of 8.75 log cfu/g, total microbes of 13.25 log cfu/g, and total mold of 4.27 log cfu/g, pH of 5.85, total lactic acid of 2.97%, Total Volatile Base (TVB) value of 153.05 mgN/100g, and water content of 59.33%. Further research will

be carried out regarding the investigation of amino acid, fatty acid, and volatile compounds from joruk.

ACKNOWLEDGEMENTS

We offer our gratitude to the Ministry of Research and Technology/National Research and Innovation Agency for their funding through Competitive Research 2019-2020

REFERENCES

- Abdel-Rahman MA, Tashiro Y, Sonomoto K. 2013. Recent advances in lactic acid production by microbial fermentation processes. *Biotechnol Adv* 31: 877-902.
- Ahillah N, Rusdanillah A, Afiana W, Sulistiani R, Mail RP. 2017. The influence of salt concentration on the fermentation of the wader fish (*Rasbora lateristriata*). *Jurnal Bioedukasi* 10: 12-17.
- Anihouvi VB, Ayernor GS, Hounhouigan JD, Akyi-Dawsoin E. 2006. Quality characteristics of lanhouin: a traditional processed fermented fish product in the Republic of Benin. *Afr J Food Agric Nutr Dev* 6: 1-15.
- Anihouvi VB, Kindossi JM, Hounhouigan JD. 2012. Processing and quality characteristics of some major fermented fish products from Africa: A critical review. *Intl Res J Biol Sci* 1: 72-84.
- AOAC. 2005. Official Methods of Analysis. Association of Official Analytical Chemists, Washington DC.
- Aristyan I, Ibrahim R, Rianingsih L. 2014. The influence of different percentages of salt toward sensory and microbiological quality of shrimp paste. *Jurnal Pengolahan dan Bioteknologi Hasil Perikanan* 3: 60-66. [Indonesian]
- Desniar, Setyaningsih I, Sumardi RS. 2012. Chemical and microbiological parameter changes and isolation of acid-producing bacteria during fermentation process of common carp (*Cyprinus carpio*) bekasam. *Jurnal Pengolahan Hasil Perikanan Indonesia* 15: 232-239. [Indonesian]
- El Sheikh AF, Ray R, Montet D, Panda S and Worawattanamateekul W. 2014. African fermented fish products in scope of risks. *Intl Food Res J* 21: 425-432
- Fatihah F, Pelealu JJ, Gugule S, Yempormase HV, Tallei TE. 2017. Quality evaluation of bakasang processed with variation of salt concentration, temperature and fermentation time. *Pak J Biol Sci* 20: 543-551. DOI: 10.3923/pjbs.2017.543.551
- Gaseem, MA. 2019. Microbiological and chemical quality of a traditional salted-fermented fish (Hout-Kasef) product of Jazan Region, Saudi Arabia. *Saudi J Biol Sci* 26: 137-140.
- Hadiwiyoto S. 1993. *Teknologi Pengolahan Hasil Perikanan*. Liberty, Yogyakarta.
- Hadiyanti MR, Wikandari PR. 2013. The effect of salt concentration and the addition of lactic acid bacteria *Lactobacillus plantarum* B1765 as starter culture for the quality product of milkfish (*Chanos chanos*) bekasam. *UNESA J Chemistry* 2: 136-143.
- Huang Z, Huang L, Xing G, Xu X, Tu C, Dong M. 2020. Effect of cofermentation with lactic acid bacteria and *K. marxianus* on physicochemical and sensory properties of goat milk. *Foods* 9: 299. DOI: 10.3390/foods9030299
- Kakati BK, Goswami UC. 2013. Characterization of the traditional fermented fish product Shidol of Northeast India prepared from *Puntius sophore* and *Setipinna phasa*. *Indian J Trad Knowl* 12: 85-90.
- Karparvar N, Safaei HG, Derakhshandeh A, Hemmati F, Mazloomi SM. 2019. Isolation and identification of lactic acid bacteria from a traditional fermented fish sauce (Mahyaveh) in Fars Province, Iran. *Intl J Nutr Sci* 4: 49-53. DOI:10.30476/IJNS.2019.81437.1005
- Khedid K, Faïd M, Mokhtari A, Soulaymani A, Zinedine A. 2009. Characterization of lactic acid bacteria isolated from the one humped camel milk produced in Morocco. *Microbiol Res* 164: 81-91. DOI: 10.1016/j.micres.2006.10.008
- Kim BK, Lee HH, Jeng MH, Shim KB. 2012. Effect of fermentation temperature and salt concentration on the quality of jack mackerel (*Trachurus japonicus*) fish sauce. *J Fish Mar Sci Educat* 24: 755-762. DOI: 10.13000/JFMSE.2012.24.6.755

- Koesoemawardani D, Herdiana N, Suharyono, Hasly MZ, Ali M. 2019. The influence of cooked rice addition on the quality of joruk, an Indonesian freshwater fermented fish. *Biosci Res* 16: 3443-3448.
- Koesoemawardani D, Hermawan YE, Herdiana N, Susilawati. 2020. Characteristic of rusip from trash fish. *J Agric Prod Technol Industr* 25: 120-128. DOI: <http://dx.doi.org/10.23960/v25i2.120-128>
- Koesoemawardani D, Marniza, Rizal S, Sella N. 2016. Addition of palm sugar in "Joruk" (fish fermentation). In: *Prosiding Seminar Nasional Pengembangan Teknologi Pertanian*. DOI: 10.25181/prosemnas.v0i0.477. [Indonesian]
- Koesoemawardani D, Rizal S, Tauhid M. 2013. Microbiological and chemical changes of rusip during fermentation. *agriTECH* 33: 265-272. DOI: 10.22146/agritech.9547
- Kusmarwati A, Heruwati ES, Utami T, Rahayu ES. 2011. Effect of the addition of *Pediococcus acidilactici* F-11 as a starter culture on the quality of rusip. *Jurnal Pascapanen dan Bioteknologi Kelautan dan Perikanan* 6: 13-26.
- Lestari S, Rinto, Huriyah SB. 2018. Increasing functionality of bekasam with *Lactobacillus acidophilus* as a starter. *Indon Fish Processing J* 21: 179-187. DOI: 10.17844/jphpi.v21i1.21596
- Majid A, Tri WA, Laras R. 2014. The influence of different salt concentration on the sensory quality and volatile compounds of anchovy paste (*Stolephorus* sp.). *Jurnal Pengolahan dan Bioteknologi Hasil Perikanan* 3: 17-24. [Indonesian]
- Maliki MK, Afrianto E, Liviawaty E, Rostini I. 2019. Effect of salt concentration with addition of lactic acid bacteria to preference levels of non-salty peda. *Global Sci J* 7: 388-393.
- Melia S, Purwati E, Kurnia YF, Pratama DR. 2019. Antimicrobial potential of *Pediococcus acidilactici* from bekasam, fermentation of sepat rawa fish (*Tricopodus trichopterus*) from Banyuasin, South Sumatra, Indonesia. *Biodiversitas* 20: 3532-3538.
- Paludan-Müller, Madsen M, Sophanodora P, Gram L, Møller PL. 2002. Fermentation and microflora of Plaasom a Thai fermented fish product prepared with different salt concentration. *Intl J Food Microbiol* 73: 61-70. DOI: 10.1016/S0168-1605(01)00688-2
- Puspita DA, Agustini TW, Purnamayati L. 2019. The effect of different concentration of salt towards the glutamic acid in catfish Bakasang powder. *Jurnal Teknologi Pangan* 3: 110-115. DOI: 10.14710/jtp.3.1.110-115. [Indonesian]
- Puspita ID, Wardani A, Puspitasari ROA, Nugraheni PS, Putra MMP, Pujiraharti S, Ustadi. 2017. Occurrence of chitinolytic bacteria in shrimp rusip and measurement of their chitin-degrading enzyme activities. *Biodiversitas* 18: 1275-1281.
- Querol A, Fleet. 2006. *Yeasts in Food and Beverages*. Springer, Brazil.
- Rizal S, Suharyono, Nurainy F, Amelia JR. 2020. The effects of low-temperature storage on the viability of *Lactobacillus casei* and the stability of antibacterial activity in green grass jelly synbiotic drinks. *Biodiversitas* 21: 3826-3831. DOI: 10.13057/biodiv/d210851
- SNI 2332.3: 2006. Determination of total plate count (ALT) in fishery products. National Standardization Agency of Indonesia, Jakarta, Indonesia.
- SNI 2332.7: 2015. Microbiological test method: Calculation of mold and yeast in fishery products. National Standardization Agency of Indonesia, Jakarta
- SNI 2716: 2016. Shrimp paste. National Standardization Agency of Indonesia, Jakarta, Indonesia.
- Thariq AS, Swastawati F, Surti T. 2014. The effect of different concentration of salt to the content of glutamic acid savoring flavor (umami) of mackerel fermented fish (*Rastrelliger neglectus*). *Jurnal Pengolahan dan Bioteknologi Hasil Perikanan* 3 (3): 104-111. [Indonesian]
- Villalobos JAM, Zamora JM, Barboza N, Garbanzo CR, Usaga J, Solano MR, Schroedter L, Widdrat AO, Gómez JPL. 2020. Multi-product lactic acid bacteria fermentations: A review. *Fermentation* 6: 1-21. DOI: 10.3390/fermentation6010023
- Widjajanti H, Arwinsyah, Munawar. 1996. Pengaruh komposisi bahan dan lama penyimpanan terhadap aspek mikrobiologi dan aspek biokimia dalam pembuatan bekasam. *Jurnal Penelitian Sains* 1: 11-15. DOI: 10.26554/jps.v0i1.222. [Indonesian]
- Yuktika S, Sutiyantri E, Dhewi ES, Martika SD, Sa'diyah RD. 2017. The influence of salt concentration on the fermentation of shrimp. *Bioedukasi* 10: 18-22. DOI: 10.20961/bioedukasi-uns.v10i2.11568. [Indonesian]
- Zuidar A, Rizal S, Widyastuti K. 2016. The effect of fish type and salt concentrations on bamboo shoots fish fermented. *Jurnal Kelitbangan* 4: 181-194.

● 20% Overall Similarity

Top sources found in the following databases:

- 13% Internet database
- Crossref database
- 11% Submitted Works database
- 15% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Mustafa A. Gassem. "Microbiological and chemical quality of a traditio...	2%
	Crossref	
2	smujo.id	1%
	Internet	
3	agritech.unhas.ac.id	<1%
	Internet	
4	mail.scialert.net	<1%
	Internet	
5	D R Pratama, E Purwati, Yuherman, S Melia. "The potential of probiotic...	<1%
	Crossref	
6	S Mayyoga, I U P Rangkuti, H Purwanto. "The effect of contact time of ...	<1%
	Crossref	
7	Z. Hanum, Yurliasni, N. Mardhiah, Z.M. Gaznur. "Antibacterial Activity ...	<1%
	Crossref	
8	Sumardianto Sumardianto, Ima Wijayanti, Fronthea Swastawati. "Chara...	<1%
	Crossref	

9	mafiadoc.com	Internet	<1%
10	"Ethnic Fermented Foods and Alcoholic Beverages of Asia", Springer S...	Crossref	<1%
11	Universitas Brawijaya on 2021-10-03	Submitted works	<1%
12	journal.upgris.ac.id	Internet	<1%
13	journalbalitbangdalamampung.org	Internet	<1%
14	"Ethnic Fermented Foods and Beverages of India: Science History and ...	Crossref	<1%
15	Microbiology Monographs, 2015.	Crossref	<1%
16	Sumartini, P W Ratrinia, K H Sitorus. "Proximate and nutrition analysis ...	Crossref	<1%
17	probiotics.wg.ugm.ac.id	Internet	<1%
18	juszzks.com.ba	Internet	<1%
19	Universidade de Sao Paulo on 2022-05-18	Submitted works	<1%
20	Universitas Hasanuddin on 2019-09-28	Submitted works	<1%

- 21 **Xue Chen, Pengcheng Dong, Ke Li, Lixian Zhu et al. "Effect of the combi...** <1%
Crossref
-
- 22 **Bahni Dhar, Binaya B. Nayak, Ranendra K. Majumdar, Amjad K. Balange...** <1%
Crossref
-
- 23 **Dessy Abdullah, Sandeep Poddar, Ramesh Prasath Rai, Endang Purwat...** <1%
Crossref
-
- 24 **N N Lubis, I Suhaidi, E Yusraini. "The effect of sauerkraut liquid with su...** <1%
Crossref
-
- 25 **Universitas Slamet Riyadi on 2020-11-07** <1%
Submitted works
-
- 26 **jaabjournal.org** <1%
Internet
-
- 27 **ncbi.nlm.nih.gov** <1%
Internet
-
- 28 **A Kusmarwati, U Hizamah, S Wibowo. " Microbiological and chemical q...** <1%
Crossref
-
- 29 **jurnal.fp.unila.ac.id** <1%
Internet
-
- 30 **CTI Education Group on 2020-09-07** <1%
Submitted works
-
- 31 **Universiti Teknologi MARA on 2018-12-23** <1%
Submitted works
-
- 32 **biblio.ugent.be** <1%
Internet

33	veterinaryworld.org	Internet	<1%
34	D Koesoemawardani, S Hidayati, Subeki. " Amino acid and fatty acid c...	Crossref	<1%
35	LU FENG, YING LI, CHANGLU MA, YANFENG TUO. " Bacterial Diversity ...	Crossref	<1%
36	Monte Sant' Angelo College on 2020-08-04	Submitted works	<1%
37	Multimedia University on 2021-06-27	Submitted works	<1%
38	Siddegowda Gopalapura Shivanne Gowda, Bhaskar Narayan, Shubha G...	Crossref	<1%
39	UNESCO-IHE Institute for Water Education on 2014-04-04	Submitted works	<1%
40	Universitas Airlangga on 2020-09-25	Submitted works	<1%
41	Universitas Jenderal Soedirman on 2018-03-18	Submitted works	<1%
42	Xu, Li-Li, Pan-Pan Hu, Xiao-Le Kong, Robert C. Hider, Tao Zhou, and Zhi...	Crossref	<1%
43	link.springer.com	Internet	<1%
44	pinpdf.com	Internet	<1%

45	repo.unand.ac.id Internet	<1%
46	mdpi.com Internet	<1%
47	scielo.br Internet	<1%
48	tandfonline.com Internet	<1%
49	Lincoln University on 2019-03-28 Submitted works	<1%
50	Sri Melia, Indri Juliyarsi, Yulianti Fitri Kurnia. "Physicochemical properti... Crossref	<1%
51	Manish Kumar Chatli, Neeraj Gandhi, Parminder Singh. "Efficacy of sod... Crossref	<1%
52	Sharon Xi Ying Chan, Nursyah Fitri, Nurul Syahidah Mio Asni, Nor Hafiz... Crossref	<1%
53	T. Wang. "Combined Application of Modified Atmosphere Packaging a... Crossref	<1%
54	"Novel Food Fermentation Technologies", Springer Science and Busine... Crossref	<1%
55	Unika Soegijapranata on 2015-03-11 Submitted works	<1%
56	iGroup on 2015-08-01 Submitted works	<1%

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded sources
- Quoted material
- Manually excluded text blocks

EXCLUDED SOURCES

smujo.id	85%
Internet	
biodiversitas.mipa.uns.ac.id	85%
Internet	
M R Hanifah, R Naufalin, R Wicaksono. "The effect of edible coating contained...	4%
Crossref	
repository.lppm.unila.ac.id	4%
Internet	
Rifda Naufalin, Rumpoko Wicaksono, Eva Triyulianingrum. "Edible coating app...	3%
Crossref	
isisn.org	2%
Internet	
atlantis-press.com	2%
Internet	
sciencegate.app	1%
Internet	

EXCLUDED TEXT BLOCKS

B I O D I V E R S I T A S Volume 22, Number 1, January 2021 Pages

Universitas Bengkulu on 2023-02-28

Microbiological, physical

repository.lppm.unila.ac.id

fermented fish product originated in Ogan Komering Ulu Timur, South Sumatra, Ind...

repository.lppm.unila.ac.id

et al. - Microbiological, physical and chemical

etheses.dur.ac.uk

BIODIVERSITAS 22 (1

Universitas Bengkulu on 2023-02-28

et al. - Microbiological, physical and chemical

etheses.dur.ac.uk

BIODIVERSITAS 22 (1

Universitas Bengkulu on 2023-02-28

Where:W : Weight of porcelain dish (g)W1 : Weight of porcelain dish

app.tisi.go.th

MATERIALS AND METHODSMaterialsThe

Arif Nuryawan, Nova O. Hutauruk, Esra Yunita S. Purba, Nanang Masruchin et al. "Properties of wood compo..."

Where:ml NaOH : Volume of NaOH usedN NaOH : Molarity of NaOHE: Equivalent...

CTI Education Group on 2020-09-07

calculated asfollow:TVB-N (mg N/100 g) = Titration (ml of 0.1 N acid) × 14

Mustafa A. Gassem. "Microbiological and chemical quality of a traditional salted-fermented fish (Hout-Kase..."

The number of colonies

Z. Hanum, Yurliasni, N. Mardhiah, Z.M. Gaznur. "Antibacterial Activity of Kefir Grain Levels on Fermented Go..."

RESULTS AND DISCUSSIONTotal Lactic Acid Bacteria

www.smujo.id

G1 (5%)G2 (10%)G3(15%)G4 (20%)G5 (25%)G6 (30

www.researchgate.net

in the same column indicated significantdifferences

www.ijat-aatsea.com

in the same column indicated significantdifferences

www.ijat-aatsea.com