

PAPER NAME

Theinfluenceofcookedriceadditiononthe
qualityofjoruk-Indonesianfreshwaterferm
entedfish.pdf

WORD COUNT

4048 Words

CHARACTER COUNT

20958 Characters

PAGE COUNT

6 Pages

FILE SIZE

842.9KB

SUBMISSION DATE

Mar 3, 2023 2:21 PM GMT+7

REPORT DATE

Mar 3, 2023 2:22 PM GMT+7

● **9% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 6% Internet database
- Crossref database
- 5% Submitted Works database
- 4% Publications database
- Crossref Posted Content database

● **Excluded from Similarity Report**

- Bibliographic material
- Manually excluded sources
- Quoted material
- Manually excluded text blocks



Available online freely at www.isisn.org

Bioscience Research

Print ISSN: 1811-9506 Online ISSN: 2218-3973

Journal by Innovative Scientific Information & Services Network



RESEARCH ARTICLE

BIOSCIENCE RESEARCH, 2019 16(4): 3443-3448.

OPEN ACCESS

The influence of cooked rice addition on the quality of *Joruk*, an Indonesian freshwater fermented fish

Dyah Koesoemawardani^{1*}, Novita Herdiana¹, Suharyono¹, Muhammad Zaenal Hasly¹, and Mahrus Ali²

¹Department of Agricultural Technology, University of Lampung, Bandar Lampung, Indonesia

²Department of Fisheries and Marine Sciences, University of Lampung, Bandar Lampung, Indonesia

*Correspondence: dyahthp@gmail.com Received 11-09-2019, Revised: 09-10-2019, Accepted: 22-10-2019 e-Published: 03-11-2019

Joruk is a fermented fish product originating from South Sumatra province, Indonesia. Traditionally it was made from freshwater fish (*Wader pari*: Cyprinidae), cooked rice, solar salt, and brown sugar. This research was aimed to acquire the concentration of cooked rice addition to produce a *joruk* product with the best properties including microbiological, chemical, and sensorial. Conducted by a Completely Randomized Design (CRD) in four replications. The treatment was the differences in concentration of cooked rice addition consisted of six different levels, i.e., 5, 10, 15, 20, 25, and 30% (w/w). Results showed that adding 20% of cooked rice provided the finest *joruk* product appearance. The characteristics were 4.92 pH, 6.92% lactic acid, 84.55 mgN/100 g volatile base, 63.30% moisture content, 4.25% ash content, 3.61% fat and 28.82% protein content, 8.61 log CFU/g of LAB, 13.74 log CFU/g total microbes, and 4.16 log CFU/g of yeast. The sensorial character of *joruk* was blackish-brown of color (7.3), slightly sour of smell (6.3), salty-runny taste (7.8), and a desirable non-intact appearance (6.5). It is suggested that *joruk* composed of high nutritional values and a potential health benefit.

Keywords: fermented fish, *Rasbora* sp., indigenous food, cooked rice addition

INTRODUCTION

Joruk is a typical fermented freshwater fish product originating from Ogan Komering Ulu Regency, South Sumatra province of Indonesia. The raw materials used in making *joruk* are freshwater fish (*Rasbora* sp: Cyprinidae), solar salt, cooked rice, and palm sugar. *Joruk* is consumed by frying or roasting to perform the distinctive aroma.

Traditionally, the addition of cooked rice in the processing of *joruk* not only provided a distinctive aroma but also stimulated the growth of beneficial microorganism such as lactic acid bacteria (LAB). One of the most advantages of fermented fish products is the high of LAB that would degrade the starch into simpler compounds and have benefits for body health (Koo et al., 2016). Most

LABs were probiotic microbes that play a beneficial role in improving digestive health. Koesoemawardhani et al., (2016) reported that *joruk* containing LABs of 10.19 log CFU/g.

The concentration of each raw material in making *joruk* was varied, and there is no processing standard. Traditionally, the amount of palm sugar added to the *joruk* fermentation process ranging from 20-50%, while cooked rice ranging from 10-20%. It causes an enormous diversities in the quality of *joruk* among the local producers. Sometimes they produced an undesirable juicy characteristic.

Koesoemawardani et al., (2016), optimized the addition of palm sugar and suggested 20% of palm sugar with the highest amount of LAB. High level of sugar (above 40%) would binding the free

water, consequently unable to be used by microbes and inhibit microbial growth (Muchtadi and Sugiyono, 2013). In addition, Nuraini et al., (2014) stated that the addition of 40% cooked rice could reduce the pH and TVB value of tilapia *bekasam*, but increase the lactic acid value, total sugar, and lysine.

There was no report concerning of cooked rice addition of the quality of *loruk*. Therefore, the purpose of this study is to determine a certain quantity of cooked rice additions for producing the finest features of the *loruk* product, including chemical, microbiological and sensorial properties.

MATERIALS AND METHODS

Research methodology

This research was conducted using a Completely Randomized Design (CRD). The treatment is the different concentration of cooked rice (N) which added to process *loruk* product were: 5% (N1), 10% (N2), 15% (N3), 20% (N4), 25% (N5), and 30% (N6) were replicate 4 times. The best treatment calculated based on De Garmo et al. (1984) by an effectiveness index (EI) among *loruk* parameters, then followed by evaluated for the total volatile base (TVB), protein, fat, ash, and sensorial properties of *loruk* product.

Joruk processing

Materials for making *loruk* consisted of freshwater fish locally called *wader pari* (*Rasbora* sp.), cooked rice (rice variety of IR 64), and palm sugar obtained from the traditional producer, and solar salt. The *wader pari* then washing and draining following by adding 10% of solar salt and 20% palm sugar. The cooked rice added according to the treatment and followed by stirring. Jars then sealed tightly and fermented for seven days.

Parameters analysis

The characteristic of the product performed by microbial and chemical analysis. The microbiological analysis was carried out by total microbes on the nutrient agar medium (INS, 2016), total yeast (INS, 2006), and lactic acid bacteria (LAB) on the MRSA medium (Fardiaz, 1992). The chemical properties including the pH value employing a pH meter according to INS (2006), lactic acid (AOAC, 2005), TVB (INS, 2009), moisture content (INS, 2006), protein and fat content (INS, 2006) and ash content determined according to Indonesian National

Standard methods (INS, 2009). The organoleptic analysis was carried out by hedonic analysis (INS, 2011). Four kinds of analysis (color, smell, taste, and appearance) were compared to a score-based test by eight semi-trained panelists for sensory evaluation.

Data analysis

The data analyzed by variance to determine whether there were differences among treatments. The similarity of data tested by the Bartlett test or by the Tuckey test and following by the Smallest Real Difference (SRD) test at the level of 5% (Hanafiah, 2011).

RESULTS AND DISCUSSION

Joruk was daily consumed by the Malay inhabitant in Southern region of Sumatra, Indonesia. Traditionally, freshwater fish were processed by fermenting and adding cooked rice for the production of *loruk*. The cooked rice addition is intended to provide a distinctive flavor and aroma. It is also a useful substrate for the growth of beneficial microbes during fermentation.

The different concentration of cooked rice showed a significant effect on the value of the moisture content of the *loruk*. The moisture content of *loruk* ranged from 62.68 to 63.70% (Table 1). The SRD test showed that the moisture content in the addition of cooked rice statistically different by 25% and 30%.

The higher the addition of cooked rice will affect the lower of moisture content because of the higher the concentration of sugar. It is indicated that the addition of more than 25% of cooked rice generated less free-water on *loruk* product. In line to Muchtadi and Sugiyono (2013), who claimed that sugar could be a preservative compound for free water binding.

In this study, the moisture content of *loruk* moisture ranged from 62.68 to 63.70%. Another study shows that the moisture content *loruk* were 56.23-64.14% (Koesoemawardani et al., 2016), while the other type of Indonesia fermented fish ranged from 72.14-74.81% (Kalista et al., 2012).

The pH value of *loruk* was inadequate pH range from 4.92 to 5.26 (Table 1). The pH value of *loruk* was in accordance with several fermented fish products, such as cyprinid *bekasam* were in range of 3.60–5.30 (Desniar et al., 2013); catfish *bekasam* were 3.99–4.41 (Kalista et al., 2012); red tilapia *bekasam* were 4.66 (Nuraini et al., 2014); and anchovies *rusip* product was 5.98 (Koesoemawardani et al., 2013).

Table 1; Moisture content, pH, and lactic acid of *joruk* after addition of cooked rice

Cooked rice addition	Characteristics		
	Moisture (%)	pH	Lactic acid (%)
N1 (5%)	63.70± 1.29 ^c	5.23± 0.14 ^b	6.48± 1.26 ^a
N2 (10%)	63.64± 1.03 ^c	5.18± 0.17 ^b	6.66± 1.39 ^a
N3 (15%)	63.55± 1.02 ^c	4.96± 0.17 ^a	6.70± 1.23 ^a
N4 (20%)	63.30± 1.29 ^{bc}	4.92± 0.16 ^a	6.92± 1.21 ^a
N5 (25%)	62.94± 1.22 ^{ab}	4.94± 0.11 ^a	6.75± 1.33 ^a
N6 (30%)	62.68± 1.18 ^a	5.26± 0.32 ^b	6.21± 1.36 ^a
SRD_{0.05}	0.45	0.21	1.32

Exp.: different character showed a statistically significant treatment on SRD 5%.

Table 2; Microbial profile of *joruk* after addition of cooked rice

Cooked rice addition	Microbial profile (log CFU/g)		
	Microbe	Yeast	LAB
N1 (5%)	13.77± 0.11 ^a	4.41± 0.03 ^{bc}	8.38± 0.28 ^a
N2 (10%)	13.58± 0.34 ^a	4.38± 0.05 ^{bc}	8.35± 0.17 ^a
N3 (15%)	13.48± 0.29 ^a	4.35± 0.17 ^{bc}	8.51± 0.17 ^a
N4 (20%)	13.74± 0.17 ^a	4.16± 0.11 ^a	8.61± 0.44 ^a
N5 (25%)	13.73± 0.11 ^a	4.28± 0.04 ^{ab}	8.53± 0.39 ^a
N6 (30%)	13.54± 0.21 ^a	4.44± 0.00 ^c	8.31± 0.31 ^a
LSD_{0.05}	0.39	0.13	0.34

Exp.: different character showed a statistically significant treatment on LSD 5%.

However, the pH values lower than shrimp fermented product, such *terasi* an Indonesian shrimp paste was 7-8 (Ali et al., 2019a), and shrimp sauce *kecalok* were 6.38-7.32 (Ali et al., 2019b).

The different concentration of cooked rice had a significant effect on the pH value during fermentation. SRD analysis showed the addition of 30% cooked rice indicated the highest pH value of the product. The addition of cooked rice affected microbial growth and acted as a carbon source for the microorganisms.

Carbohydrates hydrolyzed into simpler compounds such as dextrose, mannose, and sucrose which used by lactic acid bacteria as energy sources and produce acidic and volatile compounds. This process causes an acidic condition to the end product and lowering the pH values (Lestari et al., 2018).

Khalid (2011) stated that the optimum pH of lactic acid bacteria growth was 5.5-5.8. The addition of 30% of cooked rice can increase in pH value as 5.26. The increase in pH during fermentation due to the addition of sugar concentration as reported by Hariyadi et al., (2013), and Gianti and Evanuarini (2011).

Ariyanto et al., (2013) stated that a high initial sugar concentration would inhibit the fermentation rate because it inhibits the growth of

microorganisms. Otherwise, Kalista et al., (2012) reported that higher carbohydrates affected to lower LAB activity to produce lactic acid.

**Figure 1; Freshwater fish (*wader pari*, *Rasbora* sp.) and the *joruk* product**

The lactic acid values on *joruk* believed originating from LAB activity during fermentation degrading the carbohydrate sources from cooked rice and palm sugar. The lactic acid values ranged from 6.21–6.92%, but the variance analysis showed that the different addition of cooked rice had no significant effect on the lactic acid concentration of *joruk*.

It suspected of the low activity of some LAB in osmotic stress condition due to the high sugar content (Tsakalidou and Papadimitriou, 2011). It is the reason why the addition of different concentration of cooked rice did not affect the increase of lactic acid concentration. The seven

days fermented *loruk* product played an optimum LAB growth and affected the relatively similar concentration of lactic acid in all treatment.

We also reported the decreasing of pH value during the fermentation period from 6 to 4.92-5.26 at the end fermentation. It caused by the presence of organic acids, especially lactic acid, from the breakdown of carbohydrates by lactic acid bacteria followed by a decrease in pH value (Kalista et al., 2012; Setiarti et al., 2016).

The total number of microbes ranged from 13.48–13.77 log CFU/ g. Variance analysis showed that the addition of cooked rice did not provided significantly affect the total microbial value. It caused the processing of *loruk* occurred under spontaneous fermentation employing wide varies of microbial types such as LAB, mold, and yeast.

Koesoemawardani et al., (2016), reported that the addition of palm sugar concentration did not provide a significant effect on *loruk*. In line to *bekasam* product that the different carbohydrates sources did not significantly affect microbial growth (Desniar et al., 2013).

The total number of *loruk* LAB ranged from 8.31 – 8.61 log CFU/ g (Table 2). Some report revealed that LAB of fermented fish product ranging from 6.48 - 10.15 log CFU/ ml (Hadiyanti and Wikandari, 2013; Prianto and Djajati, 2018).

Based on the results of the study, the total value of lactic acid bacteria in the *loruk* is higher than the *bekasam* product. The microbes profile on *loruk* was dominated by LAB and considered high comparing to *bekasam* microbes. It caused the addition of both cooked rice and palm sugar during the making of *loruk* than only cooked rice addition on *bekasam* processing.

Rice and palm sugar were providing more carbon sources to fulfill the energy of microbes and will make an appropriate environment for the growth of lactic acid bacteria.

The different concentrations of cooked rice addition did not provide a different growth of LAB. It indicated that LABs were in the seven days fermented *loruk* had undergone a stationary phase and has been at the optimum growth. The addition of cooked rice would also cause too much-containing *loruk*, resulting in plasmolysis of the LABs. Maryana (2014), also noted that the addition of different carbon sources had no significant effect on the LAB, including *L. plantarum* and *L. acidophilic*. It was due to the excessive proportion of carbon sources and caused a hypertonic condition and lysis cells.

The total yeast was 4.16–4.44 log CFU/ g,

and lower than total yeast in the *rusip*, an anchovy fermented product that reported in the range of 4.57 to 5.94 CFU/ g (Koesoemawardani et al., 2013). Variance analysis showed that the addition of cooked rice had a significant effect on yeast growth (Table 2). Presumably, lactic acid bacteria have been in the stationary phase, where lactic acid bacteria will not grow and produce lactic acid, which affected pH. In this condition, the yeast will grow.

The total value of yeast is high, allegedly related to the moisture content of the *loruk*, which valued at 63.55-63.70% where in this condition, yeasts will quickly grow. Meanwhile, the total yeast value is low because the addition of rice causes the pH value to be lower so that the yeast cannot grow optimally.

The high proportion of carbon sources caused hypertonic condition and affected bacteria cells lysis (Maryana, 2014). The yeast grows because the LAB was lysis. Carbohydrates are the primary substrate for yeast growth, especially as a carbon source in the metabolic system (Lestari et al., 2018).

The fermentation of *loruk* occurred within spontaneous fermentation, which allowed the growth of all kinds of microbes due to appropriate nutrition and environments. The nutrition such as carbon, nitrogen, vitamins, and minerals are required for energy sources and compiles cell components, whereas the appropriate environments are pH, aw and temperature (Lestari et al., 2018).

Table 3; Characteristic of *loruk* treatment by adding 20% of cooked rice (N4)

Joruk characteristic	Values
TVB (mgN/100g)	84.55± 2.31
Protein (%)	28.82± 0.20
Fat (%)	3.61± 0.12
Ash (%)	26.97± 0.35
Moisture (%)	63.30± 1.29
pH	4.92± 0.16
Lactic acid (%)	6.92± 1.21
LAB (log CFU/g)	8.61± 0.44
Microbe (log CFU/g)	13.74± 0.17
Yeast/ yeast (log CFU/g)	4.16± 0.11
Organoleptic	
- Color	5.37± 1.7
- Smell	7.37± 1.8
- Taste	7.8± 0.9
- Appearance	6.5± 2.4

Determination of adequate concentration of cooked rice by chemical and microbiological

properties calculated according to an effectiveness index (EI) (data not shown) suggested that N4 treatment provide the best characteristics (Table 3). The product then analyzed for total volatile base (TVB) content, proximate, and sensory properties to display a complete quality profile of *loruk* product.

Furthermore, TVB value in this study was 84.55 mg N/ 100g. In accordant to Azizah and Wikandari (Azizah and Wikandari, 2014), was 97.30 mg.N/ 100g, as well as TVB of milkfish *bekasam* as 91.45 mg.N/ 100g (Hadiyanti and Wikandari, 2013).

The total volatile base provides the freshness of fish, the higher the TVB, the lower the level of freshness of the fish (Mueda, 2018). The increasing of TVB value due to microbial activity during fermentation, which degrading protein onto simpler and volatile compounds (Sari et al., 2018).

The protein content of *loruk* was 28.82%, fat content was 3.61% and ash of 4.25%. While the product of *bekasam* made from *seluang* fish has a protein content of 14.68%, fat and ash contents were 7.94% and 13.13% respectively (Mumtiah et al., 2014).

Sensorial analysis of *loruk* has a quality characteristic that is blackish brown with color intensity of 5.3, while the aroma has a slightly sour (7.3) and have a salty-runny taste (7.8). Aulia et al. (2018), stated salty-sour provide a distinctive flavor of the product because of the increase of amino acid content during fermentation period due to the breakdown of proteins. Furthermore, *loruk* has the appearance of non-intact (6.5) and provided whole fish appearance at the end product.

CONCLUSION

The addition 20% of cooked rice provided the best quality of *loruk* supported by chemical, microbiological, and sensorial properties. The chemical characteristics of pH 4.92, total lactic acid 6.92%, TVB 84.55 mg/ 100g, levels 63.30% moisture, 4.25% ash content, fat and protein content 3.61% and 28.82% respectively. Microbiological criteria were total LAB of 8.61 log CFU/ g, total microbes 13.74 log CFU/ g and yeast 4.16 log CFU/ g. The sensorial properties were blackish brown (7.3), slightly sour of smell (6.3), salty-runny of taste (7.8), and desirable a non-intact appearance (6.5) at the end of the product.

CONFLICT OF INTEREST

The authors declared that present study was

performed in absence of any conflict of interest.

ACKNOWLEDGEMENT

The author would like to thank the Indonesian Ministry of Research, Technology, and Higher Education for funding the research.

AUTHOR CONTRIBUTIONS

DK designed, conducted the research and write the manuscript; NH, SU, and MZA performed the research and analyzed data. MA designed experiments and reviewed the manuscript. All authors read and approved the final version.

Copyrights: © 2019 @ author (s).

This is an open access article distributed under the terms of the [Creative Commons Attribution License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

REFERENCES

- Ali M, Y Yuniarta, A Aulanni'am, and J Kusnadi. 2019a. Characteristic of *acan*: a traditional shrimp paste of Maduranese, Indonesia. IOP Conf. Ser.: Earth Environ. Sci. 230 012048. DOI: 10.1088/1755-1315/230/1/012048
- Ali M, A Kurniawan, and NM Noor. 2019b. Preliminary study on *kecalok*, an indigenous shrimp sauce from Indonesia. Squalen Bull. of Mar. and Fish. Postharvest and Biotech. 14(2): 93-101
- Ariyanto HD, Hidayatulloh F, and Murwono J. 2013. The effect of adding some sugar on alcohol productivity in the manufacture of wine made from rejected apples by using nopkor MZ.11. Jurnal Teknologi Kimia & Industri, 2(4): 226-232
- Association of Official Analytical Chemists [AOAC]. 2005. Official Methods of Analysis. (Washington DC: AOAC).
- Aulia H, Sri B, Gres M, and Andri JK. 2018. The effect of addition various concentrations of turmeric on the quality of catfish var. *Sangkuriang*. Jurnal Tadris Pendidikan Biologi, 9(1): 84-99.
- Azizah N and Wikandari PR. 2014. The using of banana peel flour in making *bekasam* with

- starter culture *Lactobacillus plantarum* B1765. UNESA Journal of Chemistry, 3(3): 138-145
- De Garmo EP, Sullivan WG and Canada JR. 1984. Engineering Economy. The 7th Edition. Macmillan Publishing Comp., New York. 264-266.
- Desniar, Rusmana I, Suwanto A, and Mubarik NR. 2013. Antimicrobial compounds produced by lactic acid bacteria originated from *bekasam*. Jurnal Akuatika, 3(2): 35-145.
- Fardiaz S. 1992. Mikrobiologi Pangan Edisi Pertama. (Jakarta: Gramedia) p283.
- Gianti I and Evanuarini H. 2011. The effect of sugar addition and time of storage on physical quality of fermented milk. Jurnal Ilmu dan Teknologi Hasil Ternak, 6(1): 28-33
- Hadiyanti MR and Wikandari PR. 2013. The effect of salt concentration and the addition of lactic acid bacteria *Lactobacillus plantarum* B1765 as a starter culture for the quality product of milkfish (*Chanos chanos*) *bekasam*. Jurnal Kimia, 2(3): 136–143.
- Hanafiah KA. 2011. *Rancangan percobaan teori dan aplikasi*. (Jakarta: Rajawali press) p238
- Haryadi, Nurliana, and Sugito. 2013. The value of pH and total of lactic acid bacteria of goat milk kefir after fermented by a various concentration of sugar and different length of incubation. Jurnal Medika Veterinaria, 7(1):4-7. DOI:10.21157/j.med.vet..v7i1.2905
- Indonesian National Standard. 2006. SNI Number: 01-2332.3; 01-2354.2; 01-2354.4; 01-2354.3; 01.2354. (Jakarta: BSN).
- Indonesian National Standard. 2009. SNI Number 2354.8:2009. (Jakarta: BSN).
- Indonesian National Standard. 2011. SNI Number 2346.2011. (Jakarta: BSN).
- Kalista A, Supriadi A, and Rachmawati SH. 2012. Catfish (*Clarias gariepinus*) *bekasam* with different sources of carbohydrates. Jurnal Fishtech, 1(1): 102–110.
- Khalid K. 2011. An overview of lactic acid bacteria. International Journal of Bioscience (IJB). 1(3): 1-13.
- Koesoemawardani D, Marniza, Rizal S and Sella N. 2016. Addition of palm sugar in *loruk* (fish fermentation). Proceeding of national seminar on Agricultural Technology Development, State Polytechnic of Lampung: 187-195
- Koesoemawardani D, Rizal S. and Tauhid M. 2013. Microbiological and chemical changes of *rusip* during fermentation. Jurnal Agritech, 33(3): 265-272.
- Koo OK, SJ Lee, KR Chung, DJ Jang, HJ Yang, and DY Kwon. 2016. Korean traditional fermented fish products: *jeotgal*. J Ethn Foods: 1-10. DOI: 10.1016/j.jef.2016.06.004
- Lestari S, Rinto, Huriah SB. 2018. The improvement of functional properties of the *bekasam* by using *Lactobacillus acidophilus* as a starter. JPHPI, 21(1): 179-187. DOI: 10.17844/jphpi.v21i1.21596.
- Maryana, D. 2014. The effect of sucrose addition on bacterial quantification in fermented whey product by using a combination of *Lactobacillus plantarum* and *Lactobacillus acidophilus*. (Makassar: Universitas Hasanuddin) p26.
- Muchtadi RT and Sugiyono. 2013. Prinsip Proses dan Teknologi Pangan. (Bogor: Alfabeta press. p320.
- Mueda RT. 2015. Physico-chemical and color characteristics of salt fermented fish sauce from anchovy *Stolephorus commersonii*. AACL Bioflux, 8(4): 565-572.
- Mumtiah ON, Kusdiyantini E, and Budiharjo A. 2014. Isolation and characterization of lactic acid bacteria, and proximate analysis of fermented *bekasam* of mujair (*Oreochromis mossambicus* Peters). Jurnal Biologi, 3(2): 20-30.
- Nuraini A, Ratna I and Laras R. 2014. The effect of different concentrations addition of cooked rice as carbohydrates sources and brown sugars to the quality *bekasam* made of red tilapia (*Oreochromis niloticus*). Jurnal Saintek Perikanan, 10(1): 19-25.
- Prianto AD, and Djajati S. 2018. Silver rasbora *bekasam* using various processed rice on microbiological and organoleptic properties. Jurnal Ilmu Pangan & Hasil Pertanian, 2(2): 107-115. DOI: 10.26877/jiphp.v2i2.3039
- Sari SI, Widiastuti I, and Lestari SD. 2018. Effects of different fermentation process on the physical and chemical characteristics of Siamese Gourami (*Trichogaster pectoralis*) fish sauce. Fishtech. 7(1): 36-48.
- Setiarto RHB, Nunuk W and Iwan S. 2016. Effect of lactic acid bacteria, fungi and yeast fermentation on the nutritional quality of sorghum flour. Agritech. 36(4): 440-449. DOI: 10.22146/agritech.16769
- Tsakalidou E and Papadimitriou K. 2011. Stress responses of lactic acid bacteria. Food Microbiology and Food Safety. (New York: Springer Science and Business Media). p530.

● 9% Overall Similarity

Top sources found in the following databases:

- 6% Internet database
- Crossref database
- 5% Submitted Works database
- 4% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Universitas Diponegoro on 2018-10-28	1%
	Submitted works	
2	Universitas Brawijaya on 2019-05-14	<1%
	Submitted works	
3	intp.fapet.ipb.ac.id	<1%
	Internet	
4	Nurul Meutia Agustiari, Sri Anggrahini, Andriati Ningrum. "Influences of...	<1%
	Crossref	
5	ecologicalprocesses.springeropen.com	<1%
	Internet	
6	journal.bio.unsoed.ac.id	<1%
	Internet	
7	garuda.ristekdikti.go.id	<1%
	Internet	
8	journal.upgris.ac.id	<1%
	Internet	

9	journals.usm.ac.id Internet	<1%
10	li01.tci-thaijo.org Internet	<1%
11	ojs.omniakuatika.net Internet	<1%
12	sdiarticle3.com Internet	<1%
13	EP Dianti, G Anjani, DN Afifah, N Rustanti, B Panunggal. " Nutrition Qual... Crossref	<1%
14	University of Lincoln on 2021-06-03 Submitted works	<1%
15	garuda.kemdikbud.go.id Internet	<1%
16	rjoas.com Internet	<1%
17	globalscientificjournal.com Internet	<1%
18	"Lactic Acid Bacteria", Springer Science and Business Media LLC, 2019 Crossref	<1%
19	Free University of Bolzano on 2022-09-13 Submitted works	<1%
20	Garcia Fontan, M.C.. "Microbiological characteristics of "androlla", a Sp... Crossref	<1%

- 21 **Mahrus Ali, Ardiansyah Kurniawan, Nuning Mahmudah Noor. "Prelimin...** <1%
Crossref
-
- 22 **jurnal.fp.unila.ac.id** <1%
Internet
-
- 23 **repo.unand.ac.id** <1%
Internet
-
- 24 **P D M H Karti, S J Manpaki, I Prihantoro. "The radio sensitivity of matur...** <1%
Crossref
-
- 25 **Universitas Brawijaya on 2019-07-31** <1%
Submitted works

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded sources
- Quoted material
- Manually excluded text blocks

EXCLUDED SOURCES

isisn.org	88%
Internet	
repository.lppm.unila.ac.id	88%
Internet	
smujo.id	6%
Internet	
biodiversitas.mipa.uns.ac.id	5%
Internet	
patpiconference.ftip.unpad.ac.id	5%
Internet	
Mahrus Ali, Yunianta, Aulanni'am, Joni Kusnadi. " Characteristic of : a traditi...	1%
Crossref	

EXCLUDED TEXT BLOCKS

Available online freely at www.isisn.orgBioscience ResearchPrint ISSN: 1811-9506...

Universitas Brawijaya on 2020-06-16

influence

biodiversitas.mipa.uns.ac.id

University of Lampung, Bandar Lampung, Indonesia

N. Herawati *, , I. F. Rivai, H. Koyama, S. Suzuki, Y. Lee. "Copper in Rice and in Soils According to Soil Type i...

Fisheries and Marine

iopscience.iop.org

2019, Revised: 09-10-2019, Accepted: 22-10-2019 e-Published

www.researchgate.net

The quality of joruk, an Indonesian freshwater fermented fish

biodiversitas.mipa.uns.ac.id

Bioscience Research, 2019 volume 16(4

Carla Vanti, Alice Panizzolo, Luca Turone, Andrew A Guccione, Francesco Saverio Violante, Paolo Pillastrini, ...

The quality of joruk, an Indonesian freshwater fermented fish

biodiversitas.mipa.uns.ac.id

Bioscience Research, 2019 volume 16(4

Carla Vanti, Alice Panizzolo, Luca Turone, Andrew A Guccione, Francesco Saverio Violante, Paolo Pillastrini, ...

The quality of joruk, an Indonesian freshwater fermented fish

biodiversitas.mipa.uns.ac.id

Bioscience Research, 2019 volume 16(4

Carla Vanti, Alice Panizzolo, Luca Turone, Andrew A Guccione, Francesco Saverio Violante, Paolo Pillastrini, ...

The quality of joruk, an Indonesian freshwater fermented fish

biodiversitas.mipa.uns.ac.id

designed experiments and reviewed themanuscript. All authors read and approved...

Universiti Sultan Zainal Abidin on 2020-05-31
