

Identification of Steroid Hormones and Fatty Acids during Gonadal Maturation of Spiny Lobster *Panulirus homarus*

Journal:	<i>Invertebrate Reproduction and Development</i>
Manuscript ID	TINV-2018-0053.R2
Manuscript Type:	Research Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Adiputra, Yudha; University of Lampung, Dept of Fisheries and Marine Science; Bogor Agricultural University, Department of Aquaculture Jr, Muhammad; Bogor Agricultural University., Department of Aquaculture Suprayudi, Muhammad; Bogor Agricultural University, Department of Aquaculture Manalu, Wasmen; Bogor Agricultural University, Department of Anatomy, Physiology, and Pharmacology Widanarni, Widanarni; Bogor Agricultural University, Department of Aquaculture
Keywords:	eyestalk ablation, fatty acid, spiny lobster, steroid hormone, thyroxine injection

SCHOLARONE™
Manuscripts

1
2
3 **1 Identification of Steroid Hormones and Fatty Acids during Gonadal Maturation of Spiny**
4
5 **2 Lobster *Panulirus homarus***
6

7
8
9
10 4 Yudha Trinoegraha Adiputra¹, Muhammad Zairin Jr.^{1*}, Muhammad Agus Suprayudi¹, Wasmen
11
12 5 Manalu², and Widanarni¹
13
14
15
16

17 7 ¹Graduate Program in Aquaculture Science, Department of Aquaculture, Faculty of Fisheries and
18
19 8 Marine Science, Bogor Agricultural University.

20
21 9 ²Department of Anatomy, Physiology, and Pharmacology, Faculty of Veterinary Medicine,
22
23 10 Bogor Agricultural University.
24
25

26
27
28 12 *Corresponding author: E-mail: zairinmz@live.com
29
30

31 13 Address: Agatis Street, Dramaga Campus, Bogor 1680 West Java, Indonesia.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 24
4
5 25
6
7
8 26 **Abstract**
9
10 27
11
12 28
13
14 29
15
16 30
17
18 31
19
20 32
21
22 33
23
24 34
25
26 35
27
28 36
29
30 37
31
32 38
33
34 39
35
36 40
37
38 41
39
40 42
41
42 43
43
44 44
45
46 45
47
48 46
49
50 47
51
52 48
53
54 49
55
56 50
57
58 51
59
60 52

Information on steroid hormones and fatty acids that play roles in lobster reproduction is still very limited although the data are indispensable to seed production in hatchery. The study was designed to identify steroid hormones and fatty acids during gonadal maturation of spiny lobster (*Panulirus homarus*). Pyrolysis GCMS was used to identify steroid hormones and fatty acids and compared their concentrations between sexes and treatments. Samples from 6 male and 18 female spiny lobsters were used with different treatments. Male spiny lobsters were treated with and without thyroxine injection. Female spiny lobsters were treated with and without eyestalk ablations during mature and immature gonad developments. Androst-5-en-17-one, 3β (androst) and estran-3-one, 17β (estran), two steroid hormones were identified at different levels of gonadal maturity of spiny lobsters. High concentrations of androst and estran were detected in the male spiny lobsters treated with thyroxine injections. **Estran has limited roles in stimulating gonadal development in male spiny lobsters.** Androst was also found in the gonad of mature female spiny lobsters, but with low concentrations. Estran showed high concentrations in female brood stock of spiny lobsters during oogenesis stages both without eyestalk ablation and with ablation of one or two eyestalks, except in the immature female gonads. It was found that stearic acid was the highest and dominant fatty acid in mature male spiny lobster. Stearic acid, oleic acid, palmitic acid, and caprylic acid were fatty acids with high concentrations in immature and mature female spiny lobsters. After 30 days in captivity, only stearic acid and oleic acid were found dominantly in **eyestalk ablated mature female** spiny lobsters.

Keywords: eyestalk ablation, fatty acid, spiny lobster, steroid hormone, thyroxine injection

1
2
3 47
4
5
6 48
7
8 49
910 **Introduction**

11
12 51 Reproduction in crustaceans is associated with endocrine process including hormonal and
13
14 52 nutritional functions (Subramoniam 2011). The roles of hormones and fatty acids in reproduction
15
16 53 are reflected in gonadal maturation technology that has been applied practically in hatcheries to
17
18 54 produce post larvae (Wilder et al. 2010). Eyestalk ablation is a conventional technique applied to
19
20 55 male and female spiny lobsters (*Panulirus homarus*) in order to accelerate gonad maturity,
21
22 56 increase feed intake, and its conversion related to reproduction, and more specifically to shorten
23
24 57 the duration from intermoult to premoult stages (Radhakrishnan & Vijayakumaran 1984;
25
26 58 Vijayakumaran & Radhakrishnan 1984; Fernandez & Radhakrishnan 2016).

27
28
29
30
31 59 Another gonadal maturation technology is the injection of sex-type steroid hormones of
32
33 60 vertebrates. Kirubakaran et al. (2005) reported two steroid hormones namely estradiol and
34
35 61 progesterone during ovarian maturation in spiny lobster. The steroid hormones play roles in
36
37 62 gonadal maturation, shorten maturation period, increase matured female, and prolong
38
39 63 fertilization in spiny lobsters (*P. interruptus*) (Nan et al. 2015).

40
41
42 64 Nandi (1967) explained that the quantitative and qualitative analysis of hormones and the release
43
44 65 of hormones from the gland were the activities that can link the important functions of steroid
45
46 66 hormone and nutrition in reproduction. Nutrients such as lipid, protein, carbohydrate, vitamin,
47
48 67 and minerals are important factors for maturation process of crustacean (Harrison 1990). Lipid
49
50 68 is a major energy source in marine invertebrate, involved in many important processes including
51
52 69 growth, molt, and reproduction (Yan et al. 2017). Nutrients requirements of brood stock can be

1
2
3 70 traced on the basis of changes in the composition of the corresponding materials when the
4
5 71 maturation process occurs in the crustacean gonads (Harrison 1990). Lack of information on
6
7
8 72 steroid hormone and fatty acids in spiny lobster gonadal maturation is due to the limited
9
10 73 utilization of this species in commercial mariculture, especially in hatchery. This study was
11
12 74 designed to identify the types and levels of steroid hormones and fatty acids during gonadal
13
14
15 75 maturation in spiny lobster.

16 17 76 **Materials and Methods**

18 19 77 *Spiny Lobster Origin and Rearing*

20
21 78 Immature and mature male and female spiny lobsters used in the experiment were obtained from
22
23
24 79 Krui, West Coast Residence, Lampung Province, Indonesia. Mature male spiny lobsters with
25
26 80 body weights range of 145-152 g, immature female spiny lobsters with body weights range of
27
28 81 126-134 g, and mature female spiny lobsters with body weights range of 148-174 g were handled
29
30
31 82 with sea sand, ice, and packed in paper box, and then transported by car about six hours to Main
32
33 83 Center for Marine Aquaculture or MCMA (Balai Besar Perikanan Budidaya Laut) in Pesawaran
34
35 84 Residence, Lampung Province, Indonesia. Permit clearance was obtained from Fish Quarantine
36
37
38 85 and Fish Quality Products Inspection Office of Lampung Province to use spiny lobsters for
39
40 86 research purposes. Four fiber illuminated plastic tanks each with the size of 200 cm x 100 cm x
41
42 87 50 cm were used for rearing of spiny lobsters. The tanks were filled with sea water with 40 cm
43
44 88 height, continuously changed, and aerated. Fresh squid or fish meats were used as feeds given
45
46
47 89 twice a day at 08 am and 05 pm with the level of 3-5 % of body weight. The experimental tanks
48
49 90 were siphoned and cleaned two times daily at 07 am and 04 pm. For a shelter of spiny lobster in
50
51 91 captivity, each tank was provided with 10 PVC pipes each with 6-inch diameter and 30 cm
52
53
54 92 length.

93 *Spiny Lobster Treatments*

94 Spiny lobster of 24 individuals consisting of 6 males and 18 females were used in the
95 experiment. Mature male spiny lobsters, each with body weight of 145-152 g, were identified or
96 marked individually by the number written on the paper covered with transparent water-proof
97 plastic and tighten into the tail by rubber band. All experimental male spiny lobsters were reared
98 in one tank. The observations for male spiny lobsters were divided into two parts.

99 Part I. The Effect of Thyroxine Injection on Gonadal Maturation of Male Spiny Lobsters

100 Two mature male spiny lobsters were used to study the effect of thyroxine hormone injection on
101 gonadal maturation. The dosages of thyroxine hormone used were 0 and 0.1 $\mu\text{g/g}$ body weight.

102 Two experimental male spiny lobsters were used in this test (each dose of thyroxine injection
103 only used one male spiny lobster). Levothyroxine sodium tablets (Thyrax N.V. Organon, Oss.
104 The Netherland) with thyroxine concentration of 100 μg in 1 tablet was used as a source of
105 thyroxine hormone in this study. Stock solution for thyroxine hormone injection was made by
106 crushing one tablet of Thyrax into powder form and dissolved in 100 μL of physiological NaCl
107 (0.9 g NaCl/100 μL sterile ddH₂O) as a stock solution and then a serial dilution was made to
108 obtain the required concentrations. Thyroxine hormone at a dose of 0 $\mu\text{g/g}$ was used 1 mL of
109 physiological NaCl for injection without thyroxine. Gonads from two male spiny lobsters
110 injected with thyroxine at doses of 0 and 0.1 $\mu\text{g/g}$ body weights, respectively, were collected
111 eight days after thyroxine injections.

112 Part II. The Effect of Thyroxine Injection on the Growth of Male Spiny Lobster

113 Four gonad mature male spiny lobsters were used to study the effect of thyroxine hormone
114 injection on the growth of male spiny lobster. The doses of thyroxine injection used were 0; 0.1;
115 0.2; and 0.5 $\mu\text{g/g}$ body weight. Thyroxine hormone stock solution and injection doses were

1
2
3 116 similar with those used in Part I above. Gonads from four male spiny lobsters injected with
4
5 117 thyroxine at doses of 0; 0.1; 0.2; and 0.5 $\mu\text{g/g}$ body weights were isolated eight days after
6
7
8 118 thyroxine injection.
9

10 119 Eighteen female spiny lobsters were used and the treatments were divided into three parts.

11 120 Part I. Effect of One Eyestalk Ablation on Immature Female Spiny Lobster

12
13
14
15 121 The first part consisted of six immature female spiny lobsters with body weight range of 122-134
16
17 122 g. The experimental female spiny lobsters were divided into two treatments *i.e.*, without eyestalk
18
19 123 ablation and with one eyestalk ablation. Each treatment consisted of three immature female spiny
20
21
22 124 lobsters.

23 24 125 Part II. Effect of One Eyestalk Ablation on Mature Female Spiny Lobster

25
26 126 The second part consisted of six mature spiny lobsters with body weight range of 148-152 g. The
27
28 127 mature female spiny lobsters were divided into two treatments *i.e.*, without eyestalk ablation and
29
30
31 128 with one eyestalk ablation. Each treatment consisted of three mature female spiny lobsters.

32 33 129 Part III. Effect of One and Two Eyestalk Ablation on Mature Female Spiny Lobster.

34
35 130 The mature spiny lobsters were treated with two stages of treatment of eyestalk ablation. This
36
37
38 131 part of experiment used six matured female spiny lobsters with body weight range of 163-174 g.
39
40 132 In the beginning of the experiment, or on day one of treatment, all of six experimental matured
41
42 133 female spiny lobsters were **eyestalk ablated**. After 30 days of one eyestalk ablation (on day 31st),
43
44
45 134 three of experimental mature female spiny lobsters previously ablated with one eyestalk were
46
47 135 continued with the ablation of the other eyestalk to obtain a two eyestalk ablation.

48
49 136 All matured female groups of experimental spiny lobsters were reared in three different tanks
50
51 137 and marking methods were used similar to those used in male spiny lobster. From the first day of
52
53
54 138 rearing or maintenance, the gonad was taken by following the changes in the three stages of

1
2
3 139 oogenesis. Practically, the condition and stage of oogenesis phases was confirmed by bending
4
5 140 the cephalothorax of female spiny lobster and highlighting it with a flask light. The changes in
6
7
8 141 the color of the gonads were mainly indicators used to determine the phases of oogenesis. After
9
10 142 the gonad reached the oogenesis phase, the spiny lobster was dissected without considering the
11
12 143 duration of maintenance to reach gonad maturity. In female spiny lobsters without eyestalk
13
14
15 144 ablation that were difficult to reach peak of oogenesis phases, feeding level was increased to 8%
16
17 145 of body weight/day and duration of rearing was extended to 14 days.

18
19 146 Ovaries were taken in each phase of three phases of oogenesis in three groups of experimental
20
21 147 female spiny lobsters. Three phases of oogenesis were primary vitellogenesis, secondary
22
23 148 vitellogenesis, and maturation that were confirmed by the histology of ovaries. Determination of
24
25 149 three phases of oogenesis and histological analysis were conducted by following the methods of
26
27 150 Subramoniam (2017a) and Shields and Boyd (2014) (Figure 1). Clove oil at a dose of 10 ml/L of
28
29 151 sea water was used as an anesthetic agent. Eyestalk ablation was conducted by cutting the
30
31 152 eyestalk by using a sterile scalpel and sanitized the cut eyestalk with iodine to protect from
32
33 153 pathogens infection. Treatment without eyestalk ablation required complete two eyes without
34
35 154 injuries. Gonad samples from male and female groups of experiments were kept in -80°C until
36
37 155 analysis with a pyrolysis GCMS.

38 156 ***Pyrolysis Gas Chromatography Mass Spectrometry Analysis***

39
40
41
42 157 Approximately 1 µg of gonad was weighed with microbalance and pyrolyzed at 400°C. The
43
44 158 products of pyrolysis were analyzed by GCMS. Gas Chromatography separations were carried
45
46 159 out with a GCMS-QP2010 (Shimadzu, Tokyo) plus instrument equipped with an rt x 5ms
47
48 160 capillary column (length of 60 cm, diameter of 0.25 mm, and thickness of 0.25µm).
49
50
51 161 Chromatographic separation was achieved by the following temperature program: 50°C for 5
52
53
54
55
56
57
58
59
60

1
2
3 162 min, then it was raised to 280°C for 50 min with pressure of 101 kPa. Helium was the carrier gas
4
5 163 at 0.85 mL/min with total flow of 46.5 mL/min and flow program mode at a linear velocity of
6
7 164 23.7 cm/sec, and purge flow of 3.0 mL/min. Split injection mode used was the ratio of 1:50.
8
9
10 165 Mass spectra was set with ion source at 200°C and interface at 280°C with solvent cut time of 1.5
11
12 166 min. Detector temperature used was 280°C and compound identification was based on
13
14 167 comparison of mass spectra with WILEY7 library database. Compounds with the highest
15
16 168 similarity (>90%) were identified as steroid hormones and fatty acids data were selected from
17
18
19 169 five ranks of compounds similarity available from library database.
20

21 **Data Analyses**

22
23
24 171 WILEY7 library database was able to identify structural and molecular weights of steroid
25
26 172 hormones and fatty acids. Due to the focus on steroid hormones and fatty acids identification,
27
28 173 two treatments on male spiny lobster used were with and without thyroxine injection. It means
29
30
31 174 the effects of thyroxine injection on growth and gonadal maturation and their doses of injection
32
33 175 were ignored. Identified steroid hormones that appeared more than one concentration were
34
35 176 presented as mean \pm SD.
36

37
38 177 Fatty acids identifications were conducted, due to their roles as main nutrients involved in
39
40 178 gonadal maturation. Further, comparisons of concentrations of those above fatty acids were
41
42 179 conducted between male and female gonads and between treatments. Fatty acids that appeared
43
44 180 more than one concentration were presented as mean \pm SD. In order to compare similar fatty
45
46 181 acids among treatments with exception to steroid hormones, concentration was grouped into
47
48
49 182 three categories *i.e.*, high (>10%), moderate (5-10%), and low (<5%).
50

51 **Results**

52
53
54 184 Pyrolysis GCMS showed the ability to identify steroid hormones and fatty acids during gonadal
55
56
57
58
59
60

185 maturation of spiny lobsters. Two steroid hormones found in this study were androst-5-en-17-
186 one,3 β (androst) and estran-3-one,17 β (estran). Pyrolysis GCMS also provided structures and
187 molecular weights of these two steroids hormones (Figure 2).

188 During gonadal maturation of male spiny lobsters, androst showed a high concentration in
189 control spiny lobsters without thyroxine injection ($2.00\pm 0.09\%$) compared to those injected with
190 thyroxine ($0.31\pm 0.00\%$). Estran was also found in gonadal mature male spiny lobsters without
191 thyroxine injection with concentrations of $0.74\pm 0.00\%$ and $1.00\pm 0.07\%$ in mature male spiny
192 lobsters with thyroxine injection (Table 1).

193 Concentrations of estran decreased in immature female spiny lobsters with one eyestalk ablation
194 and without eyestalk ablation in different stages of oogenesis. Without eyestalk ablation, the
195 immature female spiny lobsters showed an inconsistent decrease in estran concentrations within
196 different stages of oogenesis *i.e.*, primary vitellogenesis ($8.14\pm 0.50\%$), secondary vitellogenesis
197 ($3.74\pm 0.69\%$), and maturation ($4.88\pm 0.57\%$). However, immature female spiny lobsters with one
198 eyestalk ablation also showed a consistent decrease in the concentrations of estran in different
199 stages of oogenesis *i.e.*, primary vitellogenesis ($13.79\pm 2.59\%$), secondary vitellogenesis
200 ($4.61\pm 1.49\%$), and maturation stage ($3.26\pm 0.66\%$) (Table 1). Androst was not detected in
201 immature female spiny lobster.

202 Different from the immature female spiny lobster, concentrations of estran and androst were
203 detected in mature female spiny lobsters. Mature female spiny lobsters without eyestalk ablation
204 showed a pattern of increase in estran concentrations with the stage of oogenesis *i.e.*, primary
205 vitellogenesis ($0.13\pm 0.00\%$), secondary vitellogenesis ($5.30\pm 0.00\%$), and maturation stage
206 ($5.56\pm 0.81\%$). Mature female spiny lobster with one eyestalk ablation also showed an increase in
207 estran from $2.35\pm 0.66\%$ during primary vitellogenesis to $10.86\pm 3.10\%$ during maturation stage.

1
2
3 208 In this group of mature female spiny lobster with one eyestalk ablation, during secondary
4
5 209 vitellogenesis, estran and androst showed low concentrations *i.e.*, $1.37\pm 0.00\%$ and $0.37\pm 0.00\%$,
6
7
8 210 respectively (Table 1).

9
10 211 Matured female spiny lobsters with one and two eyestalk ablation after one month in captivity
11
12 212 and three stages of oogenesis showed variations of estran and androst concentrations. Mature
13
14 213 female spiny lobsters with one eyestalk ablation showed increases in the concentrations of estran
15
16 214 with the stages of oogenesis *i.e.*, primary vitellogenesis ($2.59\pm 0.00\%$), secondary vitellogenesis
17
18 215 ($3.93\pm 0.94\%$), and maturation stage ($5.17\pm 1.47\%$). Concentrations of androst were also detected
19
20 216 to increase from primary vitellogenesis ($0.84\pm 0.00\%$) to maturation stage ($1.11\pm 0.00\%$). Two-
21
22 217 **eyestalk ablated mature female spiny lobsters** showed the increase in estran concentrations
23
24 218 during oogenesis stages *i.e.*, primary vitellogenesis ($2.52\pm 0.00\%$), secondary vitellogenesis
25
26 219 ($7.07\pm 1.63\%$), and maturation stage ($7.85\pm 1.73\%$) (Table 1).

27
28
29 220 Variations of fatty acids concentrations were detected during gonadal maturation of spiny lobster
30
31 221 treated without thyroxine injection and with thyroxine injection. Results of experiment showed
32
33 222 that deposition of fatty acids in the gonads of mature male spiny lobsters was dominated by
34
35 223 stearic acid with a high concentration ($13.05\pm 2.37\%$; $10.84\pm 1.13\%$). Fatty acids with low
36
37 224 concentrations in this treatment were oleic acid ($2.83\pm 0.55\%$), pentadecanoic acid ($1.80\pm 0.00\%$),
38
39 225 cyclopropanepentanoic acid ($1.49\pm 0.00\%$), hexyl oleate ($1.35\pm 0.00\%$), and docosaenoic acid
40
41 226 ($1.83\pm 0.00\%$) (Table 2).

42
43
44 227 In immature female spiny lobsters without eyestalk ablation, fatty acids found in high
45
46 228 concentrations were stearic acid (16.47 ± 0.00 ; 22.32 ± 0.00 %), oleic acid ($25.96\pm 4.31\%$), and
47
48 229 palmitic acid ($11.33\pm 0.00\%$). Octadecatrienoic acid ($2.60\pm 0.00\%$), cyclopropanoic acid
49
50 230 ($2.59\pm 0.00\%$), prepenoic acid ($1.15\pm 0.00\%$), pentadecanoic acid ($0.80\pm 0.00\%$), nonadecanoid
51
52
53
54
55
56
57
58
59
60

1
2
3 231 acid ($1.15\pm 0.00\%$), and cyclopentaneundecanoid acid ($0.59\pm 0.00\%$) were identified with low
4
5 232 concentrations. Fatty acids with high concentrations in immature female spiny lobster with one
6
7 233 eyestalk ablation were caprylic acid ($24.08\pm 0.00\%$) and oleic acid (10.90 ± 0.85 ; $27.08\pm 8.74\%$).
8
9
10 234 Low concentration in immature female spiny lobster with one eyestalk ablation were docosenoic
11
12 235 acid ($4.89\pm 0.00\%$), hexanoic acid ($1.34\pm 0.00\%$), and undecenoic acid ($1.11\pm 0.00\%$) (Table 3).
13
14 236 In mature female spiny lobsters without eyestalk ablation, fatty acids deposition showed a great
15
16 237 variation. Fatty acids that were found with high and moderate concentrations during secondary
17
18 238 vitellogenesis and maturation stages were cyclopentaneundecanoid acid ($15.62\pm 0.00\%$), stearic
19
20 239 acid (9.63 ± 0.00 ; $16.82\pm 0.00\%$), and oleic acid ($12.35\pm 0.00\%$). Fatty acids with low
21
22 240 concentrations were octadecanoid acid ($4.50\pm 0.53\%$), prepenoic acid ($1.32\pm 0.00\%$), docosenoic
23
24 241 acid ($0.79\pm 0.00\%$), and butanoic acid ($1.53\pm 0.00\%$). Fatty acids with moderate and low
25
26 242 concentrations during primary vitellogenesis were palmitic acid ($5.78\pm 0.00\%$), oleic acid
27
28 243 ($4.98\pm 0.00\%$), ethyl palmitate ($3.32\pm 0.00\%$), and prepenoic acid ($0.07\pm 0.00\%$) (Table 3).
29
30
31 244 Mature female spiny lobsters with one eyestalk ablation showed a variety. In this group of
32
33 245 mature female spiny lobsters with one eyestalk ablation, concentrations of palmitic acid
34
35 246 (15.55 ± 0.00 ; 10.12 ± 0.00 ; $17.24\pm 0.00\%$), cyclopentaneundecanoid acid ($13.72\pm 0.00\%$), and oleic
36
37 247 acid ($11.39\pm 0.00\%$) were dominantly high. In this group of mature female spiny lobsters with
38
39 248 one eyestalk ablation, fatty acids showing moderate concentrations were cyclopropanepentanoic
40
41 249 acid ($5.40\pm 0.61\%$), eicosatrienoic acid ($4.20\pm 0.00\%$), and decanoic acid ($5.33\pm 0.00\%$). In this
42
43 250 group of mature female spiny lobsters with one eyestalk ablation, fatty acids with low
44
45 251 concentrations during secondary vitellogenesis and maturation stage were pentadecanoid acid
46
47 252 ($2.44\pm 0.83\%$), octadecanoid acid ($1.62\pm 0.00\%$), and prepenoic acid (0.38 ± 0.00 ; $0.44\pm 0.00\%$)
48
49
50
51
52
53
54 253 (Table 3).
55
56
57
58
59
60

1
2
3 254 After one month in captivity, mature female spiny lobsters with one and two eyestalk ablations
4
5 255 showed distinctly different fatty acids concentrations. In mature female spiny lobsters with one
6
7 256 and two eyestalk ablations, fatty acids showing high concentrations were stearic acid
8
9 257 (17.78 ± 0.00 ; 15.81 ± 0.00 ; 11.04 ± 0.00 ; $14.39\pm 0.00\%$) and oleic acid (16.74 ± 0.00 ; 18.80 ± 0.00 ;
10
11 258 $13.01\pm 0.00\%$) followed by a variety of moderate and low concentrations of fatty acids. In
12
13 259 contrast, fatty acids during maturation stage were dominated only by stearic acid 16.45 ± 0.00 ;
14
15 260 $11.95\pm 0.00\%$) and oleic acid (9.77 ± 0.00 ; $11.33\pm 0.00\%$) and no other fatty acid was detected
16
17 261 during maturation stage. In mature female spiny lobsters with one and two eyestalk ablations,
18
19 262 fatty acids showing moderate and low concentrations were cyclopropanepentanoic acid
20
21 263 (8.74 ± 0.00 ; 4.98 ± 1.23 ; 4.53 ± 0.75), octadecatrienoic acid (1.33 ± 0.00 ; 2.33 ± 0.00), pentadecanoic
22
23 264 acid (1.32 ± 0.00 ; 2.86 ± 0.00), and prepenoic acid (1.45 ± 0.00) (Table 3).
24
25
26
27

28 265 **Discussion**

29
30
31 266 Analytical platform of GCMS has advantages and disadvantages in determining metabolites
32
33 267 samples (Young & Alfaro 2018). The combination between platforms such as pyrolysis GCMS
34
35 268 may support more advantages that can be used to obtain broad analysis coverage including
36
37 269 hormones and fatty acids identifications with very limited sample numbers. The use of pyrolysis
38
39 270 GCMS is robust and convenient to produce the results rapidly. Pyrolysis GCMS has been applied
40
41 271 for exploring the natural active compounds from biological materials (Martinez-Balmori et al.
42
43 272 2013). The use of pyrolysis GCMS is able to find new products (Ghalibaf et al. 2017). This
44
45 273 study proved the benefits of pyrolysis GCMS in identification of steroid hormones and fatty
46
47 274 acids in different stages of gonad maturation and treatments.
48
49
50
51 275 Hormonal regulation of reproduction in spiny lobster is debatable due to the limited studies
52
53 276 previously conducted to improve spiny lobster hatchery production (Subramoniam 2017b).
54
55
56
57
58
59
60

1
2
3 277 Progesterone and 17β -estradiol are hormones recorded to affect vitellogenesis (Subramoniam &
4
5 278 Kirubakaran 2010). This present study found two new steroid hormones that had roles in gonadal
6
7 279 maturation and vitellogenesis of male and female spiny lobsters that were different from
8
9 280 vertebrate-type sex steroid hormones published so far. In general, steroids are available in two
10
11 281 main forms *i.e.*, the ecdysteroids and the vertebrate-type sex steroids (Subramoniam 2017b).
12
13 282 Distinctly, the difference in the roles of these two main steroids is ecdysteroid controls molting
14
15 283 process and gonadotrophic hormones and vertebrate-type sex hormones have main roles in
16
17 284 stimulating vitellogenesis (Subramoniam 2011).

18
19 285 The results of our present study showed that concentrations of androst and estran fluctuated with
20
21 286 sexes and stages of gonad maturity. This new steroid hormones have positive roles in gonadal
22
23 287 maturation of spiny lobster. Androst was abundantly found in male spiny lobster and estran
24
25 288 expression was much more frequent in female spiny lobster. Moreover, estran was detected in
26
27 289 male spiny lobster and androst was also detected in female spiny lobster. Both hormones were
28
29 290 detected in low concentrations and showed their limited roles in stimulating gonadal
30
31 291 development in different sexes of spiny lobsters.

32
33 292 Injections of thyroxine hormone in male spiny lobster affect androst and estran concentrations.
34
35 293 Concentrations of androst in male spiny lobsters with thyroxine injection were lower compared
36
37 294 to those without thyroxine injection. Distinctly, estran concentration was higher in male spiny
38
39 295 lobster with thyroxine injection compared to those without thyroxine injection. These results
40
41 296 showed that the roles of androst and estran were related to the thyroxine injection as was
42
43 297 indicated by the decreased androst concentration in the male gonads. These facts showed that
44
45 298 androst as a stimulator of gonad development in male spiny lobster was affected by thyroxine
46
47 299 hormone. The relation between thyroxine hormone and steroid hormone remain unclear.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 300 However, within these results it is shown that thyroxine suppressed steroid hormone synthesis. If
4
5 301 this condition continues, faster growth rate caused by thyroxine injection in male spiny lobster is
6
7 302 obviously necessary for maturation. Thus, main thyroxine function to support development in
8
9 303 vertebrate was found with comparable results. In decapoda, thyroxine hormone functions to
10
11 304 stimulate somatic growth in *Penaeus monodon* (Pillai et al. 1987) and *Macrobrachium*
12
13 305 *rosenbergii* (Roustain & Gaik 2006).

14
15
16
17 306 Moreover, the role of estran in the ovary is occurred in mature female compared to immature
18
19 307 female spiny lobsters. This condition was shown by the concentrations of estran in immature
20
21 308 female spiny lobsters even with one eyestalk ablation that were not able to increase compared to
22
23 309 the increase in its concentration in mature female spiny lobsters. The observation of
24
25 310 Quackenbush (1994) supported this result that small female lobsters at the first stage of gonad
26
27 311 maturity delayed their reproductions by allowing molting process to obtain larger sizes. During
28
29 312 vitellogenesis stages and treatments, mature female spiny lobsters showed fluctuated
30
31 313 concentrations of androst and estran. Yan et al. (2017) also showed that steroids concentrations
32
33 314 in the gonad during reproductive cycle tended to fluctuate due to biotic factors (maturation,
34
35 315 reproductive, food availability, and age) and abiotic factors (photoperiodicity, temperature, pH,
36
37 316 and dissolved oxygen). Since those biotic factors were relatively similar during the experiment,
38
39 317 in this study it was assumed that abiotic factors played roles in influencing the level of
40
41 318 vitellogenesis that further affected the expressions of androst and estran.

42
43
44
45 319 Androst and estran were detected in low concentrations in mature female with one eyestalk
46
47 320 ablation both during a short-time exposure and a long-time exposure (after one month in
48
49 321 captivity). In contrast, androst was not found in all vitellogenesis stages in mature female spiny
50
51 322 lobsters without eyestalk ablation and those with two eyestalk ablations after one month in
52
53
54
55
56
57
58
59
60

1
2
3 323 captivity. The practice of eyestalk ablation involves the losing of endogenous moult and
4
5 324 vitellogenesis inhibiting hormone (VIH) and lead to the induction of ovarian maturation (Kumar
6
7
8 325 et al.2018). It is shown that one eyestalk ablation also suppressed androst and estran during
9
10 326 vitellogenesis phases. Probably, the availability of vitellogenesis stimulating hormone (VSH)
11
12 327 reduced androst and estran into a basal level. These facts need to be clarified in the future study
13
14 328 to clarify the relation among androst, estran, VIH, and VSH. Quackenbush (1994), Subramoniam
15
16 329 and Kirubakaran (2010), and Subramoniam (2011) reported that, VIH and VSH showed
17
18 330 inhibitory and stimulatory effects on ovarian growth, vitellogenesis, and yolk regulations of
19
20 331 lobsters. In addition, it is also needed to confirm the importance of one eyestalk ablation in
21
22 332 mature spiny lobsters. In the present study it was found that female spiny lobsters without
23
24 333 eyestalk ablation could produce optimal maturation stage during treatment with high sterol diets.
25
26 334 Due to the pressure of animal right it is necessary to support the other option like feeding with
27
28 335 high cholesterol diets to replace eyestalk ablation.
29
30
31
32
33 336 In mature female spiny lobsters without eyestalk ablation and with two eyestalk ablation, estran
34
35 337 increased two folds followed by vitellogenesis stages. The peak concentration of estran was
36
37 338 reached during secondary vitellogenesis and maturation stage. It is shown that cholesterol from
38
39 339 feed is converted into estran by the hepatopancreas as a main storage and further transported to
40
41 340 the ovaries during vitellogenesis similar to those found by Fairs et al. (1989; 1990) in *Nephrops*
42
43 341 *norvegicus* and *Penaeus monodon*. It was shown that spiny lobster had steroidogenic ability to
44
45 342 produce variety of vertebrate-type steroid hormones (Kanazawa & Teshima 1971). Spiny lobster
46
47 343 is able to convert exogenous cholesterol obtained from the diet into sex steroids such as
48
49 344 progesterone, 17-hydroxyprogesterone, androstenedione, testosterone, and molting hormones
50
51
52
53
54 345 such as ecdysterone (Kanazawa & Teshima 1971; Burns et al.1984).
55
56
57
58
59
60

1
2
3 346 Fatty acids accumulations that were found during gonadal maturation of male and female spiny
4
5 347 lobsters had high variety. Feed such as squid and fish meat that were given to spiny lobster
6
7 348 during this study contained a lot of lipid. Metabolism of lipid in crustacean is located in the
8
9 349 hepatopancreas as the main site and the products were then accumulated in the muscle and the
10
11 350 ovaries (Swevers et al. 1991). Garofalaki et al. (2006) showed that fatty acid contents of the
12
13 351 muscle and hepatopancreas of *P. vulgaris* were low, but phospholipids were available abundantly
14
15 352 in these organs. Concentration of lipid in the hepatopancreas was found to decrease as a result of
16
17 353 its mobilization into the gonads during vitellogenesis and increasing lipid accumulation in the
18
19 354 ovaries at oogenesis phases (Iromo et al. 2014). These facts suggest that ovarian vitellogenin is
20
21 355 synthesized in the hepatopancreas, and later transported into the ovaries (Yan et al. 2017).
22
23
24 356 Fatty acids compounds were also found abundant and showed no single fatty acid that played a
25
26 357 dominant role in the gonadal maturation. Stearic acid is the most dominant fatty acid found in
27
28 358 mature male spiny lobster on two different treatments of thyroxine hormone. This means that
29
30 359 injection of thyroxine is not related to the elevation of stearic acid concentration in the gonads of
31
32 360 mature male spiny lobsters. Immature female spiny lobsters showed a more complex condition in
33
34 361 term of fatty acids accumulation. There are stearic acid, oleic acid, and palmitic acid in spiny
35
36 362 lobsters without eyestalk ablation. Moreover, only caprylic acid and oleic acid were dominant
37
38 363 fatty acids found in spiny lobster with one eyestalk ablation. Fatty acids depositions in the gonad
39
40 364 of female spiny lobsters with mature gonads had high varieties. Fatty acids that were also found
41
42 365 with high varieties were caprylic acid and oleic acid. Mature female spiny lobsters with one
43
44 366 eyestalk ablation had different fatty acids patterns where palmitic acid, cyclopentaneundecanoid
45
46 367 acid, and oleic acid were the most abundant in the ovaries. In addition, for a long time exposure
47
48 368 of eyestalk ablation, only two fatty acids class were found *i.e.*, stearic acid and oleic acid.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 369 This study showed a low variety of nutrients identified in immature female compared to mature
4
5 370 female spiny lobsters. Physiological changes caused by fatty acids variation are related to
6
7
8 371 hormonal changes in female brood stock (Yano 1998). It was found that fatty acids deposition in
9
10 372 the gonad was more diverse in female spiny lobsters without eyestalk ablation compared to
11
12 373 female spiny lobsters with one and two eyestalk ablation. Harrison (1990) supported the results
13
14 374 of this study that when maturation was induced by eyestalk ablation it would be able to
15
16 375 accelerate hormonal and metabolic change that would stimulate ovarian development to reach its
17
18 376 peak. It means there is a low variety in nutrient but with high concentration for specific purposes
19
20
21 377 such as egg yolk deposition.

22
23
24 378 Many new fatty acids class that were found in this study may improve our understanding and
25
26 379 knowledge to modify feed composition for brood stock reproduction in spiny lobster. In addition,
27
28 380 these results will fill the gap of information required for further development of spiny lobster
29
30
31 381 mariculture industry.

32 33 382 **Acknowledgments**

34
35 383 Doctoral Dissertation Research Grant from Ministry of Research, Technology, and Higher
36
37 384 Education of The Republic of Indonesia with contract No. 062/SP2H/LT/DRPM/2018 funded
38
39
40 385 this study.

41 42 386 **References**

43
44 387 Burns BG, Sangalang GB, Freeman HC, McMenemy M. 1984. Bioconversion of steroids by the
45
46 388 testes of the American lobster, *Homarus americanus*, *in vitro*. General and Comparative
47
48 389 Endocrinology, 54: 422-428.

49
50
51 390 Fairs NJ, Evershed RP, Quilan PT, Goad LJ. 1989. Detection of unconjugated and conjugated
52
53 391 steroids in the ovary, eggs, and haemolymph of the decapod crustacean *Nephrops*

- 1
2
3 392 *norvegicus*. General and Comparative Endocrinology, 4:199-208.
4
5 393 Fairs NJ, Quinlan PT, Goad LJ. 1990. Changes in ovarian unconjugated and conjugated steroid
6
7 394 titers during vitellogenesis in *Penaeus monodon*, Aquaculture 89: 83-99.
8
9
10 395 Fernandez F, Radhakhrisnan EV. 2016. Effect of bilateral eyestalk ablation on ovarian
11
12 396 development and moulting in early and late intermoult stages of female spiny lobster
13
14 397 *Panulirus homarus* (Linnaeus, 1758). Invertebrate Reproduction & Development, 60: 238-
15
16 398 242.
17
18
19 399 Garofalaki TF, Miniadis-Meimaroglou S, Sinanoglou VJ. 2006. Main phospholipids and their
20
21 400 fatty acid composition in muscle and cephalothorax of the edible Mediterranean crustacean
22
23 401 *Panulirus vulgaris* (spiny lobster). Chemistry and Physics of Lipids, 140: 55-65.
24
25
26 402 Ghalibaf A, Lehto J, Alén R. 2017. Fast pyrolysis of hot water-extracted and delignified silver
27
28 403 birch (*Betula pendula*) sawdust by Py-GC/MS. Journal of Analytical and Applied
29
30 404 Pyrolysis, 127: 17-22.
31
32
33 405 Harrison KE. 1990. The role of nutrition in maturation, reproduction and embryonic
34
35 406 development of decapod crustaceans: a review. Journal of Shellfish Research, 9: 1-28.
36
37
38 407 Iromo H, Zairin Jr. M, Suprayudi MA, Manalu W. 2014. Thyroxine distribution in the
39
40 408 hemolymph, hepatopancreas, ovary, sponge, and larvae of female mud crabs (*Scylla*
41
42 409 *serrata*) during ovarian maturation. Journal of Crustacean Biology, 34:760-763.
43
44
45 410 Kanazawa A, Teshima S. 1971. *In vivo* conversion of cholesterol to steroid hormones in the
46
47 411 spiny lobster, *Panulirus japonica*. Bulletin of The Japanese Society of Scientific Fisheries,
48
49 412 37: 891-898.
50
51
52 413 Kirubakaran R, Peter DM, Dharani G, Vinithkumar NV, Sreeraj G, Ravindran M. 2005. Changes
53
54 414 in vertebrate-type steroid and 5-hydroxytryptamine during ovarian recrudescence in the
55
56
57
58
59
60

- 1
2
3 415 Indian spiny lobster, *Panulirus homarus*. New Zealand Journal of Marine and Freshwater
4
5 416 Research, 39: 527-537.
6
7
8 417 Kumar V, Sinha AK, Romano N, Allen KM, Bowman BA, Thompson KR, Tidwell JH. 2018.
9
10 418 Metabolism and nutritive role of cholesterol in the growth, gonadal development, and
11
12 419 reproduction of crustaceans. Reviews in Fisheries and Aquaculture, 26: 254-273.
13
14
15 420 Martinez-Balmori D, Olivares F, Spaccini R, Aguiar KP, M, Araújo MF, Aguiar NO, Guridi F,
16
17 421 Canellas LP. 2013. Molecular characteristics of vermicompost and their relation to
18
19 422 preservation of inoculated nitrogen-fixing bacteria. Journal of Analytical and Applied
20
21 423 Pyrolysis, 104: 540-550.
22
23
24 424 Nan FH, Wu YS, Chang NC. 2015. The effect of steroid hormone feeds on the reproductive
25
26 425 biology of the spiny lobster, *Panulirus interruptus* (J.W. Randall, 1840) (Decapoda,
27
28 426 Palinura). Crustaceana, 88: 1367-1386.
29
30
31 427 Nandi J. 1967. Comparative endocrinology of steroid hormones in vertebrates. American
32
33 428 Zoologist, 7: 115-133.
34
35
36 429 Pillai SM, Verghese PU, Ravichandran P, Roy AK. 1987. Effect of thyroxine on growth and
37
38 430 moulting in *Penaeus monodon* Fabricius. Indian Journal of Animal Science, 57: 241-245.
39
40
41 431 Quackenbush LS. 1994. Lobster reproduction: a review. Crustaceana, 67: 82-94.
42
43
44 432 Radhakrishnan EV, Vijayakumaran M. 1984. Effect of eyestalk ablation in the spiny lobster
45
46 433 *Panulirus homarus* (Linnaeus): 3. On gonadal maturity. Indian Journal of Fisheries, 31:
47
48 434 209-216.
49
50
51 435 Roustaian P, Gaik LA. 2006. Effect of thyroxine immersion on larval survival, growth and
52
53 436 postlarvae production of freshwater prawn, *Macrobrachium rosenbergii* (de Man).
54
55 437 Aquaculture Research, 37:1378-1380.
56
57
58
59
60

- 1
2
3 438 Shields JD, Boyd R. 2014. Atlas of Lobster Anatomy and Histology. Virginia Institute of Marine
4
5 439 Science, Virginia.
6
7
8 440 Subramoniam T, Kirubakaran R. 2010. Endocrine regulation of vitellogenesis in lobsters. Journal
9
10 441 Marine Biology Association of India, 52: 229-236.
11
12 442 Subramoniam T. 2011. Mechanism and control of vitellogenesis in crustaceans. Fisheries
13
14 443 Science, 77: 1-21.
15
16
17 444 Subramoniam T. 2017a. Sexual Biology and Reproduction in Crustaceans. Academic Press,
18
19 445 London.
20
21
22 446 Subramoniam T. 2017b. Steroidal control of vitellogenesis in crustacean: a new understanding
23
24 447 for improving shrimp hatchery production. Proceeding Indian National. Science Academy,
25
26 448 83: 595-610.
27
28
29 449 Swevers L, Lambert JGD, De Loof A. 1991. Metabolism of vertebrate-type steroids by tissues of
30
31 450 three crustacean species. Comparative Biochemistry and Physiology, 99B: 35-41.
32
33
34 451 Vijayakumaran M, Radhakrishnan EV. 1984. Effect of eyestalk ablation in the spiny lobster
35
36 452 *Panulirus homarus* (Linnaeus): 2. On food intake and conversion. Indian Journal of
37
38 453 Fisheries, 31: 148-155.
39
40
41 454 Wilder MN, Okumura T, Tsutsui N. 2010. Reproductive mechanism in crustacea focusing on
42
43 455 selected prawn species: vitellogenin structure, processing and synthetic control. Aqua-
44
45 456 Bioscience Monographs, 3: 73-110.
46
47
48 457 Yan H, Xue M, Liu H, Wang L, Liu Q, Jiang L. 2017. Energy reserves and gonad steroid levels
49
50 458 during the reproductive cycle of Japanese mantis shrimp *Oratosquilla oratoria* De Haan,
51
52 459 1844 (Stomatopoda: Squillidae) in Pikou Bay, Dalian, China. Journal of Crustacean
53
54 460 Biology, 37: 99-108.
55
56
57
58
59
60

1
2
3 461 Yano I. 1998. Hormonal control of vitellogenesis in penaeid shrimp. In: Flagel TW, editor.
4
5 462 Advances in Shrimp Biotechnology. National Center for Genetic Engineering and
6
7 463 Biotechnology, Thailand, pp. 29-31.

8
9
10 464 Young T, Alfaro AC. 2018. Metabolomic strategies for aquaculture research: a primer. Reviews
11
12 465 in Aquaculture, 10: 26-56.

13
14
15 466

16 17 467 **Figure Legends**

18
19 468 Fig. 1 Oogenesis stages of spiny lobster (*Panulirus homarus*) ovaries used in this study.

20
21 469 pfc: primary follicular cell; fcon: fibrous connective tissue; sfc: secondary follicular
22
23 470 cell; pre: previtellogenic oocyte; lg: lipid globule; vova: vitellogenic oocyte; cg:
24
25 471 cortical granule.

26
27
28 472 Fig. 2 Structures and molecular weights of androst-5-en-17-one,3 β (androst) and estran-3-
29
30 473 one,17 β (estran). Two steroid hormones in immature and mature gonads of male

31
32
33 474
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60





Fig. 1 Oogenesis stages of spiny lobster (*Panulirus homarus*) ovaries used in this study. pfc: primary follicular cell; fcon: fibrous connective tissue; sfc: secondary follicular cell; pre: previtellogenic oocyte; lg: lipid globule; vova: vitellogenic oocyte; cg: cortical granule.

338x190mm (96 x 96 DPI)



25 Fig. 2 Structures and molecular weights of androst-5-en-17-one,3 β (androst) and estran-3-
 26 one,17 β (estran). Two steroid hormones in immature and mature gonads of male
 27 and female spiny lobsters (*Panulirus homarus*) found in this study.

28 338x190mm (96 x 96 DPI)

Table 1. Two steroid hormones androst-5-en-17-one,3 β (androst) and estran-3-one,17 β (estran) concentration (%) during gonadal maturation of spiny lobster (*Panulirus homarus*) in different treatments.

No	Sexes	Treatments	Concentration of androst (%)			Concentration of estran (%)		
			Concentration of androst (%) during oogenesis phases			Concentration of estran (%) during oogenesis phases		
			Primary vitellogenesis	Secondary vitellogenesis	Maturation	Primary vitellogenesis	Secondary vitellogenesis	Maturation
1	Male	Without thyroxine injection				2.00 \pm 0.09		0.74 \pm 0.00
2	Male	Thyroxine injection				0.31 \pm 0.00		1.00 \pm 0.07
3	Female	Immature and without eyestalk ablation	Not detected	Not detected	Not detected	8.14 \pm 1.50	3.74 \pm 0.69	4.88 \pm 0.57
4	Female	Immature and one eyestalk ablation	Not detected	Not detected	Not detected	13.79 \pm 2.59	4.61 \pm 1.49	3.26 \pm 0.66
5	Female	Mature and without eyestalk ablation	Not detected	Not detected	Not detected	0.13 \pm 0.00	5.30 \pm 0.00	5.56 \pm 0.81

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

6	Female	Mature and one eyestalk ablation	Not detected	0.37 ± 0.00	Not detected	2.35 ± 0.66	1.37 ± 0.00	10.86 ± 3.10
7	Female	Mature and one eyestalk ablation after 30 days in captivity	0.84 ± 0.00	Not detected	1.11 ± 0.00	2.59 ± 0.68	3.93 ± 0.94	5.17 ± 1.47
8	Female	Mature and two eyestalk ablation after 30 days in captivity	Not detected	Not detected	Not detected	2.52 ± 0.00	7.07 ± 1.63	7.85 ± 1.73

Table 2. Identified fatty acids concentration (%) during gonadal maturation of male spiny lobster (*Panulirus homarus*).

No	Treatments (number of sample)	Body weight (g)	Maturity	Identified fatty acids	Concentration (%)
1	Without thyroxine injection (2)	145;150	Mature	Octadecanoid acid/ Stearic acid	13.05±2.37
				Pentadecanoid acid,14-methyl-,methyl ester	1.80±0.00
				Cyclopropanepentanoic acid,2-undercyl-,methyl ester, trans	1.49±0.00
				9-Octadecenoid acid, hexyl ester/Hexyl oleate	1.35±0.00
				13-Docosenoic acid, methyl ester	1.03±0.00
2	Thyroxine injection (4)	148-152	Mature	Octadecanoid acid/Stearic acid	10.84±1.13
				9-Octadecenoid acid/ Oleic acid	2.83±0.55

Table 3. Identified fatty acids concentration (%) during gonadal maturation of female spiny lobster (*Panulirus homarus*).

No	Treatments (number of sample)	Body weight (g)	Maturity	Identified fatty acids	Concentration (%)
1	Without eyestalk ablation during primary vitellogenesis (1)	128	Immature	Octadecanoid acid/Stearic acid	22.32±0.00
				9,12,15-Octadecatrienoic acid,methyl ester	2.60±0.00
				Cyclopropanepentanoic acid,2-undecyl-,methyl ester, trans	2.59±0.00
				2-Prepenoic acid,2-methyl,2-(dimethylamino) ethyl ester	1.15±0.00
				Pentadecanoid acid,4,6,10,14-tetramethyl-,methy ester	0.80±0.00
2	Without eyestalk ablation during secondary vitellogenesis (1)	126	Immature	9-Octadecenoid acid/ Oleic acid	25.96±4.31
				Octadecanoid acid/Stearic acid	16.47±0.00
				Nonadecanoid acid, ethyl ester	1.15±0.00
				Cyclopentaneundecanoid acid	0.59±0.00
3	Without eyestalk ablation during maturation (1)	134	Immature	Hexadecanoic acid/ Palmitic acid	11.33±0.00
				9-Octadecenoid acid/ Oleic acid	4.02±0.00
4	One eyestalk ablation	132	Immature	13-Docosenoic acid,methyl ester	4.89±0.00

1							
2							
3							
4							
5	during primary						
6							
7	vitellogenesis (1)						
8							
9	5	One eyestalk ablation	125	Immature	Octanoid acid/ Caprylic acid	24.08±0.00	
10							
11		during secondary			9-Octadecenoid acid/ Oleic acid	10.90±0.85	
12							
13		vitellogenesis (1)			Hexanoic acid	1.34±0.00	
14							
15		6	One eyestalk ablation	130	Immature	9-Octadecenoid acid/ Oleic acid	27.08±8.74
16							
17		during maturation (1)			2-Undecenoic acid	1.11±0.00	
18							
19		7	Without eyestalk ablation	148	Mature	Hexadecanoic acid/ Palmitic acid	5.78±0.00
20							
21		during primary			9-Octadecenoid acid/ Oleic acid	4.98±0.00	
22							
23		vitellogenesis (1)			Hexadecanoid acid, ethyl ester/Ethyl palmitate	3.32±0.00	
24							
25					2-Prepenoic acid,2-methyl,2-(dimethylamino) ethyl ester	0.07±0.00	
26							
27		8	Without eyestalk ablation	152	Mature	Cyclopentaneundecanoid acid	15.62±0.00
28							
29		during secondary			Octadecanoid acid/Stearic acid	9.63±0.00	
30							
31		vitellogenesis (1)			9,12,15-Octadecatrienoic acid,methyl ester	4.50±0.53	
32							
33					Cyclopropanepentanoic acid,2-undercyl-,methyl ester,	2.09±0.47	
34							
35					trans		
36							
37							
38							
39							
40							
41							
42							
43							
44							
45							
46							
47							

				2-Prepenoic acid,2-methyl,2-(dimethylamino) ethyl ester	1.32±0.00
				13-Docosenoic acid, methyl ester	0.79±0.00
9	Without eyestalk ablation	151	Mature	Octadecanoic acid/Stearic acid	16.28±0.00
	during maturation (1)			9-Octadecenoid acid/ Oleic acid	12.35±0.00
				Butanoic acid,2((trifluoroacthyl)amino)-,butyl ester	1.53±0.00
				2-Prepenoic acid,2-methyl,2-(dimethylamino) ethyl ester	0.44±0.00
10	One eyestalk ablation	152	Mature	Hexadecanoic acid/ Palmitic acid	15.55±0.00
	during primary			Cyclopentaneundecanoic acid	13.72±0.00
	vitellogenesis (1)			Cyclopropanepentanoic acid,2-undercyl-,methyl ester,	5.40±0.61
				trans	
11	One eyestalk ablation	150	Mature	9-Octadecenoid acid/ Oleic acid	11.39±0.00
	during secondary			Hexadecanoic acid/ Palmitic acid	10.12±0.00
	vitellogenesis (1)			8,11,14-Eicosatrienoic acid, methyl ester	4.20±0.00
				Pentadecanoic acid,14-methyl-,methyl ester	2.44±0.83
				11-Octadecanoic acid, methy ester	1.62±0.00
				2-Prepenoic acid,2-methyl,2-(dimethylamino) ethyl ester	0.38±0.00

1						
2						
3						
4						
5	12	One eyestalk ablation	149	Mature	Hexadecanoic acid/ Palmitic acid	17.24±0.00
6						
7		during maturation (1)			Decanoic acid, 1-methylethyl ester	5.33±0.00
8						
9					6,9,12-Octadecatrienoic acid, methyl ester	2.92±0.00
10						
11					9-Octadecen-12-ynoic acid, methyl ester	2.06±0.00
12						
13					2-Prepenoic acid,2-methyl-,2-(dimethylamino)ethyl ester	1.65±0.28
14						
15					9-Octadecanoic acid, methy ester	1.58±0.00
16						
17					Cyclopropanepentanoic acid,2-undercyl-,methyl ester,	1.25±0.00
18						
19					trans	
20						
21					Pentadecanoid acid,14-methyl-,methyl ester	0.45±0.00
22						
23						
24	13	One eyestalk ablation	169	Mature	Octadecanoid acid/Stearic acid	17.78±0.00
25						
26		during primary			9-Octadecenoid acid/ Oleic acid	16.74±0.00
27						
28		vitellogenesis after 30 days			9,12,15-Octadecatrienoic acid,methyl ester	1.69±0.00
29						
30		in captivity (1)			Pentadecanoid acid,14-methyl-,methyl ester	1.33±0.00
31						
32					Phosphorothioc acid,O,O-diisopropyl ester, S-ester-N	1.11±0.00
33						
34					2-Prepenoic acid,2-methyl-,2-(dimethylamino)ethyl ester	0.90±0.00
35						
36					Octadecanoic acid, phenylmethyl ester	0.67±0.00
37						
38						
39						
40						
41						
42						
43						
44						
45						
46						
47						

1						
2						
3						
4						
5	14	One eyestalk ablation	174	Mature	9-Octadecenoid acid/ Oleic acid	18.80±0.00
6		during secondary			Octadecanoid acid/Stearic acid	15.81±0.00
7		vitellogenesis after 30 days			Cyclopropanepentanoic acid,2-undercyl-,methyl ester,	4.98±1.23
8		in captivity (1)			trans	
9					9,12,15-Octadecatrienoic acid,methyl ester	1.60±0.00
10					2-Prepenoic acid,2-methyl-,2-(dimethylamino)ethyl ester	1.33±0.00
11					Pentadecanoid acid,14-methyl-,methyl ester	0.69±0.00
12						
13						
14						
15						
16						
17						
18						
19						
20	15	One eyestalk ablation	163	Mature	Octadecanoid acid/Stearic acid	16.45±0.00
21		during maturation after 30			9-Octadecenoid acid/ Oleic acid	9.77±0.00
22		days in captivity (1)				
23						
24						
25						
26						
27	16	Two eyestalk ablation	172	Mature	Octadecanoid acid/Stearic acid	11.04±0.00
28		during primary			Cyclopentaneundecanoid acid	8.74±0.00
29		vitellogenesis after 30 days			Cyclopropanepentanoic acid,2-undercyl-,methyl ester,	4.53±0.75
30		in captivity (1)			trans	
31					12,15-Octadecadiynoic acid, methyl ester	2.33±0.00
32					2-Prepenoic acid,2-methyl-,2-(dimethylamino)ethyl ester	1.45±0.00
33						
34						
35						
36						
37						
38						
39						
40						
41						
42						
43						
44						
45						
46						
47						

1						
2						
3						
4						
5				Pentadecanoid acid,14-methyl-,methyl ester	1.32±0.00	
6						
7	17	Two eyestalk ablation	170	Mature	Octadecanoid acid/Stearic acid	14.39±0.00
8		during secondary			9-Octadecenoid acid/ Oleic acid	13.01±0.00
9		vitellogenesis after 30 days			Pentadecanoid acid,14-methyl-,methyl ester	2.86±0.00
10		in captivity (1)				
11						
12						
13						
14						
15	18	Two eyestalk ablation	168	Mature	Octadecanoid acid/Stearic acid	11.95±0.00
16		during maturation after 30			9-Octadecenoid acid/ Oleic acid	11.33±0.00
17		days in captivity (1)				
18						
19						
20						
21						
22						
23						
24						
25						
26						
27						
28						
29						
30						
31						
32						
33						
34						
35						
36						
37						
38						
39						
40						
41						
42						
43						
44						
45						
46						
47						