PAPER NAME Proceeding International Conf. on Cassa va 2017.pdf	AUTHOR purnomo
WORD COUNT 4105 Words	CHARACTER COUNT 21945 Characters
PAGE COUNT	FILE SIZE
14 Pages	2.4MB
	REPORT DATE
Mar 28, 2023 4:00 PM GMT+7	Mar 28, 2023 4:01 PM GMT+7

• 6% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 4% Internet database
- Crossref database
- 4% Submitted Works database

• Excluded from Similarity Report

- Bibliographic material
- Manually excluded sources

- 5% Publications database
- Crossref Posted Content database
- Quoted material
- Manually excluded text blocks



PROCEEDING INTERNATIONAL CONFERENCE ON CASSAVA (Sustainable Management of Renewable Resources in Tropics) Bandar Lampung, November 23rd - 24th 2017





PROCEEDING INTERNATIONAL CONFERENCE ON CASSAVA (Sustainable Management of Renewable Resources in Tropics) Bandar Lampung, November 23rd – 24th, 2017



Editors:

Erwin Yuliadi M. Syamsoel Hadi Kukuh Setiawan

Published as Collaboration with:

LPPM University of Lampung – Sungai Budi Group

December 2018

Title

Proceeding International Conference on Cassava (Sustainable Management on Renewable Resources in Tropics)

Editors

Erwin Yuliadi M. Syamsoel Hadi Kukuh Setiawan

Cover Designer

Rafika Restiningtias

Layout

Restu Paresta Putri Ulva Priska

Corrector

Ridho Akbar

Secretariat Office

Centre of Cassava Research and Development, University of Lampung JI. Soemantri Brojonegoro No. 1 Bandar Lampung 35145

All rights reserved. No part of this proceeding may be reproduced, in any form or by any means, without permission in writing from the publisher

Printed by LPPM of University of Lampung

Copyright @ 2018



CONTENTS

	Title	Page
1	Dietary Fibre Production as Co-Product of Tapioca Industry Bambang Triwiyono, Aton Yulianto, Sabirin, Budi Kusarpoko, Yanuar Sigit Pramana, and Novi Kuswardhani	1
2	The Effect of <i>ZincMicro</i> Nutrient on Root Fungi Disease Development of Cassava (<i>Manihot Esculenta</i> Crantz) in Sulusuban, Sub-District Anak Tuha, Lampung Tengah <i>Efri, R. Suharjo, M.S. Hadi, K. Setiawan, and M. Saifudin</i>	7
3	The Effect of Ethrel Treatment on The Growth and Production of Two Varieties of Cassava Plant (Manihot Esculenta Crantz) Ardian, Kukuh Setiawan, A.S. Tumanggor, and Erwin Yuliadi	15
4	Identifying Chemical Compound in Ceara Rubber Skin Which Is Potential To Be Natural Anti-Microbe By Using Gas Chromatography-Mass Spectrometry (GC- MS) <i>Bigi Undadraja and Dewi Sartika</i> .	24
5	Correlation of Whiteness and Protein Content of Modified Cassava Flour (Mocaf) Made From Different Varieties of Cassava Sri Lestari and Erliana Novitasari	28
6	Farmers Cassava Motivation To Change Partnership (Case of Farmers Sugarcane and Cassava Partnership) <i>Tubagus Hasanuddin and Via Agiesta</i>	33
7	Growth and Yield of Cassava (Manihot Esculenta Crantz) under Intercropping with Several Genotypes of Sorghum (Sorghum Bicolor [L.] Moench) M. Syamsoel Hadi, Restu Paresta, Muhammad Kamal, and Kukuh Setiawan	36
8	Attack Intensity and Its Population of Major Pests in Cassava (<i>Manihot Esculenta</i> Crantz) By Application of "ZincMicro" Fertilizer <i>Muhammad Rizki, Purnomo, M. Syamsoel Hadi, and I Gede Swibawa</i>	44
9	Using Ceara Rubber as Natural Anti-Microbe in Reducing Contamination of <i>Staphylococcus aureus, Salmonella sp., Vibrio sp.</i> and <i>Escherichia coli</i> in Mackerel Tuna Fish (<i>Euthynnus affinis</i>) <i>Widia Rini Hartari Dewi Sartika and Suharyono AS</i>	54
10	The Role of Supporting Environmental Factors on The Use of Cyber Extension By Farmers of Food Crops and Horticulture in Lampung Province Dame Trully Gultom and Sumaryo Gitosaputro	60
11	Occurrence of Mosquitocidal <i>Bacillus Thuringiensis</i> Serovar <i>Entomocidus</i> in The Soil of Indonesia <i>Akhmad Rizali</i>	65

ATTACK INTENSITY AND ITS POPULATION OF MAJOR PESTS IN CASSAVA (Manihot esculenta Crantz) BY APPLICATION OF "ZINCMICRO" FERTILIZER

Muhammad Rizki, Purnomo, M. Syamsoel Hadi, and I Gede Swibawa Department of Agrotechnology, Faculty of Agriculture, University of Lampung JI. Prof. Soemantri Brojonegoro, No. 1, Bandar Lampung 35145 E-mail : muhammadrizki748@gmail.com

ABSTRACT

The major pests that attack the cassava plant are mealybug and mite. Mealybug (*Phenacoccus*) can reduce production by 30-80%, while mite (*Tetranychus*) can reduce production by more than 50% in cassava plantation. This study aims to (1) investigate the intensity of mealybug infestation and mite on cassava with application of "Zincmicro" fertilizer; (2) investigate the population of mealybug and mite on cassava with application of "Zincmicro" fertilizer. This research was conducted in Sulusuban, Seputih Agung, Lampung Tengah, starting from august 2016 until march 2017. This research use direct observation pethod by scoring the level of pest attack on cassava plantation. The experiment showed that: 1) Application of 20 kg ha⁻¹ "Zincmicro" was significantly lower than without application "Zincmicro". Increasing the dosage of "Zincmicro" until 40 kg ha⁻¹ did not differ significantly; 2) Although there was no significant different between with and without application "Zincmicro"; 3) The population of mealybug was more abundant in the upper leaves; meanwhile, the mite population was more abundant in the lower leaves. Application of "Zincmicro".

Keywords: cassava, mealybug, mites, "zincmicro" fertilizer

INTRODUCTION

¹³assava (*Manihot esculenta* Crantz) is a useful plant. Cassava crop is widely cultivated because it can adapt in areas where water is limited. Lampung is the largest cassava production center in Indonesia. Production of cassava in Lampung Province in 2015 was reportedly reached 7,387,084 tons (Badan Pusat Statistik, 2016). Currently cassava has been developed as an agro-industry commodity, such as tapioca flour products, fermentation industry, and various food industries. Increasing demand for cassava as raw material of industry, should be fulfilled by increasing production (Wardani, 2015).

Fertilization is used to supply nutrients lost due to harvesting. Fertilizer needs must be in accordance with the amount of nutrients lost during harvest. The nutrient needs equal to the amount of nutrients lost when harvested while farmers only fertilize macro nutrients while the micro nutrients are relatively never used. The fulfillment of nutrient needs can support the health of cassava plant so that it can produce optimal and resistant to pest and disease attack. Pest and disease can decrease production until crop failure. Mealybug is one of the major pests of cassava. In 2009 to 2010 Thailand experienced crop failure due to mealybug attack. In Lampung, attack of mealybug could potentially cause crop failure (Rojanaridpiched, 2017). This study aims to investigate the intensity and the population of mealybug and mite on cassava with application of "Zinemicro" fertilizer.

MATERIALS AND METHODS

The research was conducted in Sulusuban, Seputih Agung, Lampung Tengah. Observations on population of mite and mealybug were done at Plant Pest Laboratory, Faculty of Agriculture, University of Lampung. This research was conducted for 8 months starting from August 2016 until March 2017. The materials used were cassava (clone UJ-3), "Zincmicro" fertilizer with Zn, Mn, B, Cu, Co, Mo, macro nutrients in the form of urea , SP-36, KCl, herbicides, 1 kg-plastic bags, 70% alcohol, aquades, glycerine, elastic bands and labels. While the tool used is hoe, sprayer, brush, vial bottle, binocular stereo microscope, binocular compound microscope, camera, and stationery. This research used direct observation method that is by scoring the level of pest attack on cassava cultivation. Population observation and identification were performed in the laboratory. The experiment was conducted by giving "Zincmicro" fertilizer to 12 experimental units consisting of "Zincmicro" fertilizer 0 kg ha⁻¹, 20 kg ha⁻¹ and 40 kg ha⁻¹ with 4 replications. The experimental unit is a plot of 23 m x 30 m.

			69 m	
)
	ſ	7 1U1	P2U1	P3U1
120		P1U2	P2U2	P3U2
120 111		P1U3	P2U3	P3U3
		P1U4	P2U4	P3U4
		Figure 1.	Experiment	al Layout

Information :

- P1 = Treatment 1 ("Zincmicro" fertilizer 0 kg ha⁻¹)
- P2 = Treatment 2 ("Zincmicro" fertilizer 20 kg ha⁻¹)
- P3 = Treatment 3 ("Zincmicro" fertilizer 40 kg ha⁻¹)
- 8] = Repeat 1
- U2 = Repeat 2
- U3 = Repeat 3
- U4 = Repeat 4

Cassava variety grown was UJ-3 clones. The soil was done by plowing twice with the time interval between the first plow and the second was 14 days. Cassava¹⁰ as planted with a spacing of 80 cm x 60 cm and length of cuttings of 25 cm. Fertilization was applied on 4 Week After Planting (WAP) with macro fertilizer in the form of urea 100 kg ha⁻¹, SP-36 100 kg ha⁻¹, and KCl 100 kg ha⁻¹, and "Zincmicro" fertilizer 0 kg ha⁻¹, 20 kg ha⁻¹, and 40 kg ha⁻¹. The second fertilization was conducted on 12 WAP with macro fertilizer in the form of urea 100 kg ha⁻¹, and KCl 100 kg ha⁻¹. Fertilization was done by means in addition to plants with a depth of 10 cm. Herbicide spraying was performed with systemic herbicides at 12 WAP and 24 WAP. Harvesting was done at 28 WAP. Observations were made when cassava aged 18 WAP to 26 WAP. The observation time interval was 14 days. Sampling was done by determining 10 rows of plants as subsample according to the diagonal direction on each plot. Each subsample consists of 10 samples. Determination of subsample was done randomly systematically. The first row determination was done randomly, the next row would be adjusted to the spacing and area of the experimental plot.

Observation of mite and mealybug population was conducted on one sample plant selected from subsampel line, after which the top, middle, and bottom leaves were taken with each 3 leaves. Three top, middle, and lower cassava leaves were put into plastic bags and tied with rubber bands and labeled. Leaf samples were taken to the laboratory and counted the number of individual mites and mealybug using a setereo microscope at magnification 2.5 times - 4 times.

Observation of the intensity of the attack was done visually based on the symptoms of mite and mealybug attack. Observation of the intensity of mite and mealybug attack was performed on each sample that was in the subsample. On observation of the intensity of plant damage caused by the mites was given a score of 0-5 as in Table 1. While in mealybug was scored 0-2 as in Table 2. The damage intensity was calculated by using the formula:

$$I = \Sigma ni x vi/(N x V) x 100\%$$

I = Intensity of attack

ni umber of leaves in each score category

vi = Score category (0 to^4)

N = Number of <u>leaves</u> in one plant

V = Value of the aghest score (in this case 5)

Score	The magnitude of damage	Information
	(%)	
0	0	Healthy leaves (no spotting)
1	$0 < x \le 10$	There are early yellowish spots (about 10%) on some
		lower leaves and or middle leaves.
2	$10 < x \le 20$	Yellowish spots rather (11-20%) in the lower and
		middle leaves.
3	$20 < x \le 50$	Clear damage, many yellow spots (21-50%), few areas
		that do not experience necrotic (<20%), especially the
		lower and middle leaves rather shrink, a number of
		leaves turn yellow and fall off.
4	$50 < x \le 75$	Severe damage (51-75%) in the lower and middle
		leaves, population of mites abundant and found white
		threads like cobwebs.
5	x > 75	Total leaf loss, shoots of smaller plants, more white
		threads, and death of plants.

Table 1. Scoring intensity of crop damage due to mite attack based on Indiati (2012).

RESULTS AND DISCUSSION

Intensity of Mealybug Attack (*Phenacoccus***).** The intensity of severe mealybug attack is seen in cassava plants that were dwarfed and the leaves were curly and there was a collection of mealybugs that cover the lower surface of leaves and stems in ¹² de upper leaves of the cassava plant. The highest intensity of mealybug attack was found in plots without application of "Zincmicro" fertilizer. The intensity of attack on the plot with the application of "Zincmicro" fertilizer 20 kg ha⁻¹ and 40 kg ha⁻¹ was lower than the intensity of the attack on plots given "Zincmicro" fertilizer ²⁰ kg ha⁻¹ and 40 kg ha⁻¹ was not different (Figure 2).

The intensity of attack on plant age of 18 WAP to 26 WAP did not appear to be different. In plots without application of "Zincmicro" fertilizer (P1), the intensity of the mealybug attack was 68.5% to 77.6%. In plots with application of "Zincmicro" fertilizer 20 kg ha⁻¹ (P2), the intensity of mealybug attack was 31.9% to 43.1%. In plots with application of "Zincmicro" fertilizer 40 kg ha⁻¹ (P3), the intensity of mealybug attack was 26.9% to 38.9%. The addition of "Zincmicro" fertilizer can affect the intensity of the attack of mealybug so that it is suspected to affect the cassava plant resistance.

Score	Information	Picture
0	Healthy plants, There are no symptoms of white flea pest attack on the upper leaves.	
1	Symptoms of mild attacks. There are some parts of the leaves on the plant slightly curling.	
2	Symptoms of severe attacks. At the top of the plant to be dwarfed or "bunchy top", the upper leaves will curl and stem distortion	
	$\begin{bmatrix} 100 \\ 80 \\ - \\ 60 \\ 40 \\ 20 \\ - \\ 0 \end{bmatrix}$	

Table 2. Scoring intensity of crop damage caused by pest infestation of mealybug



18 WAP 20 WAP 22 WAP 24 WAP 26 WAP

According to Catalayud *et al.* (1994) in Wardani (2015), cassava resistance is caused by differences in content of secondary compounds in cassava plants. Cyanide acid compounds are secondary compounds contained in cassava plant floem. Floem is a tissue where mealybug sucks fluid to be consumed as a nutrient. The higher of cyanide acid content in cassava, the more susceptible the variety is to the attack of mealybug (Wardani, 2015).

Population of Mealybug (*Phenacoccus*) in Lower Leaf Zone. Populations of mealybug are mostly found in the upper leaf zone. However, in severe attacks, mealybug populations could be found in the upper leaf zone. The mealybug population in the plots with aplication of "Zincmicro" 20 kg ha⁻¹ and 40 kg ha⁻¹ was lower than the mealybug population in plots without "Zincmicro" fertilizer. The mealybug population in plots with application of "Zincmicro" without "Zincmicro" fertilizers did not differ (Figure 3).



Figure 3. Diagram of the mealybug population stem in the lower leaf zone

The mealybug population at plant age of 18 WAP to 26 WAP did not appear to be differ. In plots without the application of "Zincmicro" fertilizer (P1), the population of mealybug 0 to 8 head per-leaf bottom. In the plot with application of "Zincmicro" fertilizer 20 kg ha⁻¹ (P2), the population mealybug 0 to 5 head per leaf bottom. In the plot with the application of "Zincmicro" fertilizer 40 kg ha⁻¹ (P3), the population mealybug 0 to 4 head per leaf bottom. Rainfall is a determinant factor in the dynamics of the mealybug population. Rain can cause mealybug to drift mechanically. Changes in mealybug population can also occur due to stress on plants caused by drought or lack of water. Drought will increase secondary compounds in cassava plants. This compound serves as phagostimulan for mealybug, so it can increase the population in plants (Nurhayati, 2012).

Population of Mealybug (*Phenacoccus*) in Middle Leaves Zone. The mealybug population was found in the central leaf zone more than the mealybug population in the lower leaf zone. This showed that the mealybug population strikes at the top of the leaf towards the lower leaf. The population in the plots with application of "Zincmicro" fertilizer 20 kg ha⁻¹ and 40 kg ha⁻¹ was lower than the mealybug population in plots without "Zincmicro" fertilizer. The mealybug population in plots with application of "Zincmicro" fertilizer without "Zincmicro" fertilizers did not differ (Figure 4). The mealybug population at plant age of 18 WAP to 26 WAP did not appear to be differ. In plots without application of "Zincmicro" fertilizer (P1), the population mealybug 1 to 12 head of per-leaf. In plots with application of "Zincmicro" fertilizer 40 kg ha⁻¹ (P3), the population mealybug 0 to 9 head per-leaf.



Figure 4. Diagram of the mealybug population stem in the central leaf zone

Population of Mealybug (*Phenacoccus*) in Upper Leaves Zone. When compared with the population in the middle and lower leaf zones, mealybug populations were found in the upper leaf zone. The population in the plots with application of "Zincmicro" fertilizer 20 kg ha⁻¹ and 40 kg ha⁻¹ was lower than the mealybug population in plots without "Zincmicro" fertilizer. The mealybug population in plots with application of "Zincmicro" fertilizer without "Zincmicro" fertilizers did not differ (Figure 5). The mealybug population at plant age of 18 WAP to 26 WAP did not appear to be differ. In plots without the application of "Zincmicro" fertilizer (P1) the population mealybug 1 to 20 heads per leaf top. In a plot with a "Zincmicro" fertilizer application of 20 kg ha⁻¹ (P2), the mealybug population was 4 to 12 heads per leaf. In the plot with the application of "Zincmicro" fertilizer 40 kg ha⁻¹ (P3), the population mealybug 1 to 10 heads per leaf top.



Figure 5. Diagram of the mealybug population stem in the upper leaf zone

Intensity of Mite Attack (*Tetranychus***).** The intensity of severe mite attack was seen in the leaves of the yellow crop plants and red rust around the leaf bone. In severe attacks could cause the leaves to fall. High intensity of mite attack was found in plots without application of "Zincmicro" fortilizer. The intensity of attack on the plot with application of "Zincmicro" fertilizer of 20 kg ha⁻¹ and 40 kg ha⁻¹ was lower than the intensity of attack on the plot without "Zincmicro" fertilizer. The intensity of attack on plots with "Zincmicro" fertilizer and without "Zincmicro" fertilizer did not differ (Figure 6).



Figure 6. Bar chart of mite intensity intensity

The intensity of attack on plant age of 18 WAP to 26 WAP did not look differently. In plots without application of "Zincmicro" fertilizer (P1), the intensity of the mite attack was 72.87% to 74.25%. In plot with application of "Zincmicro" fertilizer 20 kg ha⁻¹ (P2), the intensity of mite attack was 61,98% to 66,2%. In plots with application of "Zincmicro" fertilizer 40 kg ha⁻¹ (P3), the intensity of mite attack was 61.16% to 66.2%. "Zincmicro" fertilizer did not affect the intensity of mite attack. Dry climate is a factor that supports the proliferation of mites. Mites can fall mechanically due to rain (Rismunandar, 1981).

Population of Mites (*Tetranychus***) in Lower Leaf Zone.** Mites were found in the lower leaf zone. In severe attacks, mite populations can be found up to the upper leaf zone. The population of mites on plot with application of "Zincmicro" fertilizer 20 kg ha⁻¹ and 40 kg ha⁻¹ was lower than the mite population in the plot without "Zincmicro" fertilizer. The population of mites on plots with application of "Zincmicro" fertilizer without "Zincmicro" fertilizers did not differ (Figure 7). The population of mites at plant age of 18 WAP up to 26 WAP did not look differently. In plots without the application of "Zincmicro" fertilizer (P1), the population of mites 7 to 74 heads per leaf. In the plot with application of "Zincmicro" fertilizer 40 kg ha⁻¹ (P3), the mite population was 10 to 61 heads per leaf. Weather can affect mite populations. Mites populations decline at the start of the wet season and remain at a low level in winter. Maximum and minimum temperatures have a positive correlation with mite attacks (Sadeghi et al., 2016).



Figure 7. Diagram of the mites population stem in the lower leaf zone

Population of Mites (*Tetranychus***) in Middle Zone Leaves.** The population of mites found in the middle leaf zone indicated that the mite populations were mostly found in the lower leaf zone, so the development of the mites populations spreaded to the upper

leaf zone through the central leaf zone. The population of mites on plot with application of "Zincmicro" fertilizer 20 kg ha⁻¹ and 40 kg ha⁻¹ was lower than the mite population in the plot without "Zincmicro" fertilizer. The population of mites in plots given "Zincmicro" fertilizer without "Zincmicro" fertilizer did not differ (Figure 8). The population of mites at plant age of 18 WAP up to 26 WAP did not look different. In plots without application of "Zincmicro" fertilizer (P1), the population of mites was 11 to 36 heads per middle leaf. In a plot with application of "Zincmicro" fertilizer 20 kg ha⁻¹ (P2), the population of mites 4 to 27 heads per-leaf the middle. In the plot with the application of "Zincmicro" fertilizer 40 kg ha⁻¹ (P3), the population of mites 4 to 18 heads per-leaf the middle. Fertilization will affect the durability of the impulse.

This resistance characteristic rised and was driven by the existence of certain environmental conditions so that the plant was able to survive against pest attacks. Resistance of this impulse occured, among others, due to the fertilization and irrigation and other cultivation techniques. The life and development of insects was strongly influenced by the state of nutrition availability in plants, whereas the nutrient conditions of plants were affected by fertilization and irrigation. The availability and proportion of nutrients greatly affected pest life. For example Aphis louse is very sensitive to N content in plants and has a negative response to the content of K (Untung, 1993).



Figure 8. Diagram of the mites population stem in the central leaf zone

Population of Mites (*Tetranychus***) in Upper Leaf Zone.** The population of mites found in the upper leaf zone indicates that the mite populations were mostly found in the lower leaf zone so that the mites extended their attacks towards the upper leaf zone. The population of mites in the upper leaf zone was less than the population of the lower leaf zone. This was because the mites first attacked the lower leaves. The population of mites on plot with application of "Zincmicro" fertilizer 20 kg ha⁻¹ and 40 kg ha⁻¹ was lower than the mite population in the plot without "Zincmicro" fertilizer. The population of mites in plots given "Zincmicro" fertilizer and without "Zincmicro" fertilizer did not differ (Figure 9).

The population of mites at plant age of 18 WAP up to 26 WAP did not look different. In plots without the application of "Zincmicro" fertilizer (P1), the population of mites 0 to 19 heads per-leaf top. In the plot with application of "Zincmicro" fertilizer 20 kg ha⁻¹ (P2), population of mites 0 to 8 heads per leaf top. In plots with application of "Zincmicro" fertilizer 40 kg ha⁻¹ (P3), the population of mites was 0 to 5 heads per leaf.



Figure 9. Diagram of the mite population stem in the upper leaf zone

CONCLUSION

Based on this research it can be concluded that application of 20 kg ha⁻¹ "Zincmicro" was significantly lower than without application "Zincmicro". Increasing the dosage of "Zincmicro" until 40 kg ha⁻¹ did not differ significantly; although there was no significant different between with and without application "Zincmicro", application of "Zincmicro" tended to reduce mite attack intensity than without application "Zincmicro"; and the population of mealybug was more abundant in the upper leaves; meanwhile, the mite population was more abundant in the lower leaves. Application of "Zincmicro" tended to decrease the population of mealybug and mite compared to without application of "Zincmicro".

REFERENCES

- Adriani, E. 2016. Preferensi, Kesesuaian dan Parasitisme Anagyrus lopezi (De Santis) (Hymenoptera: Encrytidae) pada Berbagai Instar Mealybug Singkong, Phenacoccus manihoti Matile-Ferrero (Hemiptera: Pseudococcidae). Tesis. Fakultas Pertanian – IPB. Bogor.
- Badan Pusat Statistik. 2016. Provinsi Lampung Dalam Angka. BPS Provinsi Lampung. Lampung.
- Rojanaridpiched, C. 2017. International Conference On Cassava. Emersia Hotel, 23 24 November 2017. Lampung.
- Fadila, R. 2015. Respon Padi Sawah Varietas IF8 dan Lentera Terhadap Aplikasi Pupuk Organo Mineral dan Pupuk Hayati pada Inceptisol Situgede, Bogor. IPB. Bogor.
- Indiati. 1999. Status Tungau Merah Pada Tanaman Ubikayu. Di dalam: Rahmianna, editor. Pemberdayaan Tepung Ubikayu Sebagai Subsidi Terigu, dan Potensi Kacang-Kacangan untuk Pengayaan Kualitas Pangan. Edisi Khusus Balitkabi N. 15-1999. Malang: Balitkabi. Hlm 122-126.
- Nurhayati, A. 2012. Insidensi Cendawan Entomophthorales Pada Mealybug Pepaya dan Singkong (*Hemiptera : Pseudococcidae*) di Wilayah Bogor. IPB. Bogor.
- Sadeghi, E. S. dan R.V. Madani. 2016. The Influence of *Tetranychus urticae* Koch (Acari : Tetranychidae) Life Table and Reproductive Parameters by Applying Si on Bean at Library Condition. Department of Entomology-Arak University of Iran. Iran.
- Saputro, A. R. 2013. Biologi dan Potensi Peningkatan Populasi Mealybug Singkong. *Phenacoccus manihoti* Matile-Ferrero (*Hemiptera: Pseudococcidae*): Hama Pendatang Baru di Indonesia. Skripsi. Fakultas Pertanian – IPB. Bogor.
- Untung, K. 1993. Pengantar Pengelolaan Hama Terpadu. Gadjah Mada University Press. Yogyakarta.

Wardani, N. 2015. Mealybug Ubikayu, *Phenacoccus manihoti* Matile-Ferrero (*Hemiptera: Pseudococcidae*), Hama Invasif Baru di Indonesia. Disertasi. Fakultas Pertanian – IPB. Bogor.

. 2015. Phenacoccus manihoti Matile-Ferrero (Hemiptera : Pseudococcidae), Mealybug Invasif Baru di Indonesia. Prosiding Seminar Nasional Sains dan Inovasi Teknologi Pertanian. Balai Pengkajian Teknologi Pertanian. Lampung. 34 Hlm.

• 6% Top so	6 Overall Similarity		
 4% Cro 4%	 4% Internet database 6 Crossref database 6 Works database 7% Publications database 6% Crossref Posted Content database 7% Submitted Works database 		
TOP SC The sou display	OURCES urces with the highest number of matches withir ed.	the submission. Overlapping sources will not b	e
1	Higher Education Commission Paki Submitted works	stan on 2009-12-24	1%
2	Sarkar, Smritikana. "Growth, Yield a Publication	nd Moisture Extraction Pattern of B	<1%
3	docplayer.net		<1%
4	S W Indiati, YEB M Santi, Yusmani. Crossref	"Soybean pest control in tidal swa	<1%
5	Higher Education Commission Paki Submitted works	stan on 2010-11-24	<1%
6	ojs.uho.ac.id Internet		<1%
7	Universitas PGRI Semarang on 2020 Submitted works)-01-06	<1%
8	media.neliti.com		<1%

9	repository.unib.ac.id	<1%
10	"Abundance of Ants on Chili and Corn Intercrop Planting Techniques", I Crossref	<1%
11	protan.studentjournal.ub.ac.id	<1%
12	Robert W. Tairas, J. Pelealu, M. Tulung. "INTERAKSI VARIETAS PEPAY Crossref	<1%
13	University of Greenwich on 2021-12-03 Submitted works	<1%
14	Southeast Community College on 2020-03-22 Submitted works	<1%

 Excluded from Similarity Report Pibliographic material 	
Manually excluded sources Manually excluded text blocks	
EXCLUDED SOURCES	
repository.lppm.unila.ac.id	95%
digilib.unila.ac.id	5%
Higher Education Commission Pakistan on 2022-03-02 Submitted works	5%
Higher Education Commission Pakistan on 2010-02-22 Submitted works	4%
arccarticles.s3.amazonaws.com	4%
doaj.org Internet	3%
agraer.ms.gov.br Internet	3%
mafiadoc.com Internet	3%
scielo.br Internet	3%
Nabin Rawal, Shree Prasad Vista, Dinesh Khadka. "Nitrogen concentration, its	· ว 0/

3%

Crossref posted content

Sholihin, Sri W. Indiati, Kartika Noerwijati, Tinuk S. Wahyuni et al. "Improving t	1%
T S Wahyuni, K Noerwijati. "Cassava genotypes selection for high yield and hi Crossref	<1%
University of Minnesota System on 2022-07-20 Submitted works	<1%
Yohanis Ngongo, Tony Basuki, Bernard deRosari, Yosep Seran Mau et al. "The ^{Crossref}	<1%
I Gede Swibawa, Franciscus Xaverius Susilo, Purnomo Purnomo, Titik Nur Ae Crossref	<1%
Dicky Ashari Ramadhan, F X Susilo, Nur Yasin, I Gede Swibawa. "PENGARUH Crossref	<1%
Anthony Bellotti, Beatriz Vanessa Herrera Campo, Glenn Hyman. "Cassava Pr ^{Crossref}	<1%
EXCLUDED TEXT BLOCKS	
University of LampungJI. Soemantri Brojonegoro No. 1 Bandar Lampung 35145 eprints.undip.ac.id	
All rights reserved. No part of this "BER volume 77 issue 2 Cover and Back matter", Bulletin of Entomological Research, 2009	

Identifying Chemical Compound in Ceara Rubber Skin Which Is Potential To BeNat...

University of Minnesota System on 2021-10-18

Growth and Yield of Cassava (Manihot Esculenta Crantz

pustaka.setjen.pertanian.go.id

Jl. Prof. Soemantri Brojonegoro, No. 1, Bandar Lampung 35145

Universitas Terbuka on 2018-08-16

MATERIALS AND METHODSThe research was conducted in

repository.unib.ac.id

CONCLUSIONBased on this

repository.unib.ac.id