

PAPER NAME

**M, A, FDauzan, Jamalam L. Prociding International 2019.pdf**

AUTHOR

**Hery Novpriansyah**

WORD COUNT

**1780 Words**

CHARACTER COUNT

**9507 Characters**

PAGE COUNT

**5 Pages**

FILE SIZE

**1.9MB**

SUBMISSION DATE

**Apr 11, 2023 8:58 AM GMT+7**

REPORT DATE

**Apr 11, 2023 8:59 AM GMT+7****● 4% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 4% Publications database
- Crossref database

**● Excluded from Similarity Report**

- Internet database
- Submitted Works database
- Crossref Posted Content database
- Bibliographic material

ISBN : 978-4-909365-04-0

# **1** PROCEEDINGS of IC-GU 12 UGSAS-GU

## “6TH INTERNATIONAL WORKSHOP ON CROP PRODUCTION AND PRODUCTIVITY UNDER GLOBAL CLIMATE CHANGE”



**Editors :**

**Dr. Afandi**

**Prof. Dr. Ken Hiramatsu**

**DECEMBER 3-4, 2018**

**at FACULTY OF AGRICULTURE, LAMPUNG UNIVERSITY  
BANDAR LAMPUNG, INDONESIA**

**AGGREGATE STABILITY AND ROOT BIOMASS AFFECTED BY SOIL TILLAGE AND MULCHING IN GREEN NUT CULTIVATION (*VIGNA RADIATA* L.)**

*M. A. Fauzan, J. Lumbanraja, H. Novpriansyah, Afandi and N. Kaneko* ..... p.59

**APPLICATION of INDUCED COMPOST of CELLULOLITIC (*Aspergillus fumigatus*) AND LIGNINOLITIC (*Geotrichum* sp.) INOCULUM on The VEGETATIVE GROWTH of RED CHILI (*Capsicum annuum* L.)**

*AyuWulan Septitasari, Bambang Irawan, Zulkifli and Salman Farisi*..... p.61

**SOIL COMPACTION, WATER CONTENT, BULK DENSITY AND SOIL ROOT BIOMASS AFFECTED BY TILLAGE AND FERTILIZER ON GEDUNG MENENG SOIL UNDER GREEN BEAN GROWTH**

*Yogi Irawan, J. Lumbanraja, Nur Afni Afrianti, Afandi*..... p.62

**PERCEPTIONS OF FARMERS, EFFECTIVENESS OF FARMERS GROUP, AND DIFFUSION OF INNOVATION OF ORGANIC FARMING SYSTEM IN LAMPUNG PROVINCE**

*Tubagus Hasanuddin*..... p.65

**PRODUCTION AND HARVESTED NUTRIENT OF CASSAVA (*MANIHOT ESCULENTA* L.) AFFECTED BY COMPOST AND ITS COMBINATION WITH NPK INORGANIC FERTILIZER FOR THE 5<sup>TH</sup> PLANTING PERIOD**

*Novita Desri Wanti, Jamalam Lumbanraja, Supriatin, Sarno, Dermiyati Sugeng Triyono, and N. Kaneko*..... p.69

**SIMULATION OF CAVENDISH BANANA TRANSPORTATION**

*Debby Nuzulia Arlin, Cicih Sugianti, Siti Suharyatun, and Tamrin*..... p.72

**THE APPLICATION OF HOT WATER TREATMENT IN MANGO CV ARUMANIS**

*Cicih Sugianti and Dondy A Setyabudi*..... p.76

**HARVESTED NUTRIENT AND PRODUCTION OF CASSAVA (*Manihot esculenta*) AFFECTED BY TILLAGE AND HERBICIDE IN THE 4<sup>th</sup> PLANTING PERIOD IN GEDUNG MENENG SOIL BANDAR LAMPUNG**

*Adinda Kusuma Dewi Rachmat, Jamalam Lumbanraja, Nur Afni Afrianti, Muhajir Utomo, and N. Kaneko*..... p.80

**PRODUCTION AND HARVESTED NUTRIENTS OF SUGARCANE 1<sup>ST</sup>  
RATOON (*SACCHARUM OFFICINARUM* L.) AFFECTED BY ORGANIC AND  
INORGANIC FERTILIZER**

*Nurhidayat, Jamalam Lumbanraja, Supriatin , Sarno, Dermiyati  
and Sugeng Triyono.....*

p.83

**BIOGAS PRODUCTION FROM OIL PALM EMPTY FRUIT BUNCHES  
THROUGH DRY FERMENTATION PROCESS: PRELIMINARY RESULTS**

*Agus HARYANTO, Cicih SUGIANTI, Sugeng TRIYONO,  
and Nanda Efan APRIA.....*

p.87

**THE CURRENT STATUS OF AUTHENTICATION OF INDONESIAN  
SPECIALTY COFFEES USING UV-VISIBLE SPECTROSCOPY AND  
CHEMOMETRICS**

*Diding SUHANDY and Meinilwita YULIA.....*

p.90

**THE DIVERSITY OF ARBUSCULAR MYCORRHIZA FUNGI AT  
RHIZOSPHERE OF CASSAVA OF THAILAND CLONE CULTIVATED IN  
LAMPUNG TIMUR AND TULANG BAWANG BARAT**

*Maria Viva RINI, Kuswanta Futas HIDAYAT, Diah PURBANINGRUM,  
Annisa HASKA .....*

p.93

## Aggregate Stability and Root Biomass Affected by Soil Tillage and Mulching in Green Nut Cultivation (*Vigna radiata* L.)

M. A. Fauzan<sup>1</sup>, J. Lumbanraja<sup>2\*</sup>, H. Novpriansyah<sup>2</sup>, Afandi<sup>2</sup> and N. Kaneko<sup>3</sup>

<sup>1)</sup> Under graduate student, Faculty of Agriculture, Lampung University

<sup>2)</sup> Faculty of Agriculture, Lampung University

<sup>3)</sup> Yokohama National University, Japan.

### **Abstract**

The bare intensive tillage applied on agricultural land can drastically reduce soil quality. One of the efforts to prevent it is by applying mulch and conservation tillage. The objective of this study was to determine the effect of minimum tillage and mulch on soil aggregate stability and root biomass. This study was designed in the field using a randomized block design (RBD) with 2 factors and 4 repetitions. The first factor was tillage (minimum tillage (T1) and intensive tillage (T2)). The second factor was mulch (without mulch (M1) and 5 Mg mulch ha<sup>-1</sup> (M2)). The aggregate stability was determined using water-drop method (WDM). The results showed that the minimum tillage affected the increasing soil aggregate stability of the 6 mm and 3 mm diameter aggregate before land preparation and after harvesting, except, the 3 mm diameter aggregate before land preparation. The minimum tillage also increased root biomass during the vegetative period of the plant. The treatment of 5 Mg mulch ha<sup>-1</sup> affected in increasing the soil aggregate stability of 3 mm diameter aggregate after harvest but not the soil of 6 mm diameter aggregate. No interaction was found between the minimum tillage and 5 Mg mulch ha<sup>-1</sup> on soil aggregate stability and root biomass.

### **Introduction**

Land preparation was an important thing for any crop cultivation. Intensive tillage can damage soil structure but also stimulates soil oxidation in which the decomposition of organic matter much faster. The decrease of aggregate stability is very common in crop farming systems (Rachman and Abdurachman, 2006) since soil in intensive tillage when rain can break up the soil aggregate (Refliaty and Marpaung, 2010). Sarief (1989) reported that excretion and root penetration can stimulate or be affected by soil structure. The more roots found in the soil, the higher the root biomass results in the more stable soil structure will be. The objective of this research is to study the effect of minimum tillage and mulch on soil aggregate stability and root biomass.

### **Material and Method**

The location of the research was carried out at the Integrated Field Laboratory, Lampung University. This study in the field was designed using a randomized block design (RBD) with 2 factors and 4 replications. The first factor was tillage (minimum tillage (T1) and intensive tillage (T2)). The second factor was mulch (without mulch (M1) and 5 Mg mulch ha<sup>-1</sup> (M2)) The aggregate stability was determined using water-drop method (WDM) (McCalla, 1944). The average results of the observed values of the data were tested by the BNT test level of 5%.

### **Result and Discussions**

#### **Effect on Soil Aggregate Stability**

Table 1 showed that soil aggregate stability before land preparation was significantly higher for 6 mm diameter compare to that of intensive tillage and with no mulch, but not significantly in mm diameter. The tillage system treatment significantly reduced the stability of 6 mm and 3 mm soil aggregates after harvest. Mulching treatment was not significantly different from aggregate stability at all size and time of observation except for soil 3 mm diameter aggregates which showed significantly different observations after harvest.

#### **Effect on Roots Biomass**

Root biomass in the minimum tillage was very significantly different compared to intensive tillage during vegetative period of the plants (Table 2). This is because the condition of minimum tillage is more able to provide space for roots to grow and develop. A good root growth process will increase root biomass. In addition, it is suspected that the breakdown of roots in the soil due to intensive tillage treatment so that the roots decompose faster, while in the minimum tillage which is only treated as needed does not damage the root growth rate. Prasetyo, et al., (2014) stated that maximum tillage was an act of reversing, cutting, destroying and leveling the soil by removing plant debris that interferes with growth, while minimum tillage is a sufficient soil cultivation by maintaining

crop residues the former is still on the surface of the land.

Table 1. Aggregate stability of 6 mm and 3 mm diameter soil aggregates

Soil aggregate	Treatment	Soil aggregate stability (kinetic energy = J)	
		BL	AH
6 mm	T1	5,96 b	3,99 b
	T2	4,06 a	2,81 a
	F Test	*	*
	HSD 0,05	1,81	0,98
	M1	4,84	3,00
	M2	5,18	3,81
3 mm	T1	1,75	1,78 b
	T2	2,16	1,22 a
	F Test	ns	**
	HSD 0,01	-	0,41
	M1	1,92	1,35 a
	M2	1,99	1,65 b
F Test	ns	*	
HSD 0,05	-	0,29	

Table 2. Effect of the tillage system and mulch on root biomass in the green nut plants

Treatment	Root biomass (kg m <sup>-3</sup> )		
	BL	VP	AH
T1	0,69	3,22 b	0,48
T2	0,54	1,39 a	0,27
F Test	ns	**	ns
HSD 0,05	-	0,67	-
M1	0,61	1,98	0,40
M2	0,61	1,66	0,35
F Test	ns	ns	ns

**Effect on C- organic**

The content of total -C (%) in the tillage treatment and with mulch were not significantly different in all phases of observation . It is presumed that the source of organic matter given through mulch or from plant tissue that falls to the soil on the land had not been completely decomposed during the study period..

**Correlation Aggregate Stability with Root Biomass**

The correlation test results between aggregate stability and root biomass showed no significant correlation of the soil neither on 3 mm diameter of the aggregate nor on 6 mm

**Correlation Aggregate Stability with Total-C of Soil**

The correlation between soil aggregate stability and the total-C of 3 mm diameter was not significantly correlate but it was significantly correlate on the 6 mm diameter soil aggregate (Table 3). It was suspected that root exudates in the form of sugars and organic compounds that fill a large aggregate pore would be fragmented into smaller aggregates when the

compound decomposes. Ladd, et al. (1996) reported that organic compounds such as organic acids and polysaccharides were able to bind soil particles into smaller, more stable aggregates of soil. Organic matter (polysaccharides) decomposes faster by microorganisms is a temporary aggregate binding agent (Tisdall and Oades, 1982).

Table 3. The correlation soil aggregates 6 mm and 3 mm diameter to C- organic (%)

Variable	Correlation coefficient	
	Soil aggregate stability (kinetic energy = J)	
	6 mm	3 mm
C-organic	- 0,444 *	0,149 <sup>ns</sup>

ns = no significant effect, \* = significant effect.

**Conclusion**

- 1)The minimum tillage improved the aggregate stability of the 3 mm and 6 mm diameter aggregate on the soil before land preparation and after harvest, but not in the soil of 3 mm diameter aggregate on before land preparation. The minimum tillage increase root biomass in the vegetative period of the plant, but not in before land preparation and after harvest.
- 2) Treatment of mulch 5 Mg ha<sup>-1</sup> increased the stability of aggregate of 3 mm diameter of soil aggregate after harvest, but not for soil 3 mm diameter aggregate before land preparation. The mulch 5 Mg ha<sup>-1</sup> did not increase root biomass.
- 3) No interaction was found between the minimum tillage and 5 Mg mulch ha<sup>-1</sup> on soil aggregate stability and root biomass.

**Reference**

- 1) McCalla TM (1944) Water-drop method of determining stability of soil structure. Soil Sci. 58 (2): 117–121.
- 2) Prasetyo RA, A Nugroho dan J Moenandir (2014) Pengaruh sistem olah tanah dan berbagai mulsa organik pada pertumbuhan dan hasil tanaman kedelai (*Glycine max* (L.) Merr.) varietas grobogan. Jurnal Produksi Tanaman. 1 (6): 486-495.
- 3) Rachman A dan A Abdurachman (2006) Penetapan kemantapan agregat tanah. Balai Penelitian Tanah. Badan Penelitian dan Pengembangan Pertanian. Kementrian Pertanian, Bogor. 63-73.
- 4) Refliaty dan EJ Marpaung (2010) Kemantapan agregat ultisol pada beberapa penggunaan lahan dan kemiringan lereng. J.Hidrologi. 1(2): 35-42.
- 5) Sarief S (1989) Fisika-kimia tanah pertanian. Pustaka Buana. Bandung. 220 hlm.
- 6) Tisdal JM and Oades JM (1982) Organic matter and water-stable agregates in soils. Journal of Soil Science. 33: 141-163.

## ● 4% Overall Similarity

Top sources found in the following databases:

- 4% Publications database
- Crossref database

### TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

- 1** **Sugiatno Sugiatno, Ali Fitrah Hansyah, Rusdi Evizal, Sri Ramadiana. "Pe...** **2%**  
Crossref
- 2** **Dina Yulianti Sisca Vera, Edhi Turmudi, Eko Suprijono. "PENGARUH JA...** **<1%**  
Crossref
- 3** **Mark Anglin Harris. "Geobiotechnological Solutions to Anthropogenic ...** **<1%**  
Crossref