



ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 23 - Number 3 - March 2022

Front cover: Ricinus communis L.
(PHOTO: ROD WADDINGTON)

Published monthly

PRINTED IN INDONESIA

ISSN: 1412-033X

E-ISSN: 2085-4722



9 771412 033863



9 772085 472867

BIODIVERSITAS

Journal of Biological Diversity
Volume 23 – Number 3 – March 2022

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), **Agnieszka B. Najda** (Poland), **Ajay Kumar Gautam** (India), **Alan J. Lymbery** (Australia), **Annisa** (Indonesia), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Hassan Pourbabaie** (Iran), **Joko R. Witono** (Indonesia), **Kartika Dewi** (Indonesia), **Katsuhiko Kondo** (Japan), **Kusumadewi Sri Yulita** (Indonesia), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Mochamad A. Soendjoto** (Indonesia), **Mohamed M.M. Najim** (Srilanka), **Mohib Shah** (Pakistan), **Nurhasanah** (Indonesia), **Praptiwi** (Indonesia), **Rasool B. Tareen** (Pakistan), **Seyed Aliakbar Hedayati** (Iran), **Seyed Mehdi Talebi** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Subhash C. Santra** (India), **Sugeng Budiharta** (Indonesia), **Sugiyarto** (Indonesia), **Taufiq Purna Nugraha** (Indonesia), **Yosep S. Mau** (Indonesia)

EDITOR-IN-CHIEF:

S u t a r n o

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:

biodiversitas.mipa.uns.ac.id; smujo.id/biodiv



Society for Indonesia
Biodiversity



Sebelas Maret University
Surakarta

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or *Biodiversitas* encourages submission of manuscripts dealing with all aspects of biodiversity including plants, animals and microbes at the level of the gene, species, and ecosystem. Ethnobiology papers are also considered.

Article types The journal seeks original full-length: (1) **Research papers**, (2) **Reviews**, and (3) **Short communications**. Original research manuscripts are limited to 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles are also limited to 8,000 words, while Short communications should be less than 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). Please ensure that the manuscript is submitted using the *Biodiversitas* template, which can be found at (<https://biodiversitas.mipa.uns.ac.id/D/guidance.htm>). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Please also provide a list of five potential reviewers in your cover letter. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors are allowed to reproduce articles as long as they are not used for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance Only articles written in U.S. English are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. Manuscripts will be rejected if the content does not in line with the journal scope, does not meet the standard quality, is in an inappropriate format, contains complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biological significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in January, April, July and October.

A charge Starting on January 1, 2017, publishing costs waiver is granted to foreign (non-Indonesian) authors who first publish the manuscript in this journal, especially for graduate students from developing countries. However, other authors are charged USD 250 (IDR 3,500,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email. Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible, use the *Biodiversitas* template, and Microsoft Word based (.doc or .rtf; not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italics, except in italicised sentences. Scientific names (Genus, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. The Genus name can be shortened after first mention, except where this may generate confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial names should be avoided. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Standard chemical abbreviations can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurements** should use IS denominations, and other system should use equivalent values with the denomination of IS mentioned first. Abbreviations like g, mg, mL, etc. should not be followed by a dot.

Minus index (m-2, L-1, h-1) suggested to be used, except in things like "per-plant" or "per-plot". **Mathematical equations** can be written down in one column with text, in that case can be written separately. **Numbers** one to ten are written in words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

The **Title** of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Author name(s) should be completely written. Name and institution address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscripts written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

The **Abstract** should not be more than 200 words. Include between five and eight **Keywords**, using both scientific and local names (if any), research themes, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. Running title is about five words. The **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. A **Conclusion** should be given at the end of the discussion. **Acknowledgments** are expressed in brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest must be noted.

Figures and Tables of three pages maximum should be clearly presented. Include a label below each figure, and a label above each table (see example). Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. There is no **Appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, supplementary information can be provided on the website.

References In the text give the author names followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "cit" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order. Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). Please include DOI links for journal papers. The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158. DOI: 10.13057/biodiv/d070213

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet: Balagadde FK, Song H, Ozaki J, Collins CH, Barnett M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4:187. www.molecularsystembiology.com

The growth and yield responses of two bean cultivars to organic and inorganic nitrogen sources

DARWIN H. PANGARIBUAN^{1,✉}, ERMAWATI¹, NI MADE WIDI SURYANI², WAHYU SARA MARIA²

¹Department of Agrotechnology, Faculty of Agriculture, Universitas Lampung. Jl. Sumantri Brojonegoro No. 1, Bandar Lampung 35141, Lampung, Indonesia

²Graduate Program of Agrotechnology, Faculty of Agriculture, Universitas Lampung. Jl. Sumantri Brojonegoro No. 1, Bandar Lampung 35141, Lampung, Indonesia. Tel.: +62-721-781822, ✉email: darwin.pangaribuan@fp.unila.ac.id

Manuscript received: 15 March 2021. Revision accepted: 28 February 2022.

Abstract. Pangaribuan DH, Ermawati, Suryani NMW, Maria WS. 2022. The growth and yield responses of two bean cultivars to organic and inorganic nitrogen sources. *Biodiversitas* 23: 1677-1682. The productivity of beans is increased by selecting the best cultivar and applying fertilizers according to the needs of the plant. In addition, the application of high-yielding cultivar and N fertilizers with the right dosage could be used to increase the yield. Therefore, the objective of this study is to determine the effect of nitrogen sources on the growth and yield of two bean cultivars. A 5 × 2 factorial treatment was used in a randomized block design with three replications. The first factor is the Perkasa and Lebat-3 cultivar, the second is organic and inorganic nitrogen fertilizer, namely control, urea 100 kg ha⁻¹, as well as 10, 20, and 30% concentrations of liquid organic fertilizer from leucaena leaves. The results showed that the use of liquid organic fertilizers derived from leucaena could be used as an alternative to nitrogen fertilization. Furthermore, the 20% concentration treatment produced the highest number of trifoliolate leaves (8.81 leaves) and dry weight (4.52 g). Based on the results, the use of 20% liquid organic fertilizer with Perkasa cultivar showed the best effect on the number of pods per plant (13.38), and yield (846.67 kg ha⁻¹). The technology of liquid organic fertilizers from leucaena leaves could be recommended for bean organic production.

Keywords: Chlorophyll, liquid organic fertilizers, stomata, urea, yield

INTRODUCTION

Green bean (*Phaseolus vulgaris* L.) is one of the legume plants serving as a source of vegetable protein. The plant is widely consumed by the Indonesians due to its important role in fulfilling health needs as a nutritious food ingredient. Rihana et al. (2013), stated that beans are source of vegetable protein, rich in vitamins A, B, C and have great potential for the treatment of several diseases such as oxidative stress, cardiovascular disease, cancer, and diabetes (Camara et al. 2013). Furthermore, as the population grows, the need for fresh food and vegetables continues to also increase. Therefore, to increase the production of green beans, it is necessary to apply good cultivation techniques such as applying balanced fertilization that fulfills the nutritional needs for growth. The productivity of green beans is also increased by using improved cultivars. Meanwhile, there are numerous green beans cultivars available in the market. Ratnasari et al. (2015), stated that each cultivar has different genetic characteristics, which lead to differences in appearance, character, as well as responses to production factors.

Fertilization is one way to meet the nutritional needs of nitrogen in green beans. Meanwhile, nitrogen is an important nutrient required by green beans for the formation of chlorophyll, protoplasm, protein, and nucleic acids (Fahmi et al. 2010). Fertilization with nitrogen (N) is derived from organic and inorganic fertilizers such as urea. Furthermore, green bean nutritional need for N is higher than any other nutrients. Pahlevi et al. (2016) reported that

nitrogen affects the photosynthetic process in plants as well as the photosynthate produced. Maghfoer et al. (2018), also stated that the application of inorganic fertilizers produced large growth and plant yield. However, the continuous use of inorganic fertilizers leads to low soil fertility and productivity. Therefore, an alternative approach is needed to reduce the use of inorganic fertilizers, via organic fertilization.

Nitrogen is commonly supplied alternatively to plants using Liquid Organic Fertilizers (LOF). It is an organic fertilizer available in liquid form and contains nutrients in the form of a solution. Therefore, it is easily absorbed by plants. Furthermore, it is applied via sprinkling or spraying on the leaves or stems of plants. Organic fertilizers are made from plant waste available in the environment. In particular, liquid organic fertilizers are more beneficial because their distribution is adjustable to the needs of plants (Ginandjar et al. 2019). One example of organic matter widely used as a liquid organic fertilizer source of nitrogen is leucaena leaves (*Leucaena leucocephala*). Liquid organic fertilizer derived from this plant contains nitrogen nutrients needed by plants (Jeksen and Mutiara 2017). The N content in leucaena leaves is quite high, namely 3.84%, and is a source of organic nutrients to increase the growth and yield of green beans (Palimbungan et al. 2006). Furthermore, the application of liquid organic fertilizers increased the fresh weight of pods in green beans (Rizqiani et al. 2006), and provided the highest yield of pods and bean per plant in slam weed (*Chromolaena odorata* L.) (Duaja 2013).

Plant cultivar is a factor that greatly determines the quality of agricultural products. The use of high-yielding bean cultivar increases productivity and yield quality. Each cultivar has specific genetic characteristics and traits which lead to differences in the respective character and appearance. Examples of high-yielding beans cultivar include Perkasa and Lebat-3. The advantages of Perkasa cultivar include resistance to leaf rust disease, early maturity as well as the large and long pods (MoA 2002). In contrast, the advantages of Lebat-3 cultivar include being well adapted to the low-highlands, early maturity and has high yield potential (MoA 1999). Sinaga et al. (2017) showed that the use of cultivars had a significant effect on the growth and yield of beans. Furthermore, Duaja (2013) stated that there is an interaction between the use of bean cultivar with liquid organic fertilizer doses on the number of pods and productivity.

The fertilizer requirement by each cultivar is usually different. This is caused by individual genetic characteristics. Hence, the selection of fertilizer types and requirements need to be considered. Besides, the appropriate doses of liquid organic fertilizers also influence plant growth. Ratnasari et al. (2015) stated that variations in genetic traits of cultivar lead to different responses to environmental and production factors. There are only a few studies on the different organic and inorganic nitrogen sources. In this study, two bean cultivars were hypothesized to produce different responses to treatments with various types and doses of inorganic-and-organic-sources-based nitrogen. Therefore, the aim of the study is to determine the effect of inorganic and organic nitrogen sources on the growth and yield of two beans cultivars.

MATERIALS AND METHODS

This study was conducted at the experimental research area of Pemanggilan, Natar, Lampung province, Indonesia from January to March 2020. The soil was first cultivated using hoes, loosened to a depth of 20-30 cm, evenly processed and an experimental plot was made with a size of 3×2.5 m. Furthermore, the Perkasa and Lebat-3 cultivars were planted by drilling the planting holes, while two seeds each were planted with a spacing of 40 cm × 40 cm.

The (2×5) factorial treatment was used in a randomized block design (RBD) with three replications. The first factor was the bean cultivars, namely Perkasa and Lebat-3, while the second was the source of nitrogen, namely control, urea 100 kg ha⁻¹, as well as 10%, 20%, and 30% LOF. Furthermore, the data were analyzed using analysis of variance procedures by using Minitab 17, while the means were compared using the Honestly Significant Difference (HSD) of 5%.

The liquid fertilizer was prepared by mixing chopped leucaena leaves (1 kg of leucaena leaves: 2 liters of water), with 200 mL of molasses and 200 mL of EM4 in a drum, the mixture was then fermented for 15 days and the liquid was filtered. The inorganic fertilizers given include urea 100 kg ha⁻¹ (75 g plot⁻¹) at the beginning and 4 weeks after planting (WAP), while SP-36 200 kg ha⁻¹ (150 g plot⁻¹) and

KCL 100 kg ha⁻¹ (75 g plot⁻¹) were given at the beginning of planting. Fertilization was applied in a strip at a distance of 5 cm from the plant.

The liquid organic fertilizer was applied by spraying on the top and bottom of the plant (60:40). This was carried out once a week using a concentration of 10% (100 mL LOF solution + 900 mL water), 20% (200 mL LOF solution + 800 mL water), and 30% (300 mL LOF solution + 700 mL water). The liquid organic fertilizer analysis results showed that the N content in the *Leucaena* leaves was 0.16%, C-organic 3.08%, N-total 0.16%, P-total 0.18% and K 0.56% with a pH of 3.64 and the C/N ratio was 19.25.

Plant maintenance includes weeding manually and thinning, while the harvest was carried out in stages according to the age of each cultivar. The characteristics of beans ready for harvest include young and gloomy pod color, rough skin surface, less prominent seeds in the pods which usually make a popping sound when the pods are broken especially between 2-3 weeks after flowering.

The parameters observed include plant height (6 WAP), number of trifoliolate leaves (6 WAP), chlorophyll, carotenoids (5 WAP), number of stomata (5 WAP), soil-plant development analyses, SPAD value (5 WAP), number of pods per plant, dry weight (leaves and stem), and yield. Furthermore, the replica method modified from Paul (2017) was used to observe the stomata on the leaf surface. The leaves were first cleaned with a tissue to remove dust or dirt and then rubbed with transparent nail polish, and allowed to dry for a few minutes. A transparent strip tape measuring 1 cm x 1 cm in size was applied, smoothed, and then peeled off slowly. The sample produced was then attached to the slide and observed under a microscope.

The chlorophyll and carotenoid contents were measured using the spectrophotometric method. The green beans were crushed using a mortar and 100 mL of 70% alcohol solution was added. Furthermore, the extract was filtered and the filtrate was placed in a cuvet to measure the total chlorophyll and carotenoid contents using a spectrophotometer (Cary 100 UV-Vis). The chlorophyll and carotenoid contents were calculated using the formula (Rahimi et al. 2019):

$$\begin{aligned} \text{Chlorophyll } a &= 11,24 \times A_{662} - 2,04 \times A_{645} \\ \text{Chlorophyll } b &= 20,13 \times A_{645} - 4,19 \times A_{662} \\ \text{Chlorophyll total} &= 7,05 \times A_{662} + 18,09 \times A_{645} \\ \text{Carotenoid} &= \end{aligned}$$

$$\frac{(1000 \times A_{470} - 1,90 \times \text{clorofil } a - 63,14 \times \text{clorofil } b)}{214}$$

RESULTS AND DISCUSSION

The results showed that the Perkasa and Lebat-3 cultivar had no significantly different effect on vegetative growth namely plant height, numbers of leaves, and dry weight, while the application of inorganic-and-organic-sources-based nitrogen affected vegetative growth parameters

(Table 1). Plant height in the control was shorter than the treatment with urea, as well as 10, and 20% LOF, meanwhile, the application of N sources from urea, as well as 10, 20, and 30% LOF showed no significant differences (Table 1). Amin (2011), reported that nitrogen increased plant growth and height to produce numerous internodes which resultantly produce more leaves.

The number of trifoliolate leaves in green beans is influenced by the availability of N nutrients. This parameter increased significantly after the application of organic nitrogen sources. The numbers of trifoliolate leaves in the control were lesser compared to 10, 20, and 30% LOF. The 10% LOF treatment produced the highest number of trifoliolate leaves but was not different from the 20, and 30% LOF, as well as urea treatment. Pramitasari et al. (2016), stated that higher N absorption by plant leads to larger growth of the leaves. Furthermore, the application of LOF from leucaena leaves showed significant growth in plant height, numbers of leaves, and fresh weight of mustard greens (Pary 2015), the highest yield on the number of pakcoy leaves (Hidayat and Suharyana 2019), and increased the growth and number of fruit in tomato plants (Septirosya et al. 2019). The number of trifoliolate leaves is a plant characteristic that affects the rate of photosynthesis to capture sunlight. Sab et al. (2020) showed that a higher number of leaves would result in a higher plant dry weight. In this research, leucaena-based liquid organic fertilizers that are rich in nitrogen increased the vegetative growth including number of trifoliolate leaves and dry weight.

The application of nitrogen sources affected the dry weight of the plant. Liquid organic fertilizers 20% produced the highest dry weight but were not significantly different from urea and 10% LOF. The plant dry weight was influenced by the initial vegetative growth as this parameter is related to the number of leaves and fresh weight. The availability of sufficient nutrients in the treatment affects plant growth and yield. Arista et al. (2015) stated that nitrogen is an element that functions to increase leaf size and the percentage of protein. Meanwhile, Madusari (2019) reported that liquid organic fertilizers affect plant growth and nutrient absorption by increasing plant stem diameter and dry weight. From these data it could be said that the application of inorganic-and-organic-sources-based nitrogen gave the effect on the plant growth since early or vegetative growth.

The application of inorganic-and-organic-sources-based nitrogen affected physiological characteristics namely chlorophyll, carotenoids, number of stomata and SPAD value (greenness of leaves). However, chlorophyll and leaf carotenoids were not affected by the bean cultivar (Table 2). The chlorophyll content in the control was lower than the urea treatment, and was not different with 30% LOF. Furthermore, the chlorophyll content for the urea treatment did not differ compared to 10 and 20% LOF (Table 2). The application of urea produced a higher carotenoid content compared to the control, but was not different compared to 10 and 20% LOF treatments (Table 2). Also, the application of 10, 20, and 30% LOF showed no differences in the carotenoid content compared to the control. Bojovic

et al. (2005), stated that the leaf color of certain cultivars did not directly correlate with chlorophyll content, although Permanasari et al. (2014) stated that chlorophyll formation plays an important role in the process of photosynthesis. Wijiyanti (2019) reported that nitrogen is the main nutrient needed for the formation of chlorophyll and carotenoids. This is also in line with Razaq et al. (2017) who found that chlorophyll and carotenoids synthesis depends on optimal N availability, therefore, nitrogen plays an important role in the formation of photosynthetic pigments. Furthermore, Hendriyani et al. (2018) reported that carotenoids and chlorophyll are complementary pigments, but carotenoids are fewer in number and it play a role in the absorption of light by chlorophyll. It is clear from this research that chlorophyll and carotenoid contents are affected by the availability of N nutrients.

The application of urea produced a higher number of stomata compared to the control, while the application of urea and 20% LOF showed no differences in the number of stomata. Similarly, LOF applications of 10, and 30% also produced no difference in the numbers of stomata (Table 2). The number of stomata in the two cultivars were not affected by the application of nitrogen fertilizers, the two cultivars showed similar responses. Figure 1 showed the number of stomata as a result of treatments in two cultivars investigated. The application of inorganic fertilizers produced more stomata compared to other treatments as shown in Figure 1. Table 2 showed that urea and 20% LOF produced a higher number of stomata compared to the control, as indicated also by Figure 1. Meanwhile, the number of stomata in plants affects metabolic processes in plants such as photosynthesis. The rate of photosynthesis increases as the number of stomata rises; hence, plant productivity also increases. This is supported by Proklamasiningsih et al. (2012), which stated that the rate of photosynthesis in plants is closely related to the yield. Moreover, Putri et al. (2017), reported that stomata play an important role in photosynthesis by acting as a surface for CO₂ exchange in the leaves.

The results showed that the level of SPAD value, number of pods, and bean yield were significantly affected by the interaction of N sources and bean cultivar (Table 3). The treatment without N fertilizer had low SPAD value, while the application of nitrogen fertilizers affected the greenness of the leaves. The 30% LOF treatment with Perkasa cultivar and urea with Lebat-3 produced the highest level of SPAD value compared to the others but were still significantly the same as the other treatments with different N sources. Moreover, the SPAD value of plants treated with organic and inorganic fertilizers was higher than the control. Nitrogen is an important element that increases leaf greenness in plants (Faustina et al. 2015). Pramitasari et al. (2016) also stated that the N nutrient affects plant growth, appearance, and color hence, the plant organs turn green due to the chlorophyll content. Furthermore, Pamungkas and Supijatno (2017), reported that the level of leaf greenness indicates that the plant has sufficient nitrogen levels and good health conditions.

Table 1. Effects of treatment with different nitrogen sources and cultivar on plant height, number of trifoliolate leaves, and dry weight

Treatment	Plant height (cm)	Number of trifoliolate leaves	Dry weight (g)
Perkasa cultivar	133.60a	7.20a	3.21a
Lebat-3 cultivar	137.63a	8.11a	3.17a
HSD _{0,05}	18.47	1.43	0.92
Control	94.71c	5.57b	1.95b
Urea	171.48a	8.50ab	3.78ab
10% LOF	150.60ab	9.11a	3.13ab
20% LOF	148.02ab	8.81ab	4.52a
30% LOF	113.29bc	6.30ab	2.58ab
HSD _{0,05}	42.11	3.25	2.09

Note: In a column, common values letter(s) do not differ significantly at $p \leq 0,05$ as per HSD

Table 2. Effects of treatment with different nitrogen sources and cultivar on chlorophyll, carotenoid, and numbers of stomata


Treatment	Chlorophyll (mg g ⁻¹ FW)	Carotenoid (mg g ⁻¹ FW)	The number of stomata (mm ⁻²)
Perkasa cultivar	12.92a	1.59a	14.40a
Lebat-3 cultivar	13.24a	1.88a	14.67a
HSD _{0,05}	1.16	0.29	3.50
Control	9.68b	1.38b	8.50c
Urea	15.93a	2.35a	22.00a
10% LOF	15.56a	1.69ab	11.50bc
20% LOF	13.77a	1.76ab	17.83ab
30% LOF	10.46b	1.47b	12.83bc
HSD _{0,05}	2.64	0.67	8.05

Note: In a column, common values letter(s) do not differ significantly at $p \leq 0,05$ as per HSD

Table 3. Effects of treatment nitrogen source and cultivar on SPAD value, number of pods, and yield two cultivars bean.

Parameter	Source N	Cultivar		HSD _{0,05}
		Perkasa	Lebat-3	
SPAD value	Control	18.31 Ab	17.54Ab	9.34
	Urea	27.79 Aa	28.91 Aa	
	10% LOF	20.13 Aab	28.11 Aa	
	20% LOF	24.06 Aab	24.19 Aa	
	30% LOF	28.96 Aa	21.81 Aab	
	Number of pods	Control	5.46 Ac	
Urea		9.13 Aab	9.05 Aa	
10% LOF		9.96 Aab	12.13 Aa	
20% LOF		13.38 Aa	4.67 Bab	
30% LOF		6.92 Abc	3.75 Ab	
Yield (kg ha ⁻¹)		Control	231.11 Ac	113.33 Ab
	Urea	720.89 Aab	429.78 Aa	
	10% LOF	565.33 Ab	515.56 Aa	
	20% LOF	846.67 Aa	181.33 Bb	
	30% LOF	333.78 Ac	148.00 Ab	

Note: Mean followed by the same letter (capital letter read horizontally and small letter read vertically) do not differ significantly at $p \leq 0,05$ as per HSD

**Figure 1.** The number of stomata of Perkasa: A. Control, B. Urea, C. 10% LOF, D. 20% LOF, E. 30% LOF. The number of stomata of Lebat-3: F. control, G. Urea, H. 10% LOF, I. 20% LOF, J. 30% LOF

The application of inorganic-and-organic-sources-based nitrogen affected vegetative growth, namely number of pods and yield. Significant increase in yield (Table 3) is confirmed with better vegetative growth (Table 1 and 2). In this study, the application of 10 or 20% LOF was sufficient for the growth of green beans and was not significantly different from urea treatment. Perkasa cultivar with 20% LOF produced a statistically higher number of bean pods compared to the Lebat-3. Similarly, Perkasa cultivar with 20% LOF produced the highest number of bean pods per plant but was not significantly different from the 10% LOF and urea treatments. Furthermore, Lebat-3 cultivar with

LOF 10% produced the highest number of bean pods but was not significantly different from the urea and 20% LOF treatment.

The field observation showed that there was a symptom of N deficiency in green bean leaves without nitrogen fertilizer. This deficiency occurs in the leaves of plants without treatment (control). The symptoms include yellowish-green coloration of leaves compared to normal leaves which are fresh green. Erythrina et al. (2016) stated that nitrogen is the main nutrient in the formation of leaf color due to its important role in increasing leaf green matter and protein. Therefore, the application of nitrogen increases the green color of the leaves but when deficient, the formation of chlorophyll is reduced thereby causing the leaves to appear yellowish-green. Bojovic et al. (2009) stated that nitrogen deficiency causes a reduction in leaf greenish color and area, as well as photosynthesis due to the linear correlation between leaf chlorophyll formation and nitrogen. Furthermore, nitrogen requirement by plants is needed to be in accordance with the demand of the plant. Zainal et al. (2014) stated that the lack of Nitrogen leads to leaf chlorosis indicated by yellowing of the leaves. In contrast, excess nitrogen accelerates plant growth, especially on the stems, the leaves turn dark green.

Lebat-3 cultivar requires a lower LOF concentration compared with Perkasa to produce significant yield. The plant genetic traits in the two cultivars influenced the different responses to liquid organic fertilization. Table 3 shows that Perkasa cultivar with 20% LOF produced a higher yield compared to other treatments but was not significantly different from the 100% recommended urea treatment. The Lebat-3 cultivar with 10% LOF produced a higher yield compared to other treatments but was not significantly different from the recommended 100% urea treatment. Chaturvedi (2005), reported that nitrogen fertilization with the proper dose affects grain yield in rice and other parameters in each cultivar. Furthermore, the use of N sources for plants affects the results of photosynthesis. This is supported by Rathke (2005), which stated that plant yield is improved by increasing the source of N. In addition, nitrogen improves yield by increasing leaf area and the rate of photosynthesis, hence, high carbohydrates are produced by the plants (Chaturvedi 2005). The yield of Lebat-3 cultivar was lower compared to Perkasa but was not significantly different from the recommended 100% urea treatment and 20% LOF. Ratnasari et al. (2015), stated that the differences in the characters possessed by cultivar are caused by the varying genetic composition. Hence, each cultivar responds differently to environmental and production factors. The positive effect of liquid organic fertilizers from leucaena leaves in increasing crop yields was also found by Duaja et al. (2013) on chickpeas, Palimbungan et al. (2006) on mustard plants, Septirosya et al. (2019) on tomato plants, and Ainiya et al. (2019) on sweet corn plants. The experiment showed that the liquid organic fertilizers derived from leucaena leaves are applicable as an alternative nitrogen source in bean cultivation. This low-cost technology is easy to use by farmers and also supports organic bean production.

In conclusion, the use of plant-based nitrogen fertilization from leucaena leaves could be an alternative to nitrogen fertilization in bean plants. The 20% LOF is recommended as an adequate concentration to increase the growth and yield of green beans. Furthermore, each cultivar shows different responses to growth factors. Treatment of Perkasa (with 20% LOF) and Lebat-3 cultivar (with 10% LOF) produced the highest bean yield, respectively.

ACKNOWLEDGEMENTS

This study was funded by the Research Grant 2020 from Faculty of Agriculture, Universitas Lampung, Indonesia. The authors are grateful to Paul B Timotiwu head of LT-SIT (The Integrated Laboratory and Innovation Center of Technology, Universitas Lampung, Indonesia) for helping with pigment measurement, as well as Setyo Widagdo, Fitri Yelli, Dad Resiworo for scientific discussions.

REFERENCES

- Ainiya M, Fadli M, Despita R. 2019. Peningkatan pertumbuhan dan hasil jagung manis dengan pemanfaatan trichokompos dan POC daun lamtoro. *Agrotech Res J* 3 (2): 69-74. DOI: 10.20961/agrotechresj.v3i2.31910. [Indonesian]
- Amin MEH. 2011. Effect of different nitrogen sources on growth, yield and quality of fodder maize (*Zea mays* L.). *J Saudi Soc Agric Sci* 10: 17-23. DOI: 10.1016/j.jssas.2010.06.003.
- Arista D, Suryono, Sudadi. 2015. Efek dari kombinasi pupuk N, P dan K terhadap pertumbuhan dan hasil kacang tanah pada lahan kering Alfisol. *Jurnal Agrosains* 17 (2): 49-52. DOI: 10.20961/agsjpa.v17i2.18672. [Indonesian]
- Bojovic B, Markovic A. 2009. Correlation between nitrogen and chlorophyll content in wheat (*Triticum aestivum* L.). *Kragujevac J Sci* 31: 69-74.
- Bojovic, B, Stojanovic J. 2005. Chlorophyll and carotenoid content in wheat cultivars as a function of mineral nutrition. *Arch Biol Belgrade* 57 (4): 283-290. DOI: 10.2298/ABS0504283B.
- Camara, CRS, Urrea CA, Schlegel V. 2013. Pinto beans (*Phaseolus vulgaris* L.) as a functional food: Implications of human health. *Agriculture* 3: 90-111. DOI: 10.3390/agriculture3010090.
- Chaturvedi I. 2005. Effect of nitrogen fertilizers on growth, yield and quality of hybrid rice (*Oryza sativa*). *J Central Eur Agric* 6 (4): 611-618.
- Duaja MD, Mukhsin, Sijabat R. 2013. Analisis pertumbuhan dan hasil dua varietas buncis (*Phaseolus vulgaris* L.) pada perbedaan jenis pupuk organik cair. *Bioplantae* 2 (1): 47-54. [Indonesian]
- Duaja MD. 2013. Pengaruh jenis bahan dasar dan dosis pupuk organik cair terhadap pertumbuhan dan hasil buncis (*Phaseolus vulgaris* L.). *Bioplantae* 2 (4): 192-200. [Indonesian]
- Erythrina. 2016. Bagan warna daun: Alat untuk meningkatkan efisiensi pemupukan nitrogen pada tanaman padi. *Jurnal Litbang Pertanian* 35 (1): 1-10. DOI: 10.21082/jp3.v35n1.2016.p1-10. [Indonesian]
- Fahmi A, Syamsudin, Utami SNH, Radjaguguk B. 2010. Pengaruh interaksi hara nitrogen dan fosfor terhadap pertumbuhan tanaman jagung (*Zea mays* L.) pada tanah regosol dan latosol. *Berita Biologi* 10 (3): 297-304. [Indonesian]
- Faustina E, Sudradjat, Supijatno. 2015. Optimization of nitrogen and phosphorus fertilizer on two years old of oil palm (*Elaeis guineensis* Jacq.). *Asian J Appl Sci* 3 (3): 421-428.
- Ginandjar, S, Frasetya B, Nugraha W, Subandi M. 2019. The effect of liquid organic fertilizer of vegetable waste and planting media on growth and yield of strawberry (*Fragaria* spp.) Earlibrite Cultivar. *IOP Conf Ser Earth Environ Sci* 334: 012033. DOI: 10.1088/1755-1315/334/1/012033.

- Hendriyani IK, Nurchayati Y, Setiari N. 2018. Kandungan klorofil dan karotenoid kacang tunggak (*Vigna unguiculata* (L.) Walp.) pada umur tanaman yang berbeda. *Jurnal Biologi Tropika* 1 (2): 38-43. DOI: 10.14710/jbt.1.2.38-43. [Indonesian]
- Hidayat O, Suharyana A. 2019. Pengaruh dosis pupuk organik cair daun lamtoro terhadap pertumbuhan dan hasil tanaman pakcoy (*Brassica rapa* L.) varietas Nauli-F1. *Paspalum Jurnal Ilmu Pertanian* 7 (2): 57-63. DOI: 10.35138/paspalum.v7i2.118. [Indonesian]
- Jeksen J, Mutiara C. 2017. Analisis kualitas pupuk organik cair dari beberapa jenis tanaman Leguminosa. *Jurnal Pendidikan MIPA* 7 (2): 124-130. [Indonesian]
- Madusari S. 2019. Processing of fibre and its application as liquid organic fertilizer in oil palm (*Elaeis guineensis* Jacq.) seedling for sustainable agriculture. *J Appl Sci Adv Technol* 1 (3): 81-90.
- Maghfoer MD, Koesriharti, Islami T, Kanwal NDS. 2018. A study of the efficacy of various nutrient sources on the growth and yield of cabbage. *Agrivita J Agric Sci* 40 (1): 168-176. DOI: 10.17503/agrivita.v40i1.1721.
- MoA. 1999. Deskripsi kacang buncis varietas Lebat-3. Lampiran keputusan Menteri Pertanian No: 748/Kpts/TP.240/6/1999. [Indonesian]
- MoA. 2002. Deskripsi kacang buncis varietas Perkasa. Lampiran keputusan Menteri Pertanian No: 270/Kpts/TP.240/4/2002. [Indonesian]
- Pahlevi RW, Guritno B, Suminarti NE. 2016. Pengaruh kombinasi proporsi pemupukan nitrogen dan kalium pada pertumbuhan, hasil dan kualitas tanaman ubi jalar (*Ipomoea batatas* (L.) Lamb) varietas Cilembu pada dataran rendah. *Jurnal Produksi Tanaman* 4 (1): 16-22. [Indonesian]
- Palimbungan N, Labatar R, Hamzah F. 2006. Pengaruh ekstrak daun lamtoro sebagai pupuk organik cair terhadap pertumbuhan dan hasil produksi tanaman sawi. *Jurnal Agrisistem* 2 (2): 96-101. [Indonesian]
- Pamungkas MA, Supijatno. 2017. Pengaruh pemupukan nitrogen terhadap percabangan tanaman teh (*Camelia sinensis* (L.) O. Kuntze) untuk pembentukan bidang petik. *Buletin Agronomi* 5: 234-241. DOI: 10.29244/agrob.v5i2.16804. [Indonesian]
- Pary C. 2015. Pengaruh pupuk organik (daun lamtoro) dalam berbagai konsentrasi terhadap pertumbuhan tanaman sawi. *Jurnal Fikratuna* 7 (2): 247-255. [Indonesian]
- Paul V, Sharma L, Pandey R, Meena RC. 2017. Measurements of stomatal density and stomatal index on leaf/plant surfaces. Manual of ICAR sponsored training programme on physiological techniques to analyze the impact of climate change on crop plants. Division of Plant Physiology, IARI, New Delhi.
- Permanasari I, Irfan M, Abizar. 2014. Pertumbuhan dan hasil kedelai (*Glycine max* (L.) Merrill) dengan pemberian *Rhizobium* dan pupuk urea pada media gambut. *Jurnal Agroteknologi* 5: 29-34. [Indonesian]
- Pramitasari HE, Wardiyati T, Nawawi M. 2016. Pengaruh dosis pupuk nitrogen dan tingkat kepadatan tanaman terhadap pertumbuhan dan hasil tanaman kailan (*Brassica oleraceae* L.). *Jurnal Produksi Tanaman* 4 (1): 49-56. [Indonesian]
- Proklamasingsih E, Prijambada ID, Rachmawati D, Sancayaningsih RP. 2012. Laju fotosintesis dan kandungan klorofil kedelai pada media tanam masam dengan pemberian garam aluminium. *Agrotrop* 2 (1): 17-24. [Indonesian]
- Putri FM, Suedy SWA, Darmanti S. 2017. Pengaruh pupuk nanosilika terhadap jumlah stomata, kandungan klorofil dan pertumbuhan padi hitam (*Oryza sativa* L. cv. japonica). *Buletin Anatomi Fisiologi* 2 (1): 72-79. DOI: 10.14710/baf.2.1.2017.72-79. [Indonesian]
- Rahimi A, Moghaddam SS, Ghiyasi M, Heydarzadeh S, Ghazizadeh K, Popovic-Djordjevic J. 2019. The influence of chemical, organic and biological fertilizers on agrobiological and antioxidant properties of syrian cephalaria (*Cephalaria syriaca* L.). *Agriculture* 9: 122. DOI: 10.3390/agriculture9060122.
- Rathke GW, Christen O, Diepenbrock W. 2005. Effects of nitrogen source and rate on productivity and quality of winter oilseed rape (*Brassica napus* L.) grown in different crop rotations. *Field Crops Res* 94: 103-113. DOI: 10.1016/j.fcr.2004.11.010.
- Ratnasari D, Bangun MK, Damanik RIM. 2015. Respon dua varietas kedelai (*Glycine max* (L.) Merrill.) pada pemberian pupuk hayati dan NPK majemuk. *Jurnal Online Agroteknologi* 3 (1): 276-82. [Indonesian]
- Razaq M, Zhang P, Shen H, Salahuddin. 2017. Influence of nitrogen and phosphorus on the growth and root morphology of Acer mono. *PLoS One* 12 (2): e0171321. DOI: 10.1371/journal.pone.0171321.
- Rihana S, Heddy YBS, Maghfoer MD. 2013. Pertumbuhan dan hasil tanaman buncis (*Phaseolus vulgaris* L.) pada berbagai dosis pupuk kotoran kambing dan konsentrasi zat pengatur tumbuh Dekamon. *Jurnal Produksi Tanaman* 1 (4): 369-377. [Indonesian]
- Rizqiani NF, Ambarwati E, Yuwono NW. 2006. Pengaruh dosis dan frekuensi pemberian pupuk organik cair terhadap pertumbuhan dan hasil buncis (*Phaseolus vulgaris* L.) dataran rendah. *Ilmu Pertanian* 13 (2): 163-178. [Indonesian]
- Sab DBH, Sridhara S, Gopakkali P. 2020. Growth and yield of maize (*Zea mays* L.) as influenced by date of sowing and hybrids. *J Agric Appl Biol* 1 (2): 38-45. DOI: 10.11594/jaab.01.02.0.
- Septirosya T, Putri RH, Aulawi T. 2019. Aplikasi pupuk organik cair lamtoro pada pertumbuhan hasil tanaman tomat. *Agroscrip* 1 (1): 1-8. DOI: 10.36423/agroscrip.v1i1.185. [Indonesian]
- Sinaga AS, Guritno B, Sudiarmo. 2017. Pengaruh dosis kompos sampah rumah tangga terhadap pertumbuhan dan hasil tiga varietas buncis tipe tegak (*Phaseolus vulgaris* L.). *Jurnal Produksi Tanaman* 5 (6): 947-956. [Indonesian]
- Wijiyanti P, Hastuti ED, Haryanti S. 2019. Pengaruh masa inkubasi pupuk dari air cucian beras terhadap pertumbuhan tanaman sawi hijau (*Brassica juncea* L.). *Buletin Anatomi Fisiologi* 4 (1): 41-28. DOI: 10.14710/baf.4.1.2019.21-28. [Indonesian]
- Zainal M, Nugroho A, Suminarti NE. 2014. Respon pertumbuhan dan hasil tanaman kedelai (*Glycine max* (L.) Merrill) pada berbagai tingkat pemupukan N dan pupuk kandang ayam. *Jurnal Produksi Tanaman* 2 (6): 484-490. [Indonesian]