



## Locating-Chromatic Number of Amalgamation of Stars

Asmiati\*, H. Assiyatun & E.T. Baskoro

Combinatorial Mathematics Research Group,  
Faculty of Mathematics and Natural Sciences,  
Institut Teknologi Bandung, Jl. Ganesa 10 Bandung.  
Email: asmiati308@students.itb.ac.id

**Abstract.** Let  $G$  be a connected graph and  $c$  a proper coloring of  $G$ . For  $i=1,2,\dots,k$  define the color class  $C_i$  as the set of vertices receiving color  $i$ . The color code  $c_{\Pi}(v)$  of a vertex  $v$  in  $G$  is the ordered  $k$ -tuple  $(d(v,C_1),\dots,d(v,C_k))$  where  $d(v,C_i)$  is the distance of  $v$  to  $C_i$ . If all distinct vertices of  $G$  have distinct color codes, then  $c$  is called a locating-coloring of  $G$ . The locating-chromatic number of graph  $G$ , denoted by  $\chi_L(G)$  is the smallest  $k$  such that  $G$  has a locating coloring with  $k$  colors. In this paper we discuss the locating-chromatic number of amalgamation of stars  $S_{k,m}$ .  $S_{k,m}$  is obtained from  $k$  copies of star  $K_{1,m}$  by identifying a leaf from each star. We also determine a sufficient condition for a connected subgraph  $H \subseteq S_{k,m}$  satisfying  $\chi_L(H) \leq \chi_L(S_{k,m})$ .

**Keywords:** amalgamation of stars; color code; locating-chromatic number.

### 1 Introduction

Let  $G$  be a finite, simple, and connected graph. Let  $c$  be a proper coloring of a connected graph  $G$  using the colors  $1,2,\dots,k$  for some positive integer  $k$ , where  $c(u) \neq c(v)$  for adjacent vertices  $u$  and  $v$  in  $G$ . Thus, the coloring  $c$  can be considered as a partition  $\Pi$  of  $V(G)$  into color classes (independent sets)  $C_1, C_2, \dots, C_k$ , where the vertices of  $C_i$  are colored by  $i$  for  $1 \leq i \leq k$ . The color code  $c_{\Pi}(v)$  of a vertex  $v$  in  $G$  is the ordered  $k$ -tuple  $(d(v,C_1), \dots, d(v,C_k))$  where  $d(v,C_i) = \min\{d(v,x) \mid x \in C_i\}$  for  $1 \leq i \leq k$ . If all distinct vertices of  $G$  have distinct color codes, then  $c$  is called a *locating-coloring* of  $G$ . A *minimum locating-coloring* uses a minimum number of colors and this number is called the *locating-chromatic number* of graph  $G$ , denoted by  $\chi_L(G)$ .

---

\*Permanent address: Mathematics Departement, Faculty of Mathematics and Natural Sciences, Universitas Lampung, Jl. Brojonegoro no.1 Bandar Lampung, Lampung.  
Received January 11<sup>th</sup>, 2010, Revised July 14<sup>th</sup>, 2010, Accepted for publication July 28<sup>th</sup>, 2010.

The following definition of an amalgamation of graphs is taken from [3]. For  $i=1,2,\dots,k$ , let  $G_i$  be a graph with a fixed vertex  $v_{oi}$ . The *amalgamation*  $\text{Amal}(G_i, v_{oi})$  is a graph formed by taking all the  $G_i$ 's and identifying their fixed vertices. In this paper, we consider the amalgamation of stars. More precisely, for  $i=1,2,\dots,k$ , let  $G_i = K_{1,n_i}$ ,  $n_i \geq 1$  where  $v_{oi}$  be any leaf (a vertex of degree 1) of  $K_{1,n_i}$ . We denote the amalgamation of  $k$  stars  $K_{1,n_i}$  by  $S_{k,(n_1,n_2,\dots,n_k)}$ ,  $k \geq 2$ . We call the identified vertex as the *center* (denoted by  $x$ ), the vertices of distance 1 from the center as the *intermediate vertices* (denoted by  $l_i$ ;  $i=1,2,\dots,k$ ), and the  $j$ -th leaf of the intermediate vertex  $l_i$  by  $l_{ij}$  ( $j=1,2,\dots,m-1$ ). In particular, when  $n_i = m$ ,  $m \geq 1$  for all  $i$ , we denote the amalgamation of  $k$  isomorphic stars  $K_{1,m}$  by  $S_{k,m}$ .

The locating-chromatic number was firstly studied by Chartrand *et al.* [1]. They determined the locating-chromatic numbers of some well-known classes such as paths, cycles, complete multipartit graphs and double stars. Apart of paths and double stars, the locating-chromatic number of other trees are also considered by Chartrand *et al.* [2]. They constructed a tree of order  $n \geq 5$  with the locating-chromatic number  $k$ , where  $k \in \{3,4,\dots,n-2,n\}$ . They also showed that no tree on  $n$  vertices with locating-chromatic number  $n-1$ .

Based on the previous results, locating-chromatic number of amalgamation of stars have not been studied. Motivated by this, in this paper we determine the locating-chromatic number of amalgamation of stars.

Beside that, we also discuss the monotonicity property of the locating-chromatic number for the class of amalgamation of stars. Clearly, the locating-chromatic number of a star  $K_{1,n}$  is  $n+1$ , for any  $n$  (since all vertices must have different color codes). Since any connected subgraph  $H$  of  $K_{1,n}$  is also a star with small size, then we clearly have  $\chi_L(H) \leq \chi_L(K_{1,n})$ . However in general for any connected subgraph  $H \subseteq G$ , the locating-chromatic number of  $H$  may not be necessarily smaller or equal to the locating-chromatic number of  $G$ .

In this paper, we also investigate the monotonicity property of the locating-chromatic number for amalgamation of stars,  $S_{k,m}$ . We derive a sufficient condition for a connected subgraph  $H \subseteq S_{k,m}$  satisfying  $\chi_L(H) \leq \chi_L(S_{k,m})$ .

The following results were proved by Chartrand et al. in [1]. The set of neighbours of a vertex  $v$  in  $G$  is denoted by  $N(v)$ .

**Theorem 1.1.** *Let  $c$  be a locating-coloring in a connected graph  $G$ . If  $u$  and  $v$  are distinct vertices of  $G$  such that  $d(u,w) = d(v,w)$  for all  $w \in V(G) - \{u,v\}$ , then  $c(u) \neq c(v)$ . In particular, if  $u$  and  $v$  are non adjacent vertices of  $G$  such that  $N(u) = N(v)$ , then  $c(u) \neq c(v)$ .*

**Corollary 1.1.** *If  $G$  is a connected graph containing a vertex adjacent to  $k$  leaves of  $G$ , then  $\chi_L(G) \geq k+1$ .*

## 2 Main Results

We first prove some lemmas regarding the properties of locating-chromatic number of amalgamation of stars. From now on  $S_{k,m}$  denotes the amalgamation of  $k$  isomorphic stars  $K_{1,m}$ .

**Lemma 2.1.** *For  $k \geq 2, m \geq 2$ , let  $c$  be a proper coloring of  $S_{k,m}$ , using at least  $m$  colors. The coloring  $c$  is a locating-coloring if and only if  $c(l_i) = c(l_n), i \neq n$  implies  $\{c(l_{ij}) \mid j=1,2,\dots,m-1\}$  and  $\{c(l_{nj}) \mid j=1,2,\dots,m-1\}$  are distinct.*

*Proof.* Let  $P = \{c(l_{ij}) \mid j=1,2,\dots,m-1\}$  and  $Q = \{c(l_{nj}) \mid j=1,2,\dots,m-1\}$ . Let  $c$  be a locating-coloring of  $S_{k,m}$ ,  $k \geq 2, m \geq 2$  using at least  $m$  colors and  $c(l_i) = c(l_n)$ , for some  $i \neq n$ . Suppose that  $P = Q$ . Because  $d(l_i, u) = d(l_n, u)$  for every  $u \in V \setminus \{\{l_{ij} \mid j=1,2,\dots,m-1\} \cup \{l_{nj} \mid j=1,2,\dots,m-1\}\}$  then the color codes of  $l_i$  and  $l_n$  will be the same. So  $c$  is not a locating-coloring, a contradiction. Therefore  $P \neq Q$ .

Let  $\Pi$  be a partition of  $V(G)$  into color classes with  $|\Pi| \geq m$ . Consider  $c(l_i) = c(l_n), i \neq n$ . Since  $P \neq Q$ , there are color  $x$  and color  $y$  such that  $(x \in P, x \notin Q)$  and  $(y \in P, y \notin Q)$ . We will show that color codes for every  $v \in V(S_{k,m})$  is unique.

- Clearly,  $c_\Pi(l_i) \neq c_\Pi(l_n)$  because their color codes differ in the  $x$ th-ordinate and  $y$ th-ordinate.

- If  $c(l_{ij}) = c(l_{ns})$ , for some  $l_i \neq l_n$ , we will show that  $c_{\Pi}(l_{ij}) \neq c_{\Pi}(l_{ns})$ . We divide into two cases.

Case 1. If  $c(l_i) = c(l_n)$  then by the premise of this theorem,  $P \neq Q$ . So  $c_{\Pi}(l_{ij}) \neq c_{\Pi}(l_{ns})$ .

Case 2. Let  $c(l_i) = r_1$  and  $c(l_n) = r_2$ , with  $r_1 \neq r_2$ . Then  $c_{\Pi}(l_{ij}) \neq c_{\Pi}(l_{ns})$  because their color codes are different at least in the  $r_1$ th-ordinate and  $r_2$ th-ordinate.

- If  $c(l_i) = c(l_{nj})$ ,  $l_i \neq l_n$ , then  $c_{\Pi}(l_i)$  contains at least two components of value 1, whereas  $c_{\Pi}(l_{nj})$  contains exactly one component of value 1. Thus  $c_{\Pi}(l_i) \neq c_{\Pi}(l_{nj})$ .
- If  $c(x) = c(l_{ij})$ , then color code of  $c_{\Pi}(x)$  contains at least two components of value 1, whereas  $c_{\Pi}(l_{ij})$  contains exactly one component of value 1. Thus  $c_{\Pi}(x) \neq c_{\Pi}(l_{ij})$ .

From all above cases, we see that the color code for each vertex in  $S_{k,m}$  is unique, thus  $c$  is a locating-coloring.  $\square$

**Lemma 2.2.** *Let  $c$  be a locating coloring of  $S_{k,m}$  using  $m+a$  colors and  $H(a) = (m+a-1) \binom{m+a-1}{m-1}$ ,  $a \geq 0$ . Then  $k \leq H(a)$ .*

*Proof.* Let  $c$  be a locating-coloring of  $S_{k,m}$  using  $m+a$  colors. For fixed  $i$ , let  $c(l_i)$  be a color of intermediate vertex  $l_i$ , then the number color combinations can be used by  $\{l_{ij} \mid j=1,2,\dots,m-1\}$  is  $\binom{m+a-1}{m-1}$ . Because one color is used for coloring the center  $x$ , there are  $(m+a-1)$  colors for  $l_i$ , for every  $i=1,2,\dots,k$ . By Lemma 2.1, the maximum number of  $k$  is  $(m+a-1) \binom{m+a-1}{m-1} = H(a)$ . So  $k \leq H(a)$ .  $\square$

The main result of this paper concerns about locating-chromatic number of  $S_{k,m}$ .

**Theorem 2.1.** For  $a \geq 0, k \geq 2, m \geq 2$ , let  $H(a) = (m+a-1) \binom{m+a-1}{m-1}$ . Then,

$$\chi_L(S_{k,m}) = \begin{cases} m & \text{for } 2 \leq k \leq H(0), m \geq 3, \\ m+a & \text{for } H(a-1) < k \leq H(a), a \geq 1. \end{cases}$$


*Proof.* First, we determine the trivial lower bound. By Corollary 1.1, each vertex  $l_i$  is adjacent to  $(m-1)$  leaves, for  $i=1,2,\dots,k$ . Thus,  $\chi_L(S_{k,m}) \geq m$ .

Next, we determine the upper bound of  $\chi_L(S_{k,m})$  for  $2 \leq k \leq H(0) = m-1$ . Let  $c$  be a coloring of  $V(S_{k,m})$  using  $m$  colors. Without loss of generality, we can assign  $c(x)=1$  and  $c(l_i)=i+1$  for  $i=1,2,\dots,k$ . To make sure that the leaves will have distinct color code, we assign  $\{l_{ij} \mid j=1,2,\dots,m-1\}$  by  $\{1,2,\dots,m\} \setminus \{i+1\}$  for any  $i$ . Then, by Lemma 2.1, we have that  $c$  is a locating-coloring. Thus  $\chi_L(S_{k,m}) \leq m$ .


Next, we shall improve the lower bound for the case of  $k$  such that  $H(a-1) < k \leq H(a), a \geq 1$ . Since  $k > H(a-1)$  then by Lemma 2.2,  $\chi_L(S_{k,m}) \geq m+a$ . On the other hand if  $k > H(a)$  then by Lemma 2.2,  $\chi_L(S_{k,m}) \geq m+a+1$ . Thus  $\chi_L(S_{k,m}) \geq m+a$  if  $H(a-1) < k \leq H(a)$ .

Next, we determine the upper bound of  $\chi_L(S_{k,m})$  for  $H(a-1) < k \leq H(a), a \geq 1$ . Without loss of generality, let  $c(x)=1$  and color the intermediate vertices  $l_i$  by  $2,3,\dots,m+a$  in such a way that the number of the intermediate vertices receiving the same color  $t$  does not exceed  $\binom{m+a-1}{m-1}$ , for any  $t$ . We are able to do so because  $H(a-1) < k \leq H(a)$ . Therefore, if  $c(l_i)=c(l_n)$ ,  $i \neq n$  then we can manage  $\{c(l_{ij}) \mid j=1,2,\dots,m-1\} \neq \{c(l_{nj}) \mid j=1,2,\dots,m-1\}$ . By Lemma 2.1,  $c$  is a locating-coloring on  $S_{k,m}$ . So  $\chi_L(S_{k,m}) \leq m+a$  for  $H(a-1) < k \leq H(a)$ .  $\square$

The following figures show minimum locating-colorings on  $S_{4,6}$  and  $S_{9,3}$ .



**Figure 1** A minimum locating-coloring of  $S_{4,6}$ .



**Figure 2** A minimum locating-coloring of  $S_{9,3}$ .

Next, we discuss the monotonicity property of locating-chromatic number for the amalgamation of stars.

**Theorem 2.2** *If  $2 \leq k \leq m-1$ , then  $\chi_L(G) \leq \chi_L(S_{k,m})$  for every  $G \subseteq S_{k,m}$  and  $G \neq K_{1,m}$ .*

*Proof.* Let  $c$  be a minimum locating-coloring of  $S_{k,m}$  obtained from Theorem 2.1. Let  $G$  be any connected subgraph of  $S_{k,m}$ . Define a coloring  $c'$  on  $G$  by

preserving colors used in  $S_{k,m}$  for the corresponding vertices, namely  $c'(v') = c(v)$  if  $v$  is the corresponding vertex of  $v'$  in  $S_{k,m}$ . We show that  $c'$  is a locating-coloring of  $G$ .

If there exist  $l_i, l_n$  such that  $\{c'(l_{ij}) \mid j=1,2,\dots,r\} = \{c'(l_{nj}) \mid j=1,2,\dots,r\}$ , with  $1 \leq r \leq m-1$ , then color codes of  $l_{ij}$  and  $l_{nj}$  for every  $j=1,2,3,\dots,m$  is unique because  $c'(l_i) \neq c'(l_n)$  for every  $l_i \neq l_n$ . If  $c'(l_i) = c'(l_{nj}) \neq c'(x)$ , then the first component of  $c'_\pi(l_i)$  has value 1, whereas for  $c'_\pi(l_{nj})$  it has value 2. So color code of  $l_i$  and  $l_{nj}$  are different. Next, if  $c'(x) = c'(l_{nj})$ ,  $G \neq P_3$  then their color codes are different because  $c'(l_i) \neq c'(l_n)$  for every  $l_i \neq l_n$ . For the case  $G = P_3$ ,  $v_i \in V(P_3)$  for each  $i$  is colored by 1, 2, and 3 respectively. Because the color codes for every  $v \in V(G)$  is unique, then  $c'$  is a locating-coloring of  $G$ . So  $\chi_L(G) \leq \chi_L(S_{k,m})$  for every  $G \subseteq S_{k,m}$ ,  $G \neq K_{1,m}$ .  $\square$

Let  $S_{k,(n_1,n_2,\dots,n_k)} \subseteq S_{k,m}$ . Define  $A = \{i \mid n_i = 1\}$ . For  $k \geq m$ , we must restrict subgraphs of  $S_{k,m}$  so that satisfy monotonicity property.

**Theorem 2.3** *If  $k \geq m$  and  $|A| \leq \chi_L(S_{k,m}) - 1$  then  $\chi_L(S_{k,(n_1,n_2,\dots,n_k)}) \leq \chi_L(S_{k,m})$ .*

*Proof.* Let  $k \geq m$  and from Theorem 2.1, we have that  $\chi_L(S_{k,m}) = m + a$  for  $H(a-1) < K \leq H(a)$ ,  $a \geq 1$ . Let  $G = S_{k,(n_1,n_2,\dots,n_k)}$  be any subgraph obtained from  $S_{k,m}$  with  $1 \leq n_i \leq m$ . If  $2 \leq n_i \leq m$  for each  $i$ , then color vertices of  $G$  follow the proof of Theorem 2.1. Clearly, the coloring of  $G$  is a locating-coloring. Otherwise, we have  $n_i = 1$  for some  $i$ , and so  $|A| \geq 1$ . If  $|A| \leq \chi_L(S_{k,m}) - 1$ , then the center  $x$  is given color 1,  $l_i \in A$  for each  $i$  is colored by  $2, 3, \dots, \chi_L(S_{k,m})$ , respectively and the colors for the other vertices follow the proof of Theorem 2.1. Observe that the color codes of  $l_i$  for each  $l_i \in A$  has value 1 in the 1th-ordinate, 0 in the  $i$ th-ordinate, and 2 otherwise, these color codes are unique. For the remaining of the vertices, the color codes are also unique as proven in Theorem 2.1. As the result, the coloring of  $G$  is a locating-coloring. So  $\chi_L(S_{k,(n_1,n_2,\dots,n_k)}) \leq \chi_L(S_{k,m})$ .  $\square$

**References**

- [1] Chartrand, G., Erwin, D., Henning, M.A., Slater, P.J. & Zang. P., *The locating-chromatic number of a graph*, Bull. Inst. Combin. Appl., **36**, pp. 89-101, 2002.
- [2] Chartrand, G., Erwin, D., Henning, M.A., Slater & P.J., Zang. P., *Graph of order  $n$  with locating-chromatic number  $n-1$* , Discrete Mathematics, **269**, pp. 65-79, 2003.
- [3] Carlson, K., *Generalized books and  $C_m$ -snakes are prime graphs*, Ars Combin., **80**, pp. 215-221, 2006.